

In vivo* DNA Methylation of *Escherichia coli* DH5 α and Top10F' Strains by Bacterial Cytosine-5 Methyltransferase *M. Msp1*

Öztürk ÖZDEMİR

Cumhuriyet Üniversitesi Tıp Fakültesi Tıbbi Biyoloji ve Genetik A.B.D. Sivas-TURKEY

David HORNBY

Department of Molecular Biology and Biotechnology University of Sheffield U.K

Received: 09.04.1997

Abstract: At the chromatin level, methylated CpG dinucleotides are R.Msp1 resistant compared with nonmethylated counterparts. The DNA of two *E.coli* strains was analyzed following transformation with bacterial cytosine-5-methyltransferase gene M.Msp1 in mammalian transfection vector pcDNA3. Expression of the M.Msp1 was tested by R.Msp1 digestion. The results suggest that the DNA of both strains was fully methylated at the CCGG sequences, by the active enzyme under the control of T7 and cytomegalovirus (CMV) promoters. Methylated DNA can not be digested and it exhibits higher fragment sizes in 1% agarose gel in contrast to the untransformed cell DNAs *in vivo*.

Key Words: Bacterial MTase M.Msp1, *in vivo* DNA methylation

***Escherichia coli* DHS α ve Top10F' Hücre DNA'larının Bakteri Sitozin-5-Metiltransferaz Enzimi, *M. Msp1*, Tarafından *In vivo* Metilasyonu**

Özet: Kromatin düzeyinde metillenmiş CpG dinükleotidleri, metil grubu taşımayan nükleotidlerin aksine bir modifiye restriksiyon enzimi olan R. Msp1 enzimine dirençlidirler. Yapısında metillenmiş dinükleotid (mCpG) taşıyan bu serbest DNA aynı restriksiyon enziminin kesilmez. Bu çalışmada, metillenmiş nükleotidlerin bu dirençli olma özelliğinden yararlanarak başka bir bakteri türüne ait (*Moroxella* spp.), rekombine metiltransferaz enziminin, farklı hücrelerde benzer metilaz aktivite gösterip göstermeyeceği araştırılmak istenmiştir. Bunun için izogenik olmayan iki *E.coli* suşu (DH5 α ve Top 10F'), PCR ile 5' ucuna omurgalı nükleer transport sinyali SV40 VP1 ve markır enzim glutatyon-S-transferaz (GST) gen dizileri takılmış bakteriyel sitozin-5-spesifik metiltransferaz (MTaz) M. Msp1 rekombinant gen füzyonu ile transforme edildiler.

Transforme edilmiş her iki hücre DNA' ları R. Msp1 enziminin test edildi, rekombinant genin her iki hücre grubunda da aktif olduğu görüldü. Bir memeli gen transfer vektörü olan pcDNA₃ içinde, T7 ve sitomegalovirus (CMV) promotorlarının kontrolünde enzimin tamamen aktif olduğu, *in vivo* DH5 α ve Top10F' hücre DNA'larında CCGG dizilerini metillediği saptandı.

Anahtar Sözcükler: Bakteriyel MTaz M.Msp1, *in vivo* DNA metilasyonu

*:This study is a part of a project supported by NATO fellowship 94/B1, TÜBİTAK-TURKEY and carried out in Krebs Institute University of Sheffield, UK

Introduction

DNA methyltransferases (MTases) have long been regarded as an attractive model for studying DNA - protein interaction. Over 50 cytosine DNA MTases from organisms ranging from bacteria to mammals have been cloned and sequenced (1-3). DNA methylation is used by a wide range of prokaryotes and eukaryotes for the marking of DNA sequences and regulation of gene expression (including restriction-modification systems), DNA replication and repair. Methylation of cytosine at carbon-5 of pyrimidine ring occurs at postreplication stage of DNA synthesis by the activity of DNA methyltransferase (MTase) (4-6). Cytosine MTases transfer methyl groups from the donor S-adenosyl-L-methionine to the C-5 position of target cytosine in specifically recognized DNA sequences. It is obvious that DNA methylation is the most common form of DNA postmodification in biological systems. Many studies have shown correlations between increased gene expression and decreased DNA methylation (7-8). In mammalian cells, 3-5% of the cytosine residues in DNA are methylated. The haploid mammalian genome contains (5×10^7) CpG dinucleotides about 60% of which are methylated at the 5 position. Methylation patterns undergo sweeping reorganization during gametogenesis and early development. In addition to CpG methylation, plant DNA contains methylated bases in the trinucleotide sequence CpNpG, where N can be A, C, G or T (9). Bacterial MTases are usually found as a part of restriction-modification systems to activate in cell's defense against phage infection in prokaryotes. These enzymes share a common protein architecture. Ten conserved motifs are normally found 10-20 aa in length and a hypervariable sequence called TRD (T: threonine, R: arginine and D: aspartic acid) which is located between motifs VIII and IX (10). Two major classes of DNA MTase exist in nature. First, methylates a pyrimidine ring carbon and forms C-5 methylcytosine (M.Msp1 and M.Hha I). The second class methylates exocyclic nitrogens and forms either N⁴-methylcytosine (M.Pvu II) or N⁶-methyladenine (M.TaqI) (11-13). HhaI and HpaII DNA MTases bind DNA mismatches which methylate uracil and block DNA repair (14). Bandaru et al. proposed that HpaII MTase was mutagenic in *E. coli*. The presence of the enzyme in *E. coli* causes a substantial increase in C to T mutations at CG sites (15). M.SssI and M.HhaI MTases reveal extensive interactions with the substrate DNA backbone. The two structurally related enzymes display similar specific and nonspecific contacts with DNA while bound to their target sequences (16). Sites for cytosine methylation are known as hot spots for cytosine to thymine mutations (10). It is our claim that because of their biological importance, their sequence specificity and reaction similarity, the C-5 MTases will be the subject of many studies in the near future.

We aimed to study the gene encoding the above mentioned MTase M.Msp1 whether it is active or inactive with glutathion-S-transferase (GST) and mammalian nuclear localisation transport sequences (NLS) which were fused to the 5' end of gene. For this purpose, the recombinant plasmid, encoding the bacterial M.Msp1 protein and which was used to transform different *E. coli* strains and expression level of the fusion product, was analysed under the control of T7 and CMV promoters. It is reported here, that the protein was found to be fully active and is able to methylate the CCGG sequences of the plasmid encoding the recombinant M.Msp1 in *E.coli* strains used.

Construction of Plasmids

The aim of this study was to construct a plasmid that would direct the expression of a recombinant bacterial DNA MTase enzyme in cultured cells. A PCR based strategy was carried out for introduction of three component coding sequences into the mammalian transfection vector pcDNA₃ (Invitrogen). First, the gene, encoding the bacterial cytosine-5 specific DNA MTase *M.Msp1* was introduced into the general cloning vectors pMTL23 and pNRT2. To this construct the coding sequence for glutathion-S-transferase was added followed by a vertebrate nuclear localisation SV40 VP1 sequence both by PCR based methodology. The complete fusion protein encoding gene was finally introduced into the general purpose transfection vector pcDNA₃.

Cell Culture and Transformation

The standard method for preparing competent cells was a modification of the calcium chloride procedure (17). A single colony of *E. coli* strains (DH5 α and Top10F'), which was inoculated into 5mls LB medium and incubated overnight at 37 °C. 50 μ l of this culture, was then transferred into a further 5 mls of LB and grown until the OD₂₆₀ of the culture reached 0.6 (~2 hours). 1.5 ml samples were then centrifuged at 12,000 rpm in a microcentrifuge for 2 minutes. The supernatant was removed and the bacterial pellet resuspended in 50 μ l of ice-cold CaCl₂ (50 mM) solution. After 2 minutes centrifugation, the supernatant was removed and 100 μ l of 50 mM CaCl₂ solution was added and at the same time between 5-10 μ l of a ligation mixture or less than 100 ng (~ 1 l) of a recombinant plasmid DNAs with bacterial *M.Msp1* gene fusion were added to the competent cells, which were mixed gently and incubated on ice for 20 minutes. The cells were then heat shocked at 42 °C for 2 minutes and chilled by returning the suspension immediately onto ice. 1ml LB medium was added to each sample and subsequently incubated at 37 °C for 1 hour. Each sample was then recentrifuged at 12,000 rpm in a microcentrifuge for 2 minutes, and pellet was resuspended in a volume of 200 μ l of LB and plated out onto the appropriate media and antibiotics. DH5 α cells were plated out in agar plates containing LB medium (10 gr tryptone+5 gr yeast extract+10 gr NaCl in 1 l distilled water) with ampicillin (2 μ g/ml) and Top10F' cells were plated out in agar plates containing LB medium with ampicillin (2 μ g/ml) and neomycin (40 μ g/ml).

Preparation of DNA from Transformed Cells

Genomic and plasmid DNAs from transformed DH5 α and Top10F' were isolated from 1.5-500 mls of liquid LB culture following alkaline lysis method described in Sambrook *et al.* (18). Recombinant bacterial cultures were grown to late log phase in LB medium with suitable antibiotics and the cells were harvested at 5000 g in a Sorvall centrifuge (GS3 rotor) for 10 minutes at 4 °C. The cells were then washed in solution I (0.184g/ml CaCl₂, 0.125 M Hepes, pH: 7.15) and 2ml of freshly prepared solution of lysozyme [100 mg lysozyme and 100 μ l Tris (1M) and 10 ml H₂O]. Suspensions were subsequently extracted with phenol-chloroform three times, followed by one-time ice-cold ethanol precipitation. DNA was resuspended in an 40 μ l volume of TE buffer (10 mM Tris.Cl, 1mM EDTA, pH : 8.0). Any further purification of plasmid DNA was

carried out by adsorption onto glass beads.

Restriction Activity Assay

Restriction digestion was performed by incubating 1 µg DNA from transformed and control group cells in a total volume of 20 µl with 10-20 units of restriction enzyme. Digests were carried out at the suppliers recommended temperatures (usually 37 °C) for 2-24 hours, using specific buffers supplied with the enzymes. The restriction patterns were then analyzed by 1 % agar gel electrophoresis.

Methylation Activity Assay

As a control pUC19, and recombinant plasmids that include NLS +GST + M.Msp1 gene were digested with R.Msp1 and R.HpaII for detection of activity of recombinant gene in *E.coli* strains. Digestion mixture (20 µl) included; 5 µl plasmid DNA + 2 µl R.Msp1 or R.HpaII restriction endonuclease (RE) buffers + 2 µl of RNase (10 µl/ml) + 1 µl R.Msp1 or R.HpaII RE + 9 µl distilled water. The reaction mixture was incubated at 37 °C for 3 hours. Bacteriophage λ DNA digested with HindIII and EcoRI was used as DNA molecular weight standards and unmethylated pUC19 DNA, transformed DH5α and Top10F' plasmid DNAs were spotted onto 1% agar gels and stained with ethidium bromide (EtBr), (4 µg/ml). The amount of DNA present was quantified by comparing the UV induced fluorescence emitted by EtBr molecules intercalated into the sample DNA with that of a series of markers spotted onto the same gel.

Results

In order to establish the effect of bacterial cytosine-5 M.Msp1 MTase on DH5α and Top10F' cells DNA, we have constructed a fusion of a bacterial cytosine-specific DNA MTase with a vertebrate nuclear targeting signal (NLS-SV40 VP1) and the marker enzyme glutathione-S-transferase (GST). Approximately 100 ng of pNRT2 encoding M.Msp1 gene as template and a 52 mer top strand, a 39 mer bottom strand as primers were used and PCR was set up according to the protocol which was outlined in the expand long template PCR system package (Techne) and explained in materials methods (Figure 1).

These DNA methyltransferase genes were separately cloned into the nonisogenic *E.coli* strains DH5α and Top10F' respectively. Transformed and control group colonies were selected with appropriate antibiotics (Figure 2).

These recombinant plasmids were stably maintained under selective growth conditions. DNA methylation was monitored at different times after induction by determining the susceptibility of the DNA to R.Msp1 digestion. After isolation of small scale (mini- prep) plasmid DNAs, from transformed *E.coli* strains that included NLS+GST+M.Msp1 recombinant gene, were digested with R.Msp1 for detection of methylation activity of fusion protein. This showed that production of bacterial methyltransferase can be expressed by the T7 and CMV promoters. The

fusion protein was found fully active and able to methylate CCGG sequences of the plasmids encoding the *M.Msp1* in *E.coli* strains used. The multicopy plasmid DNAs appeared to be completely methylated. We always observed uncut, high-molecular-weight DNA fragments on 1 % agar gels (Figure 3 and 4).




Figure 1. PCR Product Lane 1:HindIII and EcoRI digested (DNA marker Lanes 2-3 : (2000 bps, NLS+ GST+ *M.Msp1* gene fusion

Discussion

Approximately 90% of methylated Cs is followed on its 3' side by a G residue. Conveniently, this sequence forms part of the recognition sequence CCGG for two restriction enzymes *R.Msp1* and *R.HpaII* which differ in their ability to digest in this sequence when the C is methylated. Hence, if DNA is digested with either *R.Msp1* and *R.HpaII*, both enzymes will give the same pattern of bands only if all the C residues within the recognition sites are unmethylated (19).

We were able to express monospecific (enzyme modifying a single DNA sequence) cytosine-5 DNA MTase at high levels in two *E.coli* strains as functional fusions with glutathione-S-transferase and vertebrate nuclear targeting signal SV40 VP1. First we constructed the bacterial cytosine specific DNA *M.Msp1* MTase gene fusion including a nuclear localisation signal and glutathione-S-transferase and placed it under control of the T7 and CMV promoters (Figure 1).




Figure 2. *E.coli* DH5 α and Top10F' cells were transformed with PCR product (M.Msp1 gene fusion) and plated out in LB medium and selected with ampicillin for DH5 α , ampicillin and neomycin for Top10F'. After mini-prep DNA isolation, DNAs were digested with Hind III and EcoRI restriction endonucleases for selection of fragments carrying the fusion gene (~ 2000 bps). Lanes Lane 1, 2 and 3 : Positive DH5 α colonies Lane 4 : λ DNA digested with Hind III and EcoRI Lanes 5, 7, 8, 9 and 10 : Positive Top10F' colonies Lane 6 : Negative Top10F' colony




Figure 3. Agar gel electrophoresis (1%) of ethidium bromide stained plasmid DNA from *E.coli* DH5 α cells transformed with bacterial cytosine -5- specific DNA MTase M.Msp1 gene fusion .Lanes Lane1: Transformed cells' DNA digested with 10 U of R.Msp1. DNAs were resistant to the enzyme digestion and no fragmentations was found Lane 2 : HindIII and EcoRI digested λ DNA marker Lane 3 : pUC19 plasmid DNA digested with 10 U of R.Msp1.

After transformation of and establishment in two different non-isogenic *E.coli* cells, with the PCR product of the fusion genes construct, we have found that genes with CpG islands have a significant probability of becoming inactivated by methylation. The results indicate that the primary cause of preferential cutting is specific methylation of the CCGG sites in both strains. Expression of the M.Msp1 gene fusion was tested by the cognate restriction enzyme R.Msp1. Enzyme was active in both strains and methylated DNA could not be cut. Methylated DNA exhibited higher fragment sizes in 1% agar gel (Figure 3 and 4). Resistance to R.Msp1 digestion appeared to require at least 8 hours to achieve sufficient methylation to yield high-molecular-

weight protected fragments. This result suggests that methylation is non-random and certain large regions are inaccessible to bacterial C-5 MTase action within the two *E. coli* DH5 α and Top10F' strains. Therefore, foreign MTase genes with regulated expression and vertebrate nuclear targeting signal SV40 VP1 could be useful as an *in vivo* probe to test gene expression and chromatin organization.




Figure 4. Annealing profiles of transformed *E. coli* Top10F' and untransformed pUC19 plasmids DNA digested with 10 U of R.Msp1. The bacterial cytosine-5 MTase M.Msp1 fusion protein was active in all transformed strains and prevented the DNA fragmentation contrary to the restriction activity of cognate enzyme R.Msp1 (which can not cleave if the external cytosine is methylated in its recognition site CCGG). Lanes 1 and 4: Transformed *E. coli* Top10F' plasmid DNAs, no fragmentation. Lane 2: HindIII and EcoRI digested λ DNA marker. Lane 3: pUC19 plasmid DNA digested with 10 U of R.Msp1, high fragmentation.

References

1. Razin A, Riggs AD: DNA methylation and gene function. *Science* 210: 604 - 610, 1980.
2. Cheng X, Kumar S, Posfai JW and Roberts RJ: Crystal structure of the HhaI methyltransferase complexed with S-adenosyl-L-methionine. *Cell* 74: 299 - 307, 1993.
3. Wu JC and Santi DV: Kinetics and catalytic mechanism of HhaI methyltransferase. *J Biol Chem* 262: 4778-4786, 1987.
4. Antequera F, Boyes J and Bird A: High levels of de novo methylation and altered chromatin structure at CpG islands in cell lines. *Cell* 62: 502 -514, 1990.
5. Boyes J, Bird A: DNA methylation inhibits transcription indirectly via a methyl CpG binding protein. *Cell* 64:1123 -1134, 1991.
6. Brandeis M, Ariel M and Cedar H: Dynamics of DNA methylation during development. *BioEssays* 45: 709-713, 1993.
7. Sturm K, Taylor JH: Distribution of 5-methylcytosine in DNA of somatic and germline cells from bovine tissues. *Nucl Acids Res* 9: 4537-4546, 1981.

8. Laird PW, Grusby LJ, Faxeli A, Dickinson SL, Jung WE, Li E, Weinberg RA and Jaenisch R : Suppression of intestinal neoplasia by DNA hypomethylation . *Cell* 81: 197-205, 1995.
9. Clark SJ, Harrison J and Frammer M : CpNpG methylation in mammalian cells. *Nature Genet* 10: 20-27,1995.
10. Gabbara S Sheluho D and Bhagwat A : Cytosine methyltransferase from E.coli in which active site cysteine is replaced with serine is partially active. *Biochem J*(in press).
11. Wronski MA von , and Brent TP : Effect of 5-azacytidine on expression of the human DNA repair enzyme O6-methylguanine - DNA methyltransferase . *Carcinogenesis* 15: 577-582, 1994.
12. Gabbara S, and Bhagwat A : The mechanism of inhibition of DNA (cytosine-5-) methyltransferase by 5-azacytidine is likely to involve methyltransfer to the inhibitor. *Biochem J* 307 : 87-92, 1995 .
13. Lauster R, Trautner TA, Noyer-Weinder M: Cytosine-specific type II DNA methyltransferase. A conserved enzyme core with variable target-recognizing domains. *J Mol Biol* 206: 305-312 , 1989.
14. Yang AJ, Shen JC, Zingg JM, Mi S, and Jones PA: HhaI and HpaII DNA methyltransferases bind DNA mismatches, methylate uracil and block DNA repair. *Nucl Acids Res* 23 (8) : 1380-1387, 1995.
15. Bandaru B, Wyszyski M, and Bhagwat A : HpaII methyltransferase is mutagenic in E.coli *J of Bacteriol* 177 (10): 2950-2952 ,1995.
16. Renbaum P, and Razin A : Footprint analysis of M.SssI and M.HhaI methyltransferases reveals extensive interaction with the substrate DNA backbone . *J Mol Biol* 284: 19-26 , 1995.
17. Chen C and Okoayama H: High efficiency transformation of mammalian cells by plasmid DNA. *Mol Cell Biol* 7: 2745, 1987.
18. Sambrook J, Fritsch EF, Maniatis T : Edition 2 (*Molecular Cloning, A laboratory manuel* (Cold Spring Harbor Laboratory Press, Cold Spring Horbor 1989.
19. Ayté J, Gómez GG, and Hegardt FG : Methylation of regulatory region of the mitochondrial 3-hydroxy-3-methylglutaryl-CoA syntase gene leads to its transcriptional inactivation . *Biochem J* 295: 807-812, 1992.