

Second Hankel determinant for certain subclasses of bi-univalent functions involving Chebyshev polynomials

Halit ORHAN¹, Evrim TOKLU^{2,*}, Ekrem KADIOĞLU³

¹Department of Mathematics, Faculty of Science, Atatürk University, Erzurum, Turkey

²Department of Mathematics, Faculty of Education, Ağrı İbrahim Çeçen University, Ağrı, Turkey

³Department of Mathematics, Faculty of Science, Atatürk University, Erzurum, Turkey

Received: 16.07.2017

Accepted/Published Online: 03.05.2018

Final Version: 24.07.2018

Abstract: In this paper our purpose is to find the upper bound estimate for the second Hankel determinant $|a_2a_4 - a_3^2|$ for functions defined by convolution belonging to the class $\mathcal{N}_\sigma^{\mu,\delta}(\lambda, t)$ by using Chebyshev polynomials.

Key words: Univalent function, bi-univalent function, coefficient bounds, Chebyshev polynomial, Hankel determinant, convolution

1. Introduction

Let \mathcal{A} denote the class of functions $f(z)$ analytic in the open unit disk $\mathbb{U} := \{z \in \mathbb{C} : |z| < 1\}$ and normalized by

$$f(z) = z + \sum_{n \geq 2} a_n z^n. \quad (1.1)$$

Because of the Koebe one-quarter theorem it is well known that every univalent function $f \in \mathcal{A}$ has an inverse $f^{-1} : f(\mathbb{U}) \rightarrow \mathbb{U}$ satisfying

$$f^{-1}(f(z)) = z, \quad (z \in \mathbb{U})$$

and

$$f(f^{-1}(w)) = w, \quad (|w| < 1/4).$$

Moreover, it is easy to check that the inverse function has the series expansion of the form

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots, \quad w \in f(\mathbb{U}). \quad (1.2)$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in \mathbb{U} if both f and its inverse $g = f^{-1}$ are univalent in \mathbb{U} . Let σ denote the class of bi-univalent functions in \mathbb{U} given by (1.1). For a brief history of functions in the class σ , and also various other properties of the bi-univalent function, one can see recent works [2, 8, 12, 21, 26] and the references therein.

*Correspondence: evrimitoklu@gmail.com

2010 AMS Mathematics Subject Classification: 30C45

Some of the prominent and well-examined subclasses of univalent functions of class \mathcal{S} are those of the class $\mathcal{S}^*(\alpha)$ of starlike functions of order α in \mathbb{U} and the class $\mathcal{K}(\alpha)$ of convex functions of order α in \mathbb{U} . Moreover, by means of the analytic descriptions, we have

$$\mathcal{S}^*(\alpha) := \left\{ f : f \in \mathcal{A} \text{ and } \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha; \quad z \in \mathbb{U}; \quad 0 \leq \alpha < 1 \right\}$$

and

$$\mathcal{K}(\alpha) := \left\{ f : f \in \mathcal{A} \text{ and } \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha; \quad z \in \mathbb{U}; \quad 0 \leq \alpha < 1 \right\}.$$

For $0 \leq \alpha < 1$, a function $f \in \sigma$ is in the class $\mathcal{S}_\sigma^*(\alpha)$ of bi-starlike function of order α , or $\mathcal{K}_\sigma(\alpha)$ of bi-convex function of order α if both f and its inverse f^{-1} are, respectively, starlike or convex functions of order α .

We say that $f \in \mathcal{A}$ is subordinate to the function $g \in \mathcal{A}$ in \mathbb{U} , written $f(z) \prec g(z)$, if there exists a Schwarz function w , analytic in \mathbb{U} , with $w(0) = 0$ and $|w(z)| < 1$, and such that $f(z) = g(w(z))$.

For $f(z)$ given by (1.1) and $\Theta(z)$ defined by

$$\Theta(z) = z + \sum_{n \geq 2} \theta_n z^n, \quad (\theta_n \geq 0), \tag{1.3}$$

the Hadamard product (or convolution) $(f * \Theta)(z)$ of the functions $f(z)$ and $\Theta(z)$ is defined by

$$(f * \Theta)(z) = z + \sum_{n \geq 2} a_n \theta_n z^n = (\Theta * f)(z). \tag{1.4}$$

Next, we consider the function

$$\begin{aligned} f_\delta(z) &= \int_0^z \left(\frac{1+r}{1-r} \right)^\delta \frac{1}{1-r^2} dr \\ &= z + \delta z^2 + \frac{1}{3}(2\delta^2 + 1)z^3 + \dots \\ &= z + \sum_{n \geq 2} b_n(\delta)z^n, \quad (\delta > 0, \quad z \in \mathbb{U}). \end{aligned} \tag{1.5}$$

It is worth mentioning that if $\delta < 1$, then $zf'_\delta(z)$ is starlike with two slits. Moreover, we can see that since $zf'_1(z)$ is the Koebe function, all the functions f_δ are univalent and convex in \mathbb{U} . For more detail about the function $f_\delta(z)$ one can refer to [27]. If we put the function $f_\delta(z)$ defined by (1.5) in for the function $\Theta(z)$ given by (1.3) in the equality (1.4), we have

$$h_\delta(z) = (f * f_\delta)(z) = z + \sum_{n \geq 2} a_n b_n(\delta)z^n = (f_\delta * f)(z). \tag{1.6}$$

In 1976, Noonan and Thomas [20] defined the q th Hankel determinant of f given by (1.1) for integers $n \geq 1$ and $q \geq 1$ by

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \dots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \dots & a_{n+q-2} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n+q-1} & a_{n+q-2} & \dots & a_{n+2q-2} \end{vmatrix}, \quad (a_1 = 1).$$

This determinant has been investigated by several authors in the literature [13, 20]. For instance, this determinant is useful in showing that a function of bounded characteristic in \mathbb{U} , i.e. a function that is a ratio of two bounded analytic functions with its Laurent series around the origin having integral coefficients, is rational [9]. Moreover, it is important to mention that the Hankel determinants $H_2(1) = a_3 - a_2^2$ and $H_2(2) = a_2a_4 - a_3^2$ are well known as Fekete–Szegő and second Hankel determinant functionals, respectively. In 1969, the Fekete–Szegő problem for the classes \mathcal{S}^* and \mathcal{K} was investigated by Keogh and Merkes [17]. Recently, many authors have discussed upper bounds for the Hankel determinant of functions belonging to various subclasses of univalent functions (see [1, 11, 18] and the references therein). Very recently, the upper bounds of $H_2(2)$ for the classes $\mathcal{S}_\sigma^*(\alpha)$ and $\mathcal{K}_\sigma(\alpha)$ were investigated by Deniz et al. [12]. Later, the works were extended by Orhan et al. [23, 24] and Altınkaya and Yalçın [4].

Chebyshev polynomials, which are used by us in this study, play an important role in many branches of mathematics, especially in numerical analysis (see [10]). We know that there are several kinds of Chebyshev polynomials. In particular, we shall introduce the first and second kind of polynomials, $T_n(x)$ and $U_n(x)$. For a brief history of the Chebyshev polynomials of first kind $T_n(x)$ and second kind $U_n(x)$ and their numerous uses in different applications, one can refer [3, 14, 15].

The most remarkable kinds of the Chebyshev polynomials are the first and second kinds, and in the case of real variable x on $(-1, 1)$ they are defined by

$$T_n(x) = \cos(n \arccos x) \quad U_n(x) = \frac{\sin[(n + 1) \arccos x]}{\sin(\arccos x)} = \frac{\sin[(n + 1) \arccos x]}{\sqrt{1 - x^2}}.$$

Now we consider the function that is the generating function of a Chebyshev polynomial:

$$G(t, z) = \frac{1}{1 - 2tz + z^2}, \quad t \in \left(\frac{1}{2}, 1\right), z \in \mathbb{U}.$$

It is well known that if $t = \cos \theta$, $t \in (-\pi/3, \pi/3)$, then

$$\begin{aligned} G(t, z) &= 1 + \sum_{n \geq 1} \frac{\sin(n + 1)\theta}{\sin \theta} z^n \\ &= 1 + 2 \cos \theta z + (3 \cos^2 \theta - \sin^2 \theta) z^2 + \dots, \quad (z \in \mathbb{U}). \end{aligned}$$

That is, in view of [28], we can write

$$G(t, z) = 1 + U_1(t)z + U_2(t)z^2 + U_3(t)z^3 + \dots, \quad t \in \left(\frac{1}{2}, 1\right), z \in \mathbb{U}, \tag{1.7}$$

where $U_n(t)$ stands for the second kind of Chebyshev polynomials. From the definition of the second kind of Chebyshev polynomials, we easily arrive at $U_1(t) = 2t$. Also, it is well known that we have the following recurrence relation:

$$U_{n+1}(t) = 2tU_n(t) - U_{n-2}(t),$$

for all $n \in \mathbb{N}$. From here, we can easily obtain

$$U_1(t) = 2t, \quad U_2(t) = 4t^2 - 1, \quad U_3(t) = 8t^3 - 4t, \quad U_4(t) = 16t^4 - 12t^2 + 1, \dots \tag{1.8}$$

Definition 1 For $\lambda \geq 1$, $\mu \geq 0$, $\delta \geq 1$ and $t \in (1/2, 1]$, a function $h_\delta \in \sigma$ given by (1.6) is said to be in class $\mathcal{N}_\sigma^{\mu, \delta}(\lambda, t)$ if the following subordinations hold for all $z, w \in \mathbb{U}$:

$$(1 - \lambda)\left(\frac{h_\delta(z)}{z}\right)^\mu + \lambda h'_\delta(z)\left(\frac{h_\delta(z)}{z}\right)^{\mu-1} \prec G(z, t) \tag{1.9}$$

and

$$(1 - \lambda)\left(\frac{k_\delta(w)}{w}\right)^\mu + \lambda k'_\delta(w)\left(\frac{k_\delta(w)}{w}\right)^{\mu-1} \prec G(w, t), \tag{1.10}$$

where the function $k_\delta = h_\delta^{-1}$ is defined by (1.2).

Obviously, for $\delta = 1$, we get that $\mathcal{N}_\sigma^{\mu, 1}(\lambda, t) = \mathcal{N}_\sigma^\mu(\lambda, t)$. It is important to mention that the class $\mathcal{N}_\sigma^\mu(\lambda, t)$ was introduced and investigated by Bulut et al. [6]. They also discussed initial coefficient estimates and Fkete–Szegő bounds for the class $\mathcal{N}_\sigma^\mu(\lambda, t)$ and its subclasses, given in the following remark.

Remark 1 (i) For $\delta = 1$ and $\mu = 1$, we get the class $\mathcal{N}_\sigma^{1, 1}(\lambda, t) = \mathcal{B}_\sigma(\lambda, t)$ consisting of functions $f \in \sigma$ satisfying the condition

$$(1 - \lambda)\frac{f(z)}{z} + \lambda f'(z) \prec G(z, t)$$

and

$$(1 - \lambda)\frac{g(w)}{w} + \lambda g'(w) \prec G(w, t)$$

where the function $g = f^{-1}$ is defined by (1.2). This class was introduced and studied by Bulut et al. [7] (see also [19]).

(ii) For $\delta = 1$ and $\lambda = 1$, we obtain the class $\mathcal{N}_\sigma^{\mu, 1}(1, t) = \mathcal{B}_\sigma^\mu(t)$ consisting of bi-Bazilevič functions:

$$f'(z) \left(\frac{f(z)}{z}\right)^{\mu-1} \prec G(z, t)$$

and

$$g'(w) \left(\frac{g(w)}{w}\right)^{\mu-1} \prec G(w, t),$$

where the function $g = f^{-1}$ is defined by (1.2). This class was introduced and studied by Altınkaya and Yalçın [5].

(iii) For $\delta = 1$, $\mu = 1$, and $\lambda = 1$, we have the class $\mathcal{N}_\sigma^{1, 1}(1, t) = \mathcal{B}_\sigma(t)$ consisting of functions f satisfying the condition

$$f'(z) \prec G(z, t)$$

and

$$g'(w) \prec G(w, t),$$

where the function $g = f^{-1}$ is defined by (1.2).

(iv) For $\delta = 1$, $\lambda = 1$, and $\mu = 0$, we have the class $\mathcal{N}_\sigma^{0,1} = \mathcal{S}_\sigma^*(t)$ satisfying the condition

$$\frac{zf'(z)}{f(z)} \prec G(z, t)$$

and

$$\frac{wg'(z)}{g(w)} \prec G(w, t),$$

where the function $g = f^{-1}$ is defined by (1.2).

Let us take a look at some lemmas that are very useful in building our main results.

Let \mathcal{P} denote the class of analytic functions p in \mathbb{U} such that $p(0) = 1$ and $\operatorname{Re}(p(z)) > 0$, $z \in \mathbb{U}$. We know that this class is usually called the Carathéodory class.

Lemma 1 (see [25]) *If the function $p \in \mathcal{P}$ is given by the following series:*

$$p(z) = 1 + c_1z + c_2z^2 + c_3z^3 + \dots, \tag{1.11}$$

then the sharp estimate given by

$$|c_n| \leq 2 \quad (n = 1, 2, 3, \dots) \tag{1.12}$$

holds true.

Lemma 2 [16] *If the function $p \in \mathcal{P}$ is given by the series (1.11), then*

$$\begin{aligned} 2c_2 &= c_1^2 + x(4 - c_1^2), \\ 4c_3 &= c_1^3 + 2(4 - c_1^2)c_1x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z, \end{aligned}$$

for some x and z with $|x| \leq 1$ and $|z| \leq 1$.

In the present investigation, we seek the upper bound for the second Hankel determinant for functions h_δ belonging to the class $\mathcal{N}_\sigma^{\mu,\delta}(\lambda, t)$ by making use of the Chebyshev polynomial expansions and the Hadamard product. Also, we give some remarkable consequences related to the class $\mathcal{N}_\sigma^{\mu,\delta}(\lambda, t)$.

2. Main results

Theorem 1 *Let $h_\delta \in \sigma$ of the form (1.6) be in $\mathcal{N}_\sigma^{\mu,\delta}(\lambda; t)$. Then*

$$|a_2a_4 - a_3^2| \leq \begin{cases} \varphi(2^-, t), & \chi_1 \geq 0 \quad \chi_2 \geq 0 \\ \frac{36t^2}{(2\delta^2+1)^2(2\lambda+\mu)^2}, & \chi_1 \leq 0 \quad \chi_2 \leq 0 \\ \max \left\{ \frac{36t^2}{(2\delta^2+1)^2(2\lambda+\mu)^2}, \varphi(2^-, t) \right\}, & \chi_1 > 0 \quad \chi_2 < 0 \\ \max \{ \varphi(c_0, t), \varphi(2^-, t) \}, & \chi_1 < 0 \quad \chi_2 > 0 \end{cases},$$

where

$$\begin{aligned} \varphi(2^-, t) &= \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + \mu)^2} + \frac{\chi_1 + 9\chi_2}{6\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(2\lambda + \mu)^2(\lambda + \mu)^4}, \\ \varphi(c_0, t) &= -\frac{27\chi_2^2}{8\chi_1\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(2\lambda + \mu)^2(\lambda + \mu)^4} \\ &\quad + \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + \mu)^2}, \quad c_0 = \sqrt{\frac{-18\chi_2}{\chi_1}} \end{aligned}$$

and

$$\begin{aligned} \chi_1 &= 18(\lambda + \mu)^3 \left(3\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu) - (2\delta^2 + 1)^2(2\lambda + \mu)^2 \right) U_1^2(t) \\ &\quad + (2\lambda + \mu)^2 U_1(t) |\Omega_{\lambda, \mu, \delta}(t)| - 9(\lambda + \mu)^2(2\lambda + \mu) U_1(t) \left((3\lambda + \mu)(8\delta^4 - 4\delta^2 + 5) U_1^2(t) \right. \\ &\quad \left. + 4(2\delta^2 + 1)^2(2\lambda + \mu)(\lambda + \mu) U_2(t) \right), \\ \chi_2 &= \left[(2\lambda + \mu)(3\lambda + \mu) (8\delta^4 - 4\delta^2 + 5) U_1^3(t) + 4(2\delta^2 + 1)^2(\lambda + \mu)(2\lambda + \mu)^2 U_1(t) U_2(t) \right. \\ &\quad \left. + (\lambda + \mu) U_1^2(t) (2(2\delta^2 + 1)^2(2\lambda + \mu)^2 - 12\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)) \right] (\lambda + \mu)^2, \end{aligned}$$

$$\Omega_{\lambda, \mu, \delta}(t) = 18(2\delta^2 + 1)^2(\lambda + \mu)^3 U_3(t) - U_1^3(t)(3\lambda + \mu) \left(3(2\delta^2 + 1)^2(\mu^2 + 3\mu - 4) + 54\delta(\delta^3 + 2\delta) \right).$$

Proof Let $h_\delta \in \mathcal{N}_{\sigma}^{\mu, \delta}(\lambda, t)$. Then we have

$$(1 - \lambda) \left(\frac{h_\delta(z)}{z} \right)^\mu + \lambda h'_\delta(z) \left(\frac{h_\delta(z)}{z} \right)^{\mu-1} = G(t, u(z)) \tag{2.1}$$

and

$$(1 - \lambda) \left(\frac{k_\delta(w)}{w} \right)^\mu + \lambda k'_\delta(w) \left(\frac{k_\delta(w)}{w} \right)^{\mu-1} = G(t, v(w)) \tag{2.2}$$

where $p_1, p_2 \in \mathcal{P}$ and defined by

$$p_1(z) = \frac{1 + u(z)}{1 - u(z)} = 1 + c_1 z + c_2 z^2 + c_3 z^3 + \dots \tag{2.3}$$

and

$$p_2(w) = \frac{1 + v(w)}{1 - v(w)} = 1 + d_1 w + d_2 w^2 + d_3 w^3 + \dots \tag{2.4}$$

It follows from (2.3) and (2.4) that

$$u(z) = \frac{p_1(z) - 1}{p_1(z) + 1} = \frac{1}{2} \left[c_1 z + \left(c_2 - \frac{c_1^2}{2} \right) z^2 + \left(c_3 - c_1 c_2 + \frac{c_1^3}{4} \right) z^3 + \dots \right] \tag{2.5}$$

and

$$v(w) = \frac{p_2(w) - 1}{p_2(w) + 1} = \frac{1}{2} \left[d_1 w + \left(d_2 - \frac{d_1^2}{2} \right) w^2 + \left(d_3 - d_1 d_2 + \frac{d_1^3}{4} \right) w^3 + \dots \right]. \tag{2.6}$$

Using (2.5) together with (2.6), and taking $G(z, t)$ as given in (1.7), we get that

$$G(t, u(z)) = 1 + \frac{U_1(t)}{2} c_1 z + \left[\frac{U_1(t)}{2} \left(c_2 - \frac{c_1^2}{2} \right) + \frac{U_2(t)}{4} c_1^2 \right] z^2 + \left[\frac{U_1(t)}{2} \left(c_3 - c_1 c_2 + \frac{c_1^3}{4} \right) + \frac{U_2(t)}{2} c_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{U_3(t)}{8} c_1^3 \right] z^3 + \dots \tag{2.7}$$

and

$$G(t, v(w)) = 1 + \frac{U_1(t)}{2} d_1 w + \left[\frac{U_1(t)}{2} \left(d_2 - \frac{d_1^2}{2} \right) + \frac{U_2(t)}{4} d_1^2 \right] w^2 + \left[\frac{U_1(t)}{2} \left(d_3 - d_1 d_2 + \frac{d_1^3}{4} \right) + \frac{U_2(t)}{2} d_1 \left(d_2 - \frac{d_1^2}{2} \right) + \frac{U_3(t)}{8} d_1^3 \right] w^3 + \dots \tag{2.8}$$

By considering (2.1), (2.7) and (2.2), (2.8), when some elementary calculations are done, we get that

$$(\lambda + \mu) a_2 b_2(\delta) = \frac{U_1(t)}{2} c_1, \tag{2.9}$$

$$(2\lambda + \mu) \left[a_3 b_3(\delta) + (\mu - 1) \frac{a_2^2 b_2^2(\delta)}{2} \right] = \frac{U_1(t)}{2} \left(c_2 - \frac{c_1^2}{2} \right) + \frac{U_2(t)}{4} c_1^2, \tag{2.10}$$

$$(3\lambda + \mu) \left[a_4 b_4(\delta) + (\mu - 1) a_2 a_3 b_2(\delta) b_3(\delta) + (\mu - 1)(\mu - 2) \frac{a_2^3 b_2^3(\delta)}{6} \right] = \frac{U_1(t)}{2} \left(c_3 - c_1 c_2 + \frac{c_1^3}{4} \right) + \frac{U_2(t)}{2} c_1 \left(c_2 - \frac{c_1^2}{2} \right) + \frac{U_3(t)}{8} c_1^3, \tag{2.11}$$

and

$$-(\lambda + \mu) a_2 b_2(\delta) = \frac{U_1(t)}{2} d_1, \tag{2.12}$$

$$(2\lambda + \mu) \left[(\mu + 3) \frac{a_2^2 b_2^2(\delta)}{2} - a_3 b_3(\delta) \right] = \frac{U_1(t)}{2} \left(d_2 - \frac{d_1^2}{2} \right) + \frac{U_2(t)}{4} d_1^2, \tag{2.13}$$

$$(3\lambda + \mu) \left[(\mu + 4) a_2 a_3 b_2(\delta) b_3(\delta) - (\mu + 4)(\mu + 5) \frac{a_2^3 b_2^3(\delta)}{6} - a_4 b_4(\delta) \right] = \frac{U_1(t)}{2} \left(d_3 - d_1 d_2 + \frac{d_1^3}{4} \right) + \frac{U_2(t)}{2} d_1 \left(d_2 - \frac{d_1^2}{2} \right) + \frac{U_3(t)}{8} d_1^3. \tag{2.14}$$

Using (2.9) along with (2.12), we find that

$$c_1 = -d_1 \tag{2.15}$$

and

$$a_2 = \frac{U_1(t)}{2\delta(\lambda + \mu)}c_1. \tag{2.16}$$

Now, from (2.10), (2.13), and (2.16), we obtain that

$$a_3 = \frac{3}{2\delta^2 + 1} \left[\frac{U_1^2(t)}{4(\lambda + \mu)^2}c_1^2 + \frac{U_1(t)}{4(2\lambda + \mu)}(c_2 - d_2) \right]. \tag{2.17}$$

Also, subtracting (2.14) from (2.11) and using (2.16) together with (2.17), we get that

$$a_4 = \frac{3}{\delta^3 + 2\delta} \left[\left(\frac{U_1(t) - 2U_2(t) + U_3(t)}{8(3\lambda + \mu)} - \frac{(\mu^2 + 3\mu - 4)U_1^3(t)}{48(\lambda + \mu)^3} \right) c_1^3 + \frac{U_1(t)}{4(3\lambda + \mu)}(c_3 - d_3) \right. \\ \left. + \frac{5U_1^2(t)}{16(\lambda + \mu)(2\lambda + \mu)}c_1(c_2 - d_2) + \frac{U_2(t) - U_1(t)}{4(3\lambda + \mu)}c_1(c_2 + d_2) \right]. \tag{2.18}$$

Thus, we can easily determine that

$$|a_2a_4 - a_3^2| = \left| \frac{U_1(t)\Delta_{\lambda,\mu,\delta}(t)}{96\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(\lambda + \mu)^4}c_1^4 - \frac{9U_1^2(t)}{16(2\delta^2 + 1)^2(2\lambda + \mu)^2}(c_2 - d_2)^2 \right. \\ \left. + \frac{3U_1(t)[U_2(t) - U_1(t)]}{8\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)}c_1^2(c_2 + d_2) + \frac{3U_1^2(t)}{8\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)}c_1(c_3 - d_3) \right. \\ \left. + \frac{U_1^3(t)[15(2\delta^2 + 1)^2 - 36\delta(\delta^3 + 2\delta)]}{32\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(\lambda + \mu)^2(2\lambda + \mu)}c_1^2(c_2 - d_2) \right|, \tag{2.19}$$

where

$$\Delta_{\lambda,\mu,\delta}(t) = 18(2\delta^2 + 1)^2(\lambda + \mu)^3((U_1(t) - 2U_2(t) + U_3(t)) \\ - U_1^3(t)(3\lambda + \mu)(3(2\delta^2 + 1)^2(\mu^2 + 3\mu - 4) + 54\delta(\delta^3 + 2\delta))).$$

In view of Lemma 2 and (2.15), we write

$$c_2 - d_2 = \frac{4 - c_1^2}{2}(x - y), \tag{2.20}$$

$$c_2 + d_2 = c_1^2 + \frac{4 - c_1^2}{2}(x + y), \tag{2.21}$$

$$c_3 - d_3 = \frac{c_1^3}{2} + \frac{(4 - c_1^2)c_1}{2}(x + y) - \frac{(4 - c_1^2)c_1}{4}(x^2 + y^2) + \frac{4 - c_1^2}{2} [(1 - |x|^2)z - (1 - |y|^2)w], \tag{2.22}$$

for some x, y and z, w with $|x| \leq 1, |y| \leq 1, |z| \leq 1,$ and $|w| \leq 1$. Next, we will need to plug the last three equations given by (2.20), (2.21), and (2.22) into the Hankel functional given by equation (2.19). Also, with

the help of an application of the triangle inequality, we have

$$\begin{aligned} |a_2a_4 - a_3^2| &\leq \frac{U_1(t) |\Omega_{\lambda,\mu,\delta}(t)|}{96\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(\lambda + \mu)^4} c_1^4 + \frac{3U_1^2(t)c_1(4 - c_1^2)}{8\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} \\ &+ \left[\frac{(15(2\delta^2 + 1)^2 - 36\delta(\delta^3 + 2\delta)) U_1^3 c_1^2(4 - c_1^2)}{64\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(\lambda + \mu)^2(2\lambda + \mu)} + \frac{3U_1(t)U_2(t)c_1^2(4 - c_1^2)}{16\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} \right] (|x| + |y|) \\ &+ \left[\frac{3U_1^2(t)c_1^2(4 - c_1^2)}{32\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} - \frac{3U_1^2(t)c_1(4 - c_1^2)}{16\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} \right] (|x|^2 + |y|^2) \\ &+ \frac{9U_1^2(t)(4 - c_1^2)^2}{64(2\delta^2 + 1)^2(2\lambda + \mu)^2} (|x| + |y|)^2, \end{aligned}$$

where

$$\Omega_{\lambda,\mu,\delta}(t) = 18(2\delta^2 + 1)^2(\lambda + \mu)^3U_3(t) - U_1^3(t)(3\lambda + \mu) \left(3(2\delta^2 + 1)^2(\mu^2 + 3\mu - 4) + 54\delta(\delta^3 + 2\delta) \right).$$

Since class \mathcal{P} is invariant under the rotations, by (1.12) we may suppose without loss of generality that $c_1 := c \in [0, 2]$. Therefore, for $\eta = |x| \leq 1$ and $\zeta = |y| \leq 1$, we obtain

$$|a_2a_4 - a_3^2| \leq \kappa_1 + \kappa_2(\gamma_1 + \gamma_2) + \kappa_3(\gamma_1^2 + \gamma_2^2) + \kappa_4(\gamma_1 + \gamma_2)^2 = \psi(\gamma_1, \gamma_2)$$

where

$$\begin{aligned} \kappa_1 &= \frac{U_1(t) |\Omega_{\lambda,\mu,\delta}(t)|}{96\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(\lambda + \mu)^4} c^4 + \frac{3U_1^2(t)c(4 - c^2)}{8\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} \geq 0, \\ \kappa_2 &= \left[\frac{(15(2\delta^2 + 1)^2 - 36\delta(\delta^3 + 2\delta)) U_1^3 c^2(4 - c^2)}{64\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(\lambda + \mu)^2(2\lambda + \mu)} + \frac{3U_1(t)U_2(t)c^2(4 - c^2)}{16\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} \right] \geq 0, \\ \kappa_3 &= \frac{3U_1^2(t)c(c - 2)(4 - c^2)}{32\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)} \leq 0, \\ \kappa_4 &= \frac{9U_1^2(t)(4 - c^2)^2}{64(2\delta^2 + 1)^2(2\lambda + \mu)^2} \geq 0, \quad \frac{1}{2} < t < 1. \end{aligned}$$

Now let us consider the closed square $\mathbb{S} = \{(\gamma_1, \gamma_2) : 0 \leq \gamma_1 \leq 1, 0 \leq \gamma_2 \leq 1\}$. In that case, all that we need to do is to maximize the function $\psi(\gamma_1, \gamma_2)$ in the closed square \mathbb{S} for $c \in [0, 2]$. Since $\kappa_3 \leq 0$ and $\kappa_3 + 2\kappa_4 \geq 0$ for all $t \in (\frac{1}{2}, 1)$ and $c \in (0, 2)$, we conclude that

$$\psi_{\gamma_1\gamma_1}\psi_{\gamma_2\gamma_2} - (\psi_{\gamma_1\gamma_2})^2 < 0, \quad \text{for all } \gamma_1, \gamma_2 \in \mathbb{S}.$$

Thus, the function ψ cannot have a local maximum in the interior of the square \mathbb{S} . Now we investigate the maximum of ψ on the boundary of the square \mathbb{S} .

For $\gamma_1 = 0$ and $0 \leq \gamma_2 \leq 1$ (similarly $\gamma_2 = 0$ and $0 \leq \gamma_1 \leq 1$), we get

$$\psi(0, \gamma_2) = \phi(\gamma_2) = \kappa_1 + \kappa_2\gamma_2 + (\kappa_3 + \kappa_4)\gamma_2^2.$$

Next, we are going to be dealing with the following two cases separately.

Case 1 Let $\kappa_3 + \kappa_4 \geq 0$. In this case for $0 < \gamma_2 < 1$, and any fixed c with $0 \leq c < 2$ and for all t with $\frac{1}{2} < t < 1$, it is clear that $\phi'(\gamma_2) = 2(\kappa_3 + \kappa_4)\gamma_2 + \kappa_2 > 0$, that is, $\phi(\gamma_2)$ is an increasing function. Hence, for fixed $c \in [0, 2)$ and $t \in (1/2, 1)$, the function $\phi(\gamma_2)$ attains a maximum at $\gamma_2 = 1$ and

$$\max(\phi(\gamma_2)) = \phi(1) = \kappa_1 + \kappa_2 + \kappa_3 + \kappa_4.$$

Case 2 Let $\kappa_3 + \kappa_4 < 0$. Since $\kappa_2 + 2(\kappa_3 + \kappa_4) \geq 0$ for $0 < \gamma_2 < 1$, any fixed c with $0 \leq c < 2$, and all $t \in (1/2, 1)$, it is clear that $\kappa_2 + 2(\kappa_3 + \kappa_4) < 2(\kappa_3 + \kappa_4)\gamma_2 + \kappa_2 < \kappa_2$ and so $\phi'(\gamma_2) > 0$. Hence, for fixed $c \in [0, 2)$ and $t \in (\frac{1}{2}, 1)$, the function $\phi(\gamma_2)$ attains a maximum at $\gamma_2 = 1$.

For $\gamma_1 = 1$ and $0 \leq \gamma_2 \leq 1$ (similarly $\gamma_2 = 1$ and $0 \leq \gamma_1 \leq 1$), we get

$$\psi(1, \gamma_2) = \Psi(\gamma_2) = (\kappa_3 + \kappa_4)\gamma_2^2 + (\kappa_2 + 2\kappa_4)\gamma_2 + \kappa_1 + \kappa_2 + \kappa_3 + \kappa_4.$$

Thus, from the above cases of $\kappa_3 + \kappa_4$, we get that

$$\max \Psi(\gamma_2) = \Psi(1) = \kappa_1 + 2\kappa_2 + 2\kappa_3 + 4\kappa_4.$$

Since $\phi(1) \leq \Psi(1)$ for $c \in (0, 2)$ and $t \in (\frac{1}{2}, 1)$, we obtain

$$\max(\psi(\gamma_1, \gamma_2)) = \psi(1, 1)$$

on the boundary of the square \mathbb{S} . Thus, the maximum of ψ occurs at $\gamma_1 = 1$ and $\gamma_2 = 1$ in the closed square \mathbb{S} .

Let a function $\varphi : [0, 2] \rightarrow \mathbb{R}$ defined by

$$\varphi(c, t) = \max(\psi(\gamma_1, \gamma_2)) = \psi(1, 1) = \kappa_1 + 2\kappa_2 + 2\kappa_3 + 4\kappa_4 \tag{2.23}$$

for fixed values of t . Substituting the values of $\kappa_1, \kappa_2, \kappa_3$, and κ_4 in the function φ defined by (2.23) yields

$$\varphi(c, t) = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + \mu)^2} + \frac{\chi_1 c^4 + 36\chi_2 c^2}{96\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(2\lambda + \mu)^2(\lambda + \mu)^4},$$

where

$$\begin{aligned} \chi_1 = & 18(\lambda + \mu)^3 \left(3\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu) - (2\delta^2 + 1)^2(2\lambda + \mu)^2 \right) U_1^2(t) \\ & + (2\lambda + \mu)^2 U_1(t) |\Omega_{\lambda, \mu, \delta}(t)| - 9(\lambda + \mu)^2(2\lambda + \mu) U_1(t) \left((3\lambda + \mu)(8\delta^4 - 4\delta^2 + 5) U_1^2(t) \right. \\ & \left. + 4(2\delta^2 + 1)^2(2\lambda + \mu)(\lambda + \mu) U_2(t) \right), \\ \chi_2 = & \left[(2\lambda + \mu)(3\lambda + \mu) (8\delta^4 - 4\delta^2 + 5) U_1^3(t) + 4(2\delta^2 + 1)^2(\lambda + \mu)(2\lambda + \mu)^2 U_1(t) U_2(t) \right. \\ & \left. + (\lambda + \mu) U_1^2(t) (2(2\delta^2 + 1)^2(2\lambda + \mu)^2 - 12\delta(\delta^3 + 2\delta)(\lambda + \mu)(3\lambda + \mu)) \right] (\lambda + \mu)^2. \end{aligned}$$

Assuming that $\varphi(c, t)$ has a maximum value in an interior of $c \in [0, 2]$, by elementary calculation, we obtain that

$$\varphi'(c, t) = \frac{\chi_1 c^3 + 18\chi_2 c}{24\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(2\lambda + \mu)^2(\lambda + \mu)^4}.$$

We will examine the sign of the function $\varphi'(c, t)$ depending on the different cases of the signs of χ_1 and χ_2 as follows:

1. Let $\chi_1 \geq 0$ and $\chi_2 \geq 0$, and then $\varphi'(c, t) \geq 0$, so $\varphi(c, t)$ is an increasing function. Therefore,

$$\begin{aligned} \max \{\varphi(c, t) : c \in (0, 2)\} = \varphi(2^-, t) &= \frac{\chi_1 + 9\chi_2}{6\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(2\lambda + \mu)^2(\lambda + \mu)^4} \\ &+ \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + \mu)^2}. \end{aligned} \tag{2.24}$$

That is, $\max \{\max \{\psi(\gamma_1, \gamma_2) : 0 \leq \gamma_1, \gamma_2 \leq 1\} : 0 < c < 2\} = \varphi(2^-, t)$.

2. Let $\chi_1 \leq 0$ and $\chi_2 \leq 0$, and then $\varphi'(c, t) \leq 0$, so $\varphi(c, t)$ is an decreasing function on the interval $(0, 2)$. Therefore,

$$\max \{\varphi(c, t) : c \in (0, 2)\} = \varphi(0^+, t) = 4\kappa_4 = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + \mu)^2}. \tag{2.25}$$

3. Let $\chi_1 > 0$ and $\chi_2 < 0$, and then $c_0 = \sqrt{\frac{-18\chi_2}{\chi_1}}$ is a critical point of the function $\varphi(c, t)$. We suppose that $c_0 \in (0, 2)$, and since $\varphi''(c_0, t) > 0$, c_0 is a local minimum point of the function $\varphi(c, t)$. That is, the function $\varphi(c, t)$ cannot have a local maximum.

4. Let $\chi_1 < 0$ and $\chi_2 > 0$, and then c_0 is a critical point of the function $\varphi(c, t)$. We assume that $c_0 \in (0, 2)$. Since $\varphi''(c_0, t) < 0$, c_0 is a local maximum point of the function $\varphi(c, t)$ and the maximum value occurs at $c = c_0$. Therefore,

$$\max \{\varphi(c, t) : c \in (0, 2)\} = \varphi(c_0, t) \tag{2.26}$$

where

$$\varphi(c_0, t) = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + \mu)^2} - \frac{27\chi_2^2}{8\chi_1\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + \mu)(2\lambda + \mu)^2(\lambda + \mu)^4}.$$

Thus, from (2.24) to (2.26), the proof of **Theorem 1** is completed. □

Now we would like to draw attention to some remarkable results obtained for some values of λ , μ , and δ in **Theorem 1**.

Corollary 1 Let $h_\delta \in \sigma$ of the form (1.6) be in $\mathcal{B}_\sigma^\delta(\lambda, t)$. Then

$$|a_2 a_4 - a_3^2| \leq \begin{cases} \varphi(2^-, t), & \chi_3 \geq 0 \quad \chi_4 \geq 0 \\ \frac{36t^2}{(2\delta^2 + 1)^2(2\lambda + 1)^2}, & \chi_3 \leq 0 \quad \chi_4 \leq 0 \\ \max \left\{ \frac{36t^2}{(2\delta^2 + 1)^2(2\lambda + 1)^2}, \varphi(2^-, t) \right\}, & \chi_3 > 0 \quad \chi_4 < 0 \\ \max \{ \varphi(c_0, t), \varphi(2^-, t) \}, & \chi_3 < 0 \quad \chi_4 > 0 \end{cases},$$

where

$$\varphi(2^-, t) = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + 1)^2} + \frac{\chi_3 + 9\chi_4}{6\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + 1)(2\lambda + 1)^2(\lambda + 1)^4},$$

$$\varphi(c_0, t) = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2\lambda + 1)^2} - \frac{27\chi_4^2}{8\chi_3\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3\lambda + 1)(2\lambda + 1)^2(\lambda + 1)^4},$$

and $c_0 = \sqrt{\frac{-18\chi_4}{\chi_3}}$, $\chi_3 = \chi_1(\lambda, \mu = 1, \delta; t)$, $\chi_4 = \chi_2(\lambda, \mu = 1, \delta; t)$.

Taking $\lambda = 1$ in **Theorem 1**, we get the following result.

Corollary 2 Let $h_\delta \in \sigma$ of the form (1.6) be in $\mathcal{B}_\sigma^{\mu, \delta}(t)$. Then

$$|a_2a_4 - a_3^2| \leq \begin{cases} \varphi(2^-, t), & \chi_5 \geq 0 \quad \chi_6 \geq 0 \\ \frac{36t^2}{(2\delta^2+1)^2(2+\mu)^2}, & \chi_5 \leq 0 \quad \chi_6 \leq 0 \\ \max \left\{ \frac{36t^2}{(2\delta^2+1)^2(2+\mu)^2}, \varphi(2^-, t) \right\}, & \chi_5 > 0 \quad \chi_6 < 0 \\ \max \{ \varphi(c_0, t), \varphi(2^-, t) \}, & \chi_5 < 0 \quad \chi_6 > 0 \end{cases},$$

where

$$\varphi(2^-, t) = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2 + \mu)^2} + \frac{\chi_5 + 9\chi_6}{6\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3 + \mu)(2 + \mu)^2(1 + \mu)^4},$$

$$\varphi(c_0, t) = \frac{9U_1^2(t)}{(2\delta^2 + 1)^2(2 + \mu)^2} - \frac{27\chi_6^2}{8\chi_5\delta(\delta^3 + 2\delta)(2\delta^2 + 1)^2(3 + \mu)(2 + \mu)^2(1 + \mu)^4},$$

and $c_0 = \sqrt{\frac{-18\chi_6}{\chi_5}}$, $\chi_5 = \chi_1(\lambda = 1, \mu, \delta; t)$, $\chi_6 = \chi_2(\lambda = 1, \mu, \delta; t)$.

Putting $\lambda = 1$ and $\mu = 1$ in **Theorem 1**, we find the following result.

Corollary 3 If $h_\delta \in \mathcal{B}_\sigma^\delta(t)$ is of the form (1.6), then

$$|a_2a_4 - a_3^2| \leq \begin{cases} \varphi(2^-, t), & \frac{1}{2} < t \leq t_0 \\ \varphi(c_0, t), & t_0 < t < 1 \end{cases},$$

where

$$\varphi(2^-, t) = \frac{3t^2((2\delta^2 + 1)^2 - t(5\delta^4 + 2\delta^2 + 2))}{(\delta^2 + 2)(2\delta^3 + \delta)^2},$$

$$\begin{aligned} \varphi(c_0, t) = & - \frac{t(\nabla_{(4,20,-3);(22,40,64)}^{(3,1,-1)})^2}{\delta(2\delta^2 + 1)^2(\delta^3 + 2\delta)(\nabla_{(-4,4,-3);(22,40,64)}^{(3,1,-1)} + 6t|t(5\delta^4 + 2\delta^2 + 2) - (2\delta^2 + 1)^2|)} \\ & + \frac{4t^2}{(2\delta^2 + 1)^2}, \end{aligned}$$

$\nabla_{(a,b,c);(d,e,f)}^{(m,n,r)}(\delta; t) = \nabla_{(a,b,c);(d,e,f)}^{(m,n,r)} = m(1 + 2\delta^2)^2 + nt(a + bt^2 + ct^4) + rt^2(d + et^2 + ft^4)$. Moreover, the value of t_0 is the root of equation $\chi_1 = 0$ for $\lambda = \mu = 1$ and $\frac{1}{2} < t < 1$.

Setting $\delta = 1$ in **Corollary 3**, we obtain the following result.

Corollary 4 If $h_\delta \in \mathcal{B}_\sigma(t)$ is of the form (1.6), then

$$|a_2a_4 - a_3^2| \leq \begin{cases} t^2(1-t^2), & \frac{1}{2} < t \leq t_{0_1} \\ \frac{t(260t^4+84t^3-139t^2-18t+9)}{8(18t^3+42t^2-17t-9)}, & t_{0_1} < t < 1 \end{cases},$$

where the value of t_{0_1} , which is approximately $t_{0_1} = 0.603615$, is the root of equation $\chi_1 = 0$ for $\lambda = \mu = \delta = 1$ and $\frac{1}{2} < t < 1$.

Next, taking $\lambda = 1$ and $\mu = 0$ in Theorem 1, we arrive at the following result.

Corollary 5 If $h_\delta \in \mathcal{S}_\sigma^{*,\delta}(t)$ is of the form (1.6), then

$$|a_2a_4 - a_3^2| \leq \begin{cases} \varphi(2^-, t), & \frac{1}{2} < t \leq t_{0_2} \\ \varphi(c_0, t), & t_{0_2} < t < 1 \end{cases},$$

where

$$\begin{aligned} \varphi(2^-, t) &= \frac{8t^2((2\delta^2 + 1)^2 - 6t^2(\delta^2 - 1)^2)}{(2\delta^3 + \delta)^2(\delta^2 + 2)}, \\ \varphi(c_0, t) &= -\frac{t(\nabla_{(-2,10,1);(23,20,56)}^{(-2,-1,1)})^2}{\delta(2\delta^2 + 1)^2(\delta^3 + 2\delta)(\nabla_{(4,-2,7);(23,20,56)}^{(-4,1,2)} - 8t|6t^2(\delta^2 - 1)^2 - (2\delta^2 + 1)^2|)} \\ &\quad + \frac{9t^2}{(2\delta^2 + 1)^2}. \end{aligned}$$

Moreover, the value of t_{0_2} is the root of equation $\chi_1 = 0$ for $\lambda = 1$, $\mu = 0$, and $\frac{1}{2} < t < 1$.

Now, taking $\delta = 1$ in Corollary 5, we attain the following result.

Corollary 6 If $h_\delta \in \mathcal{S}_\sigma^*(t)$ is of the form (1.6), then

$$|a_2a_4 - a_3^2| \leq \begin{cases} \frac{8t^2}{3}, & \frac{1}{2} < t \leq \frac{7+\sqrt{401}}{44} \\ t^2 + \frac{t(2+t-11t^2)^2}{3(-4-7t+22t^2)}, & \frac{7+\sqrt{401}}{44} < t < 1 \end{cases}.$$

When $\delta = 1$, we note that the results given above coincide with the results in [22].

References

- [1] Ali RM, Lee SK, Ravichandran V, Supramanian S. The Fekete-Szegő coefficient functional for transforms of analytic functions. Bull Iranian Math Soc 2009; 35: 119-142.
- [2] Ali RM, Lee SK, Ravichandran V, Supramanian S. Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions. Appl Math Lett 2012; 25: 344-351.
- [3] Altınkaya Ş, Yalçın S. Chebyshev polynomial coefficient bounds for a subclass of bi-univalent functions. arXiv: 1605.08224v2.
- [4] Altınkaya Ş, Yalçın S. Upper bound second Hankel determinant for bi-Bazilevič. Mediterr J Math 2016; 13: 4081-4090.

- [5] Altınkaya Ş, Yalçın S. On the Chebyshev polynomial coefficient problem of bi-Bazilevič function. *Commun Fac Sci Univ Ank Series A1* 2017; 66: 289-296.
- [6] Bulut S, Magesh N, Abirami C. A comprehensive class of analytic bi-univalent functions by means of Chebyshev polynomials. *J Fract Calc Appl* 2017; 8: 32-39.
- [7] Bulut S, Magesh N, Balaji VK. Initial bounds for analytic and bi-univalent functions by means of Chebyshev polynomials. *J Class Anal* 2017; 11: 83-89.
- [8] Çağlar M, Orhan H, Yağmur N. Coefficient bounds for new subclasses of bi-univalent functions. *Filomat* 2013; 27: 1165-1171.
- [9] Cantor DG. Power series with the integral coefficients. *B Am Math Soc* 1963; 69: 362-366.
- [10] Chihara TS. *An Introduction to Orthogonal Polynomials*. New York, NY, USA: Gordon and Breach Science Publisher, 1978.
- [11] Deekonda VK, Thoutreedy R. An upper bound to the second Hankel determinant for functions in Mocanu class. *Vietnam J Math* 2015; 43: 541-549.
- [12] Deniz E, Çağlar M, Orhan H. Second Hankel determinant for bi-starlike and bi-convex functions of order β . *Appl Math Comput* 2015; 271: 301-307.
- [13] Dienes P. *The Taylor Series: An Introduction to the Theory of Functions of a Complex Variable*. New York, NY, USA: Dover, 1957.
- [14] Doha EH. The first and second kind Chebyshev coefficients of the moments of the general-order derivative of an infinitely differentiable function. *Int J Comput Math* 1994; 51: 23-35.
- [15] Dziok J, Raina RK, Sokól J. Applications of Chebyshev polynomials to classes of analytic functions. *C R Math Acad Sci Paris* 2015; 353: 433-438.
- [16] Grenander U, Szegő G. *Toeplitz Forms and Their Applications*. Berkeley, CA, USA: University California Press, 1958.
- [17] Keogh FR, Merkes EP. A coefficient inequality for certain classes of analytic functions. *P Am Math Soc* 1969; 20: 8-12.
- [18] Lee K, Ravichandran V, Supramaniam S. Bounds for the second Hankel determinant of certain univalent functions. *J Inequal Appl* 2013; 281: 1-17.
- [19] Mustafa N. Upper bound for the second Hankel determinant of certain subclass of analytic and bi-univalent functions. *arXiv-1702.06826*.
- [20] Noonan JW, Thomas DK. On the second Hankel determinant of a really mean p -valent functions. *T Am Math Soc* 1976; 223: 337-346.
- [21] Orhan H, Magesh N, Balaji VK. Fekete-Szegő problem for certain classes of Ma-Minda bi-univalent functions. *Afr Math* 2016; 27: 889-897.
- [22] Orhan H, Magesh N, Balaji VK. Second Hankel determinant for certain class of bi-univalent functions defined by Chebyshev polynomials. *Asian-Eur J Math* 2017. doi: 10.1142/S1793557119500177.
- [23] Orhan H, Magesh N, Yamini J. Bounds for the second Hankel determinant of certain bi-univalent functions. *Turk J Math* 2016; 40: 679-687.
- [24] Orhan H, Toklu E, Kadioğlu E. Second Hankel determinant problem for k -bi-starlike functions. *Filomat* 2017; 31: 3897-3904.
- [25] Pommerenke C. *Univalent Functions*. Gottingen, Germany: Vandenhoeck and Ruprecht, 1975.
- [26] Srivastava HM, Mishra AK, Gochhayat P. Certain subclasses of analytic and bi-univalent functions. *Appl Math Lett* 2010; 23: 1188-1192.
- [27] Trimble SY. A coefficient inequality for convex univalent functions. *P Am Math Soc* 1975; 48: 266-267.
- [28] Whittaker T, Watson GN. *A Course Of Modern Analysis*. Cambridge, UK: Cambridge University Press, 1996.