

Crustacean Biodiversity of *Padina pavonia* (L.) Facies Along the Aegean Coasts of Turkey

Fevzi KIRKIM, Ahmet KOCATAŞ, Tuncer KATAĞAN
Ege University, Fisheries Faculty, Hydrobiology Section, 35100, Bornova, İzmir - TURKEY

Murat SEZGİN
Ondokuz Mayıs University, Fisheries Faculty, Hydrobiology Section, 57000, Sinop - TURKEY
e-mail: msezgin@omu.edu.tr

A. Suat ATEŞ
Onsekiz Mart University, Fisheries Faculty, Hydrobiology Section, 18000, Çanakkale - TURKEY

Received: 24.03.2004

Abstract: This research was carried out to determine the crustacean species associated with *Padina pavonia* facies distributed in the upper-infralittoral zone of the Aegean Sea coasts of Turkey and their bioecological features. The investigations were performed at depths of 2-5 m in 13 different stations chosen in the Aegean Sea in June and July, 1995. As a result of the study, a total of 3279 specimens belonging to 85 species were identified. Of these, *Ampithoe ramondi* was commonest with a dominance value of 21.47%, followed by *Elasmopus pocillimanus* with 13.94% and *Erichthonius brasiliensis* with 4.61%.

Key Words: Diversity, *Padina pavonia*, Crustacea, Aegean Sea, Turkey

Türkiye'nin Ege Denizi Kıyıları *Padina pavonia* (L.) Fasiesinin Crustacea Çeşitliliği

Özet: Bu araştırma Türkiye'nin Ege Denizi kıyılarının üst infralittoral zonunda dağılım gösteren *Padina pavonia* fasiesinin Crustacea türlerini tespit etmek amacıyla yürütülmüştür. Araştırmalar 1995 yılının haziran ve temmuz aylarında Ege Denizi'nde seçilen 13 farklı istasyonda 2-5 m derinliklerde gerçekleştirilmiştir. Sonuç olarak 85 türe ait toplam 3279 birey tanımlanmıştır. Bunlardan *Ampithoe ramondi* % 21,47' lik dominansi değeri ile en yaygın tür olurken, bunu % 13,94 ile *Elasmopus pocillimanus* ve % 4,61 ile *Erichthonius brasiliensis* izlemektedir.

Anahtar Sözcükler: Diversite, *Padina pavonia*, Crustacea, Ege Denizi, Türkiye

Introduction

Photophilic algae represent a substrate with a particular structure populated by a great number of animals. The mass of these photophil organisms makes a marked contribution to raising the productivity of the marine coast (Tiganus, 1972).

Padina pavonia is a representative of brown algae (Phaeophyceae), usually located on rocky substrates, various shell bottoms and coral fragments in shallow waters, and show a wider distribution in unpolluted environments. Peres and Picard (1964) indicated that *P. pavonia* facies usually occur in shallow waters due to its tolerance to variations in edaphic factors. In the Aegean Sea *P. pavonia* is a member of the genus *Padina*, which is

represented by 4 species (*P. boergesenii*, *P. pavonia*, *P. tenuis* and *P. gymnospora*) in the Mediterranean Sea (European Register of Marine Species, 2003).

Bellan-Santini (1969) investigated the *P. pavonia* facies in unpolluted and calm waters of Marseilles Bay in the Mediterranean using qualitative and quantitative methods. Russo (1997) studied the epifauna associated with some algae species and *P. pavonia* on the coasts of Cyprus. Studies on the macrobenthic fauna inhabiting algae facies of the coastal hard substratum, especially *P. pavonia*, in the Turkish Aegean Sea are scarce. Studies were conducted by Kocataş (1978), Ergen (1980), Ergen et al. (1985), Ergen et al. (1994), and Öztürk and Ergen (2000), the majority of which concentrated on Mollusca and Polychaeta groups in the vicinity of İzmir Bay. The

most specific study on crustacean fauna within *P. pavonia* facies was carried out by Kocataş (1976).

Therefore, samplings in this study were made over a wider geographical range capable of representing the Turkish Aegean Sea coasts, and we aimed to examine the crustacean fauna associated with *P. pavonia* facies based on qualitative and quantitative data.

Materials and Methods

In order to determine crustacean assemblages within *P. pavonia* facies, samplings were performed at 13 different localities (1. Saros Bay-Güneyli, 2. Çanakkale-Monument, 3. Altınoluk, 4. Ayvalık, 5. Dikili, 6. Foça, 7. Urla, 8. Çeşme, 9. Sığacık-Seferihisar, 10. Kuşadası, 11. Bodrum, 12. Datça, and 13. Marmaris, from North to South) in the upper-infralittoral zone of the Aegean Sea.

Samples were collected according to the methodology proposed by Bellan-Santini (1969), and a 400 cm² unit

area was sampled for *P. pavonia* facies. For this purpose, a metal frame (20 x 20 cm) covered with a bag made up of a plankton net was used.

The *P. pavonia* roots and leaves within the metal frame were excavated using a spatula, and the material collected was preserved in 4% formalin for further analysis back in the laboratory. The samples were washed through a 1 mm sieve and the crustacean specimens were sorted. The extracted fauna was separated into taxonomic groups, identified and counted under a stereomicroscope. Groups were identified and listed according to the revisions given by Bacescu (1951) (Cumacea), Riggio (1973) (Tanaidacea), Giordani-Soika (1950), Holdich (1968; 1970) (Isopoda), Ruffo (1982, 1989, 1993, 1998) (Amphipoda) and Zariquiey Alvarez (1968), D'Udekem D'Acoz (1996) and Falciai and Minervini (1996) (Decapoda).

To elucidate the community structure, Soyer's (1970) frequency index (f %), Bellan-Santini's (1969)




Figure 1. Map of study area.

quantitative dominance index (DI %), Shannon-Weaver's (1949) diversity index (H^1), Pielou's (1975) evenness index and Bray-Curtis's (1957) similarity index were calculated.

The frequency index of a particular species was estimated by

$f = m / M \times 100$, where m = number of stations where the species was found and M = number of all stations.

The dominance index of a certain species was estimated by

$DI = m / M \times 100$, where m = individual number of a species in the stations and M = total individual numbers of all species.

The Shannon-Weaver diversity index was estimated by

$$H^1 = - \sum_{i=1}^n \log p_i \cdot \log_2 p_i$$

$$p_i = \frac{S}{N}$$

where S = total individual number of a species and N = total individual numbers of all species.

The Pielou evenness index was estimated by

$$J^1 = \frac{H^1}{\log_2 S}$$

where H^1 = Shannon index value; S = species number

The Bray-Curtis similarity index was estimated by

$$S_{jk} = 100 \{1 - \frac{|\sum |y_{ij} - y_{ik}|}{\sum |y_{ij} + y_{ik}|}\}$$

Results

As a result of the study, carried out at 13 different stations along the Turkish Aegean Sea coast, a total of 3279 individuals belonging to 85 species (2 Cumacea, 2 Tanaidacea, 14 Isopoda, 18 Decapoda and 49 Amphipoda) were recorded (Figure 2, Table 1).

The highest number of species was observed at station 6 (Foça) with 32 species, was followed by station 7 (Urla-Karantina Island) with 27 species and station 9 (Siğacık) with 25 species, and the lowest number was at station 12 (Datça) with 10 species. The highest number of specimens was at station 3 with 430 specimens, whereas the lowest value was recorded at station 4 (Ayvalık) (Figure 3).

The differences among the stations are in accordance with their substrate heterogeneity. At some stations *P. pavonia* constitutes a poor facies and its vicinity is surrounded with naked stones. However, at stations with both species and specimen richness at high values *P. pavonia* comprises dense facies and is covered with dense algae facies such as *Cystoseira* spp. and *Halopteris* spp. Crustacean species occur in dense populations in such biotopes.

Shannon-Weaver diversity index values (H^1) among the sampling stations did not show significant difference, and these values ranged between 4.32 and 2.56. The highest DI value was at station 6 (Foça) and the lowest DI value at station 8 (Çeşme) (Figure 3). The evenness index (J^1) values mainly ranged between 0.69 and 0.89, revealing that the distribution of species at the stations is regular. However, at station 8 (Çeşme), *Ampithoe*




Figure 2. Numbers of species and specimens belonging to Crustacea groups.

Table. List of species, numbers of individuals at stations, values of dominance and abundance.

Species	Stations													Σ	f %	DI%
	1	2	3	4	5	6	7	8	9	10	11	12	13			
Total individual	216	278	117	430	214	258	404	262	154	374	202	133	237	3279		
Total species	18	21	16	23	14	32	27	11	22	25	21	10	20	85		
Cumacea																
<i>Bodotria scorpioides</i> (Montagu, 1804)		2		1						2				23.08	0.15	
<i>Cumella limicola</i> Sars, 1879	3	2	1		29	1			3		1		1	61.54	1.25	
Tanaidacea																
<i>Apseudes robustus</i> G.O. Sars, 1882				1										7.69	0.03	
<i>Leptochelia savignyi</i> (Kroyer, 1842)	1	1	3		1	4			1		9		1	61.54	0.64	
Isopoda																
<i>Carpas stebbingi</i> (Monod, 1939)	2	3				5	36	12	17	13	5	18	10	76.92	3.69	
<i>Bopyrus squillarum</i> Latreille, 1802						1							2	15.38	0.09	
<i>Cymodoce emarginata</i> Leach, 1818	6	3	2	3	4	3				21				53.85	1.28	
<i>Cymodoce spinosa</i> (Risso, 1816)					4		5							15.38	0.27	
<i>Cymodoce truncata</i> Leach, 1814					4			3			2	3		30.77	0.37	
<i>Cymodoce tuberculata</i> Costa in Hope, 1851						3							3	15.38	0.18	
<i>Dynamene edwardsi</i> (Lucas, 1849)						5					4		6	23.08	0.46	
<i>Dynamene magnitorata</i> Holdich, 1968				3		2								15.38	0.15	
<i>Dynamene torelliae</i> Holdich, 1968						22	31		6	7				30.77	2.01	
<i>Joeropsis brevicornis</i> subsp. <i>littoralis</i> Amar, 1949		7				12							7	23.08	0.79	
<i>Uromunna petiti</i> (Amar, 1948)									2					7.69	0.06	
<i>Paranthura costana</i> Bate & Westwood, 1868				3					1					15.38	0.12	
<i>Synisoma capito</i> (Rathke, 1837)		4		2		1								23.08	0.21	
<i>Synisoma appendiculata</i> (Risso, 1816)	3												3	15.38	0.18	
Amphipoda																
<i>Ampelisca pseudospinimana</i> Bellan-Santini & Kaim-Malka, 1977						2								7.69	0.06	
<i>Amphilocheus neapolitanus</i> Della Valle, 1893							11		1	17		12		30.77	1.25	
<i>Ampithoe ferox</i> (Chevreux, 1902)						13								7.69	0.40	
<i>Ampithoe helleri</i> Karaman, 1975			2											7.69	0.06	
<i>Ampithoe ramondi</i> Audouin, 1826	55	55	22	130	50	50	101	90	27	47	27	25	25	100.00	21.47	
<i>Aora spinicornis</i> Afonso, 1976				3		7								15.38	0.30	
<i>Apherusa bispinosa</i> (Bate, 1857)	1		11											15.38	0.37	
<i>Apherusa chierieghinii</i> Giordani-Soika, 1950				17							7			15.38	0.73	
<i>Apherusa vexatrix</i> Krapp-Schickel, 1979						10			17	45				23.08	2.20	
<i>Atylus guttatus</i> (Costa, 1851)					8		7			30				23.08	1.37	
<i>Atylus massiliensis</i> Bellan-Santini, 1975										22				7.69	0.67	
<i>Caprella acanthifera</i> Leach, 1814						3				25				15.38	0.85	
<i>Caprella grandimana</i> Mayer, 1882		7								22				15.38	0.88	
<i>Caprella rapax</i> Mayer, 1890	9	27		22	17		17	12					14	53.85	3.60	
<i>Colomastix pusilla</i> Grube, 1861		3												7.69	0.09	
<i>Corophium acherusicum</i> Costa, 1851		7												7.69	0.21	
<i>Corophium</i> sp.			2											7.69	0.06	
<i>Cymadusa crassicornis</i> (Costa, 1857)		47	13	30										23.08	2.74	
<i>Dexamine spiniventris</i> (Costa, 1853)	1	32	25	32	26	9	17	27	15	12	5	31		92.31	7.08	
<i>Dexamine spinosa</i> (Montagu, 1813)						17	15			22		11		30.77	1.98	
<i>Elasmopus affinis</i> Della Valle, 1893											67			7.69	2.04	
<i>Elasmopus brasiliensis</i> (Dana, 1855)												6		7.69	0.18	
<i>Elasmopus pocillimanus</i> (Bate, 1862)	80		15	120	50	20	40		12	30	40		50	76.92	13.94	
<i>Erichthonius brasiliensis</i> (Dana, 1855)				22		12	22	75	12	8				46.15	4.61	
<i>Erichthonius punctatus</i> (Bate, 1857)						12	12							15.38	0.73	
<i>Eusiroides dellavallei</i> Chevreux, 1899			3											7.69	0.09	
<i>Gammarella fucicola</i> (Leach, 1814)			1	12										15.38	0.40	
<i>Guernea coalita</i> (Norman, 1868)		12	3	11				9						30.77	1.07	
<i>Hyale schmidtii</i> (Heller, 1866)			8				12			6			11	30.77	1.13	

Table (contuened)

Species	Stations													Σ	
	1	2	3	4	5	6	7	8	9	10	11	12	13		
<i>Hyale stebbingi</i> Chevreux, 1888											4			7.69	0.12
<i>Leucothoe spinicarpa</i> (Abildgaard, 1789)						4			1					15.38	0.15
<i>Lysianassa caesarea</i> Ruffo, 1987	10			10		4			4					30.77	0.85
<i>Lysianassa costae</i> (Milne-Edwards, 1830)		5			5					7			2	30.77	0.58
<i>Maera grossimana</i> (Montagu, 1808)						7								7.69	0.21
<i>Maera inaequipes</i> (Costa, 1857)						6	15			15			60	30.77	2.93
<i>Melita palmata</i> (Montagu, 1804)									6	6				15.38	0.37
<i>Microdeutopus algicola</i> Della Valle, 1893	20													7.69	0.61
<i>Microdeutopus chelifera</i> (Bate, 1862)			2								3			15.38	0.15
<i>Microdeutopus gryllotalpa</i> Costa, 1853		20												7.69	0.61
<i>Microdeutopus sporadhi</i> Myers, 1969												2		7.69	0.06
<i>Microdeutopus stationis</i> Della Valle, 1893						11							12	15.38	0.70
<i>Orchomene humilis</i> (Costa, 1853)									2					7.69	0.06
<i>Peltocoxa marioni</i> Catta, 1875												2		7.69	0.06
<i>Peltocoxa mediterranea</i> Schiecke, 1977						1								7.69	0.03
<i>Pereinotus testudo</i> (Montagu, 1808)	5	7			12	6	7			7				46.15	1.34
<i>Phtisica marina</i> Slabber, 1749		17									2			15.38	0.58
<i>Podocerus variegatus</i> Leach, 1814									13		1	23	13	30.77	1.52
<i>Stenothoe monoculoides</i> (Montagu, 1815)			4											7.69	0.12
<i>Stenothoe tergestina</i> Nebeski, 1881	8	16					15							23.08	1.19
Decapoda															
<i>Acanthonyx lunulatus</i> (Risso, 1816)							1		1	3	7		2	38.46	0.43
<i>Achaeus gracilis</i> O. G. Costa, 1839				1										7.69	0.03
<i>Athanas nitescens</i> (Leach, 1814)	6				3	3	1	1			1		1	53.85	0.49
<i>Eriphia verrucosa</i> (Forskål, 1775)							1							7.69	0.03
<i>Galathea bolivari</i> Zariquiey- Alvarez, 1950									2		1			15.38	0.09
<i>Hippolyte leptocerus</i> (Heller, 1863)								7	26		2	7		30.77	1.28
<i>Hippolyte</i> sp.	2													7.69	0.06
<i>Hippolyte garciaraso</i> D'Udekem d'Acoz, 1996								2						7.69	0.06
<i>Liocarcinus pusillus</i> (Leach, 1815)		1									1			15.38	0.06
<i>Lysmata seticaudata</i> (Risso, 1816)						1	17		6	2			7	38.46	1.01
<i>Pagurus chevreuxi</i> (Bouvier, 1896)				1					1		7			23.08	0.27
<i>Palaemon longirostris</i> (H. Milne Edwards, 1838)					1		4	4	4					30.77	0.40
<i>Pilumnus hirtellus</i> (Linnaeus, 1761)	1		2				1				1			30.77	0.15
<i>Pilumnus hirsutus</i> Stimpson, 1858										1				7.69	0.03
<i>Pirimela denticulata</i> (Montagu, 1808)	3						1							15.38	0.12
<i>Pisa hirticortis</i> (Herbst, 1804)				1										7.69	0.03
<i>Pisidia longimana</i> (Risso, 1816)			2		1	5	3			2			7	46.15	0.61
<i>Sirpus zariquieyi</i> Gordon, 1953				1			1							15.38	0.06




Figure 3. Numbers of species and specimens at stations.

ramondi and *Erichthonius brasiliensis* (Amphipoda), which are represented by 90 and 75 specimens respectively, lowered the evenness index value (Figure 4).

For the relative importance of crustacean species sampled from the 13 stations, the Soyer frequency index was computed and 9 species were designated as very common, 19 species as common and 57 species as rare. Four species, which had the highest values and are very common in *P. pavonia* facies, are *Ampithoe ramondi* (100%), *Dexamine spiniventris* (92.3%), *Elasmopus pocillimanus* and *Carpas stebbingi* (76.9%).

Apeudes robustus, *Uromunna petiti*, *Ampelisca pseudospinimana*, *Ampithoe helleri*, *Atylus massilensis*, *Colomastix pusilla*, *Corophium acherisicum*, *Corophium* sp., *Elasmopus affinis*, *E. brasiliensis*, *Eusiroides dellavallei*, *Hyale stebbingi*, *Maera grossimana*, *Microdeutopus algicola*, *M. gryllotalpa*, *M. sporadhi*, *Orchomene humilis*, *Peltocoxa marioni*, *P. mediterranea*,

Stenothoe monoculoides, *Acheus gracilis*, *Eriphia verrucosa*, *Hippolyte* sp., *Hippolyte garciaraso*, *Pilumnus hirsutus* and *Pisa hirticortis* were recorded from only 1 of the 13 stations and are rare species of this community. *A. ramondi* was the most dominant crustacean species with 771 specimens (21.48%) followed by *E. pocillimanus* with 513 specimens (13.9%) and *E. brasiliensis* 163 (4.6%) at *P. pavonia* facies (Figure 6).

According to results of the Bray-Curtis similarity index, 2 stations (station 1 and 5) with a value of 58%, and stations 7 and 8, and 10 and 12 with a value of 50% shared the same groups (Figure 7).

Discussion

A total of 3279 individuals belonging to 85 crustacean species were recorded from 13 different *P. pavonia* facies along the Aegean Sea coast of Turkey.




Figure 4. Comparison of evenness and diversity index values.




Figure 5. Dispersion of species as a result of 3 frequency index group values.




Figure 6. Dominance values of species.




Figure 7. Similarity among the stations (Bray-Curtis).

In studies previously carried out on *P. pavonia* facies Bellan-Santini (1969) reported 115 species (25 algae and 90 animal species) from Marseilles Bay in the Mediterranean Sea. Ergen et al. (1985) recorded 91 species and 1156 individuals belonging to different benthic macrofauna groups at Urla (station 7 in this study) and recorded the rate of crustacean species as 31.9%. Kocataş (1978) indicated that 28 out of 48 amphipod species were recorded in *P. pavonia* communities from the rocky substrates in the upper-infralittoral zone of İzmir Bay. Kocataş (1978) reported that 65 crustacean species, from a total of 231 species, were recorded in *P. pavonia* facies during his investigations into benthic forms of rocky shores in İzmir Bay. Ergen (1980) recorded 48 Polychaeta species in *P. pavonia* facies during his study to 5 facies in İzmir Bay. Ergen et al. (1994) reported 57 crustacean species on soft and hard bottoms including *P. pavonia* facies at

Gencelli Harbour (Aliğa-İzmir) Öztürk and Ergen (2000) reported 19 gastropod species in *P. pavonia* facies of 24 stations located on the Turkish coast of the Aegean Sea. When all the previous studies mentioned above are considered, our study includes the richest number of crustacea species.

A comparison of the sampling stations, in terms of the northern Aegean (1-8) and southern Aegean (8-13) revealed that the crustacean species diversity is somewhat higher in the southern Aegean Sea. This can be explained by the wider range of sampling stations that represent various habitats, and consequently by *P. pavonia* facies being exposed to different hydrographical conditions. In addition, the Aegean Sea, serving as a transition zone between the Mediterranean and Black Sea, also impacts on the wide distribution of marine organisms (Kocataş and Bilecik, 1992).

References

- Bacescu, M. 1951. Cumacea Fauna R.P.R. IV, Bucureşti, 94 p.
- Bellan-Santini, D., Karamani G., Krapp-Schickel, G., Ledoyer, M., Myers, A.A., Ruffo, S. and Schiecke, U. 1982. Gammaridea (Acanthonozomatidae to Gammaridae). In: Sandro Ruffo (ed.), The Amphipoda of the Mediterranean, Part I, Mémoires de l'Institut Océanographique, Monaco, 13: 364 p.
- Bellan-Santini, D. 1969. Contribution à l'étude des peuplement infralittoraux sur substrat rocheux (Etude qualitative et quantitative de la franch Superiere), Recherche Travaux Station Marine Endoume, France, 63 (47): 9-294.
- Bellan-Santini, D., Diviacco, G., Krapp-Schickel, G., Myers, A.A. and Ruffo, S. 1989. Gammaridea (Haustoriidae to Lysianassidae). In: Sandro Ruffo (ed.), The Amphipoda of the Mediterranean, Part II, Mémoires de l'Institut Océanographique, Monaco, 13: 365-576.
- Bellan-Santini, D., Karaman, G.S., Krapp-Schickel, G., Ledoyer, M. and Ruffo, S. 1993. Gammaridea (Melphidippidae to Talitridae) Ingolfiellidae, Caprellidae, In: Sandro Ruffo (ed.), The Amphipoda of the Mediterranean, Part III, Mémoires de l'Institut Océanographique, Monaco, 13: 577-813.

- Bellan-Santini, D., Karaman, G.S., Ledoyer, M., Myers, A.A., Ruffo, S. and Vader, W. 1998. Localities and Map, Addenda to Parts 1-3, Key to Families, Ecology, Faunistics and Zoogeography, Bibliography, Index. In: The Amphipoda of the Mediterranean, Sandro Ruffo (ed.). Part 4, Mémoires de l'Institut Océanographique, Monaco, 13: 815-959.
- Bray, J.R. and Curtis, J.T. 1957. An ordination of the upland forest communities of South Wisconsin. Ecological Monographs, 27: 325-347.
- D'Udekem D'Acoz, C. 1996. The Genus *Hippolyte* Leach, 1814 (Crustacea: Decapoda: Caridea: Hippolytidae) in the East Atlantic Ocean and the Mediterranean Sea, with a checklist of all species in the genus, Zool. Verh., 30 ix. : 1-133.
- European Register of Marine Species. 2003 Algae: brief check list, accessible via E.R.M.S Home Page on WWW.).
- Ergen, Z. 1980. İzmir Körfezi'nde üst infralittoral zonun bazı fasieslerinde Polychaeta'nın karşılaştırmalı olarak incelenmesi, TÜBİTAK VII. Bilim Kongresi, Kuşadası, Aydın, 275-284.
- Ergen, Z., Kocataş, A. and Katağan, T. 1985. Evolution Des Peuplements a *Padina pavonia* dans le Golfe d'İzmir (Turquie). Rapp. Comm. Int. Mer Médit., 29(5): 317-319.
- Ergen, Z., Kocataş, A., Katağan, T. and Çınar, M.E. 1994. Gencelli Limanı (Aliağa-İzmir) Bentik Faunası, E.Ü. Fen Fakültesi Dergisi Seri B, 16(2): 1047-1059.
- Falciari, L. and Minervini, R. 1996. Guide des Homards, Crabes, Longoustes, Crevettes et Autres Crustacés Décapodes d' Europe, Delachaux et Niestle SA, Lausanne-Paris, 287 p.
- Giordani-Soika, A. 1950. Tanaidacei degli Isopodi marini della laguna di Venezia, Archiv. Oceanogr. Limnologia, 7(2-3): 213-238.
- Holdich, D.M. 1968. Reproduction, growth and bionomics of *Dynamene bidentata* (Crustacea: Isopoda), J. Zool., London, 156: 137-153.
- Holdich, D.M. 1970. The distribution and habitat preferences of the Afro-European species of *Dynamene* (Crustacea: Isopoda), I. Nat. Hist., 4: 419-438.
- Kocataş, A. and Bilecik, N. 1992. Ege Denizi ve Canlı Kaynakları, T.C. Tarım ve Köyişleri Bakanlığı, Su Ürünleri Araştırma Enstitüsü Müdürlüğü, Bodrum, Seri A, 7: 1-88.
- Kocataş, A. 1976. Note préliminaire sur les Amphipodes recueillis dans les horizons supérieurs de l'étage infralittoral rocheux du golfe d'İzmir (Turquie). Téthys 7(2-3): 235-239.
- Kocataş, A. 1978. İzmir Körfezi kayalık sahillerinin bentik formları üzerinde kalitatif ve kantitatif araştırmalar, Ege Üniversitesi Fen Fakültesi Monografiler Serisi, İzmir, 12: 1-93.
- Öztürk, B. and Ergen, Z. 2000. Les Archéogastéropodes (Mollusca-Gastropoda) du littoral Turc de la Mer Egée. Acta Adriat., 41(2): 59-70.
- Peres, J.M. and Picard, J. 1964. Nouveau manuel de bionomie benthique de la Mer Méditerranée. Rec. Trav. Sta. Mar. Endoume 31(47): 5-137.
- Pielou, E.C. 1975. Ecological Diversity. Wiley-Inter Science Publ., London.
- Riggio, S. 1993. I Tanaidacei dei Mari Italiani: Quadro Delle Conoscenze. Bull. Mus. Civ. St. Nat. Verona, 20(2): 583-698.
- Russo, R.A. 1997. Epifauna living on sublittoral seaweeds around Cyprus, Hydrobiologia 344: 169-179.
- Shannon, C.E. and Weaver, V. 1949. A mathematical theory of communication, Univ. Pres, Illinois, Urbana: 101-117.
- Soyer, J. 1970. Bionomie benthique du plateau continental de la côte catalane française. III. Les peuplements de Copepodes harpacticoides (Crustacea), Vie et Milieu, 21: 337-511.
- Tiganuş, V. 1972. Ecologic observations on the Fauna Associated to the Cystoseira Belt along the Romanian Black Sea Coast. Cercetări Marine, I.R.C.M., 4: 153-167.
- Zariquiey Alvarez, R. 1968. Crustáceos Decápodos Ibéricos, investigación pesquera, 32, 510 p.