

Seasonal dynamics and spatial distribution of *Dactylogyrus crucifer* Wagener, 1857 on the gills of roach (*Rutilus rutilus* L.) from Lake Sapanca, Turkey

Erhan SOYLU^{1,*}, Bahaddin RÜZGAR², Meral SOYLU¹

¹Marmara University, Vocational School of Technical Sciences, Fisheries Department, İstanbul - TURKEY

²Marmara University, School of Banking and Insurance, İstanbul - TURKEY

Received: 25.08.2008

Abstract: This study determined the seasonal dynamics and spatial distribution of the monogenean parasite *Dactylogyrus crucifer* on the gills of roach *Rutilus rutilus* from Lake Sapanca, Turkey. A total of 183 specimens of roach between January and November 2005 were examined in monthly intervals. The highest mean intensity was recorded in March (227.1 ± 87.14) and the lowest in November (13.3 ± 8.2). Fifty three *R. rutilus* specimens were therefore investigated for spatial distribution of *D. crucifer* in March 2006. No significant differences in distribution were found according to any of the spatial determinants.

Key words: Roach, monogenean, seasonal occurrence, spatial distribution, Lake Sapanca

Sapanca Gölü (Türkiye)'nde Kızılöz Balığı (*Rutilus rutilus* L.)'nin solungaçlarında *Dactylogyrus crucifer* Wagener, 1857 in mevsimsel dinamikleri ve mekansal dağılımı

Özet: Bu çalışmada, Sapanca Gölünde *Rutilus rutilus*'un solungaçlarındaki monogenean parazit *Dactylogyrus crucifer*'in mevsimsel dinamikleri ve mekansal dağılımı belirlendi. 2005 yılı Ocak ve Kasım ayları arasında toplam 183 kızılöz balığı ile aylık olarak çalışılarak parazitin en yüksek ortalama yoğunluğu Mart'ta (227.1 ± 87.14) ve en düşük değeri Kasım'da (13.3 ± 8.2) bulundu. Bu nedenle 2006 yılı Mart ayında *D. crucifer*'in mekansal dağılımı araştırıldı. Mekansal dağılım yönünden hiçbir anlamlı fark bulunamadı.

Anahtar sözcükler: Kızılöz, monogenean, mevsimsel varlık, mekansal dağılım, Sapanca Gölü

Introduction

The spatial distribution of gill parasites has been investigated by several authors (El Hafidi et al., 1998; Dzika, 1999; Chapman et al., 2000; Matejusova et al., 2002; Simkova et al., 2002; Gutierrez and Martorelli,

1999; Turgut et al., 2006). The microhabitats and coexistence of *Dactylogyrus* species of *Rutilus rutilus* have also been investigated (Koskivaara et al., 1992; Koskivaara and Valtonen, 1992; Bagge and Valtonen, 1999; Simkova et al., 2000, 2002). Various factors have

* E-mail: esoylu@marmara.edu.tr

been reported to affect parasite populations on gills. Seasons and water temperature affect the occurrence and abundance of monogenoids (Koskivaara et al., 1991; Simkova et al., 2001) and the trophic status of a lake determines the presence and abundance of certain parasites (Wisniewski, 1958). Physiological and immunological factors of the fish host may determine the degree of infection (Dobson and May, 1987). Some fish develop immunity against the *Dactylogyrus* species (Paperna, 1964). Site specificity in monogeneans may result from physico-chemical requirements (Chapman et al., 2000). Variations in water current over the gill surface, or which arch has the greatest area, can affect the hemibranch preference of monogenoids (Gutierrez and Martorelli, 1999).

The present study investigated the monogenean parasite *D. crucifer* on the gills of *R. rutilus* from Lake Sapanca to determine its seasonal occurrence and spatial distribution in monospecific infections.

Materials and methods

A total of 183 *R. rutilus* were collected from the oligotrophic Lake Sapanca using gill nets once per month between January and November 2005 and examined for the occurrence of *D. crucifer*. Fish were transported to the laboratory alive in the lake water and killed immediately prior to examination. Their length and sex were recorded. The gill arches were removed fresh from the left and right side. All soft parts of a gill arch were isolated and all Dactylogyrid specimens were counted and identified using sclerotised parts of the parasites and the reproductive organs as per Gussev (Gussev, 1985).

In March 2006, the month in which the intensity of infection was found to be highest, 53 *R. rutilus* specimens (total length 19.0-28.0 cm) were examined for the spatial distribution of *D. crucifer* in monospecific infections. Parasites from the left and right gill arches were collected after fixation of the arches in 4% formalin solution. Each gill arch was placed in a Petri dish filled with distilled water. The left and right gill arches were numbered as I to IV, with gill arch I being nearest to the operculum. The 2 hemibranches of each gill arch were designated anterior and posterior. Each hemibranch was divided into 6 sectors, with proximal and distal parts being divided into dorsal, medial, and ventral areas (Figure).




Figure. Numbering of left branchial arch. 1, 2, 3 distal parts, 4, 5, 6 proximal parts.

The parasites were collected from each sector separately and the species of each identified on the basis of chitinized parts (haptor, copulatory organ). Microhabitats on the gill arch were classified in accordance with the following: El Hafidi et al. (1998) and Dzika (1999). The prevalence and mean intensities of the parasite species were determined according to Bush et al. (1997). The distribution of *D. crucifer* was analyzed by nonparametric statistics tests, Kruskal Wallis ANOVA Test, and followed by Mann Whitney U tests to determine the parasite population differences between the gill arches and between the segments.

Results

Of the 183 roach specimens (total length range: 17.0-36.0 cm; mean 23.42 ± 3.60) collected between January and November 2005, 149 were found to be infected with *D. crucifer* (prevalence 81.4%). During this period 14,549 specimens of *D. crucifer* were examined. The mean intensity and mean abundance of infection was detected as 95.8 ± 62.7 and 90.0 ± 66.2 , respectively. Statistical data of the collected fish are shown in Table 1. The mean intensities in male and female fish were found to be 75.90 ± 66.65 and 83.97 ± 76.58 , respectively, and there were no

Table 1. Number of uninfected and infected fish, prevalence, range, and mean intensity of *Dactylogyrus crucifer* on the gills of *Rutilus rutilus*.

| Months | Uninfected fish number | Infected fish number | | Prevalence of infection (%) | Intensity of infection min-max | Mean intensity of infection \pm SD |
|--------|------------------------|----------------------|--------|-----------------------------|--------------------------------|--------------------------------------|
| | | male | female | | | |
| Jan | 14 | 7 | 4 | 44.0 | 5-48 | 26.2 \pm 10.05 |
| Feb | 10 | 5 | 4 | 47.3 | 12-40 | 24.2 \pm 10.97 |
| Mar | | 2 | 7 | 100.0 | 123-387 | 227.1 \pm 87.14 |
| Apr | 5 | 7 | 6 | 72.2 | 36-170 | 113.3 \pm 39.78 |
| May | | 6 | 9 | 100.0 | 30-152 | 100.7 \pm 45.34 |
| Jun | | 8 | 5 | 100.0 | 73-236 | 128.8 \pm 48.64 |
| Jul | | 8 | 10 | 100.0 | 31-102 | 68.4 \pm 19.51 |
| Aug | | 9 | 8 | 100.0 | 41-208 | 92.3 \pm 41.45 |
| Sep | | 6 | 11 | 100.0 | 29-213 | 109.3 \pm 54.02 |
| Oct | | 6 | 11 | 100.0 | 33-195 | 149.7 \pm 55.74 |
| Nov | 5 | 5 | 5 | 66.6 | 1-27 | 13.3 \pm 7.78 |

significant differences of infection between male and female fish ($P > 0.05$). Although *D. crucifer* were the dominant parasite, *D. sphyrna* and *D. vistulae* were observed in May, June, and July only, but they were low in number and therefore not taken into account.

In the 53 roach specimens investigated in March 2006, to determine the spatial distribution of *D. crucifer*, a total of 4514 *D. crucifer* individuals were recorded. The total numbers in each sector of each

branchial arch are given in Table 2. The mean intensity of the parasites was 98.8.

An evaluation of the Kruskal Wallis test found no significant differences in distribution according to any of the spatial determinants. As a result, Mann Whitney U tests revealed no meaningful result. However, most of the differences in spatial distribution were considerable. Of the *D. crucifer*, 56.2% were in the anterior hemibranches and 48.8%

Table 2. Total numbers of *Dactylogyrus crucifer* in each sector of each branchial arch of *Rutilus rutilus* (n = 53).

| ARCH | | Sector of Branchial Arch | | | | | | TOTAL |
|-------|---|--------------------------|-----|-----|-----|------|-----|-------|
| | | 1 | 2 | 3 | 4 | 5 | 6 | |
| I | L | 45 | 159 | 83 | 144 | 151 | 91 | 673 |
| | R | 38 | 91 | 38 | 129 | 121 | 83 | 500 |
| II | L | 105 | 98 | 106 | 76 | 151 | 128 | 664 |
| | R | 128 | 151 | 91 | 38 | 113 | 90 | 611 |
| III | L | 91 | 136 | 83 | 151 | 212 | 98 | 771 |
| | R | 53 | 106 | 76 | 91 | 144 | 68 | 538 |
| IV | L | 106 | 114 | 53 | 45 | 83 | 30 | 431 |
| | R | 61 | 68 | 38 | 23 | 98 | 38 | 326 |
| TOTAL | | 627 | 923 | 568 | 697 | 1073 | 626 | 4514 |

in the posterior ($P = 0.114 > 0.05$) and 56.2% were in the left gill arches and 43.8 % in the right ($P = 0.149 > 0.05$). According to gill arch number, 16.8% were in gill arch IV and around 28% in average in the other 3 ($P = 0.177 > 0.05$). One difference, which was almost significant, was between the dorsal, medial, and ventral segments, where 44.2% were in the medial segment and around 28% in average were in the others ($P = 0.084 > 0.05$). This was reflected by the numbers in the sectors; 20.4% and 23.8% in (medial) sectors 2 and 5, respectively, and around 14% in average in the rest ($P = 0.013 > 0.05$). However, the difference between the numbers of *D. crucifer* in the proximal and distal parts was small; 53.1% and 46.9%, respectively, ($P = 0.386 > 0.05$), Table 3.

Discussion

Few studies on the distribution of monogenean species of the inland water fish of Turkey have been conducted. Özer and Öztürk (2005) studied *Dactylogyrus cornu* on the vimba (*Vimba vimba*

tanella) in the Sinop region of Turkey. Soylu (2007) studied the seasonal occurrence and site selection of *Paradiplozoon homoion* on the gills of *Pseudophoxinus antalyae*. The metazoan parasites of *R. rutilus* in Lake Sapanca have been previously investigated and *D. crucifer*, *D. sphyrna*, *D. vistulae*, *Diplostomum* sp, *Tylodelphys clavata*, *Palaeorchis incognitis*, *Aspidogaster limacoides*, glochidia of mollusks, and *Piscicola geometra* were recorded by Karabiber (2006). In the present study, 3 *Dactylogyrus* species, *D. crucifer*, *D. sphyrna*, and *D. vistulae*, were found on roach from the oligotrophic Lake Sapanca, although only *D. crucifer* prevailed throughout the year. Kogteva (1957) found *D. crucifer* on the gills of rudd *Scardinius erythrophthalmus*, roach *R. rutilus*, and white bream *Blicca bjoerkna* from the USSR. As well as *R. rutilus*, both *S. erythrophthalmus* and *B. bjoerkna* are present in Lake Sapanca, but *D. crucifer* has previously been reported only on the gills of *R. rutilus* (Soylu, 1991). In the present study *D. sphyrna* and *D. vistulae* were found in limited numbers in May, June, and July.

Table 3. Distribution of *Dactylogyrus crucifer* on the gill apparatus of *Rutilus rutilus*, single species infections.

| | P Kruskal-Wallis | No of <i>D. crucifer</i> | % of <i>D. crucifer</i> |
|----------------------|---------------------|-----------------------------|----------------------------|
| Left side | P = 0.149 > 0.05 | 2539 | 56.2 |
| Right side | | 1975 | 43.8 |
| Gill arch I | P = 0.177 > 0.05 | 1173 | 26.0 |
| Gill arch II | | 1275 | 28.2 |
| Gill arch III | | 1309 | 29.0 |
| Gill arch IV | | 757 | 16.8 |
| Dorsal segment | P = 0.084 > 0.05 | 1265 | 28.0 |
| Medial segment | | 1996 | 44.2 |
| Ventral segment | | 1253 | 27.8 |
| Proximal part | P = 0.386 > 0.05 | 2396 | 53.1 |
| Distal part | | 2118 | 46.9 |
| Sector 1 | P = 0.013 > 0.05 | 627 | 13.9 |
| Sector 2 | | 923 | 20.4 |
| Sector 3 | | 568 | 12.6 |
| Sector 4 | | 697 | 15.4 |
| Sector 5 | | 1073 | 23.8 |
| Sector 6 | | 626 | 13.9 |
| Anterior hemibranch | P = 0.114 > 0.05 | 2536 | 56.2 |
| Posterior hemibranch | | 1978 | 48.8 |

Koskivaara and Valtonen (1992) found *D. crucifer*, *D. nanus*, *D. suecicus*, *D. microcanthus*, *D. similis*, *D. cabollario*, *D. fallax*, *D. sphyrna*, and *D. vistulae* on roach from 3 small and interconnected but limnologically different lakes in central Finland. They reported finding all these dactylogyrids except for *D. vistulae* on roach in the oligotrophic Lake Peurunka. They claimed that *D. crucifer*, *D. nanus*, and *D. suecicus* are mainly found in large numbers in oligotrophic lakes. They identified *D. sphyrna* and *D. vistulae* as a satellite species originating from less common cyprinid species found in the same area (Koskivaara et al., 1991).

Ten dactylogyrid species from roach in the Morava river basin in the Czech Republic were reported by Kadlec et al. (2003). Simkova et al. (2001) found 6 dactylogyrid species on the same host individual and recorded *D. crucifer*, *D. nanus*, *D. rutilii*, and *D. suecicus* as common dactylogyrids on the gills of roach. However, on the roach in the present study from Lake Sapanca, the dactylogyrid diversity was far less rich.

The maximum mean intensity of *D. crucifer* on the gills of *R. rutilus* in Lake Sapanca was found in March. We found no significant difference in left-right preference. Similarly, others have found no significant differences in monogenean numbers between left and right gill arches of host fish (El Hafidi et al., 1998; Dzika, 1999). *Pseudodactylogyrus anguillae*, however, has been reported to prefer the left side of European eel, whereas *P. bini* was found mostly on the right side (Buchmann, 1988).

Parasites have been reported to prefer sheltered parts of the gill. Wooten (1974), studying ruffle respiratory currents, reports significantly higher water flowrates over the middle 2 gills, the ventral segments, and the distal parts of the gill filaments, but no significant difference in flowrates over anterior and posterior gill hemibranches. Oncomiracidia attach to the outside surface of the gill filaments and may migrate to the area of the arch sheltered from the direct effect of respiratory current (Paling, 1968). A great volume of water in the gill ventilating current passes over the 2nd and 3rd gill arches, spreading the cercariae onto the anterior gills, after which they actively select the preferred habitat (Madhavi, 1986). The findings of the present study were consistent with these reports, in that the greatest tendency, although not significant, was for the *D. crucifer* to prefer the more sheltered proximal parts.

This study found that in *R. rutilus*, *D. crucifer* was more abundant on the medial segments, anterior hemibranches, and on left side than on other microhabitats of the gills, although the differences were not very distinct. It also found that *D. crucifer* settled on almost all the sectors and that it was the dominant parasite species throughout the year, which means that there were no interspecific interactions. Ramasamy et al. (1985) suggest that the microhabitat extension can also occur in cases of high parasite density and intraspecific competition for space. This intraspecific competition was evident in the present study where *D. crucifer* was high in density and spread over almost all the gill arches and sectors.

References

- Bagge, A.M. and Valtonen, E.T. 1999. Development of monogenean communities on the gills of roach fry (*Rutilus rutilus*). *Parasitology* 118: 479-487.
- Buchmann, K. 1988. Spatial distribution of *Pseudodactylogyrus anguillae* and *P. bini* (Monogenea) on the gills of the European eel, *Anguilla anguilla*. *Journal of Fish Biology* 32: 801-802.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. and Shostak, A.W. 1997. Parasitology meets ecology on its own term: Margolis et al. revised. *Journal of Parasitology* 83: 575-583.
- Chapman, L.J., Lanciani, C.A. and Chapman, C.A. 2000. Ecology of a diplozoan parasite on the gills of the African cyprinid *Barbus neumayeri*. *African Journal of Ecology* 38: 312-320.
- Dobson, A.P. and May, R.M. 1987. The effects of parasites on fish populations-theoretical aspects. *International Journal of Parasitology* 17: 363-370.
- Dzika, E. 1999. Microhabitat of *Pseudodactylogyrus anguillae* and *P. bini* (Monogenea: dactylogyridae) on the gills of large-size European eel, *Anguilla anguilla* from Lake Gaj, Poland. *Folia Parasitologica* 46: 33-367.
- El Hafidi, F., Berrada-Rkhami, O., Benazzou, T. and Gabrion, C. 1998. Microhabitat distribution and coexistence of Microcotylidae (Monogenea) on the gills of striped mulled *Mugil cephalus*: change or competition? *Parasitological Research* 84: 315-320.

- Gussev, A. V. 1985. Key to the parasites of the freshwater fish fauna of the USSR II, Ed. O. N. Bauer Izdat Nauka, Leningrad. Keys to the Fauna of the USSR Vol. 143 pp. 424, Parasitic Monogeneans.
- Gutierrez, P. and Martorelli, S.R. 1999. Niche preferences and spatial distribution of Monogenea on the gills of *Pimelodus maculatus* in Rio de la Plata (Argentina). *Parasitology* 119: 183-188.
- Kadlec, D., Simkova, A., Jarkovsky, J. and Gelnar, M. 2003. Parasite communities of freshwater fish under flood conditions. *Parasitology Research* 89: 272-283.
- Karabiber, T. 2006. Metazoan parasites of roach (*Rutilus rutilus* L., 1758) from Lake Sapanca. Marmara Üniv. Institute in Pure and Applied Science. Msc Thesis. 45 pp. (in Turkish).
- Kogteva, E.P. 1957. Fish parasites from Pskov-Chud water reserve. In *Parasites and Diseases of Fish*. Bulletin of the All-union Scientific Research Institute of Lake and River Fisheries. Ed. G.K. Petrushevskii (English).
- Koskivaara, M., Valtonen, E.T. and Prost, M. 1991. Dactylogyrids on the gills of roach in central Finland features of infection and species composition. *International Journal for Parasitology* 21: 565-572.
- Koskivaara, M. and Valtonen, E. T. 1992. *Dactylogyrus* (Monogenea) communities on the gills of roach in three lakes in Central Finland. *Parasitology* 104: 263-272.
- Koskivaara, M., Valtonen, E.T. and Vouri, K.M. 1992. Microhabitat distribution and coexistence of *Dactylogyrus* species (Monogenea) on the gills of roach. *Parasitology* 104: 273-281.
- Madhavi, R. 1986. Distribution of metacercariae of *Centrocestus formosanus* (Trematoda: Heterophyidae) on the gills of *Aplocheilichthys panchax*. *Journal of Fish Biology* 29: 685-690.
- Matejusova, I., Simkova, A., Sasal, P. and Gelnar, M. 2002. Microhabitat distribution of *Pseudodactylogyrus anguillae* and *P. bini* among and within gill arches of the European eel (*Anguilla anguilla* L.). *Parasitology Research* 89: 290-296.
- Özer, A. and Öztürk, T. 2005. *Dactylogyrus cornu* Linstow, 1878 (Monogenea) infestation on the vimba (*Vimba vimba tanella* Nordmann, 1840) caught in the Sinop region of Turkey in relation to the host factors. *Turkish Journal of Veterinary and Animal Sciences* 29: 1119-1123.
- Paling, J.E. 1968. A method of estimating the relative volumes of water flow over the different gills of a freshwater fish. *Journal of Experimental Biology* 48: 533-544.
- Paperna, I. 1964. Competitive exclusion of *Dactylogyrus extensus* by *Dactylogyrus vastator* (Trematoda, Monogenea) on the gills of Reared Carp. *Journal of Parasitology* 50: 94-98.
- Ramasamy, P., Ramalingam, K., Hana, R.B.E. and Halton, D.W. 1985. Microhabitats of gill parasites (Monogenea and Copepoda) of Teleost (*Scomberoides* spp.). *International Journal for Parasitology* 15: 385-397.
- Simkova, A., Desdevises, Y., Gelnar, M. and Morand, S. 2000. Coexistence of nine gill ectoparasites (*Dactylogyrus*: Monogenea) parasitizing the roach (*Rutilus rutilus* L.) history and present ecology. *International Journal of Parasitology* 30: 1077-1088.
- Simkova, A., Gelnar, M. and Sasal, P. 2001. Aggregation of congeneric parasites (Monogenea: *Dactylogyrus*) among gill microhabitats within one host species (*Rutilus rutilus*). *Parasitology* 123: 599-607
- Simkova, A., Ondrackova, M., Gelnar, M. and Morand, S. 2002. Morphology and coexistence of congeneric ectoparasite species: reinforcement of reproductive isolation? *Biological Journal of the Linneans Society* 76: 125-135.
- Soylu, E. 1991. Monogeneans of some fishes from Lake Sapanca. İstanbul University, Institute of Marine Sciences and Geography Bulletin 8: 145-156.
- Soylu, E. 2007. Seasonal occurrence and site selection of *Paradiplozoon homoion* (Bychowsky and Nagibina, 1959) on the gills of *Pseudophoxinus antalyae* Bogutskaya, 1992 from Kepez-Antalya, Turkey. *European Association of Fish Pathologists Bulletin* 27: 71-74.
- Turgut, E., Shinn, A. and Wootten, R. 2006. Spatial distribution of *Dactylogyrus* (Monogenea) on the gills of the host fish. *Turkish Journal of Fisheries and Aquatic Sciences* 6: 93-98.
- Wisniewski, W. L. 1958. Characterisation of the parasitofauna of an eutrophic lake (Parasitofauna of the biocoenosis of Druznno Lake-part I.). *Acta Parasitologica Polonica*, 6: 1-64.
- Wootten, R. 1974. The spatial distribution of *Dactylogyrus amphibothrium* on the gills of ruffe *Gymnocephalus cernua* and its relation to the relative amounts of water passing over the parts of the gills. *Journal of Helminthology* 48: 167-174.