

## Nucleolar organizer region distribution in *Nannospalax ehrenbergi* (Nehring, 1898) (Rodentia: Spalacidae) from Iraq

Yüksel COŞKUN<sup>1</sup>, Nursel AŞAN BAYDEMİR<sup>2\*</sup>, Alaettin KAYA<sup>1</sup>, Ayşe Merve KARÖZ<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Science, Dicle University, Diyarbakır, Turkey

<sup>2</sup>Department of Biology, Faculty of Arts and Sciences, Kırıkkale University, Kırıkkale, Turkey

Received: 04.04.2013 • Accepted: 08.10.2013 • Published Online: 17.01.2014 • Printed: 14.02.2014

**Abstract:** The distribution of nucleolar organizer regions (NORs) in *Nannospalax ehrenbergi* from Iraq was determined for the first time. In the karyotype of *N. ehrenbergi* the diploid number, fundamental number, and fundamental autosomal number were 52, 76, and 72, respectively. NORs occur in the telomeric regions of submetacentric and acrocentric autosome pairs.

**Key words:** Ag-NOR banding, chromosomal form, Palestine mole rat, Iraq

Blind mole rats (family Spalacidae) are highly adapted to subterranean life and occupy dry and arid grassland regions in the eastern Mediterranean region and Europe (Topachevskii, 1969; Musser and Carleton, 2005). According to Nevo et al. (1995), Musser and Carleton (2005), and Kryštufek and Vohralik (2009), the subfamily Spalacinae consists of either a single genus, *Spalax*, or 2 genera, *Spalax* and *Nannospalax* (Topachevskii, 1969; Lyapunova et al., 1974; Gromov and Baranova, 1981; Zima and Kral, 1984). Musser and Carleton (2005) accepted the name *Nannospalax* as a synonym of *Spalax*. However, recently Németh et al. (2009), Kryštufek and Vohralik (2009), and Kandemir et al. (2012) discussed the taxonomic status of the taxon and decided to use the *Nannospalax* species as a superspecies. According to Kryštufek and Vohralik (2009) and Kandemir et al. (2012), the genus *Nannospalax* is represented by 3 superspecies in Turkey: *N. leucodon* in Turkish Thrace, *N. xanthodon* all over Anatolia, and *N. ehrenbergi* in southeastern Anatolia.

*Nannospalax ehrenbergi* was first described by Nehring in 1898 from Jaffa, Israel (Nehring, 1898). This species is distributed in southeastern Turkey, northern Iraq, Syria, Lebanon, Jordan, Israel, and North Africa (Topachevskii, 1969; Savic and Nevo, 1990; Musser and Carleton, 2005; Kryštufek and Vohralik, 2009). However, Seçkin and Coşkun (2006) determined that *N. ehrenbergi* is preferred by *Asio otus* as one of the mammalian diets from Diyarbakır Province, southeastern Turkey. Although more than 30 chromosomal forms were recorded from Turkey for the genus *Nannospalax*

(Coşkun, 2003; Aşan and Yağcı, 2006; Sözen et al., 2006b, 2011, 2013; Coşkun et al., 2009; Kandemir et al., 2012; Matur et al., 2013), and in addition several different karyotypes of *N. ehrenbergi* were described from across the distribution area (Coşkun 2004; Coşkun et al., 2006; Sözen et al., 2006a), only 1 chromosomal form was presented by Coşkun et al. (2012) from Mosul, Iraq.

So far G-, C-, and AgNOR-banding of different chromosomal forms of *Nannospalax xanthodon* and *N. leucodon* were achieved from Turkey by various authors (Ivanitskaya et al., 1997, 2008; Gülkaç and Küçükdumlu, 1999; Arslan and Bölükbaş, 2010; Arslan et al., 2011a, 2011b; Aşan Baydemir et al., 2013; Arslan et al., 2013; Arslan and Zima, 2013). However, only a few banded karyotypes of *N. ehrenbergi* were determined (Ivanitskaya et al., 1997; Gülkaç and Küçükdumlu, 1999). The aim of the present study was to present the nucleolar organizer region (NOR) distribution of the Palestine mole rat, *Nannospalax ehrenbergi*, from Iraq for the first time.

Two male specimens of *Nannospalax ehrenbergi* from the southwest of Mosul, Iraq (36°10'N, 43°03'E and 36°08'N, 43°01'E), were karyotyped. Mitotic metaphases were obtained from the bone marrow as described by Lee and Elder (1980). NORs were detected by the technique of Howell and Black (1980). At least 20 well-spread and Ag-NOR banded metaphase plates were photographed. The voucher specimens and the slides are deposited in the Department of Biology, Faculty of Science, Dicle University.

\* Correspondence: nurselasan@yahoo.com

The karyotype had  $2n = 52$ ,  $NF = 76$ , and  $NFa = 72$ . The chromosome set consisted of 11 meta-/submetacentric autosome pairs (nos. 1–11), and 14 acrocentric pairs decreasing in size (nos. 12–25). The X chromosome was a large metacentric chromosome while the Y was a small acrocentric. No secondary constriction was detected in the examined metaphase plates (Figure 1).


NORs were located only on 3 pairs of autosomes, at the telomeric regions of the short arms of 1 large (no. 2) and 1 small submetacentric chromosome (no. 11) pair and 1 acrocentric pair (no. 25) (Figure 2).

Five chromosomal forms ( $2n = 48, 52, 54, 56$ , and  $58$ ) were recorded for *N. ehrenbergi* by various authors from southeastern Anatolia (Ivanitskaya et al., 1997; Sözen et al., 1999, 2006; Coşkun et al., 2006, 2009; Kankılıç et al., 2007). According to Ivanitskaya et al. (1997), the most widely distributed chromosomal form of *N. ehrenbergi* is  $2n = 52$  and  $NFa = 72$ . Coşkun et al. (2012) stated that the chromosomal set of the Iraqi population of *Nannospalax ehrenbergi* was in accordance with the  $2n = 52$  population distributed in southeastern Anatolia.


Ag-NOR banded karyotypes of the  $2n = 52$  chromosomal form of *N. ehrenbergi* from Turkey were achieved by Ivanitskaya et al. (1997), and Gülkaç and Küçükdumlu (1999). Ivanitskaya et al. (1997) determined terminally located NORs in 3 pairs of biarmed autosomes whereas Gülkaç and Küçükdumlu (1999) recorded NORs in 2 pairs of large submetacentric autosomes. In addition to the Ag-NOR banded chromosomes recorded by Ivanitskaya et al. (1997) and Gülkaç and Küçükdumlu (1999), NORs were determined in an acrocentric pair in Iraqi specimens. As a consequence, the number of NOR-bearing chromosomes of *N. ehrenbergi* may facilitate future cytotaxonomic studies in the subfamily Spalacinae. However, molecular studies will also determine the genetic diversity of the genus *Spalax* in Turkey (Kankılıç et al., 2005; Arslan et al., 2010; Kandemir et al., 2012; Kankılıç and Gürpınar, 2013).

#### Acknowledgment

We thank the anonymous reviewers for their valuable contributions the manuscript.



**Figure 1.** Conventionally stained karyotype of *Nannospalax ehrenbergi* in Mosul province (Al-Jurn), Iraq (male).



**Figure 2.** Ag-NOR stained metaphase plate of *Nannospalax ehrenbergi* (arrows indicate NOR-bearing chromosomes).

## References

- Arslan A, Akan Ş, Zima J (2011a). Variation in C-heterochromatin and NOR distribution among chromosomal races of mole rats (Spalacidae) from Central Anatolia, Turkey. *Mamm Biol* 76: 28–35.
- Arslan A, Arisoy A, Zima J (2013). The chromosome banding pattern in two cytotypes (2n = 36 and 38) of blind mole rats from Turkey (Mammalia: Spalaxidae). *Zool Middle East* 59: 95–100.
- Arslan A, Bölükbaş F (2010). C-heterochromatin and NORs distribution of mole rat, *Nannospalax xanthodon* from Aksaray, Turkey. *Caryologia* 63: 398–404.
- Arslan E, Gülbağçe E, Arıkoğlu H, Arslan A, Buzan EV, Krystufek B (2010). Mitochondrial divergence between three cytotypes of the Anatolian mole rat, *Nannospalax xanthodon*. *Zool Middle East* 50: 27–34.
- Arslan A, Toyran K, Gözütok S, Yorulmaz T (2011b). C- and NOR stained karyotypes of mole rat, *Nannospalax xanthodon* (2n = 54) from Kırıkkale, Turkey. *Turk J Biol* 35: 655–661.
- Arslan A, Zima J (2013). The banded karyotype of the 2n = 58 chromosomal race of mole rats from Erzincan, Turkey. *Folia Zool* 62: 19–23.
- Aşan N, Yağcı T (2006). Karyotype and hair scale structure of *Nannospalax leucodon* (Nordmann, 1840) from Central Anatolia (Rodentia: Spalacidae). *Turk J Zool* 32: 125–130.
- Aşan Baydemir N, Yağcı T, Çakır Ş (2013). Karyological studies of the 2n = 60 cytotype of *Nannospalax nehringi* from Central Anatolia. *Caryologia*. 66: 49–53.
- Coşkun Y (2003). A study on the morphology and karyology of *Nannospalax nehringi* (Satunin, 1898) (Rodentia: Spalacidae) from Northeast Anatolia, Turkey. *Turk J Zool* 27: 171–176.
- Coşkun Y (2004). Morphological and karyological characteristics of *Nannospalax ehrenbergi* (Nehring, 1898) (Rodentia: Spalacidae) from Hatay province, Turkey. *Turk J Zool* 28: 205–212.
- Coşkun Y, El Namee A, Kaya A (2012). Karyotype of *Nannospalax ehrenbergi* (Nehring, 1898) (Rodentia: Spalacidae) in the Mosul Province, Iraq. *Hystrix* 23: 75–78.
- Coşkun Y, Kaya A, Yürümez G (2009). Chromosomal forms of the mole rat *Nannospalax nehringi* (Satunin, 1898) from the Van Lake Basin in Eastern Turkey. *Zool Middle East* 48: 17–24.
- Coşkun Y, Ulutürk S, Yürümez G (2006). Chromosomal diversity in mole-rats of the species *Nannospalax ehrenbergi* (Rodentia: Spalacidae) from South Anatolia, Turkey. *Z Saugetierkd* 71 : 244–250.
- Gromov IM, Baranova GI (1981). Catalogue of Mammals in USSR. Leningrad: Nauka.
- Gülkaç D, Küçükdumlu İ (1999). Variation in the nucleolus organizer regions (NORs) in two mole rat species (*Spalax leucodon* and *Spalax ehrenbergi*). *Turk J Biol* 23: 153–158.
- Howell WM, Black DA (1980). Controlled silver staining for nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 36: 1014–1015.
- Ivanitskaya E, Coşkun Y, Nevo E (1997). Banded karyotypes of mole rats (*Spalax*, Spalacidae, Rodentia) from Turkey: a comparative analysis. *J Zool Syst Evol Research* 35: 171–174.
- Ivanitskaya E, Sözen M, Rashkovetsky L, Matur F, Nevo E (2008). Discrimination of 2n = 60 *Spalax leucodon* (Spalacidae, Rodentia) in Turkey by means of classical and molecular cytogenetic techniques. *Cytogenet Genome Res* 122: 139–149.
- Kandemir İ, Sözen M, Matur F, Kankılıç T, Martinkova N, Çolak F, Özkurt Ş, Çolak E (2012). Phylogeny of species and cytotypes of mole rats (Spalacidae) in Turkey inferred from mitochondrial cytochrome b gene sequences. *Folia Zool* 61: 25–33.
- Kankılıç T, Çolak E, Çolak R, Yiğit N (2005). Allozyme variation in *Spalax leucodon* Nordmann, 1840 (Rodentia: Spalacidae) in the area between Ankara and Beyşehir. *Turk J Zool* 29: 377–384.
- Kankılıç T, Gürpınar C (2014). Revised classification design of the Anatolian species of *Nannospalax* (Rodentia: Spalacidae) using a RFLP analysis. *Turk J Zool* 38: 67–78.
- Kankılıç T, Kankılıç T, Çolak R, Çolak E, Karataş A (2007). Karyological comparison of populations of the *Spalax leucodon* Nordmann 1840 superspecies (Rodentia: Spalacidae) in Turkey. *Zool Middle East* 42: 15–24.
- Kryštufek B, Vohralik V (2009). Mammals of Turkey and Cyprus. Rodentia II. Cricetinae, Muridae, Spalacidae, Calomyscidae, Capromyidae, Hystricidae, Castoridae. Koper, Slovenia: Knjižnica Annales Majora, University of Primorska Science and Research Centre.
- Lee MR, Elder FF (1980). Yeast stimulation of bone marrow mitosis for cytogenetic investigations. *Cytogenet Cell Genet* 26: 36–40.
- Matur F, Çolak F, Ceylan T, Sevidik M, Sözen M (2013). Chromosomal evolution of the genus *Nannospalax* (Palmer 1903) (Rodentia, Muridae) from western Turkey. *Turk J Zool* 37: 470–487.
- Musser GG, Carleton MD (2005). Family Muridae. In: Wilson DE, Reeder DM, editors. *Mammal Species of the World: A Taxonomic and Geographic Reference*. 3rd ed. Washington, DC: Smithsonian Institution Press, pp. 1189–1531.
- Nehring A (1898). Über mehrere neue *Spalax* Arten. *Sitzungsber. d. Gesellsch. Naturforsch Freunde z Berlin* 10: 151–183.
- Németh A, Révay T, Hegyeli Z, Farkas J, Czabán D, Rózsás A, Csorba G (2009). Chromosomal forms and risk assessment of *Nannospalax* (superspecies *leucodon*) (Mammalia: Rodentia) in the Carpathian Basin. *Folia Zool* 58: 349–361.
- Savic I, Nevo E (1990). The Spalacidae: Evolutionary history, speciation and population biology. In: Reig O, editor. *Evolution of Subterranean Mammals at the Organismal and Molecular Levels*. New York: Alan R Liss, Inc., pp. 129–153.
- Seçkin S, Coşkun Y (2006). Mammalian remains in the pellets of the long-eared owls (*Asio otus*) in Diyarbakır province. *Turk J Zool* 30: 271–278.

- Sözen M, Çataklı K, Eroğlu F, Matur F, Sevindik M (2011). Distribution of chromosomal forms of *Nannospalax nehringi* (Satunin, 1898) (Rodentia: Spalacidae) in Çankırı and Çorum provinces, Turkey. Turk J Zool 35: 367–374.
- Sözen M, Çolak F, Sevindik M, Matur F (2013). Cytotypes of *Nannospalax xanthodon* (Satunin, 1898) (Rodentia: Spalacidae) from western Anatolia. Turk J Zool 37: 462–469.
- Sözen M, Çolak E, Yiğit N, Özkurt Ş, Verimli R (1999). Contributions to the karyology and taxonomy of the genus *Spalax* Gldenstaedt, 1770 (Mammalia: Rodentia) in Turkey. Z Saugetierkd 64: 210–219.
- Sözen M, Matur F, Çolak E, Özkurt Ş, Karataş A (2006a). Some karyological records and a new chromosomal form for *Spalax* (Mammalia: Rodentia) in Turkey. Folia Zool 55: 247–256.
- Sözen M, Sevindik M, Matur F (2006b). Karyological and some morphological characteristics of *Spalax leucodon* Nordmann, 1840 (Mammalia: Rodentia) superspecies around Kastamonu province, Turkey. Turk J Zool 30: 205–219.
- Topachevskii VA (1969). Fauna of the USSR. Mammals: Mole Rats, Spalacidae. Vol. 3, No. 3. Leningrad: Academia Nauka.