

Herpetofauna of the Lake District around Burdur

Oğuzhan EGE¹, Batuhan Yaman YAKIN², Cemal Varol TOK^{2*}

¹Department of Biology, Zoology Section, Faculty of Sciences, Ege University, İzmir, Turkey

²Department of Biology, Zoology Section, Faculty of Sciences and Arts, Çanakkale Onsekiz Mart University, Terzioğlu Campus, Çanakkale, Turkey

Received: 03.02.2015 • Accepted/Published Online: 27.07.2015 • Printed: 30.11.2015

Abstract: This study investigated the amphibian and reptilian fauna of the Lake District around Burdur, Turkey. A total of 21 amphibian and reptile species were identified, consisting of two anurans, two turtles, one tortoise, seven lizards, and nine snakes. A zoogeographical assessment was also carried out in which the chorotypes and the IUCN criteria of the species were added.

Key words: Burdur, Lake District, herpetofauna, Amphibia, Reptilia, chorotype, zoogeography

In the surveys compiled by Baran (1986) and Demirsoy (1997a, 1997b) of studies of the amphibians and reptiles in Turkey up to recent times, it was found that researchers had mostly either investigated species one by one or else compared samples from different regions. However, there has been an increase in the number of studies detailing all herpetofauna in a specific region. In general, collecting specimens from different localities in specific regions is more complicated. Thus, it is more important to determine and describe the entire amphibian and reptilian species of a specific region (Baran, 1980, 1984; Yılmaz, 1984, 1989; Arıkan, 1988, 1994; Tok, 1993, 1995; Kumlutaş et al., 1998, 1999, 2001, 2004a, 2004b, 2011; Özdemir, 1998; Özdemir and Baran, 2002; Ilgaz and Kumlutaş, 2005; Kaya, 2005; Hür et al., 2008; Bulut, 2010; Afsar and Tok, 2011; Bulut et al., 2011; Uysal, 2011; Afsar et al., 2012; Tok et al., 2012a, 2012b; Cihan and Tok, 2014; Özcan and Üzümlü, 2014; Tok and Çiçek, 2014). Furthermore, some regions have been granted status as legally protected areas in order to conserve their natural richness. In particular, Lake Burdur has been singled out for conservation by the Ramsar Convention on Wetlands of International Importance as a Wildlife Protection Area, as both an important area for amphibian, reptile, bird, and mammalian species and an important wetland. The lack of a comprehensive study on the herpetofauna of this area up to now enhances the importance of this study.

The aim of this study was to determine the amphibian and reptilian species of Burdur Province, part of the Turkish Lake District in southwestern Anatolia, and to

make a zoogeographical assessment by determining the distributions of those species.

The study area, the province of Burdur, covers an area of 6887 km² between 36°53'N–37°50'N and 29°24'E–30°53'E. The area is formed of tectonic and karstic sedimentary basins between the Western Taurus Mountains. Because of the six lakes (Karataş, Salda, Yarıklı, Akgöl, Gölhisar, and Burdur) within the tectonic sedimentary basins, this region is usually called the Lake District. The name “Burdur” applies to both the province and its regional capital, as well as the lake nearby. In this study, it generally denotes the province, which includes the six lakes.

In order to determine the herpetofauna of the study area, a total of 131 amphibian and reptilian specimens were collected between September 1999 and May/June 2000 from the vicinity of Burdur in field trips. The specimens were identified by the literature (Başoğlu and Baran, 1977, 1980; Özeti and Yılmaz, 1994). The preserved specimens are in the collection of the Ege University Zoology Department, İzmir (ZDEU). Turtles and tortoises were released after examination. Evaluation of the current conservation status of the species was made. Zoogeographical assessments were carried out in consideration of the species' origin by classifying them into major chorotypes according to Vigna Taglianti et al. (1999) and Sindaco et al. (2000).

Species were captured using different methods. Aquatic species were caught by hand or with a scoop, and terrestrial species by hand or by use of a net.

As a result of the research, 2 amphibian and 19 reptilian species were found inhabiting the research area.

* Correspondence: cvtok@comu.edu.tr

The localities in which specimens were collected are given in Figure 1 and the Table.

Herpetological examination of the Lake District around Burdur shows that the species in this area are arranged into eight major chorotypes, two for amphibians and seven for reptiles, according to the classification of Vigna Taglianti et al. (2000) (Table). The most common chorotype in the area is E-Mediterranean (six species, 28.6%), and then respectively Turano-Mediterranean (four species, 19%), Turano-Europeo-Mediterranean (three species, 14.4%), SW-Asiatic (three species, 14.4%), Central Asiatic-European (2 species, 9.5%), Euro-Siberian (1 species, 4.7%), SW-Anatolian Endemic (1 species, 4.7%), and Mediterranean (1 species, 4.7%) (Vigna Taglianti et al., 1999; Sindaco et al., 2000) (Figure 2).

The following species were determined in our study area: *Bufo variabilis* (Pallas, 1769), *Pelophylax caralitanus* (Arikan, 1988), *Mauremys rivulata* (Valenciennes, 1833), *Emys orbicularis* (Linnaeus, 1758), *Testudo graeca* (Linnaeus, 1758), *Mediodactylus kotschyi* (Steindachner, 1870), *Stellagama stellio* (Linnaeus, 1758), *Lacerta trilineata* (Bedriaga, 1886), *Anatololacerta danfordi* (Gunther, 1876), *Ophisops elegans* (Ménétriés, 1832), *Trachylepis aurata*

(Linnaeus, 1758), *Typhlops vermicularis* (Merrem, 1820), *Dolichophis caspius* (Gmelin, 1789), *Dolichophis jugularis* (Linnaeus, 1758), *Eirenis modestus* (Martin, 1838), *Malpolon insignitus* (Geoffroy de St-Hilaire, 1809), *Natrix natrix* (Linnaeus, 1758), *Natrix tessellata* (Laurenti, 1768), and *Telescopus fallax* (Fleischmann, 1831). We have also included a lizard species that was seen but not collected (*Blanus strauchi* Bedriaga, 1884), and a slide taken by a previous researcher (hydrologist Semih Üçüncü) was used for a snake species [*Zamenis situla* (Linnaeus, 1758)]. The most common amphibians were *Bufo variabilis* (Pallas, 1769) and *Pelophylax caralitanus* (Arikan, 1988). For reptiles, the most common were two lizard species: *Ophisops elegans* (Ménétriés, 1832) and *Anatololacerta danfordi* (Gunther, 1876).

According to IUCN Red List data, 17% of the species are LC (Least Concern), one of them is VU (Vulnerable) (*Testudo graeca* Linnaeus 1758), two of them are NT (Near Threatened) [(*Pelophylax caralitanus* Arikan, 1988 and *Emys orbicularis* (Linnaeus, 1758)], and one of the species (*Bufo variabilis* Pallas, 1769) is DD (Data Deficient), due to its taxonomical status being uncertain and a lack of information about its distribution (IUCN, 2013) (Table).




Figure 1. The localities from which specimens were collected in Burdur Province (1. İbecik, 2. Gölhisar Lake, 3. Kibyra, 4. Yapraklı Dam, 5. Yamadıburnu, 6. Gölhisar, 7. Karamanlı Dam, 8. Salda Lake, 9. Yeşilova, 10. Düden Road, 11. Taşpınar, 12. Bayındır, 13. Sazak Village, 14. Yarışlı Village, 15. Yarışlı Lake, 16. Karataş Lake, 17. Akören, 18. Karakent, 19. Örencik, 20. Yarı Village, 21. Kuruçay, 22. Suderesi, 23. İnsuyu, 24. Sagalassos, 25. Ağlasun Plain, 26. Beşkonak, 27. Kocaaliler-Beşkonak road, 28. Kocaaliler).

Table. Chorotype classification, localities, and conservation status of the IUCN criteria for amphibian and reptilian species in the study area (IUCN: International Union for the Conservation of Nature and Natural Resources, VU: Vulnerable, NT: Near Threatened, DD: Data Deficient, LC: Least Concern).

	Species	Localities	Chorotypes	IUCN
1	<i>Bufo variabilis</i> (Pallas, 1769)	8, 11, 18, 23, 17	Turano-European-Mediterranean	DD
2	<i>Pelophylax caralitanus</i> Arikan, 1988	2, 5, 7, 8, 10, 12, 13, 16, 17, 22, 25, 28	Euro-Siberian	NT
3	<i>Emys orbicularis</i> (Linnaeus, 1758)	2, 15	Turano-European-Mediterranean	NT
4	<i>Mauremys rivulata</i> (Valenciennes, 1833)	5, 28	Turano-European-Mediterranean	LC
5	<i>Testudo graeca</i> Linnaeus, 1758	2, 5, 8, 9, 10, 16, 17, 20, 27, 28	Turano-Mediterranean	VU
6	<i>Trachylepis aurata</i> (Linnaeus, 1758)	2, 13	SW-Asiatic	LC
7	<i>Anatololacerta danfordi</i> Gunther, 1876	10, 14, 24	SW-Anatolian Endemic	LC
8	<i>Lacerta trilineata</i> Bedriaga, 1886	4, 6, 9, 10, 18, 25, 28	E-Mediterranean	LC
9	<i>Ophisops elegans</i> Ménétriés, 1832	3, 8, 10, 14, 15, 16	E-Mediterranean	LC
10	<i>Mediodactylus kotschy</i> (Steindachner, 1870)	8	E-Mediterranean	LC
11	<i>Stellagama stellio</i> (Linnaeus, 1758)	2, 12, 13, 14, 18	E-Mediterranean	LC
12	<i>Blanus trauchi</i> Bedriaga, 1884	24	E-Mediterranean	LC
13	<i>Typhlops vermicularis</i> (Merrem, 1820)	17, 18	Turano-Mediterranean	LC
14	<i>Dolicophis caspius</i> (Gmelin, 1789)	5, 8, 20	Turano-Mediterranean	LC
15	<i>Dolicophis jugularis</i> Linnaeus, 1758	10	SW-Asiatic	LC
16	<i>Eirenis modestus</i> (Martin, 1838)	5, 14, 15, 18	SW-Asiatic	LC
17	<i>Malpolon insignitus</i> (Geoffroy De St-Hilaire, 1809)	26	Mediterranean	LC
18	<i>Telescopus fallax</i> Fleischmann, 1831	21	Turano-Mediterranean	LC
19	<i>Natrix natrix</i> (Linnaeus, 1758)	4, 9, 11, 17	Central Asiatic-European-Mediterranean	LC
20	<i>Natrix tessellata</i> (Laurenti, 1768)	8, 10	Central Asiatic-European	LC
21	<i>Zamenis situla</i> (Linnaeus, 1758)	1	E-Mediterranean	LC




Figure 2. Distribution of major chorotypes of amphibian and reptilian species.

Threats to these species should also be mentioned here. These include dangers such as overharvesting, human disturbance, incorrect agricultural practices, disease, natural disasters, and the degradation and loss of habitat (Cox et al., 2006).

It is hoped that the herpetological information presented here concerning the herpetofauna of the lakes of Burdur, which is a part of the Lake District, an important

wetland, will encourage more comprehensive studies in the future.

Acknowledgments

This study forms part of the MSc thesis of Oğuzhan Ege, supervised by Prof Dr C Varol Tok, and was funded by the Research Council (BAP) of Ege University, İzmir, Turkey.

References

- Afsar M, Ayaz D, Afsar B, Tok CV (2012). Camili Biyosfer Rezerv Alanı'nın (Borçka, Artvin, Türkiye) Herpetofaunası. Anadolu Üniversitesi Bilim ve Teknoloji Dergisi 2: 41–49.
- Afsar M, Tok CV (2011). The herpetofauna of the Sultan Mountains (Afyon-Konya-Isparta), Turkey. Turk J Zool 35: 491–501.
- Arıkan H (1988). On a new form of *Rana ridibunda* (Anura: Ranidae) from Turkey. İstanbul Üniv Fen Fak Mecmuası 53: 81–87.
- Arıkan H, Özeti N, Çevik İE, Tosunoğlu M (1994). *Rana ridibunda caralitana* (Anura: Ranidae)'nin Göller Bölgesi'nde Dağılışı. Turk J Zool 18: 141–145 (in Turkish).
- Baran İ (1980). Doğu ve Güneydoğu Anadolu'nun Kaplumbağa ve Kertenkele faunası. Ege Üniv Fen Fakültesi Dergisi Seri B 4: 203–219 (in Turkish).
- Baran İ (1984). İzmir ve Bodrum Arasındaki Adalarımızın Herpetofaunasının Taksonomik Araştırılması. Doğa Bilim Dergisi Seri A 8: 43–52 (in Turkish).
- Baran İ (1986). Bibliographie der Amphibien und Reptilien der Türkei. In: Kasperek M, editor. Zoologische Bibliographie der Türkei. Heidelberg, Germany: Kasperek, pp. 79–118 (in German).
- Başoğlu, M, Baran İ (1977). Türkiye Sürüngenleri, Kısım I. Kaplumbağa ve Kertenkeleler. Bornova, İzmir, Turkey: Ege University Faculty of Science Book Series (in Turkish).
- Başoğlu M, Baran İ (1980). Türkiye Sürüngenleri, Kısım II, Yılanlar. Bornova, İzmir, Turkey: Ege University Faculty of Science Book Series (in Turkish).
- Başoğlu M, Özeti N, Yılmaz İ (1994). Türkiye Amfibileri. Bornova, İzmir, Turkey: Ege University Faculty of Science Book Series (in Turkish).
- Bulut A (2010). Gökçeadadaki (Çanakkale, Türkiye) amfibi ve sürüngenlerin taksonomisi ve ekolojisi. MSc, Çanakkale Onsekiz Mart University, Çanakkale, Turkey (in Turkish).
- Bulut A, Tok CV, Yakın BY (2011). Gökçeadadaki (Çanakkale, Türkiye) Amfibi ve Sürüngenlerin Ekolojisi. In: X. Ulusal Ekoloji ve Çevre Kongresi, 4–7 October 2011, Çanakkale, Turkey (in Turkish).
- Cihan D, Tok CV (2014). Herpetofauna of the vicinity of Akşehir and Eber (Konya, Afyon), Turkey. Turk J Zool 38: 234–241.
- Cox N, Chanson J, Stuart S, compilers (2006). The Status and Distribution of Reptiles and Amphibians of the Mediterranean Basin. Gland, Switzerland and Cambridge, UK: IUCN.
- Demirsoy A (1997a). Türkiye Omurgalıları - Türkiye Omurgalı Faunasının Sistematik ve Biyolojik özelliklerinin Araştırılması ve Koruma Önlemlerinin Saptanması - AMFİBİLER. Ankara, Turkey: Çevre Bakanlığı, Çevre Bakanlığı Genel Müdürlüğü (in Turkish).
- Demirsoy A (1997b). Türkiye Omurgalıları - Türkiye Omurgalı Faunasının Sistematik ve Biyolojik, Özelliklerinin Araştırılması ve Koruma Önlemlerinin Saptanması - SÜRÜNGENLER. Ankara, Turkey: Çevre Bakanlığı, Çevre Bakanlığı Genel Müdürlüğü (in Turkish).
- Hür H, Uğurtaş İH, İşbilir A (2008). The amphibian and reptile species of Kazdağı National Park. Turk J Zool 32: 359–362.
- Ilgaz K, Kumlutaş Y (2005). The amphibian and reptile species of İğneada (Kırklareli) and its vicinity. Pak J Biol Sci 8: 558–560.
- Kaya S (2005). Çanakkale ili ve civarının herpetofaunası. MSc, Çanakkale Onsekiz Mart University, Çanakkale, Turkey (in Turkish).
- Kumlutaş Y, Durmuş SH, Ilgaz Ç (2011). Kaş-Kekova Özel Çevre Koruma Bölgesi'nin Herpetofaunası. Anadolu Doğa Bilimleri Dergisi 2: 28–34.
- Kumlutaş Y, Ilgaz Ç, Durmuş SH (2001). Herpetofauna of Spil Mountain (Manisa) and its vicinity: results of field surveys. Anadolu Univ J Sci Tech 2: 63–66.
- Kumlutaş Y, Öz M, Durmuş H, Tunç MR, Özdemir A, Düşen S (2004a). On some lizard species of the western Taurus range. Turk J Zool 28: 225–236.
- Kumlutaş Y, Özdemir A, Ilgaz Ç, Tosunoğlu M (2004b). The amphibian and reptile species of Bozdağ (Ödemiş). Turk J Zool 28: 317–319.
- Kumlutaş Y, Tok CV, Türkozan O (1998). The herpetofauna of Ordu-Giresun region. Turk J Zool 22: 199–201.
- Kumlutaş Y, Tosunoğlu M, Göçmen B (1999). Karadeniz Bölgesi *Rana ridibunda* (Anura: Ranidae) Populasyonları Üzerinde Morfolojik Araştırma. Turk J Zool 23: 801–806 (in Turkish).
- Özcan S, Üzümlü N (2014). The herpetofauna of Madran Mountain (Aydın, Turkey). Turk J Zool 38: 108–113.
- Özdemir A (1998). Murat Dağı (Kütahya-Uşak) herpetofaunasının araştırılması. MSc, Dokuz Eylül University, İzmir, Turkey (in Turkish).

- Özdemir A, Baran İ (2002). Research on the herpetofauna of Murat Mountain (Kütahya-Uşak). *Turk J Zool* 26: 189–195.
- Sindaco R, Venchi A, Carpaneto G, Bologna M (2000). The reptiles of Anatolia. A checklist and zoogeographical analysis. *Biogeographia* 21: 441–554.
- Tok CV (1993). Reşadiye (Datça) yarımadası herpetofaunası üzerine taksonomik ve biyolojik araştırmalar. PhD, Ege University, Bornova, İzmir, Turkey (in Turkish).
- Tok CV (1995). Reşadiye (Datça) Yarımadası'nın Herpetofaunası. *Turk J Zool* 19: 119–121 (in Turkish).
- Tok CV, Çicek K (2014). Amphibians and reptiles in the province of Çanakkale (Marmara Region, Turkey) (Amphibia; Reptilia). *Herpetozoa* 27: 65–76.
- Tok CV, Gürkan M, Yakın BY (2012a). Ağı Dağı'nın Herpetofaunası. In: Kazdağları III. National Symposium, 24–26 May 2012, Çanakkale, Turkey, pp. 59–63 (in Turkish).
- Tok CV, Yakın BY, Gürkan M (2012b). Kirazlı (Çanakkale) Civarının Herpetofaunası. In: Kazdağları III. National Symposium, 24–26 May 2012, Çanakkale, Turkey, p. 621 (in Turkish).
- Uysal İ (2011). Kavak Deltası (Saroz körfezi, Çanakkale-Türkiye)'nin herpetofaunası ve mevcut türlerin habitat seçimleri. MSc, Çanakkale Onsekiz Mart University, Çanakkale, Turkey (in Turkish).
- Vigna Taglianti A, Audisia P, Biondi M, Bologna MA, Carpaneto GM, De Biase A, Fattorini S, Piattella E, Sindaco R, Venchi A et al. (1999). A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. *Biogeographia* 20: 31–59.
- Yılmaz İ (1984). Trakya Kuyruksuz Kurbağaları Üzerine Morfolojik ve Taksonomik Bir Araştırma (Anura: Discoglossidae, Pelobatidae, Bufonidae, Hylidae, Ranidae). *Doğa Bilim Dergisi* 8: 244–264 (in Turkish).
- Yılmaz İ (1989). Kuzey Anadolu Amfibilerinin Yayılışı üzerine bir çalışma (Amphibia: Urodela, Anura). *Doğa Bilim Dergisi* 13: 130–140 (in Turkish).