

## Predation of a scorpion (Scorpiones: Buthidae) by an assassin bug (Heteroptera: Reduviidae) in the Brazilian Atlantic Forest

André Felipe de Araujo LIRA<sup>1\*</sup>, Victor Luiz Nascimento de ARAÚJO<sup>2</sup>, Cleide Maria Ribeiro de ALBUQUERQUE<sup>2</sup>

<sup>1</sup>Post Graduate Program in Animal Biology, Department of Zoology, Center of Biological Sciences, Federal University of Pernambuco, Cidade Universitária, Recife, PE, Brazil

<sup>2</sup>Department of Zoology, Center of Biological Sciences, Federal University of Pernambuco, Cidade Universitária, Recife, PE, Brazil

Received: 17.04.2015 • Accepted/Published Online: 05.09.2015 • Final Version: 05.02.2016

**Abstract:** Litter-dwelling arthropods comprise about three-fourths of the total animal biomass in tropical forests. These invertebrates are involved in many interspecific interactions, from mutualism to predation. We report herein the predation of a scorpion by an immature assassin bug (Harpactorini) during a nocturnal manual search for scorpions in a fragment of the Brazilian Atlantic Forest. The specimens were found 15 cm above the ground on a seedling, and the prey was two-fold larger than the predator. The assassin bug had its rostrum inserted into the pleura of a juvenile *Tityus pusillus* Pocock, 1893 scorpion, between the first and second segments of the mesosoma; when disturbed, the predator jumped to the leaf litter without releasing its prey. To the best of our knowledge, this is the first report of juvenile predator-prey interactions between a heteropteran and a scorpion in this biome.

**Key words:** Harpactorini, intraguild predation, natural history, *Tityus pusillus*

Scorpions are top predators, capturing a great variety of prey exerting direct influence on the energy flow through predation of different invertebrates (Polis, 1990; McCormick and Polis, 1990). However, scorpions are also subjected to predation by vertebrates and invertebrates (Polis et al., 1981). Predation of scorpions by invertebrates, mainly ants, coleopterans, heteropterans, chilopods, solpugids, other scorpions, and spiders, causes a high rate of mortality among juveniles and smaller species of scorpions (Cloudsley-Thompson, 1960; Williams, 1966; Cloudsley-Thompson, 1977; McCormick and Polis, 1990; Punzo, 1998; Melic, 2000; Moreno-González and Hazzí, 2012; Stevenson and Stohlgren, 2015). Here, we report the first case of scorpion predation by a juvenile assassin bug (Heteroptera, Reduviidae, Harpactorinae, Harpactorini), during a nocturnal collection of scorpions in the Brazilian Atlantic Forest; this is the second reported case of scorpion predation in this biome (Lira and Costa, 2014).

Because of the difficulty in identifying juveniles, the assassin bug was classified to the tribe level, Harpactorini. The immature assassin bug preyed on a juvenile *Tityus pusillus* Pocock, 1893 (Scorpiones: Buthidae), which is the most abundant and widely found scorpion in the northeastern region of the Brazilian Atlantic Forest

(Lourenço, 2002; Porto et al., 2010; Lira and Albuquerque, 2014; Lira et al., 2015) and is capable of causing injuries to humans (Albuquerque et al., 2009). This scorpion is a sedentary predator commonly found in the leaf litter layers; its abundance increases in the dry season, and it shows a positive relationship with dry litter mass (Lira et al., 2013, 2015; Lira and DeSouza, 2014).

Despite the smaller size (8.6 mm) of the predator in comparison with the size of its prey (14.3 mm), the assassin bug managed to insert its rostrum into the pleura of the scorpion, between the first and second segments of the mesosoma (Figure). The predation was observed in Reserva Biológica de Saltinho (08°43'43"S, 35°10'39.8"W), Tamandaré, southern coast of Pernambuco state, Brazil. The reserve has a total area of 562.25 ha composed of the Ombrophilous Dense Forest (Lisboa et al., 2011; ICMBio, 2015). In previous studies of other Atlantic Forest fragments, *T. pusillus* has been observed only as the predator of spiders, crickets, cockroaches, and moths (Lira pers. obs.). The specimens of our study were found on a seedling (~45 cm), on which the assassin bug was oriented vertically 15 cm above the ground with its head facing upward. When disturbed, the predator jumped to the leaf litter with the scorpion in its mouthparts without releasing

\* Correspondence: andref.lira@gmail.com



**Figure.** Nymph of assassin bug (Harpactorini) preying a *Tityus pusillus* Pocock, 1893 juvenile.

its prey. Both animals were collected, and the voucher materials have been deposited in the Arachnological Collection of the Universidade Federal de Pernambuco.

The predation of *T. pusillus* by an assassin bug that belongs to Harpactorini described in this study differs from previously reported predations by animals that belong to Heteroptera; adult, not juvenile, scorpions that belong to Buthidae have been reported to exhibit such behavior (Stevenson and Stohlgren, 2015). An adult *Microtomus purcis* (Drury, 1782) (Heteroptera: Reduviidae) was found feeding on juvenile and adult *Centruroides hentzi* (Banks, 1900) (Scorpiones: Buthidae) under a pine bark in the coastal plain of the United States. Both species are characteristically bark-dwelling species (Horn and Hanula, 2002; Stevenson et al., 2012), and the interactions between *M. purcis* and *C. hentzi* were attributed to the similar microhabitats of both species (Stevenson and Stohlgren, 2015).

Nevertheless, both studies agree about the predator-prey size for assassin bugs that feed on larger prey. It is important to note that assassin bugs are the largest group

of predatory hemimetabolous insects (~6800 species) and one of the more numerous animal predators (Hwang and Weirauch, 2012). Assassin bugs that exhibit extraoral digestion, such as the specimen reported in the present study, may consume relatively large prey, allowing them to prey on a large variety of animals as predicted by Cohen (1995). Assassin bugs use venomous saliva that paralyzes their prey within a short duration of time, after which the bugs use their forelegs to hold the prey and suck the body juices (Evangelin et al., 2014). In addition, some members of the tribe Harpactorini have developed a novel predation strategy called sticky trap predation, as they have sticky glands on the legs that secrete a sticky substance used to capture their prey (Barth, 1952; Ambrose, 1999; Weirauch, 2006; Zhang and Weirauch, 2013).

Despite their capacity to capture larger prey, nonhematophagous assassin bugs prefer to capture smaller prey. Cogni et al. (2002) investigated the influence of prey size on predation success in the assassin bug *Zelus longipes* (Linnaeus, 1767) (Heteroptera: Reduviidae). According to these researchers, the fact that successful attacks were more frequent in small prey reduces the risk of injury to the predator. In fact, scorpions can be dangerous prey because of their venom, and the predator can easily become potential prey. Thus, the venomous saliva of assassin bugs is very useful for capturing dangerous prey such as scorpions.

The observation reported in this study extends the list of invertebrate predators of scorpions, and, to our knowledge, it is the first report of an assassin bug preying on a scorpion in the Brazilian Atlantic Forest. Further studies are required to obtain more and detailed information on predator-prey interactions between assassin bugs and scorpions and to understand the ecological implication of this interaction.

### Acknowledgments

We are very grateful to Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the doctoral scholarship to AFAL. We are also very grateful to Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for permission to use the area to collect samples. We are indebted to Dr Hécio Gil-Santana, who identified the assassin bug.

### References

- Albuquerque CMR, Porto TJ, Amorim, MLP, Santana Neto PL (2009). Scorpionism caused by *Tityus pusillus* Pocock, 1893 (Scorpiones; Buthidae) in State of Pernambuco. *Rev Soc Bra Med Trop* 42: 206–208.
- Ambrose DP (1999). *Assassin Bugs*. Enfield, NH, USA: Science Publishers.
- Barth R (1952). Die Fangdrüsen an den Beinen von *Zelus (Diplocodus) leucogrammus* (Perty). *Zool Jahrb* 73: 323–336 (in German).
- Cloudsley-Thompson JL (1960). Notes on Arachnida, 35 - A scorpion eaten by a beetle. *Entomol Mon Mag* 95: 223.
- Cloudsley-Thompson JL (1977). Adaptational biology of Solifugae (Solpugida). *Bull Br Arachnol Soc* 4: 61–71.

- Cogni R, Freitas AVL, Amaral Filho BF (2002). Influence of prey size on predation success by *Zelus longipes* L. (Het., Reduviidae). *J Appl Entomol* 126: 74–78.
- Cohen AC (1995). Extra-oral digestion in predaceous terrestrial arthropoda. *Annu Rev Entomol* 40: 85–103.
- Evangelin G, Horne B, Muthupandi M, John WS (2014). Venomous saliva of non-haematophagous reduviid bugs (Heteroptera: Reduviidae): A review. *Biolife* 2: 615–626.
- Horn S, Hanula JL (2002). Life history and habitat associations of the Broad Wood Cockroach, *Parcoblatta lata* (Blattaria: Blattellidae), and other native cockroaches in the coastal plain of South Carolina. *Ann Entomol Soc Am* 95: 665–671.
- Hwang WS, Weirauch C (2012). Evolutionary history of assassin bugs (Insecta: Hemiptera: Reduviidae): insights from divergence dating and ancestral state reconstruction. *PLoS One* 7: e45523.
- ICMBio (2015). Instituto Chico Mendes de Conservação da Biodiversidade. Available from: <http://www.icmbio.gov.br/portal> (Accessed 21 March 2015).
- Lira AFA, Albuquerque CMR (2014). Diversity of scorpions (Chelicerata: Arachnida) in the Atlantic Forest in Pernambuco, northeastern Brazil. *Check List* 10: 1331–1335.
- Lira AFA, Costa AA (2014). First record of a brown widow spider *Latrodectus geometricus* Koch, 1841 (Araneae, Theridiidae) feeding scorpion (Scorpiones, Bothriuridae) in a Brazilian Atlantic Forest. *Braz J Biol* 74: 1011.
- Lira AFA, DeSouza AM (2014). Microhabitat use by scorpion species (Arachnida: Scorpiones) in the montane Atlantic Rain Forest, Brazil. *Rev Ibér Aracnol* 24: 107–108.
- Lira AFA, Rego FNAA, Albuquerque CMR (2015). How important are environmental factors for the population structure of co-occurring scorpion species in a tropical forest? *Can J Zool* 93: 15–19.
- Lira AFA, Souza AM, Silva Filho AAC, Albuquerque CMR (2013). Spatio-temporal microhabitat use by two co-occurring species of scorpions in Atlantic rainforest in Brazil. *Zoology* 116: 182–185.
- Lisboa EBF, Moura GJB, Melo IVC, Andrade EVE, Figuerêdo Júnior JM (2011). Aspectos ecológicos de *Hypsiboas semilineatus* (Spix, 1824) (Amphibia, Anura, Hylidae) em fragmento de Mata Atlântica, nordeste do Brasil. *RICA* 2: 21–30 (article in Portuguese with an abstract in English).
- Lourenço WR (2002). *Scorpions of Brazil*. Paris, France: Les Éditions de l'If.
- McCormick SJ, Polis GA (1990). Prey, predators, and parasites. In: Polis GA, editor. *The Biology of Scorpions*. Stanford, CA, USA: Stanford University Press, pp. 294–320.
- Melic A (2000). El género *Latrodectus* Walckenaer, 1805 en la península Ibérica (Araneae: Theridiidae). *Rev Ibér Aracnol* 1: 13–30 (article in Spanish with an abstract in English).
- Moreno-González JA, Hazzi NA (2012). Intraguild predation case: *Tityus forcipula* Gervais, 1843 (Scorpiones, Buthidae) feeding on *Chactas vanbenedeni* Gervais, 1843 (Scorpiones, Chactidae) in Colombia. *Rev Ibér Aracnol* 20: 117–120.
- Polis GA (1990). *The Biology of Scorpions*. Stanford, CA, USA: Stanford University Press.
- Polis GA, Sissom WD, McCormick SJ (1981). Predators of scorpions: field data and a review. *J Arid Environ* 4: 309–326.
- Porto TJ, Brazil TK, Lira da Silva RM (2010). Scorpions, state of Bahia, northeastern Brazil. *Check List* 6: 292–297.
- Punzo F (1998). *The Biology of Camel Spiders (Arachnida, Solifugae)*. Boston, MA, USA: Kluwer Academic Publisher.
- Stevenson DJ, Greer G, Elliott MJ (2012). The distribution and habitat of *Centruroides hentzi* (Banks) (Scorpiones, Buthidae) in Georgia. *Southeast Nat* 11: 589–598.
- Stevenson DJ, Stohlgren KM (2015). Predation on the scorpion *Centruroides hentzi* (Banks) (Scorpiones: Buthidae) by the assassin bug *Microtomus purcis* (Drury) (Insecta: Hemiptera: Reduviidae). *Southeast Nat* 14: N1–N4.
- Weirauch C (2006). Observations on the sticky trap predator *Zelus luridus* Stal (Heteroptera: Reduviidae: Harpactorinae), with the description of a novel gland associated with the female genitalia. *Denisia* 50: 1169–1180.
- Williams SC (1966). Burrowing habitat of the scorpion *Anuroctonus phaeodactylus* (Wood) (Scorpionida: Vaejovidae). *Proc Calif Acad Sci* 34: 419–428.
- Zhang G, Weirauch C (2013) Sticky predators: a comparative study of sticky glands in harpactorine assassin bugs (Insecta: Hemiptera: Reduviidae). *Acta Zool* 94: 1–10.