

The Morphology of *Vipera ammodytes transcaucasiana* (Reptilia, Viperidae) Specimens Collected from Murgul (Artvin, Turkey)

Bilal KUTRUP

Karadeniz Technical University, Science Faculty, Biology Department, Trabzon–TURKEY

Received: 19.10.1998

Abstract: The morphological characteristics and distribution of *Vipera ammodytes transcaucasiana*, which is a member of the *Vipera kaznakovi* group, were studied. The transcaucasian horned viper specimens were captured from Murgul, Artvin, in Turkey.

This species differs from all other vipers in the Near and Middle East in respect of its pronounced horn. It is similar to *Vipera pontica* in the greenish–yellowish tip of the tail, but differs in having more apicals.

Key Words: Reptilia, Squamata, Viperidae, *Vipera ammodytes transcaucasiana*, Erenköy, Morphology.

Murgul (Artvin)'dan Yakalanan *Vipera ammodytes transcaucasiana* (Reptilia, Viperidae)'nın Morfolojisi

Özet: Bu çalışmada *Vipera kaznakovi* grubunun bir üyesi olan *Vipera ammodytes transcaucasiana*'nın morfolojik karakterleri ve yayılışı araştırılmıştır. Boynuzlu engerek olarak bilinen *Vipera ammodytes transcaucasiana* örnekleri Artvin iline bağlı Murgul ilçesinden yakalanmıştır.

Bu engerek oldukça belirgin boynuzu ile yakın ve orta doğudaki engerek türlerinden farklıdır. Kuyruk ucunun sarımsı yeşil olması ile *Vipera pontica*'ya benzer fakat apical sayısının daha fazla olması ile ondan ayrılır.

Anahtar Sözcükler: Reptilia, Squamata, Viperidae, *Vipera ammodytes transcaucasiana*, Erenköy, Morfoloji.

Introduction

The first study of zoogeographic and systematic characteristics in Turkish vipers was conducted by Başoğlu (1). Further studies have been carried out by Baran (2), Başoğlu & Baran (3), Tuncer (4), Böhme & Joger (5) and Tok & Kumlutaş (6).

The first report of *Vipera ammodytes transcaucasiana* in Turkey was from the vicinity of Borçka, in the province of Artvin (7). Eiselt & Baran (8) described two viper specimens from Köseadağ, near Zara, in the province of Sivas, as *Vipera ammodytes transcaucasiana*. Nilson & Andren (9) reported this taxon from Ordu, Zonguldak, Adapazarı and Konya. Teynie (10) stated that a specimen caught in the İstanbul area resembled *Vipera ammodytes transcaucasiana*. Tok & Kumlutaş (6) described a specimen of *Vipera ammodytes transcaucasiana*, which was caught in Perşembe (central Black Sea region).

Recent studies of the systematics of vipers in the Caucasus region have shown that the taxonomy of these species is rather complex. It has been reported that there are two viper groups in this region, *Vipera kaznakovi*,

and *Vipera ursini* (11, 12, 13). *Vipera ammodytes transcaucasiana* has been shown to be an important species in the *Vipera kaznakovi* group in north-eastern Turkey and the adjacent Transcaucasia region since 1990 (14).

To date there have been few *Vipera ammodytes transcaucasiana* specimens caught in the province of Artvin. The first specimen in the region was examined by Derjugin (7). Then, Nilson (9) reported that he saw one *Vipera ammodytes transcaucasiana* in Borçka in 1988.

It can be concluded that there has been little substantial research carried out on *Vipera ammodytes transcaucasiana* in the eastern Black Sea region, and only a few specimens have been caught in this region. The aim of this study was to define new locations and to capture more *Vipera ammodytes transcaucasiana* specimens.

Material and Method

This research was predominantly a study of external morphology. Field trips were conducted in different parts



Figure 1. Dorsal view of an adult male.

Vipera ammodytes transcaucasiana in Turkey have shown that specimens caught from Istanbul, Adapazarı, Bursa and Kuşadası did not belong to the species *Vipera ammodytes taranscaucasiana*. They were more similar to *Vipera ammodytes meridionalis* or *Vipera ammodytes montandani* (6).

Although this subspecies was reported by Nilson (9) in his study carried out in the central Black Sea area, Baran did not mention it in his study conducted in the same region. In recent years, Tok & Kumlutaş (6) captured a specimen of *Vipera ammodytes transcaucasiana* from Perşembe (central Black Sea region). Accordingly, we thought that *Vipera ammodytes transcaucasiana* could be found throughout the eastern Black Sea region. For this reason, the research area was extended towards the west of Çoruh Valley.

Within the area covered by the study, it was only Çoruh Valley (from the village of Murgul, Erenköy) where specimens of *Vipera ammodytes transcaucasiana* were found. The main shelter for this subspecies was hazelnut orchards which are common to all the regions along the Black Sea coast. However it was determined that the drier habitat of Erenköy is better suited to *Vipera ammodytes transcaucasiana* than the habitat in Trabzon and Rize, which have a more humid climate. *Vipera ammodytes transcaucasiana* does not like a humid climate, like *Vipera kaznakovi*.

Vipera ammodytes transcaucasiana has a pronounced horn on the snout, and this is a distinguishing characteristic for the different forms of *Vipera ammodytes*. Similar head morphology is also found in *Vipera pontica* and *Vipera darevskii*, but *Vipera*



Figure 2. Dorsal view of an adult female.

ammodytes transcaucasiana is identical in terms of its long horn. The snout shape seems to be an evolutionary trait that is more pronounced in this viper than is normally found in Caucasian viper populations. Table 2 shows that the Erenköy specimens had a pronounced horn similar to the Borçka specimen. There was no mention of the upturned snout in the Perşembe

specimens, but the Sivas specimens had 2 or 3 scales in front of the pronounced horn (2). The number of scales in the same area in the Erenköy specimens were: 2 in the female and 3 in the male, as in the Sivas specimens. The Perşembe specimens show similarities to the Erenköy specimens in terms of rostral index. However, The Perşembe and Sivas specimens had lower ventrals than

Table 2. Variation in scalitions characters in different isolated populations of *Vipera ammodytes transcaucasiana*.

Locality	Ventrals	Subcaudals	Crawnscales	Loreals	Apicals	Canthals	Rostral index	Sublabials
Borçka*	150-162	36-39	45-58	6-7	10-12	1-1.5	-	-
Erenköy	154-157	34-38	44-48	7-7	8-12	2-2	1.33-1.56	9-10
Perşembe	150-153	33-37	-	-	-	-	1.28	11-11
Sivas	150-154	37-39	-	-	-	-	1.30-1.50	11-11

(*) Billing et al. (14).



Figure 3. Dorsal view of the head of an adult male.

the Erenköy and Borçka specimens (Table 2). Thus, the Erenköy and Borçka specimens are quite alike in terms of these characteristics.

It is essential that more studies be carried out in order to determine the distribution of this viper in Turkey. Since there still are some questions left unanswered, we think it is necessary to continue work on this topic in order to obtain a more comprehensive picture of the issue discussed in this paper.

Unlike *Vipera kaznakovi*, *Vipera ammodytes transcaucasiana* is not under protection. As a result, it is in danger, because foreign tourists have started to come to the Çoruh Valley to collect specimens.

Finally, the vipers in the Çoruh valley, especially *Vipera ammodytes transcaucasiana*, should be studied in more detail and placed under protection like *Vipera kaznakovi*.

References

1. Başoğlu, M., Some varieties of *Vipera berus* from the extreme north-eastern Anatolia. Rev. Fac. Sci. Univ. Istanbul, Ser. B, 12: 182-190, 1947.
2. Baran, I., Türkiye Yılanlarının Taksonomik Revizyonu ve Coğrafik Dağılımları. TBTK yayınları no: 309, T.B.A.G. Seri no: 9, Ankara, 1-177s., 1976.
3. Başoğlu, M., Baran, I., Türkiye Sürüngenleri, Kısım II. Yılanlar, Ege Univ. Fen Fak. Kitaplar Ser. no: 81, İzmir, 218s., 1980.
4. Tuncer, S., Investigation of *Vipera kaznakovi* (Hopa Snake) as Predator. International Symposium on Biological Control, Antalya, Turkey, 27-30 November 1990, pp. 23-26, 1990.
5. Böhme, W., Joger, U., Eine neue Art des *Vipera berus* complex aus der Türkei. Amph. Rept. Vol. 4, pp. 265-271, 1983.
6. Tok, C.V., Kumlutaş, Y., On specimens of *Vipera ammodytes taranscaucasiana* (Viperidae) from Perşembe, Black Sea region of Turkey. Zoology in the Middle East 13, 47-50, 1996.
7. Derjugin, K., Matériaux pour l'herpétologie du sud-ouest du Transcaucase et des environs de Trapezonde. Ann. Mus. Zool. St.-Petersbourg 6: 84-111, St. Petersburg, 1901.
8. Eiselt, J., Baran, I., Ergebnisse zoologischer Sammelreisen in der Türkei. Viperidae, Annalen des Naturhistorischen Museum Wien 74: Vienna, 357-369, 1970.
9. Nilson, G., Andren, C., The mountain vipers of the Middle East the *Vipera xanthina* complex (Reptilia, Viperidae) Bonner Zoologische Monografien. Vol. 20, 1-90, 1988.
10. Teynie, A., Observations herpétologiques en Turquie. II. Partie., Bulltin Societe Herp. France 58, 20-29, 1991.

11. Vedmedeja, V., Orlov, N. L., Tuniev, B., On taxonomy of three vipers species of the *Vipera kaznakovi* complex. Proc. Zool. Inst. Leningr. 157, pp. 265-271, 1986.
12. Orlov, N., Tuniev, B., The recent areas, their possible genesis and the phylogeny of three viper species of the Eurosiberian group of the *Vipera kaznakovi* complex in the Caucasus. Proc. Zool. Inst. Leningr. 157, pp. 107-135, 1986.
13. Nilson, G., Tuniev, S., Orlov, N., Hoggren, M., Andren, C., Systematic of the vipers of the Caucasus : Polymorphism or Sibling Species. Asiatic Herpetological Research, Vol. 6, pp. 1-26, 1995.
14. Billing, H., Nilson, G., Satler, B., *Vipera pontica* sp. n., a new viper species in the *Vipera kaznakovi* group (Reptilia, Viperidae) from north-eastern Turkey and adjacent Transcaucasiana. Zoologia Scripta, Vol. 19, no: 2, pp. 227-231, 1990.
15. Baran, I., Yılmaz, I., Kete, R., Kumlutaş, Y., Durmuş, H., Batı ve Orta Karadeniz Bölgesinin Herpetofaunası. Doğa Zooloji Der., 16: 275-288, 1992.