

Isolation and Characterization of Aerobic Denitrifiers from Agricultural Soil

Ebru ÇELEN, Mehmet Akif KILIÇ

Department of Biology, Faculty of Arts and Science, Akdeniz University, Dumlupınar Bulvarı,
07058 Kampus, Antalya - TURKEY

Received: 13.08.2003

Abstract: Denitrification is generally considered an anaerobic process. However, in recent years it has been shown that bacteria can also reduce nitrate to nitrite under aerobic conditions. The characterization of biologically available nitrogen forms and their biological cycling mechanisms is important for ecological and agricultural implications. In this study, aerobic nitrate reducers were isolated from greenhouse soil. Using a nitrate reduction assay, it was found that 39 out of 60 isolates can reduce nitrate to nitrite under aerobic conditions. Five of those 39 isolates were selected for further characterization of their aerobic nitrate reduction activity. Four of those 5 isolates were found to have periplasmic nitrate reductase activity using MV⁺ as a non-biological electron donor. Nitrite production of these isolates under aerobic conditions was determined using different nitrogen forms as sole nitrogen sources. It was found that nitrite accumulation in all isolates was higher when nitrate was the sole nitrogen source than when nitrate+ammonium was the sole nitrogen source. No nitrite accumulation was observed when ammonium was the sole nitrogen source. This study suggests that when ammonium exists together with nitrate, aerobic nitrate reduction does not lead to high nitrite accumulation.

Key Words: denitrification, aerobic denitrification, aerobic nitrate reduction, periplasmic nitrate reductase.

Tarım Toprağından Aerobik Denitrifikasyon Bakterilerin İzolasyonu ve Karakterizasyonu

Özet: Denitrifikasyon genelde anaerobik proses olarak kabul edilir. Ancak, son yıllarda aerobik koşullarda da nitratı nitrite indirgeyen bakteriler gösterilmiştir. Azot formlarının ve azotun biyolojik dönüşüm mekanizmalarının bilinmesi ekolojik ve tarımsal önem taşır. Bu çalışmada, tarım toprağından aerobik denitrifikasyon yapan bakteriler izole edildi. 60 izolattan 39'nun aerobik koşullarda nitratı nitrite indirgeyebildiği tespit edildi. 39 izolattan beşi aerobik nitrat redüksiyon aktivitesinin belirlenmesi için seçildi. Biyojik olmayan elektron vericisi MV⁺ kullanarak beş izolattan dördünün periplazmik nitrat redüktaz aktivitesine sahip olduğu bulundu. Bu izolatların farklı azot kaynaklarında, aerobik koşullarda nitrit üretme özellikleri belirlendi. İzolatların nitratlı ortamda, amonyak + nitratlı ortama kıyasla kültürlerinde yüksek seviyede nitrit biriktiği, azot kaynağı amonyak olan ortamda ise, izolatların hiçbirinin nitrit biriktirmediği gözlemlendi. Sonuç olarak, aerobik nitrat redüksiyonunda nitratla birlikte amonyağın varlığı yüksek seviyede nitrit birikimine neden olmamaktadır.

Anahtar Sözcükler: Denitrifikasyon, aerobik denitrifikasyon, aerobik nitrat redüksiyonu, periplazmik nitrat redüktaz.

Introduction

Denitrification is widely considered to be the conversion of nitrate to its reduced forms and its final products are usually in gas form. For this reason, denitrification is considered to be a process that removes bioavailable nitrogen forms from the Earth's surface (1). Since the loss of nitrate from agricultural soil reduces crop yield as well as affecting the fauna and flora on and in the soil, denitrification is not a desirable process from an agricultural point of view. Moreover, excess nitrate or nitrite in soil or water ecosystems leads to environmental

and health problems (2). This excess nitrate in water or soil can be converted to its acid forms, which consequently reduce soil and water quality (3,4). In addition to ecological problems, excess nitrate also stimulates denitrification, resulting in the loss of bioavailable nitrogen used by plants in the form of the gaseous nitrogen compounds N₂O and N₂ (5). Despite these detrimental effects, denitrification is beneficial in sewage treatment because it converts NO₃ to its gaseous forms, thus reducing the amount of available nitrogen in the sewage.

Denitrification is originally described as an anaerobic respiratory process in which the first step is catalysis by membrane-bound nitrate reductase (1,6). However, recently it has been shown that denitrification can also take place under aerobic conditions and a periplasmic nitrate reductase seems to be the major enzyme in aerobic denitrification (7,8).

There are 3 nitrate reductases in bacteria that can perform the reduction of nitrate to nitrite under different physiological conditions. Membrane-bound respiratory nitrate reductase (NAR) can generate metabolic energy by using nitrate as a terminal electron acceptor during anaerobic denitrification. Cytoplasmic assimilatory nitrate reductase (NAS) can convert nitrate to ammonium, which can then be incorporated into biomolecules. The third enzyme is a dissimilatory nitrate reductase (NAP), which exists in the periplasm of some Gram-negative bacteria and can reduce nitrate to nitrite under both aerobic and anaerobic conditions (9 -12).

NAP has been isolated from many Gram-negative bacteria (8, 13-17) and its gene has been identified in all members of proteobacteria, predominantly in γ -enterobacteria (18). Isolations from soil and fresh water indicated that aerobic nitrate reduction is widespread in nature (19).

NAP activity is not repressed by either ammonium or oxygen. The NAP system contains 4 to 7 genes, clustered in *nap* operon, and the *napABC* genes are essential for *in vivo* nitrate reduction. In some bacteria the *nap* genes are located on endogenous plasmids (e.g., *Rhodobacter capsulatus*) and others on chromosomes (e.g., *Escherichia coli*). *Nap A* gene analysis of the pseudomonas isolates indicated that some strains that have NAP genes and express activity are incapable of aerobic nitrate reduction (20).

Different physiological functions have been proposed for this enzyme. The NAP seems not to be primarily involved in nitrate assimilation or anaerobic respiration, although nitrite generated by NAP can be used as a nitrogen source or as a substrate for anaerobic denitrification depending on the organisms (12). It is more likely that NAP is involved in aerobic denitrification and/or the maintenance of an optimal redox potential. The physiological function of NAP may vary in different organisms or even in the same bacterium under different metabolic conditions.

Since enzymes responsible for nitrate reduction are located in different parts of the cell, the activity of nitrate reductases is determined using non-physiological electron donors such as benzyl viologen (BV⁺) and methyl viologen (MV⁺) dyes. Both the membrane-bound (NAR) and the periplasmic nitrate reductase (NAP) can utilize reduced BV⁺ and MV⁺ as electron donors. However, the cell membrane is permeable to BV⁺, whereas MV⁺ is relatively incapable of crossing the cell membrane. Therefore, BV⁺ and MV⁺ are used to detect the activity of nitrate reductase enzymes whose active sites are located in the cytoplasm or in the periplasm, respectively (8,14,17,21,22).

Understanding the conversion of nitrate in agricultural soil is of ecological and economical importance. Since nitrite production is usually considered to be an anaerobic process and the extent of aerobic nitrate reduction is not fully known, the effect of aerobic nitrite production on the global nitrogen cycle, environment and agriculture is not known. The aim of this study was to isolate and characterize aerobic nitrate reducers from agricultural soil.

Materials and Methods

Isolation of Bacteria

Soil samples were taken from the area around vine tree roots (approximately 20-30 cm diameter area of the body) in 4 randomly chosen sections of a greenhouse at Akdeniz University Agricultural Faculty. A total of 12 samples were collected from 5 cm below the surface of the soil using a sterile metal cylindrical tool (45 cm in length and 5 cm in diameter). All soil samples were collected in a sterile glass container and delivered to the laboratory within 20 min.

Soil samples were sieved using a sterile sieve with pores of 6 mm. Ten grams of sieved sample was dissolved in 100 ml of dissolving buffer (0.85% NaCl and Tween 80). After appropriate dilutions, samples were inoculated to both defined agar medium (30 mM succinate, 10 mM NH₄Cl, 14 g K₂HPO₄, 2.7 g KH₂PO₄, 2.5 g NaCl, 5ml 2% MgSO₄ solution, 4 mM FeSO₄ and 50 mM fungicide, pH 7.2) (19). Plates were incubated at 30 °C for 4 days. The number of bacteria in the soil sample was calculated by counting colonies forming in the defined agar plates (23). The morphological and biochemical characterization of isolates was performed according to Bergey's Manual of Bacteriology (24).

Nitrate Reduction Assay

The ability of colonies to reduce nitrate to nitrite was determined using the nitrate reductase assay modified from 19. The isolates were grown on defined medium plates for 2 days at 30 °C. One hundred micro liters of 1% sodium nitrate was added to the colonies on plates. After 10 min incubation at room temperature, 100 µl of the nitrate detection reagent (containing 2 parts 4% w/v sulfanilamide in 20% conc. HCl and 1 part 0.08% w/v naphthylethylenediamine in 1% conc. HCl) was added and the occurrence of a reddish pink color indicated the existence of nitrite, and therefore aerobic reduction of nitrate to nitrite by bacteria.

Nitrite Assay

The nitrite concentration in culture supernatants was estimated by colorimetric determination (22,25). Isolates were grown in defined liquid medium with different nitrogen sources (either 10 mM of NO₃ or NO₃ and NH₃ or NH₃) by shaking at 150 rpm at 30 °C for 12-16 h depending on the time needed for the isolates to reach their late logarithmic phase. Cells were harvested by centrifugation at 6000 rpm for 3 min and 1 ml of supernatant was taken into 3 ml cuvettes. Then 0.02 ml of nitrite determination reagent was added to the same cuvettes that were left at room temperature for 20 min and the color was measured spectrophotometrically at 540 nm against the blank.

Periplasmic Nitrate Reductase Activity

Nitrate reduction activity in whole cells was measured by following the oxidant-dependent oxidation of dithionite-reduced methyl viologen (MV⁺) (7,19). A single colony of the isolates was inoculated in nutrient broth medium and grown overnight at 30 °C under shaking. The cultures were harvested and the pellets were washed twice with 10 mM phosphate buffer (pH 7.4). The cell suspensions were used to inoculate 50 ml of defined medium containing 10 mM NaNO₃ in 250 ml conical flasks and were incubated in shaken flask culture at 30 °C for 12-16 h depending on the time needed for the isolates to reach their late logarithmic phase. For the periplasmic nitrate reductase activity, the isolates were washed twice with 10 mM phosphate buffer (pH 7.4) to avoid any nitrate contamination from the medium. The

cells of the isolates were diluted in 2 ml of the same buffer in a 3 ml cuvette and the concentration of cells in this cuvette was adjusted to give an O.D.= 0.4 at 600 nm. Two milliliters of the cell suspension adjusted to final conc. was injected into a 3 ml rubber sealed cuvette. After adding 20 µl of 100 mM methyl viologen to the same cuvette using a Hamilton syringe, air from the cuvette was removed with oxygen-free nitrogen over 10 min. Aliquots of a freshly prepared, concentrated solution of sodium dithionite were added to the cuvette with the Hamilton syringe until a steady-state absorbance at 600 nm of 1.5-2.0 was obtained. The reaction was started by injecting nitrate through the rubber septum to a final conc. of 10 mM. The resulting absorbance decrease was measured at 600 nm and recorded every 5 s. All assays were performed in a water jacketed chamber maintained at 30 °C. The enzyme activity was determined using the formula: $\epsilon = A / c \times d \text{ cm mol}^{-1}$ (ϵ for MV⁺ is 13 mM⁻¹ cm⁻¹) (26).

Results and Discussion

The number of bacteria grown on defined agar plates at 30 °C for 2 days was 1 x 10⁶ cfu/g soil. Sixty isolates, which formed colonies on defined agar plates, were selected for nitrate reductase activity. Thirty-nine of the 60 tested isolates were found to reduce the added nitrate to nitrite under aerobic conditions in the test system according to the formation of a reddish pink color on the plates. In the same test system, no nitrite was observed in 21 out of those 60 tested isolates. Nitrate reduction in the test system by isolates is attributed to the activity of periplasmic nitrate reductase enzyme (9,19) because under these experimental conditions, 1) activity of NAS is repressed by ammonium that already exists in the growth medium (27); 2) NAR cannot reduce nitrate to nitrite under oxic conditions (9,28,29). However, nitrite seen on the test system might be the result of heterotrophic nitrification of ammonium (30) that exists on the growth medium but the absence of color formation on the control plates rules out this possibility.

The morphological and biochemical properties of those 39 isolates found to reduce nitrate were characterized and 5 of those 39 characterized isolates' biochemical properties are given in Table 1. Since the physiological function of the periplasmic nitrate reductase in nitrogen fixing bacteria (*Rhizobium* spp. and

Table 1. Biochemical and morphological properties of isolates.

| Isolates/properties | A11 | C2 | C8 | D3 | E2 |
|----------------------|-----|----|----|----|----|
| Gram | - | - | - | - | - |
| Catalase | + | - | - | + | - |
| Oxidase activity | + | + | + | + | + |
| Glucose fermentation | - | - | - | - | - |
| Nitrogen fixation | - | + | + | - | + |
| Chemolithotroph | - | + | + | + | + |
| Starch hydrolysis | - | - | - | - | - |
| Arginine dihydrolase | + | - | - | + | - |
| Gelatin hydrolysis | - | - | - | - | - |
| Fluorescent pigment | - | - | - | + | - |
| Denitrification | - | + | + | - | + |

Azospirillum brasilense Sp245) was recently found to be uniquely different from its general assumed functions (8,17), 3 of 5 isolates were selected from the nitrogen fixing bacterial group. Although these nitrogen fixing isolates seem to be the same according to their properties given in Table 1, their colony morphology on defined agar medium and growth on aerobic nitrogen fixing liquid medium indicate that they differ from each other.

The 5 isolates selected were grown aerobically in a defined liquid medium with different nitrogen sources (either 10 mM of NO_3 or NO_3 and NH_3 or NH_3) as a sole nitrogen source. It was found that nitrite began to accumulate after 2 and 8 h of incubation of the isolates when NO_3 or NO_3 and NH_3 was the sole nitrogen source. Maximum nitrite accumulations for the isolates are given in Table 2.

When NO_3 is the sole nitrogen source in the growth medium, bacteria with assimilatory nitrate reductase can reduce nitrate to nitrite and then nitrite is reduced to ammonium. Therefore, nitrite accumulation in the NO_3 growth medium can be attributed to NAS activity. On the other hand, NAP may be responsible for the nitrite accumulation in these cultures as it is an oxygen insensitive periplasmic dissimilatory enzyme and its

product can be readily excreted from cells. In the light of this study, it is not clear whether NAS can produce this excess amount of nitrite or whether NAP contributes significantly to nitrate reduction, but nitrite accumulation in these cultures deserves to be characterized.

Nitrite accumulation in the NO_3 and NH_3 growth medium can be attributed to neither NAS nor NAR as they are repressed by ammonium and oxygen, respectively. Therefore, nitrite accumulation under oxic conditions suggests that nitrite produced by the isolates A11, C2, C8, D3 and E2 is likely to be a result of their NAP activity (Table 2).

No nitrite was determined at any stage of growth when NH_3 was the sole nitrogen source. Some denitrifying bacteria can produce nitrite by oxidation of ammonium through heterotrophic nitrification (30). Although it is not known whether the isolates can produce nitrite by heterotrophic nitrification, it is clear that nitrite does not accumulate as a result of this mechanism.

In order to show that the isolates have NAP activity, reduced MV^+ was used as a non-biological electron donor. Table 3 shows that A11, C2, C8 and E2 have NAP enzyme activity. No NAP activity was observed in the isolate D3. This result suggests that either the isolate D3 has no NAP enzyme or its NAP activity cannot be

Table 3. Whole-cell MV^+ dependent periplasmic nitrate reductase activities.*

| Isolate | nmol/min/mg |
|---------|-------------|
| A11 | 8 |
| C2 | 16 |
| C8 | 16 |
| D3 | 0 |
| E2 | 20 |

* Units are nanomoles of MV^+ electron donor oxidized per minute per milligram cell of dry weight.

Table 2. Maximum nitrite accumulation for the isolates when they were grown aerobically in NO_3 or NH_3 and NO_3 medium as a sole nitrogen source.

| Isolate | NO_3 medium (nitrite μM) | $\text{NH}_3 + \text{NO}_3$ medium (nitrite μM) | Nitrite accumulation Assimilation/Dissimilation |
|---------|--|--|--|
| A11 | 1000 | 5 | 200 |
| C2 | 1400 | 300 | 5 |
| C8 | 15 | 5 | 3 |
| D3 | 220 | 30 | 7 |
| E2 | 450 | 10 | 45 |

determined with this assay. Three of the isolates that showed NAP activity belong to free aerobic nitrogen fixing bacteria. Recently, Bedzyk et al. (1999) and Steenhoudt et al. (2001) have shown that nitrogen fixing bacteria, *Rhizobium* spp. and *Azospirillum brasilense* Sp245, also have the NAP enzyme.

NO₃ and NO₂ are 2 important nitrogen forms for organisms and their environment. Denitrifying bacteria are used to remove nitrate from polluted waters (31). The aerobic nitrogen fixer chemolithotrophic isolates (C2, C8 and E2) found to reduce nitrate to nitrite under aerobic conditions can also reduce nitrate under anaerobic conditions (Table 1). Therefore, these isolates can be used in the treatment of waters that are poor in organic materials.

The study suggests that when ammonium exists together with nitrate, aerobic nitrate reduction does not generally lead to high nitrite accumulation. However, when nitrate exists as the sole nitrogen source, nitrite can accumulate in large amounts in aerobic zones of the soil. Since nitrite is not a preferred nitrogen form for plants and leads to soil acidification and, consequently, loss of

cations, the use of nitrate-only fertilizers may not be as beneficial as expected in terms of soil productivity. Furthermore, nitrite production as a result of aerobic denitrification suggests that where crop yield is concerned, it is also important to consider the loss of NO₃, which is readily available to plants, from agricultural soil by aerobic nitrate reduction, as well as by anaerobic denitrification.

Acknowledgments

This work was supported by Akdeniz University's Scientific Research Project Fund.

Corresponding author:

Mehmet Akif KILIÇ

Department of Biology,

Faculty of Arts and Sciences, Akdeniz University

Dumlupınar Bulvarı, 07058 Kampus,

Antalya - TURKEY

mkilic@akdeniz.edu.tr

References

- Zumft WG. Cell biology and molecular basis of denitrification. *Microbiol and Mol Bio Rew* 61: 533-616, 1997.
- Lisa AC, Todoroff K, Shaw GM. Maternal exposure to nitrate from drinking water and diet and risk for neural tube defects. *Am J Epidemiol* 153: 325-31, 2001.
- Aber JD. Nitrogen cycling and nitrogen saturation in temperate forest ecosystems. *Trends Ecol Evol* 7: 220-224, 1992.
- Schulze ED. Air-pollution and forest decline in a spruce (*Picea abies*) forest. *Science* 244: 776-783, 1989.
- Lloyd D, Boddy L, Davies K. Persistence of bacterial denitrification capacity under aerobic conditions: the rule rather than the exception. *FEMS Microbiol Ecol* 45: 185-190, 1987.
- Ferguson SJ. Denitrification and its control. *Antonie van Leeuwenhoek* 66: 89-110, 1994.
- Bell LC, Richardson DJ, Ferguson SJ. Periplasmic and membrane-bound respiratory nitrate reductase in *Thiosphaera pantotropha*. *FEBS Lett* 265: 85-87, 1990.
- Bedzyk L, Wang T, Ye RW. The periplasmic nitrate reductase in *Pseudomonas* sp. strain G-179 catalyzes the first step of denitrification. *J Bact* 181: 2802-2806, 1999.
- Berks BC, Ferguson SJ, Moir JWB, Richardson DJ. Enzymes and associated electron transport systems that catalyse the respiratory reduction of nitrogen oxides and oxyanions. *Biochim Biophys Acta* 1232: 97-173, 1995.
- Cole JA. In the nitrogen and sulphur cyclus. In: Cole JA and Ferguson SJ eds. *Assimilatory and dissimilatory reduction of nitrate to ammonia*. Cambridge University Press; pp. 281-327, 1988.
- Hochstein LI, Tomlinson GA. The enzymes associated with denitrification. *Ann Rev Microbiol* 42: 231-261, 1988.
- Moreno-Vivian C, Cabello P, Martinez-Luque M et al. Prokaryotic nitrate reduction: Molecular properties and functional distinction among bacterial nitrate reductases. *J Bact* 181: 6573-6584, 1999.
- McEwan AG, Wetzstein HG, Meyer O et al. The periplasmic nitrate reductase of *Rhodobacter capsulatus*; purification, characterisation and distinction from a single reductase for trimethylamine-N-oxide, dimethylsulphoxide and chlorate. *Arch Microbiol* 147: 340-345, 1987.
- Siddiqui RA, Warnece-Eberz U, Hengsberger A et al. Structure and function of a periplasmic nitrate reductase in *Alcaligenes eutrophus* H16. *J Bact* 175: 5867-5876, 1993.
- Berks BC, Richardson DJ, Robinson C et al. Purification and characterization of the periplasmic nitrate reductase from *Thiosphaera pantotropha*. *Eur J Biochem* 220: 117-124, 1994.
- Carter JP, Richardson DJ, Spiro S., Isolation and characterisation of a strain of *Pseudomonas putida* that can express a periplasmic nitrate reductase. *Arch Microbiol* 163: 159-166, 1995.

17. Steenhoudt O, Keijers V, Okon Y, Vanderleyden J. Identification and characterization of a periplasmic nitrate reductase in *Azospirillum brasilense* Sp245. *Arch Microbiol* 175: 344-352, 2001.
18. Richardson DJ, Berks BC, Russell DA et al. Functional, biochemical and genetic diversity of prokaryotic nitrate reductases. *Cell Mol Life Sci* 58: 165-178, 2001.
19. Carter JP, Hsiao Y, Spiro S, Richardson DJ. Soil and sediment bacteria capable of aerobic nitrate respiration. *Appl Environ Microbiol* 61: 2852-2858, 1995.
20. Flanagan DA, Gregory LG, Carter JP et al. Detection of genes for periplasmic nitrate reductase in nitrate respiring bacteria and in community DNA. *FEMS Microbiol Lett* 177: 263-270, 1999.
21. Sears HJ, Sawers G, Berks BC et al. Control of periplasmic nitrate reductase gene expression (napEDABC) from *Paracoccus pantotrophus* in response to oxygen and carbon substrates. *Microbiol* 146: 2977-2985, 2000.
22. Kilic MA. Physiological and biochemical studies of the periplasmic nitrate reductase from a *Pseudomonas* species. MSc, University of East Anglia, 1995.
23. Wollum AG. Cultural methods for soil microorganisms. In: Page AL, Miller RH, Keeney DR eds. *Methods of soil analysis part II*. American Society of Agronomy; pp. 1159, 1982.
24. Krieg RN, Holt GJ. *Bergey's manual of systematic bacteriology*. Williams and Wilkins, 1, United States of America, 1984.
25. Coleman JK, Cornish-Bowden A, Cole JA. Purification and properties of nitrate reductase from *Escherichia coli* K12. *Biochem J* 175: 483-493, 1978.
26. Uchumira H, Enjoji H, Seki T et al. Nitrate reductase-formate dehydrogenase couple involved in the fungal denitrification by *Fusarium oxysporum*. *J Biochem* 131: 579-586, 2002.
27. McCarty GW, Bremner JM. Regulation of assimilatory nitrate reductase activity in soil by microbial assimilation of ammonium. *Proc Natl Acad Sci* 89: 453-456, 1992.
28. Nicholls DG, Ferguson SJ. *Bioenergetics 2*. Academic Press, London: 1992.
29. Sears HJ, Ferguson SJ, Richardson DJ, Spiro S. The identification of a periplasmic nitrate reductase in *Paracoccus denitrificans*. *FEMS Microbiol Lett* 113: 107-112, 1993.
30. Robertson L, Kuenen JG. Heterotrophic nitrification in *Thiosphaera pantotropha*: Oxygen uptake and enzyme studies. *J Gen Microbiol* 134: 857-863, 1988.
31. Okmen GK, Algur FO. Farklı karbon kaynaklarının ve C/N oranlarının mikrobiyal denitrifikasyon üzerine etkileri. *Türk J Biol* 24: 533-542, 2000.