

1-1-2021

## Concomitant accumulations of ions, osmoprotectants and antioxidant system-related substances provide salt tolerance capability to succulent extreme-halophyte *Scorzonera hieraciifolia*

CANSU ALTUNTAŞ

RABİYE TERZİ

Follow this and additional works at: <https://dctubitak.researchcommons.org/botany>



Part of the [Botany Commons](#)

---

### Recommended Citation

ALTUNTAŞ, CANSU and TERZİ, RABİYE (2021) "Concomitant accumulations of ions, osmoprotectants and antioxidant system-related substances provide salt tolerance capability to succulent extreme-halophyte *Scorzonera hieraciifolia*," *Turkish Journal of Botany*: Vol. 45: No. 4, Article 7. <https://doi.org/10.3906/bot-2102-7>

Available at: <https://dctubitak.researchcommons.org/botany/vol45/iss4/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals.

## Concomitant accumulations of ions, osmoprotectants and antioxidant system-related substances provide salt tolerance capability to succulent extreme-halophyte *Scorzonera hieraciifolia*

Cansu ALTUNTAŞ<sup>ORCID</sup>, Rabiye TERZİ<sup>ORCID</sup>

Department of Biology, Faculty of Science, Karadeniz Technical University, Trabzon, Turkey

Received: 04.02.2021 • Accepted/Published Online: 22.05.2021 • Final Version: 14.07.2021

**Abstract:** Halophytes adapting to live in salinized areas can activate some tolerance mechanism through signal compounds to cope with salinity. However, the role of co-activity of signal compounds in salt tolerance of halophytes has not yet fully understood. We researched the changes in signal compounds involved in the salt tolerance mechanism, including inorganic ions, osmoprotectants and substances related to the antioxidant system in *Scorzonera hieraciifolia* with fleshy shoots extreme-halophyte. The levels of calcium, magnesium, proline, soluble sugar, hydrogen peroxide, superoxide, ascorbate and glutathione increased when thickness of shoot tissues enhanced under excess salinity. There were 3.3-fold, 5-fold, 8-fold and 10-fold enhancements in the levels of inorganic ions ( $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ ),  $\text{H}_2\text{O}_2$ , ascorbate and glutathione in the shoots treated with excess salinity, respectively. Contents of sodium, potassium and chlorine, and antioxidant enzyme activities also increased in the salinized shoots. The increases in antioxidant enzyme activities were consistent with increases in their protein contents according to Western blot analysis. The results suggest that extraordinary salt tolerance capacity in *Scorzonera hieraciifolia* can improve by modulated accumulations of signal compounds, particularly calcium, magnesium, osmoprotectants, reactive oxygen species and antioxidant substances. Moreover, massive induction of antioxidant enzymes can make strong contributions to salinity tolerance of *S. hieraciifolia*.

**Key words:** Antioxidant system, extreme-halophyte, inorganic ion, osmoprotectant, salt tolerance, signal compound

### 1. Introduction

Salt stress, one of the abiotic stresses, has become a common problem for crops all over the world, especially in recent years. Salinity can cause osmotic and ionic imbalance, functional and structural protein damage and membrane injury in plants. However, plants can promote some tolerance mechanism linked with signal transduction, reactive oxygen species (ROS) production, induction of antioxidants, synthesizing of osmoprotectants, controlling ion absorption, and expression of salt responsive genes and transcription factor to cope with salinity (Gupta and Huang, 2014; Mishra and Tanna, 2017; Rubnawaz et al., 2020). Halophytes, salt-tolerant or salt-resistant plants may also exhibit physiological and morphological changes such as leaf shedding and succulent structures (Mishra and Tanna, 2017). The tolerance degrees of halophytes to salinity could vary greatly between species (Akyol et al., 2020). Some halophytic species grow well at low salinity level while extreme halophytes can survive even at salt concentrations higher than 500 mM (Inan et al., 2004). For instance, crops such as sugar beet, date palm and barley

that could survive at 85 mM NaCl irrigation water are sometimes considered halophytes while extreme halophyte *Salicornia bigelovii* could survive at up to 1300 mM NaCl, the high end of salt tolerance (Akyol et al., 2020).

Besides, signal compounds, which are members of signaling networks, participate in the growth and developmental responses of plants to unfavorable environmental factors. For instance, it has been recently reported that ascorbic acid (ASC) could contribute to the propagation of ROS/ $\text{Ca}^{2+}$  signals and act an extracellular signaling role in plant tissue (Makavitskaya et al., 2018). As a key-redox signaling component, glutathione (GSH) can interact with other signaling molecules including ROS and activate various defense mechanisms against unfavorable stress factors (Ghanta and Chattopadhyay, 2011). Similarly, some osmoprotectants such as proline and soluble sugars play signal role in abiotic stress tolerance of maize seedlings (Hayat et al., 2012; Altuntaş et al., 2019). On the other hand, osmoprotectants accumulated in plants during water deficiency and salinity help plants to avoid ion toxicity and maintain water uptake under the stress conditions without

\* Correspondence: rabiyyeterzi@yahoo.com.tr

preventing the normal metabolic processes. Like organic solutes (soluble sugars, proline, betaine, glycerol, and other low molecular weight metabolites), inorganic ions ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Cl}^-$ ) also have an osmoprotectant property and play a key role in osmotic adjustment to maintain water uptake (Zhou and Yu, 2009; Chen and Jiang, 2010). Some organic osmoprotectants and calcium, an inorganic osmoprotectant, act a signaling role in plants. Although, putative second messenger function of magnesium has been currently reported in an animal cell (Stangherlin and O'Neill, 2018), no signaling role of magnesium and other inorganic ions in responses of plants to abiotic stresses has been clarified yet.

The osmotic regulation by accumulation of inorganic anions and cations in the vacuole is the most important feature of succulent halophytes (Shabala and Mackay, 2011). It was reported that detrimental effects of increased  $\text{Na}^+$  and  $\text{Cl}^-$  contents were prevented by enhanced uptake of ions such as  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{K}^+$  in extreme halophyte two grasses (Mangalassery et al., 2017). Conversely, Ahmad et al. (2013) noted that  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{K}^+$  contents declined while  $\text{Na}^+$  concentration was increasing in shoot of *Salicornia persica*. They suggested that succulent halophytic *Salicornia persica* might use sodium ion for acceleration of water uptake under water shortage to regulate sodium concentration in cellular spaces. On the other hand, in plant cells, ensuring potassium hemostasis in a salinity environment is a key factor in detecting capability of salt tolerance. Moreover, the plants maintaining cytosolic  $\text{K}^+$ / $\text{Na}^+$  ratio can have salt tolerance capacity (Liang et al., 2018). In the light of these records, the contribution of variations in the mineral ion levels to the salinity tolerance in halophytes exposed to high salt concentrations is not well understood.

High salt concentrations in the living environment can cause hyperosmotic stress as well as ion imbalance in plants. As a result of these primary effects, oxidative stress can occur as a secondary effect (Gupta and Huang, 2014). However, the induction of antioxidant defense system consisting of enzymatic and nonenzymatic components maintains balance of reactive oxygen species within the cell. Well managed oxidative stress significantly contributed to the tolerance capacity of salinized plants by keeping cellular ROS levels in balance (Gupta and Huang, 2014).

Some studies have conducted on the salinity tolerance mechanism of succulent halophytes or extreme halophytes. For instance, Zeng et al. (2018) investigated some antioxidant enzyme activities in succulent halophyte *Carpobrotus rosii* suggested that salt stress resulted in significant increases in activities of major antioxidant enzymes, such as ascorbate peroxidase (APX), catalase (CAT) and glutathione reductase (GR) and decrease in

superoxide dismutase (SOD) activity in the mesophyll tissue. Likewise, extreme-halophyte *Thellungiella parvula*, a halophytic relative of *Arabidopsis* (*Arabidopsis thaliana*), could regulate ion hemostasis by providing osmotic adjustment and alleviate hazardous effects of excess salinity by inducing antioxidant system (Uzilday et al., 2015). It was also recorded that proline content enhanced in succulent extreme-halophyte, *Salvadora persica* under  $\text{NaCl}$  stress conditions (Parida et al., 2016). However, more researches should be performed on signal compounds to explain the salt tolerance mechanism of succulent-extreme halophytes.

*Scorzonera hieraciifolia* is an endemic halophyte species, mainly distributed in salt habitats in Irano-Turanian regions. It has been currently reported that *S. hieraciifolia* is a medicinal plant having in vitro antioxidant and anti-inflammatory activity (Sarı et al., 2019). We have observed fleshy shoots in *S. hieraciifolia* in the habitat. Understanding of the salt tolerance mechanism of *S. hieraciifolia*, a medicinal succulent extreme-halophyte, is important to provide a contribution to cultivation of medicinal plants in salinized areas.

The aim of the present study is to elucidate the signaling compounds involved in responses of the succulent extreme-halophytes to salt stress, including osmotic regulation and induction of antioxidant system. Firstly, extreme halophyte, *Thellungiella parvula* (syn. *Eutrema parvulum*), closely related to *Arabidopsis* (Uzilday et al., 2015), is used in the current research to determine whether *S. hieraciifolia* is an extreme-halophyte. Based on the measurement of some stress parameters indicating stress damage such as relative water content, lipid peroxidation and  $\text{H}_2\text{O}_2$  content by comparing to *Thellungiella parvula* under different salinity conditions (0, 150, 300, 450 and 600 mM), it was determined that *S. hieraciifolia* was an extreme-halophyte. We hypothesized that extraordinary salt tolerance of succulent extreme-halophytes can be provided by combined induction of multiple signal compounds especially calcium, magnesium, osmoprotectants, reactive oxygen species and antioxidant substances. We also tested the hypothesis that induction of antioxidant system can make strong contributions to salt stress tolerance of *S. hieraciifolia*.

## 2. Materials and methods

### 2.1. Plant materials and sterilization

*Scorzonera hieraciifolia* L. achenes were collected from Sivas Tödürge Lake (İran-Turan Region, Turkey) and brought to the laboratory. *Thellungiella parvula* (synonym, *Eutrema parvulum*) seeds gently requested and acquired from Prof. Dr. İsmail TURKAN, Ege University, İzmir, TURKEY. The mature achenes of *S. hieraciifolia* were carefully separated from their pappus bristles. The achenes

were stratified for 2 days at 4°C in petri dishes containing double-layer wet filter paper. After that, the achenes were treated with 10% (v/v) sodium hypochlorite (NaOCl) for sterilization, and then washing process was performed three times with sterile distilled water.

To determine whether *S. hieraciifolia* is an extreme halophyte plant, we compared *S. hieraciifolia* with the model plant for extreme halophytes, *T. parvula*, closely related to *Arabidopsis thaliana*. Similarly, *T. parvula* seeds were treated with 10% (v/v) NaOCl solution for sterilization followed by three rinsing stages with sterile distilled water.

## 2.2. Media, culture conditions and stress treatments

The sterilized *S. hieraciifolia* achenes and *T. parvula* seeds aseptically germinated at MS medium (pH = 5.8) (Murashige and Skoog, 1962) which is sterilized by autoclaving for 15 min at 121°C and 1 atm pressure. The nutrient basal medium is consisted of MS including vitamins (4.95 g L<sup>-1</sup>), 1% sugar (w/v) and 0.8% agar (w/v). This basal medium was supplemented with growth regulators including naphthalene acetic acid (NAA, 0.1 mg L<sup>-1</sup>), kinetin (1 mg L<sup>-1</sup>), isopentenyl adenine (2IP, 2 mg L<sup>-1</sup>) at different developmental stage of the plants. *S. hieraciifolia* and *T. parvula* plants were grown until their shoots develop (for 45 days and 35 days, respectively), at 24±2 °C, on light intensity 400–430 µmol photons m<sup>-2</sup>s<sup>-1</sup> and 16h light/8 h dark photoperiod. Subsequently, *S. hieraciifolia* and *T. parvula* plants were exposed to salt stress through 0, 150, 300, 450 and 600 mM NaCl supplements to MS media for 7 days. Salt concentrations were gradually enhanced by 150 mM NaCl supplements with 2 days intervals until reaching the last salinity levels (150, 300, 450 or 600 mM). Thus, gradually increasing salt concentrations were applied to the plants with transferring of the plants from previous salt concentration to the next concentration. Five different NaCl treatment groups (0, 150, 300, 450 and 600 mM NaCl) were constituted for *S. hieraciifolia* and *T. parvula*. Morphological observations and following measurements and analysis were conducted.

## 2.3. Stress determination parameters

To determine whether *S. hieraciifolia* is an extreme halophyte, we comparatively observed visual symptoms such as shoot chlorosis, wilting of shoots and plant death in *S. hieraciifolia* and *T. parvula* under salinity. Also, we determined changes in some stress parameters such as relative water content (RWC), lipid peroxidation and H<sub>2</sub>O<sub>2</sub> content.

### 2.3.1. Relative water content

The relative water content of *S. hieraciifolia* and *T. parvula* shoots were measured according to Castillo (1996). After shoot fresh weight (FW) was noted, the samples were incubated at 4°C for 16 h into distilled water. Shoots were

then blotted and shoot turgid weight (TW) was measured. To measure the shoot dry weight (DW), samples were continuously incubated in an oven at 80 °C until achieving a constant dry weight. The dry weights were determined. The shoot RWC was calculated using the equation:

$$\text{RWC (\%)} = (\text{FW} - \text{DW}) / (\text{TW} - \text{DW}) \times 100$$

### 2.3.2. Lipid peroxidation

The levels of thiobarbituric acid reactive substances (TBARS) were determined to assess the membrane damages in *S. hieraciifolia* and *T. parvula* shoots (Heath and Packer, 1968) ( $\epsilon = 155/(\text{mM cm})$ ). Trichloroacetic acid 0.1% (w/v) (TCA) was used for homogenization of the samples. Thiobarbituric acid in 20% TCA 0.5% (w/v) was added to the supernatants. After the reaction, the absorbances were read at 532 and 600 nm.

### 2.3.3. H<sub>2</sub>O<sub>2</sub> content

To determine hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) content, *S. hieraciifolia* and *T. parvula* samples were extracted with 5% cold TCA (w/v) with activated charcoal. After the extract was centrifuged, 10 mM potassium phosphate (K<sub>2</sub>HPO<sub>4</sub>) buffer (pH = 7.0) and 1 M potassium iodide (KI) were added to 1 mL supernatant. The absorbance was then read at 390 nm (Velikova et al., 2000).

## 2.4. Succulence measurements

We measured succulence degree, shoot thickness and plant area to determine the shoot succulence in extreme-halophyte *S. hieraciifolia*. The shoot succulence degree (SSD) was performed according to Qi et al. (2008). After measuring fresh weights and dry weights of *S. hieraciifolia*, the shoot succulence degree was expressed as the ratio of these two values to each other.

To determine shoot thickness (ST), all shoots of each treatment were transversally cut with the help of a razor blade in *S. hieraciifolia*. Subsequently, widths of the shoots were measured at several points under light microscope (Olympus, CX21, Olympus Corporation, Tokyo, Japan) with ocular meter and it was calculated by averaging these values.

The images of each plant area (PA) were determined by "ImageJ" Software Program (NIH, Bethesda, Maryland, USA). In this program, the images had to be in a certain format (8-bit grayscale, 16-bit gray scale, etc.). Image-Type 8-bit option that is appropriate to *S. hieraciifolia* was selected, and the scale was adjusted by entering the reference range 1 cm from the Analyze-Set Scale option. The area calculation for each shoot was performed from the Analyze-Tools-ROI Manager step.

## 2.5. Determination of osmoprotectants

Changes in total soluble sugar and proline contents in *S. hieraciifolia* exposed to enhanced salinity were determined.

### 2.5.1. Soluble sugar

To detect soluble sugar content, dry samples of *S. hieraciifolia* (0.1 g) were homogenized with 5 mL 70%



ethanol, and then it was boiled at 80 °C for 3 min and centrifuged at 10,000 × g for 5 min. For spectrophotometric measurement, 900 µL of distilled water was added to 100 µL of the supernatant and diluted. 5% phenol (1 mL) was added to this mixture and 96% sulfuric acid (5 mL) was added and then the mixture was homogenized by vortex. The mixture was cooled to room temperature and the absorbance of the mixture was recorded at 490 nm (Dubois, 1956).

### 2.5.2. Proline

Proline detection method was performed according to Bates et al. (1973). Oven dried shoots (0.1 g) were homogenized with 3% sulfosalicylic acid (5mL). The supernatants (1mL) were transferred to test tubes and mixed with equal volumes of glacial acetic acid and ninhydrin reagent. After that, the tubes were kept at 100 °C for 1 h. The test tubes were placed in an ice bath. The samples were mixed by a vortex after toluene (3mL) was added to the samples. The absorbance was measured at 520 nm on a UV-VIS spectrophotometer.

### 2.6. Inorganic ions

Using dry extracts of the shoots, anion-cation analyzes were performed with the Dionex ICS-5000 system (Haddad, 1997). Analysis of chloride ion content was performed with Dionex Ion Pac AS 9 HC (4×250 mm) separation column and 10 mM sodium carbonate fluid phase (flow rate: 1 mL/min). Analysis of the cation ion contents (Na<sup>+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>, Ca<sup>2+</sup>) was carried out with Dionex Ion Pac CS 12 A (3×150 mm) column and 20 mM meta sulfonic acid conductor phase (flow rate: 0.5 mL/min). The cation suppressor column Dionex CERS 500 (2 mm) and anion suppressor column Dionex AERS 500 (4 mm) were used. Samples were given to the column in 20 µL volumes by auto-sampler. Thermo Scientific conductivity detector was used in the system and calculations were made with Chromeleon Software System. (Thermo Fisher Scientific, Waltham, USA)

## 2.7. Antioxidant system-related substances

### 2.7.1. ROS levels

To determine content of superoxide by spectrophotometrically, using the tetrazolium salt XTT {(2, 3-bis (2-methoxy-4-nitro-5-sulphophenyl)-5-[(phenylamino) carbonyl]-2H-tetrazolium hydroxide} (Schopfer et al., 2001). The shoots of *S. hieracifolia* (0.5 g) were cut into small pieces and were vacuum-infiltrated for 20 min with 5 mL of 10 mM Na-citrate buffer (pH = 7.0) containing 500 µM XTT, with/without 3.5 U ml<sup>-1</sup> superoxide dismutase in dark. The samples were kept in this buffer for 2 h. The increase in XTT reduction in the shoots was recorded at 470 nm in a spectrophotometer.

### 2.7.2. Histochemical detection of H<sub>2</sub>O<sub>2</sub>

As mentioned above, the production of H<sub>2</sub>O<sub>2</sub> was determined both spectrophotometrically, and it was

visualized in vivo by 3,3'-diaminobenzidine (DAB) staining methods (Daudi et al., 2012). *S. hieracifolia* shoots were exposed to DAB prepared in 0.05% Tween 20 (v/v) and 10 mM sodium phosphate buffer (pH = 7.0). The shoots were then placed in test tubes and incubated at 80–100 rpm in a shaker. After incubation, the shoots were boiled in water at 90–95 °C for 15 ± 5 min in bleach solution; ethanol: acetic acid: glycerol (3: 1: 1). Color changes were observed in the shoots.

### 2.7.3. Ascorbate and glutathione contents

Ascorbate concentration was determined according to Liso et al. (1984). *S. hieracifolia* shoots (0.25 gr) were homogenized with 5 mL, 5% (w/v) m-phosphoric acid and the extract was centrifuged at 10,000 × g for 4 min. Sample (70 µL) was added to 3 ml of reaction medium containing 0.1 M citrate-0.2 M phosphate buffer (pH = 6.2). The initial absorbance was recorded at 265 nm and then the ascorbate concentration was determined by reading the reduction of 5 min after the addition of two units of ascorbate oxidase to the reaction medium. After ascorbate oxidation was completed, ascorbate oxidase was inhibited with 10 mM sodium azide. Dithiothreitol (DTT) (2.5 mM) was added to the medium and following reduction (3 min) with DTT, the absorbance was recorded again at 265 nm.

Glutathione content was determined according to the total glutathione assay kit (Northwest Life Science Specialties, LLC.) according to the instructions of the producer firm. Glutathione was measured using a reaction mixture containing 250 mM K<sub>2</sub>HPO<sub>4</sub>/KH<sub>2</sub>PO<sub>4</sub> (pH 7.5), 200 µM NADPH, 600 µM DTNB, 25 µL extract and 0.3 U GR. The change in absorbance was observed at 412 nm for 3 min. GSH concentration was calculated on the standard graph obtained by using GSH at 0-5 µM concentrations.

### 2.7.4. Antioxidant enzyme assays and protein determination

For superoxide dismutase, catalase and guaiacol peroxidase extractions, the shoot tissues (0.1 g) were homogenized with a 50 mM sodium phosphate buffer (pH = 7.8) with 1 mM ethylenediaminetetraacetic acid (EDTA) and polyvinylpyrrolidone (1%). For determination of APX activity, 2 mM of ascorbate was added into the sodium phosphate buffer. The samples were centrifuged at 15,000 × g for 15 min.

Superoxide dismutase activity was determined by Beauchamp and Fridovich (1971). Reaction was started by the addition of 2 µM riboflavin to 1 mL reaction medium (50 mM potassium phosphate buffer (pH = 7.8), 0.1 mM EDTA, 13 mM L-methionine, 75 µM nitro blue tetrazolium (NBT), and 50 µL extract). The absorbance was recorded at 560 nm after the mixture was exposed to white light at 375 µmol m<sup>-2</sup> s<sup>-1</sup> for 10 min. One unit (U) of SOD activity was defined as the amount of enzyme needed to bring about 50% inhibition of the NBT photoreduction rate.

Catalase activity was determined by according to Bergmeyer and Grafl (1983). The reduction in reaction time was measured at 240 nm for 5 min, containing 50 mM potassium phosphate buffer (pH = 7.0), 30 mM H<sub>2</sub>O<sub>2</sub> and 20 µL enzyme extract. Catalase activity was calculated using the 39.4 mM<sup>-1</sup>cm<sup>-1</sup> epsilon coefficient for H<sub>2</sub>O<sub>2</sub>. Ascorbate peroxidase was measured by the method of Nakano and Asada (1987). APX activity was determined by measuring at 290 nm in 1 mL reaction mixture containing 50 mM potassium phosphate buffer (pH = 7.0), 250 µM ascorbate, 5 mM H<sub>2</sub>O<sub>2</sub> and 20 µL enzyme extract. ( $\epsilon = 2.8 \text{ mM}^{-1}\text{cm}^{-1}$ ).

Guaiacol peroxidase activity was measured by increase in absorbance at 470 nm (25°C,  $\epsilon = 26.6 \text{ mM}^{-1}\text{cm}^{-1}$ ) in a 100 mM potassium phosphate buffer (pH = 7.0) containing 0.1 mM EDTA, 5 mM guaiacol, 15 mM H<sub>2</sub>O<sub>2</sub> and 50 µL of enzyme extract (Urbanek et al., 1991).

Glutathione reductase activity was determined spectrophotometrically (Foyer and Halliwell, 1976). GR was assayed by the fall in absorbance at 340 nm as NADPH was oxidized. The assay contained 50 mM Tris-HCl (pH 7.8), 150 µM NADPH, 500 µM oxidized glutathione (GSSG) and 50 µL extract. The activity of GR was calculated using an extinction coefficient of 6.22 mM<sup>-1</sup>cm<sup>-1</sup> for NADPH at 340 nm.

Antioxidant enzyme activities were presented on a protein basis. Detection of protein content was performed according to method of Bradford (1976), using BSA as a standard.

#### 2.7.5. Western blot analysis of antioxidant enzymes

Protein extractions of SOD, CAT, glutathione peroxidase (GPX; E.C. 1.11.1.9) and GR were performed using 4X PEB (protein extraction buffer, AS08 300 Agrisera Inc.) according to the manufacturer firm's instructions (for more details, see Shin et al., 2013). Samples were dissolved in an equal volume of sample buffer (2X Laemmli sample buffer 1610737 Bio-Rad) and heated at 95 °C for 5 min. Separation of total proteins (30 µg) by 12% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE, (TGX Stain-Free Precast Protein Gels, 4568095 Bio-Rad) was performed by electrophoresis (Mini-PROTEAN Tetra Cell system, 165800 Bio-Rad) at room temperature. The marker (Precision Plus Protein Western C Blotting Standards, 1610376 Bio-Rad) was used to determine the molecular weight of the proteins. Proteins separated in gel electrophoresis were transferred to the PVDF membrane (Trans-Blot Turbo Mini PVDF Transfer Packs, 1704156 Bio-Rad) using the Trans-Blot Turbo Transfer System (1704155 Bio-Rad). After transfer, the membrane was blocked at 4 °C with 2.5% milk powder in TBS. Fe-SOD Chloroplastic Fe-Dependent Superoxide Dismutase (A S06 125) for SOD, Glutathione Peroxidase Chloroplastic (A S04 055) for GPX, and, Catalase (A S09 501) for CAT were

primary used antibodies and they incubated overnight at 4 °C. After incubation, the membranes were incubated with goat anti-rabbit IgG-HRP secondary antibodies (AS09 602 Agrisera) for 4 h at room temperature. The density of the scanned protein bands was calculated with Image Lab Software (1709690, Bio-Rad Laboratories, Inc., California, USA).

#### 2.8. Statistical analysis

All experiments were carried out five times with five biological replicates. All results were presented as means  $\pm$  standard deviation. All physiological data were processed with one-way analysis of variance ( $\alpha = 0.05$ ) using the SPSS Ver. 15.0 software for Microsoft Windows (SPSS Inc., Chicago, IL, USA). Mean differences were determined with the Duncan multiple comparison test at  $\alpha = 0.05$ .

### 3. Results

#### 3.1. Determination of extreme halophyte characteristic of *S. hieraciifolia*

##### 3.1.1. Visual symptom changes


It was observed that *S. hieraciifolia* grown on the MS media could survive under the excess salt stress condition (600 mM NaCl) though shoot chlorosis and wilting. Conversely, *T. parvula* could survive at the 450 mM NaCl treatment but shoot senescence and plant death was observed at the 600 mM NaCl treatment. Additionally, shoot number increased in the 300 mM NaCl treatment in comparison with the 0 mM NaCl treatment in *S. hieraciifolia* while the shoot length was decreasing. Therefore, optimal growth was observed at 300 mM NaCl in *S. hieraciifolia*. At the same way, *T. parvula* displayed optimum growth at 300 mM NaCl (Figure 1 A-D).

##### 3.1.2. Effects of salinity on relative water content

The relative water content (%) of *S. hieraciifolia* shoots significantly enhanced at all levels of salinity in comparison with the 0 mM NaCl treatment. High increases in RWC were determined in the 450 and 600 mM NaCl treatments. Conversely, in *T. parvula*, the RWC value decreased while salt concentration was increasing. The highest decrease was determined the shoots treated with 450 mM NaCl in *T. parvula* (Table 1).

##### 3.1.3. Effects of salinity on lipid peroxidation

Lipid peroxidation enhanced in the shoots treated with 300, 450 and 600 mM NaCl, while 150 mM NaCl treatment had no significant effect in comparison with 0 mM NaCl in *S. hieraciifolia*. However, in *T. parvula*, TBARS content was increased in all salt treatments compared to the 0 mM NaCl treatment and the highest TBARS content was observed in the 450 mM salt treatment. As mentioned above, the shoots of *T. parvula* lost their vitality in the 600 mM NaCl treatment. The rates of increase of TBARS content in both plants were same (1.1-fold) in the 300 mM



**Figure 1.** Morphological alterations in the shoots of *Scorzonera hieraciifolia* and *Thellungiella parvula* treated with 0, 150, 300, 450 and 600 mM NaCl (A, B). *S. hieraciifolia* grown at 300 mM NaCl (C). *S. hieraciifolia* grown at 600 mM NaCl (D).

**Table 1.** Alterations in some stress-determining parameters in the shoots of *Scorzonera hieraciifolia* and *Thellungiella parvula* treated with 0, 150, 300, 450 and 600 mM NaCl. Means  $\pm$  SD. Means followed by different letters are significantly different according to Duncan's test at  $p \pm 0.05$ .

NaCl	<i>Scorzonera hieraciifolia</i>			<i>Thellungiella parvula</i>		
(mM)	RWC (%)	TBARS ( $\mu\text{molg}^{-1}\text{DW}$ )	$\text{H}_2\text{O}_2$ ( $\mu\text{molg}^{-1}\text{DW}$ )	RWC (%)	TBARS ( $\mu\text{molg}^{-1}\text{DW}$ )	$\text{H}_2\text{O}_2$ ( $\mu\text{molg}^{-1}\text{DW}$ )
0	73.51 $\pm$ 0.8 d	2.8 $\pm$ 0.09 d	6.3 $\pm$ 0.01 e	64.23 $\pm$ 0.1 a	6.1 $\pm$ 0.9 d	6.2 $\pm$ 0.01 d
150	76.44 $\pm$ 0.8 c	2.7 $\pm$ 0.08 d	8.7 $\pm$ 0.01 d	54.64 $\pm$ 0.2 b	8.2 $\pm$ 0.9 c	11 $\pm$ 0.01 c
300	82.84 $\pm$ 0.8 b	3.2 $\pm$ 0.08 c	11.5 $\pm$ 0.05 c	53.12 $\pm$ 0.5 c	9.5 $\pm$ 0.5 b	15.1 $\pm$ 0.02 b
450	92.07 $\pm$ 0.6 a	8.3 $\pm$ 0.25 b	28.0 $\pm$ 0.05 b	49.35 $\pm$ 0.2 d	23.2 $\pm$ 0.6 a	51.3 $\pm$ 0.01 a
600	94.1 $\pm$ 1.0 a	9.1 $\pm$ 0.12 a	32.2 $\pm$ 0.07 a	NA	NA	NA

DW: dry weight. NA: non-alive

NaCl treatments where observed the optimum growth. However, there were 2.9-fold and 3.8-fold increases in *S. hieraciifolia* and *T. parvula* in the 450 mM NaCl treatments, respectively (Table 1).

#### 3.1.4. Effects of salinity on $\text{H}_2\text{O}_2$ content

$\text{H}_2\text{O}_2$  contents gradually increased in *S. hieraciifolia* and *T. parvula* while salt concentration was increasing.  $\text{H}_2\text{O}_2$  contents and the rates of the increases observed in *T. parvula* were higher than those of *S. hieraciifolia*. For example, there were 2.4-fold increase in *T. parvula* and 1.8-fold in *S. hieraciifolia* in the 300 mM NaCl treated shoots as compared with non-saline conditions. Likewise, there were 8.3-fold and 4.4-fold enhancements in  $\text{H}_2\text{O}_2$  contents of the shoots treated with 450 mM NaCl in *T. parvula* and *S. hieraciifolia*, respectively (Table 1).

#### 3.2. Determination of shoot succulence in *S. hieraciifolia*

Fleshy shoots were observed at 300 mM and higher salt concentrations in *Scorzonera hieraciifolia*. Accordingly, the shoot succulence degree increased in the 300 mM NaCl treatment in comparison with the non-stressed shoots. The succulence degrees were high in the 450 and 600 mM NaCl treatments (Table 2). Likewise, the shoot thickness increased depending on salt concentration. The highest increase in the shoot thickness was observed in the 600 mM NaCl treated shoots. As compared with the 0 mM NaCl treatment, there were 2.4-fold and 9-fold enhancements in the thicknesses of shoots treated with 300 mM and 600 mM NaCl, respectively (Table 2).

The salt treatments also increased the plant area. The highest plant area was detected in plants treated with

**Table 2.** Changes in parameters related to shoot succulence in *S. hieraciifolia*. Means ± SD. Means followed by different letters are significantly different according to Duncan's test at  $p \pm 0.05$ . SSD: shoot succulence degree, ST: shoot thickness, PA: plant area.

NaCl (mM)	SSD	ST ( $\mu\text{m}$ )	PA ( $\text{cm}^2 \text{ plant}^{-1}$ )
0	46 ± 2.5 c	39.2 ± 0.3 e	2.31 ± 0.6 d
150	47 ± 1.1 c	52.8 ± 0.5 d	3.81 ± 0.4 c
300	69 ± 1.4 b	92.5 ± 4.1 c	12.5 ± 2.1 a
450	86 ± 2.5 a	311 ± 4.8 b	8.08 ± 2.1 b
600	88 ± 2.7 a	351 ± 7.3 a	6.28 ± 0.8 b

300 mM NaCl (Table 2). There was 5.4-fold and 2.7-fold enhancements in the plant areas treated with 300 mM and 600 mM NaCl, respectively (Table 2).

### 3.3. Effects of salinity on osmoprotectants

#### 3.3.1. Soluble sugar and proline contents

Both total soluble sugar and proline contents increased by the all salt treatments compared to the non-stressed shoots in *Scorzonera hieraciifolia*. Moreover, the highest increases in the contents of proline (2.6-fold) and total soluble sugar (1.4-fold) were observed in the 600 mM salt treated seedlings in comparison with the 0 mM NaCl treatment. There were 1.2-fold and 1.3-fold increases in proline and total soluble sugar contents of the shoots treated with 300 mM NaCl, respectively (Figure 2A, B).

#### 3.4. Ion contents


$\text{K}^+$  ion concentration increased at all NaCl concentrations in comparison with the non-stressed shoots.  $\text{K}^+$  content was high in the 150 and 300 mM NaCl treated shoots.  $\text{Na}^+$  and  $\text{Cl}^-$  contents increased gradually with the enhanced NaCl concentrations as compared to 0 mM NaCl treatment. The highest  $\text{Na}^+$  and  $\text{Cl}^-$  contents were detected in the shoots exposed to 600 mM NaCl.  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ion contents significantly increased in all salt treatments as compared with non-saline condition. The highest increases in  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  contents were determined in the 600 mM NaCl treated shoots and the rate of increase of both  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  was recorded by 3.3-fold (Table 3).

### 3.5. Effects of salinity on antioxidant system-related signal compounds

#### 3.5.1. ROS levels

Superoxide ( $\text{O}_2^{\cdot-}$ ) level in the shoot of *S. hieraciifolia* increased in parallel with the increasing salt level. The highest superoxide level was detected in the shoots treated with 600 mM NaCl (Figure 3).

The changes in  $\text{H}_2\text{O}_2$  level in the shoot tissues of *S. hieraciifolia* was also determined with the observation of light brown by DAB staining. We observed that all salt



**Figure 2.** Alterations in osmoregulation related signal compounds, total soluble sugar (A) and proline (B), in the shoots of *S. hieraciifolia*. Vertical bars represent standard deviations. Different letters denote significant differences among all treatments at  $p < 0.05$ .

treatments increased  $\text{H}_2\text{O}_2$  level in comparison with the non-stressed shoots (Figure 4).

#### 3.5.2. Ascorbate and glutathione contents

The total ascorbate concentration no significantly changed in the shoots treated with 150 mM NaCl. However, the 300 mM NaCl and further salt concentrations (450 and 600 mM) significantly increased the total ascorbate content in comparison with 0 mM NaCl treatment. As compared to the 0 mM NaCl treatment, there were 2.2-fold and 8-fold increases in ASC content of the shoots treated with 300 mM and 600 mM NaCl, respectively (Figure 5A).

Total glutathione content gradually enhanced while salt concentration was increasing in *S. hieraciifolia*. There were 3.4-fold and 10-fold increases in GSH content of the shoots treated with 300 mM and 600 mM NaCl in comparison with the 0 mM NaCl treatment respectively (Figure 5B).


#### 3.5.3. Effects of salinity on antioxidant enzyme activities

The activities SOD and APX significantly enhanced while NaCl concentrations were increasing. The highest



**Table 3.** Alterations in ion contents in the shoots of *S. hieraciifolia*. Means  $\pm$  SD. Means followed by different letters are significantly different according to Duncan's test at  $p \pm 0.05$ .

NaCl (mM)	K <sup>+</sup>	Na <sup>+</sup>	Ca <sup>++</sup>	Mg <sup>++</sup>	Cl <sup>-</sup>
0	198.6 $\pm$ 0.7 d	5.01 $\pm$ 0.01 e	8.72 $\pm$ 0.4 e	0.37 $\pm$ 0.03 d	1.76 $\pm$ 0.2 e
150	373.1 $\pm$ 10 a	46.7 $\pm$ 0.06 d	23.6 $\pm$ 0.9 b	1.18 $\pm$ 0.06 a	18.9 $\pm$ 2.2 d
300	364.1 $\pm$ 0.5 a	76.9 $\pm$ 0.1 c	22.5 $\pm$ 0.3 c	0.94 $\pm$ 0.01 b	24.2 $\pm$ 1.2 c
450	269.9 $\pm$ 0.9 c	103.6 $\pm$ 0.9 b	20.6 $\pm$ 0.4 d	0.82 $\pm$ 0.03 c	26.3 $\pm$ 0.1 b
600	288.9 $\pm$ 3.5 b	144.5 $\pm$ 1.8	28.7 $\pm$ 0.5 a	1.22 $\pm$ 0.04 a	28.6 $\pm$ 1.5 a




**Figure 3.** Alterations in superoxide level in the shoots of *S. hieraciifolia*. Vertical bars represent standard deviations. Different letters denote significant differences among all treatments at  $p < 0.05$ .

increases in the activities of SOD and APX were recorded in the 600 mM NaCl treatment (Figure 6 A). The catalase activity increased in the 150 and 300 mM NaCl treatments but it decreased in the 450 and 600 mM NaCl treatments (Figure 6 B). There was no significant change in GPOD activity in the shoots treated with 150 mM NaCl, while the activity increased in 300 mM, (1.7-fold), 450 mM (3.4-fold) and 600 mM (4-fold) NaCl treatments as compared to the 0 mM NaCl treatment (Figure 6 B). As for glutathione reductase, the activity increased under all salt stress conditions. The rate of the increase was high (2.5-fold) in the high (600 mM) salt treatment (Figure 6 B).

#### 3.5.4. Effect of salinity on antioxidant enzyme contents

The increases in the SOD activity were consistent with increases in Fe-SOD contents in *S. hieraciifolia*. Fe-SOD content increased in all salt treatments compared to 0 mM NaCl treatment and, Fe-SOD content was high in the shoots treated with 450 and 600 mM NaCl (Figure 7). The protein content of catalase distinctively enhanced in the shoots treated with 300 mM NaCl. Similar to the CAT activity, there was a decrease trend in the shoots




**Figure 4.** Alterations in H<sub>2</sub>O<sub>2</sub> content detected by DAB staining in the shoots of *S. hieraciifolia*.

exposed to 450 mM NaCl. Also, CAT content did not change significantly in 150 and 600 mM salt treatments (Figure 6). Glutathione peroxidase content increased in all salt treatments in comparison with the non-stressed shoots. GPX content was especially high in the 300 mM NaCl and upper salt treatments (Figure 7). GR content increased in the shoots treated with 150, 300 and 450 mM NaCl compared to the non-stressed shoots. There was a significant increase in the GR protein content in the 300 mM treatment in *S. hieraciifolia* (Figure 7).

#### 4. Discussion

We detected that *S. hieraciifolia* displayed optimal growth at the 300 mM NaCl concentration, as similar to *Salvadora persica*, stem-succulent extreme-halophyte, reported by Aghaleh et al. (2009). *S. hieraciifolia* was able to survive, even at the 600 mM NaCl concentration in spite of chlorosis. Likewise, Aghaleh et al. (2009) recorded that succulent halophyte, *Salvadora persica* was able to survive at 600 mM salt, which is a dose higher than the concentration of salt in seawater. *T. parvula* also displayed optimal growth at the 300 mM NaCl but it was unable to tolerate the 600 mM NaCl concentration in MS media. Accordingly, Uzilday et al. (2015) showed that *T. parvula* grown in a media including peat moss, vermiculite and




**Figure 5.** Alterations in non-enzymatic antioxidant signal compounds, ascorbate (A) and glutathione (B), in the shoots of *S. hieraciifolia*. Vertical bars represent standard deviations. Different letters denote significant differences among all treatments at  $p < 0.05$ .

soil mix could withstand 300 mM NaCl treatment. These results indicated that the maximum salt concentration that could be tolerated by halophytes may depend on the plant species, the presence of succulent structures, severity of the stress treatments and media exposed to the stress.

The relative water content increased in accordance with increased salt concentrations in *S. hieraciifolia* with fleshy shoots. Conversely, RWC value decreased with increasing salt concentration in *T. parvula*. Our data in *T. parvula* is consistent with previous reports that RWC and osmotic potential decreased in extreme halophyte, *Salsola crassa* (Yildiztugay et al., 2014). Increase in RWC indicated that succulent extreme-halophytes could dilute excess salt in their succulent structures and they could promote a mechanism of salt tolerance.

Amount of thiobarbituric acid reactive substances that is the product of lipid peroxidation, and ROS levels in plant tissues are important parameters indicating the stress damage of plants (Gupta and Huang, 2014). Uzilday et al. (2015) reported no significant change in membrane damage in shoots of *T. parvula* grown in saline soil (300 mM NaCl). However, our findings obtained in MS media showed that TBARS content and  $\text{H}_2\text{O}_2$  level increased by enhanced salt concentrations in *T. parvula*, and the rates




**Figure 6.** Alterations in antioxidant enzyme activities, superoxide dismutase, ascorbate peroxidase (A), glutathione reductase, catalase and guaiacol peroxidase (B), in the shoots of *S. hieraciifolia*. Vertical bars represent standard deviations. Different letters denote significant differences among all treatments at  $p < 0.05$ .

of the increases in *T. parvula* were higher than those of *S. hieraciifolia*. Likewise, Uzilday et al. (2015) reported that  $\text{H}_2\text{O}_2$  content significantly increased by the 300 mM NaCl treatment but no changed by low salt (50 and 200 mM NaCl) treatments in *T. parvula*. Visual symptom changes and RWC, TBARS and  $\text{H}_2\text{O}_2$  content data showed that *S. hieraciifolia* can be capable of tolerating high salt conditions and thus it can have characteristic properties of extreme-halophytes like *T. parvula*.

We also determined shoot succulence in *S. hieraciifolia* by measuring changes in the shoot succulence degree, shoot thickness and plant area under increased salinity conditions. Consistent with our findings, Parida et al. (2016) reported that succulence degree increased when salinity was increasing in extreme-halophyte, *Salvadora persica*. Increased shoot thickness and plant area due to increased salt concentration supported the idea that *S. hieraciifolia* displayed a succulent shoot structure under salt stress.

Plants supply osmotic adjustment via two processes under high saline environment: ion accumulation in the



**Figure 7.** Alterations in antioxidant enzyme contents, Fe-superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase, in the shoots of *S. hieraciifolia*. Vertical bars represent standard deviations. Different letters denote significant differences among all treatments at  $p < 0.05$ .

vacuole and synthesis of compatible solutes in cytosol (Zhou and Yu, 2009; Chen and Jiang, 2010; Shabala and Mackay, 2011; Boughalleb et al., 2020). Our results showed that proline content and total soluble sugars significantly increased in *S. hieraciifolia* shoots at all levels of salinity. Accordingly, Parida et al. (2016) recorded that proline content increased in succulent extreme-halophyte, *Salvadora persica* under NaCl stress conditions. Liang et al. (2018) suggested that the soluble sugar content of Arabidopsis overexpressing salt-related wheat *TaSST* gene was significantly higher than that of wild-type and the transgenic plants resisted external salt stress by accumulating soluble sugars. The regulated accumulation of total soluble sugar and proline with signaling functions may contribute to the osmotic adjustment to continue water uptake in *S. hieraciifolia* under excess salinity.

Our findings showed that  $K^+$  content increased in all salt treatments, and the increases were high in the 150 and 300 mM NaCl treated shoots. Levels of  $Na^+$  and  $Cl^-$  ions also increased parallel with the increase in NaCl concentrations in the shoots. Moreover, we observed 3.3-fold increases in both  $Ca^{2+}$  and  $Mg^{2+}$  contents of the 600 mM NaCl treated shoots (Table 3). The same increase rate pointed out that  $Mg^{2+}$  may have a secondary messenger function similar to  $Ca^{2+}$  and affect uptake of other inorganic ions under salt stress in plants. Magnesium also led to the increase in the osmotic potential in plant tissues to maintain water uptake from medium to the tissues (Ahmad et al., 2013). Therefore, in the present study, the possible signaling role of  $Mg^{2+}$  reported earlier in animals has been expressed for the first time in plants. Additionally, based on the increases in  $Na^+$ ,  $K^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$  and  $Cl^-$  contents, we can say that *S. hieraciifolia* can provide osmotic adjustment by using inorganic ions and induce water uptake under excess salinity.

Our results showed that increased salinity enhanced the amounts of superoxide. However, it was known that cellular antioxidant system could provide an important contribution to cellular redox homeostasis through the activation of non-enzymatic or enzymatic antioxidants in plants subjected to salt stress (Gupta and Huang, 2014). In our study, there were 2.2-fold and 8-fold increases in ASC content of the shoots treated with 300 mM and 600 mM NaCl, respectively. We observed that like ASC, GSH content increased by salt stress and there were the 3.4-fold and 10-fold increases in the 300 mM and 600 mM NaCl treatments, respectively. Coherently with our results, Yildiztugay et al. (2014) reported that ROS level and total ASC and total GSH contents increased in extreme halophyte *Salsola crassa*. Therefore, the antioxidant signal compounds ASC and GSH could activate the signaling network to stimulate the antioxidant system responses to provide salt tolerance to *S. hieraciifolia*.

Antioxidant enzymes such as SOD activity was increased under salt stress conditions in *S. hieraciifolia*. There were 1.6-fold and 2.4-fold enhancements in SOD activity of the shoots treated with 300 mM and 600 mM NaCl, respectively. Increase in SOD activity was reported in *T. parvula* during the salt stress (Uzilday et al., 2015). CAT activity enhanced in the shoots treated with 150 mM and 300 mM NaCl although the activity declined under high salt stress conditions. Similar to our results, Parida and Jha (2010) reported that CAT activity decreased under high salinity (400 mM and 600 mM NaCl) in succulent extreme-halophyte *Salicornia brachiata*. Moreover, GPOD activity increased by 1.7-fold at 300 mM NaCl and 4-fold at 600 mM NaCl in *S. hieraciifolia*. The activity of GR playing an important role in controlling endogenous  $H_2O_2$  content through a redox cycle containing glutathione and

ascorbate (Gupta and Huang, 2014), increased under salt stress in *S. hieraciifolia*. There were 1.3-fold and 2.5-fold enhancements in the GR activities in the 300 mM and 600 mM NaCl treatments, respectively. Also, APX activity showed significant stimulation in all treatments of salinity and, the activity increased by 1.3-fold and 1.5-fold at 300 mM and 600 mM NaCl, respectively. In the light of these findings, we can say that succulent extreme-halophyte, *S. hieraciifolia* can improve their salt tolerance capacity through the powerful antioxidant enzyme system.

We also detected here the changes in the protein contents of some antioxidant enzymes. As known, in higher plants, there are three major families of SOD according to the protein fold and the metal cofactor: manganese SOD (Mn-SOD), iron SOD (Fe-SOD), and copper/zinc SOD (Cu/Zn-SOD) (Wang et al., 2016). Fe-SOD and GPX protein contents increased in the 300 mM NaCl treatments in comparison with the un-stressed treatment and the highest increases were observed in the 600 mM NaCl treatment. CAT protein content was high in the 300 mM NaCl treatment. Also, GR protein content enhanced in all salt treatments except 600 mM NaCl treatment. There could be essential protective roles of antioxidant enzyme contents in conjunction with their activities in the scavenging processes. The results point out the fact that the antioxidant enzyme system act as a defense arsenal for building an adaptive mechanism under high salinity in succulent extreme-halophyte, *S. hieraciifolia*.

## 5. Conclusion

Our results showed that succulent *Scorzonera hieraciifolia* with antioxidant and anti-inflammatory activity may

be an extreme salt tolerant that could survive at excess salinity (600 mM NaCl) conditions. *S. hieraciifolia* displays optimal growth at 300 mM NaCl in the MS media. *S. hieraciifolia*, can display shoot succulence by up-taking water effectively. The combined accumulations of signal compounds involved in osmoregulation such as proline, total soluble sugar, calcium and magnesium can provide the exceptional salt tolerance to *S. hieraciifolia*. Also, ASC can contribute activation of ROS and Ca<sup>2+</sup> signaling under salt stress in succulent extreme-halophyte *Scorzonera hieraciifolia*. The modified accumulation of ASC, GSH and other signaling compounds may trigger the propagation of salt tolerance mechanism. Therefore, the exceptional salt tolerance of succulent extreme-halophyte *S. hieraciifolia* is achieved by regulated accumulation of combined signal compounds involved in osmoregulation and induction of antioxidant system. Concomitant increases in antioxidant enzyme activities, antioxidant enzyme contents and the shoot succulence in the salinized plants showed that *S. hieraciifolia* might use the comprehensive strategy when coping with salinity.

## Acknowledgments

This work was supported by the Research Unit of Karadeniz Technical University (Project No: FDK-2018-7712). We also thank Prof. Dr. İsmail TURKAN for providing us with the seeds of *Thellungiella parvula*.

## Author contributions

CA performed the all experiments and wrote the manuscript; RT analyzed all data, read and edited the manuscript.

## References

- Aghaleh M, Niknam V, Ebrahimzadeh H, Razavi K (2009). Salt stress effects on growth, pigments, proteins and lipid peroxidation in *Salicornia persica* and *S. europaea*. *Biologia Plantarum* 53 (2): 243-248. doi: 10.1007/s10535-009-0046-7
- Ahmad ST, Sima NAKK, Mirzaei HH (2013). Effects of sodium chloride on physiological aspects of *Salicornia persica* growth. *Journal of Plant Nutrition* 36 (3): 401-414. doi: 10.1080/01904167.2012.746366
- Akyol TY, Yılmaz O, Uzılday B, Uzılday RÖ, Türkan İ (2020). Plant response to salinity: an analysis of ROS formation, signaling, and antioxidant defense. *Turkish Journal of Botany* 44 (1): 1-13. doi: 10.3906/bot-1911-15
- Altuntaş C, Sezgin A, Demiralay M, Terzi R, Sağlam A et al. (2019). Application of sucrose modulates the expressions of genes involved in proline and polyamine metabolism in maize seedlings exposed to drought. *Biologia Plantarum* 63: 247-252. doi: 10.32615/bp.2019.028
- Arnon DI (1949). Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* 24 (1): 1. doi: 10.1104/pp.24.1.1
- Bates LS, Waldren RP, Teare ID (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil* 39 (1): 205-207. doi: 10.1007/bf00018060
- Beauchamp C, Fridovich I (1971). Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry* 44 (1): 276-287. doi: 10.1016/0003-2697(71)90370-8
- Bergmeyer HU, Graßl M (1983). *Methods of Enzymatic Analysis*, Vol. 2. 3rd ed. Weinheim, Germany: Verlag Chemie, pp. 267-268.
- Boughalleb F, Abdellaoui R, Mahmoudi M, Bakhshandeh E (2020). Changes in phenolic profile, soluble sugar, proline, and antioxidant enzyme activities of *Polygonum equisetiforme* in response to salinity. *Turkish Journal of Botany* 44 (1): 25-35. doi: 10.3906/bot-1908-2



- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72 (1-2): 248-254. doi: 10.1016/0003-2697(76)90527-3
- Castillo FJ (1996). Antioxidative protection in the inducible CAM plant *Sedum album* L. following the imposition of severe water stress and recovery. *Oecologia* 107 (4): 469-477. doi: 10.1007/BF00333937
- Chen H, Jiang JG (2010). Osmotic adjustment and plant adaptation to environmental changes related to drought and salinity. *Environmental Reviews* 18 (NA): 309-319. doi: 10.1139/A10-014
- Daudi A, Cheng Z, O'Brien JA, Mammarella N, Khan S et al. (2012). The apoplastic oxidative burst peroxidase in *Arabidopsis* is a major component of pattern-triggered immunity. *The Plant Cell* 24 (1): 275-287. doi: 10.1105/tpc.111.093039
- Dubois M, Gilles KA, Hamilton JK, Rebers PT, Smith F (1956). Colorimetric method for determination of sugars and related substances. *Analytical Chemistry* 28 (3): 350-356. doi: 10.1021/ac60111a017
- Foyer CH, Halliwell B (1976). The presence of glutathione and glutathione reductase in chloroplasts: a proposed role in ascorbic acid metabolism. *Planta* 133 (1): 21-25. doi: 10.1007/BF00386001
- Ghanta S, Chattopadhyay S (2011). Glutathione as a signaling molecule: another challenge to pathogens. *Plant Signaling and Behavior* 6 (6): 783-788. doi: 10.4161/psb.6.6.15147
- Gupta B, Huang B (2014). Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. *International Journal of Genomics* doi: <http://dx.doi.org/10.1155/2014/701596>
- Haddad PR (1997). Comparison of ion chromatography and capillary electrophoresis for the determination of inorganic ions. *Journal of Chromatography A* 770 (1-2): 281-290. doi: 10.1016/S0021-9673(96)01085-0
- Hayat S, Hayat Q, Alyemeni MN, Wani AS, Pichtel J et al. (2012). Role of proline under changing environments: a review. *Plant Signaling Behavior* 7 (11): 1456-1466. doi: 10.4161/psb.21949
- Heath RL, Packer L (1968). Photoperoxidation in isolated chloroplasts: I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics* 125 (1): 189-198. doi: 10.1016/0003-9861(68)90654-1
- Inan G, Zhang Q, Li P, Wang Z, Cao Z et al. (2004). Salt cress. A halophyte and cryophyte *Arabidopsis* relative model system and its applicability to molecular genetic analyses of growth and development of extremophiles. *Plant Physiology* 135 (3): 1718-1737. doi: 10.1104/pp.104.041723
- Liang W, Ma X, Wan P, Liu L (2018). Plant salt-tolerance mechanism: A review. *Biochemical and Biophysical Research Communications* 495 (1): 286-291. doi: 10.1016/j.bbrc.2017.11.043
- Liso R, Calabrese G, Bitonti MB, Arrigoni O (1984). Relationship between ascorbic acid and cell division. *Experimental Cell Research* 150 (2): 314-320. doi: 10.1016/0014-4827(84)90574-3
- Makavitskaya M, Svistunenko D, Navaselsky I, Hryvusevich P, Mackievic V et al. (2018). Novel roles of ascorbate in plants: induction of cytosolic Ca<sup>2+</sup> signals and efflux from cells via anion channels. *Journal of Experimental Botany* 69 (14): 3477-3489. doi: 10.1093/jxb/ery056
- Mangalassery S, Dayal D, Kumar A, Bhatt K, Nakar R et al. (2017). Pattern of salt accumulation and its impact on salinity tolerance in two halophyte grasses in extreme saline desert in India. *Indian Journal of Experimental Biology* 55 (8): 542-548.
- Mishra A, Tanna B (2017). Halophytes: potential resources for salt stress tolerance genes and promoters. *Frontiers in Plant Science* 8: 829. doi: 10.3389/fpls.2017.00829
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum* 15 (3): 473-497. doi: 10.1111/j.1399-3054.1962.tb08052.x
- Nakano Y, Asada K (1987). Purification of ascorbate peroxidase in spinach chloroplasts; its inactivation in ascorbate-depleted medium and reactivation by monodehydroascorbate radical. *Plant and Cell Physiology* 28 (1): 131-140. doi: 10.1093/oxfordjournals.pcp.a077268.
- Parida AK, Jha B (2010). Antioxidative defense potential to salinity in the euhalophyte *Salicornia brachiata*. *Journal of Plant Growth Regulation* 29 (2): 137-148. doi: 10.1007/s00344-009-9129-0
- Parida AK, Veerabathini SK, Kumari A, Agarwal PK (2016). Physiological, anatomical and metabolic implications of salt tolerance in the halophyte *Salvadora persica* under hydroponic culture condition. *Frontiers in Plant Science* 7: 351. doi: 10.3389/fpls.2016.00357
- Rubnawaz S, Kayani W, Mahmood R, Mirza B (2020). Enhanced stress tolerance in transformed *Ajuga bracteosa* Wall. ex Benth. regenerants by upregulated gene expression of metabolic pathways. *Turkish Journal of Botany* 44 (4): 410-426. doi: 10.3906/bot-2002-13
- Qi CH, Chen M, Song J, Wang BS (2009). Increase in aquaporin activity is involved in leaf succulence of the euhalophyte *Suaeda salsa*, under salinity. *Plant Science* 176 (2): 200-205. doi: 10.1016/j.plantsci.2008.09.019
- Sarı A, Şahin H, Özsoy N, Çelik B Ö (2019). Phenolic compounds and in vitro antioxidant, anti-inflammatory, antimicrobial activities of *Scorzonera hieraciifolia* Hayek roots. *South African Journal of Botany* 125: 116-119. doi: 10.1016/j.sajb.2019.07.009
- Schopfer P, Plachy C, Frahry G (2001). Release of reactive oxygen intermediates (superoxide radicals, hydrogen peroxide, and hydroxyl radicals) and peroxidase in germinating radish seeds controlled by light, gibberellin, and abscisic acid. *Plant Physiology* 125 (4): 1591-1602. doi: 10.1104/pp.125.4.1591
- Shabala S, Mackay A (2011). Ion transport in halophytes. *Advances in Botanical Research* 57: 151-199. doi: 10.1016/B978-0-12-387692-8.00005-9
- Sharma P, Dubey RS (2005). Drought induces oxidative stress and enhances the activities of antioxidant enzymes in growing rice seedlings. *Plant Growth Regulation* 46 (3): 209-221. doi: 10.1007/s10725-005-0002-2

- Shin SY, Kim MH, Kim YH, Park HM, Yoon HS (2013). Co-expression of monodehydroascorbate reductase and dehydroascorbate reductase from *Brassica rapa* effectively confers tolerance to freezing-induced oxidative stress. *Molecules and Cells* 36 (4): 304-315. doi: 10.1007/s10059-013-0071-4
- Stangherlin A, O'Neill JS (2018). Signal transduction: magnesium manifests as a second messenger. *Current Biology* 28 (24): R1403-R1405. doi: 10.1016/j.cub.2018.11.003
- Urbanek H, Kuzniak-Gebarowska E, Herka K (1991). Elicitation of defense responses in bean leaves by *Botrytis cinerea* polygalacturonase. *Acta Physiologiae Plantarum* 13 (1): 43-50.
- Uzilday B, Ozgur R, Sekmen AH, Yildiztugay E, Turkan I (2015). Changes in the alternative electron sinks and antioxidant defence in chloroplasts of the extreme halophyte *Eutrema parvulum* (*Thellungiella parvula*) under salinity. *Annals of Botany* 115 (3): 449-463. doi: 10.1093/aob/mcu184
- Velikova V, Yordanov I, Edreva A (2000). Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Science* 151 (1): 59-66. doi: 10.1016/S0168-9452(99)00197-1
- Wang M, Zhao X, Xiao Z, Yin X, Xing T et al. (2016). A wheat superoxide dismutase gene *TaSOD2* enhances salt resistance through modulating redox homeostasis by promoting NADPH oxidase activity. *Plant Molecular Biology* 91 (1-2): 115-130. doi: 10.1007/s11103-016-0446-y
- Xin Z, Browse J (2000). Cold comfort farm: the acclimation of plants to freezing temperatures. *Plant, Cell and Environment* 23 (9): 893-902. doi: 10.1046/j.1365-3040.2000.00611.x
- Yildiztugay E, Ozfidan-Konakci C, Kucukoduk M (2014). The role of antioxidant responses on the tolerance range of extreme halophyte *Salsola crassa* grown under toxic salt concentrations. *Ecotoxicology and Environmental Safety* 110: 21-30. doi: 10.1016/j.ecoenv.2014.08.013
- Zeng F, Shabala S, Maksimović JD, Maksimović V, Bonales-Alatorre E et al. (2018). Revealing mechanisms of salinity tissue tolerance in succulent halophytes: a case study for *Carpobrotus rossi*. *Plant, Cell and Environment* 41 (11): 2654-2667. doi: 10.1111/pce.13391
- Zhou Q, Yu BJ (2009). Accumulation of inorganic and organic osmolytes and their role in osmotic adjustment in NaCl-stressed vetiver grass seedlings. *Russian Journal of Plant Physiology* 56 (5): 678-685. doi: 10.1134/S1021443709050148