

1-1-2021

Hermitian-Toeplitz determinants for functions with bounded turning

VIRENDRA KUMAR

NAK EUN CHO

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

KUMAR, VIRENDRA and CHO, NAK EUN (2021) "Hermitian-Toeplitz determinants for functions with bounded turning," *Turkish Journal of Mathematics*: Vol. 45: No. 6, Article 20. <https://doi.org/10.3906/mat-2101-104>

Available at: <https://journals.tubitak.gov.tr/math/vol45/iss6/20>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Hermitian–Toeplitz determinants for functions with bounded turning

Virendra KUMAR¹ , Nak Eun CHO^{2,*} 

¹Department of Mathematics, Ramanujan College, University of Delhi, Kalkaji, New Delhi–110019, India

²Department of Applied Mathematics, Pukyong National University, Busan–48513, South Korea

Received: 25.01.2021

Accepted/Published Online: 20.10.2021

Final Version: 29.11.2021

Abstract: There is a rich literature on estimation of second and third Hankel determinants for normalised analytic functions in geometric function theory. It is also, therefore, natural to explore the concept of the Hermitian–Toeplitz determinants for such functions. In this paper, the sharp lower and upper estimations for third-order Hermitian–Toeplitz determinant for functions with bounded turning of order α , are obtained.

Key words: Analytic functions, functions with bounded turning of order α , Hermitian–Toeplitz determinant

1. Introduction

Finding the sharp estimates on the coefficient's functionals has been one of the major research area of geometric function theory since the advent of the Bieberbach conjecture for normalised univalent functions and then theory of the univalent functions developed around this conjecture. Later, among the coefficient functionals, the major area have been the estimation of bound on the Fekete–Szegő functional(1933) and the Hankel determinants. Although the estimation of bound on Hankel determinant started during 1960's, since the Bieberbach conjecture was unsolved, not so many researchers took interest in investigating the bound on the Hankel determinants, except a few articles [17, 28]. In the last few years investigation of the Hankel determinant gained much attention and brief survey of those work until 2013 can be found in the introduction of the paper [24]. Much recent history of development in this direction can be found in [1, 3, 4, 7, 9, 13, 16, 26, 31]. Wide variety of applications of Toeplitz–plus–Hankel systems arise in linear filtering theory, discrete inverse scattering, and discretization of certain integral equations arising in mathematical physics [30]. The paper [6] gives the higher-order asymptotic formulas for the eigenvalues of large Hermitian–Toeplitz matrices with moderately smooth symbols which trace out a simple loop on the real line and related applications in physics. The determinant of Hermitian–Toeplitz matrices finds its applications in signal processing [29], see also [20].

The class of normalised analytic functions of the form $f(z) = z + a_2z^2 + a_3z^3 + a_4z^4 + \dots$ defined on the open unit disk $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ is represented by the symbol \mathcal{A} . The collection of functions in \mathcal{A} which are univalent also is denoted by \mathcal{S} . Recently, Ali et al. [2] introduced symmetric Toeplitz determinant $T_{q,n}(f)$

*Correspondence: necho@pknu.ac.kr

2010 *AMS Mathematics Subject Classification*: 30C45, 30C50

defined by

$$T_{q,n}(f) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & a_n & \cdots & a_{n+q-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n+q-1} & a_{n+q-2} & \cdots & a_n \end{vmatrix}$$

and estimated the bounds on $T_{2,n}(f), T_{3,1}(f), T_{3,2}(f)$ and $T_{2,3}(f)$ for certain subclasses of analytic functions. Actuated by above work, recently Cudna et al. [12] considered Hermitian–Toeplitz determinants with its entries as coefficients of a normalised analytic function as follows:

$$H_{q,n}^T(f) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ \bar{a}_{n+1} & a_n & \cdots & a_{n+q-2} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{a}_{n+q-1} & \bar{a}_{n+q-2} & \cdots & a_n \end{vmatrix}, \text{ where } a_n \in \mathbb{C}.$$

Further, they investigated the sharp lower and upper bound for third order Hermitian–Toeplitz determinants for the classes of starlike and convex functions of order α ($0 \leq \alpha < 1$). It should be noted that the determinant $H_{q,n}^T(f)$ is rationally invariant. Moreover, if a_n are real, then $H_{q,n}^T(f)$ is Hermitian and therefore, the determinant $H_{q,n}^T(f)$ is a real number, see [12]. From the above definition, it is easy to verify that $H_{2,1}^T(f) = 1 - |a_2|^2$ and

$$H_{3,1}^T(f) = \begin{vmatrix} 1 & a_2 & a_3 \\ \bar{a}_2 & 1 & a_2 \\ \bar{a}_3 & \bar{a}_2 & 1 \end{vmatrix} = 2 \operatorname{Re}(a_2^2 \bar{a}_3) - 2|a_2|^2 - |a_3|^2 + 1.$$

Kumar et al. [22] gave a generalisation to the results investigated in [12] by investigating those results for Janowski starlike and convex functions. Kumar [20] investigated lower and upper bounds on the second and third order Hermitian–Toeplitz determinants for certain subclasses of close-to-convex functions. Some more results in this direction may be found in [10, 21]. The book [5] edited by Bättcher and Grudsky contains a chapter on Toeplitz matrices which describes the condition on a Toeplitz matrix with entries from a given sequence of complex numbers under which the Toeplitz matrix induces a bounded operator on space certain space. For more details on Toeplitz matrices one refer to [8, 23]. A review on Toeplitz matrices is also available in [15].

Now we consider the class of analytic functions f whose derivative have positive real part of order α ($0 \leq \alpha < 1$) i.e. $\operatorname{Re} f'(z) > \alpha$ ($z \in \mathbb{D}$). The collection of such functions is denoted by $\mathcal{R}(\alpha)$. The class is important in the sense that the functions in this class are univalent. More precisely, the condition $\operatorname{Re} f'(z) > 0$ ($z \in \mathbb{D}$) gives an important sufficient condition for univalence of normalised analytic functions, see [14]. The p -valent analogue of this class was considered in [18, 19] and investigated the bounds on second and third Hankel determinants. A generalisation of their work was done by Cho et al. [11]. The work reported in the papers [12] and [2] inculcates us to estimate sharp lower and upper bounds on the third-order Hermitian–Toeplitz determinant for functions in the class $\mathcal{R}(\alpha)$.

Now we introduce the class of functions with positive real part which is going to be an important tool for calculating the bounds. Let $\mathcal{P}(\alpha)$ denote the class of analytic functions of the form $p(z) = 1 + p_1z + p_2z^2 +$

$p_3z^3 + \dots$ with $p(0) = 1$ and $\operatorname{Re} p(z) > \alpha$ ($0 \leq \alpha < 1$). It is easy to see that $f \in \mathcal{R}(\alpha)$ if and only if $p \in \mathcal{P}(\alpha)$. Also, let $\mathcal{P}(0) =: \mathcal{P}$.

Lemma 1.1 [25] *If $p(z) = 1 + p_1z + p_2z^2 + p_3z^3 + \dots \in \mathcal{P}$, then*

$$2p_2 = p_1^2 + (4 - p_1^2)\zeta, \quad p_1 \geq 0 \tag{1.1}$$

for some $\zeta \in \mathbb{C}$ such that $|\zeta| \leq 1$.

2. Third order Hermitian–Toeplitz determinant

It is well-known that for $f(z) = z + a_2z^2 + a_3z^3 + \dots \in \mathcal{R}(\alpha)$, the sharp bound $|a_2| \leq 1 - \alpha$ holds. Now the sharp upper bound for $H_{2,1}^T(f) = 1 - |a_2|^2$ is naturally 1, however, the sharp lower bound is $1 - (1 - \alpha)^2$ i.e. $\alpha(2 - \alpha)$. The equality in the upper bound holds for the function

$$\tilde{f}_0(z) = z. \tag{2.1}$$

Equality in lower bound holds in case of the function

$$\tilde{f}_1(z) = \int_0^z \left((1 - \alpha) \frac{1 - t}{1 + t} + \alpha \right) dt = z - (1 - \alpha)z^2 + \frac{2(1 - \alpha)}{3}z^3 + \dots \tag{2.2}$$

We state this result as a theorem:

Theorem 2.1 *Let $f(z) = z + a_2z^2 + a_3z^3 + \dots \in \mathcal{R}(\alpha)$. Then the following sharp estimations hold:*

$$(2 - \alpha)\alpha \leq H_{2,1}^T(f) \leq 1.$$

Example 2.2 *For the function $\tilde{f}_0(z) = z, z \in \mathbb{D}$, we have $a_2 = a_3 = a_4 = 0$ and clearly $f \in \mathcal{R}$. Thus we have $H_{2,1}^T(f) = 1$. The function \tilde{f}_1 is also an example in the class $\mathcal{R}(\alpha)$ which gives $H_{2,1}^T(f) = \alpha(2 - \alpha)$ which is the lower bound in case of Theorem 2.1.*

We now investigate the sharp lower and upper bound on third order Hermitian–Toeplitz determinant.

Theorem 2.3 *Let $f(z) = z + a_2z^2 + a_3z^3 + \dots \in \mathcal{R}(\alpha)$. Then the following sharp estimations hold:*

$$1 \geq H_{3,1}^T(f) \geq \begin{cases} \frac{(2\alpha^2 - 7\alpha + 1)^2}{8(3\alpha - 1)}, & 0 \leq \alpha < \frac{1}{18}; \\ -\frac{1}{9}(2\alpha + 1)(6\alpha^2 - 10\alpha + 1), & \frac{1}{18} \leq \alpha < 1. \end{cases}$$

Proof Since $f(z) = z + a_2z^2 + a_3z^3 + \dots \in \mathcal{R}(\alpha)$, it follows that there exists $p(z) = 1 + p_1z + p_2z^2 + p_3z^3 + \dots \in \mathcal{P}$, such that

$$f'(z) = (1 - \alpha)p(z) + \alpha. \tag{2.3}$$

Comparing the coefficients of similar power terms in (2.3), we have

$$a_2 = \frac{(1 - \alpha)p_1}{2} \quad \text{and} \quad a_3 = \frac{(1 - \alpha)p_2}{3}. \tag{2.4}$$

With the aid of (2.4), we have

$$\begin{aligned} H_{3,1}^T(f) &= 1 + 2 \operatorname{Re}(a_2^2 a_3) - 2|a_2|^2 - |a_3|^2 \\ &= 1 + \frac{1}{6}(1 - \alpha)^3 \operatorname{Re} p_1^2 \bar{p}_2 - \frac{1}{2}(1 - \alpha)^2 |p_1|^2 - \frac{1}{9}(1 - \alpha)^2 |p_2|^2. \end{aligned}$$

It is well-known that the class $\mathcal{P}(\alpha)$ and $\mathcal{R}(\alpha)$ are rotationally invariant. For this reason, we shall limit ourselves to a consideration of nonnegative value of p_1 . Also since $|p_1| \leq 2$, we can very much assume $0 \leq p_1 \leq 2$ and let $p_1^2 =: x$, and so $x \in [0, 4]$. Also throughout this proof we also use the notation $y =: |\zeta| \in [0, 1]$ wherever be needed. Now replacing p_2 with its equivalent expression using (1.1) from Lemma 1.1 and simplifying, we get

$$\begin{aligned} H_{3,1}^T(f) &= 1 + \frac{(1 - \alpha)^3}{12} p_1^4 - \frac{p_1^2}{2} (1 - \alpha)^2 + \frac{(1 - \alpha)^3}{12} p_1^2 (4 - p_1^2) \operatorname{Re} \zeta \\ &\quad - \frac{(1 - \alpha)^2}{36} (|p_1|^4 + (4 - p_1^2)|\zeta|^2 + 2p_1^2(4 - p_1^2) \operatorname{Re} \zeta). \end{aligned} \tag{2.5}$$

In view of this fact and assumptions, we can write (2.5), as

$$\begin{aligned} H_{3,1}^T(f) &= 1 + \frac{(2 - 3\alpha)(1 - \alpha)^2}{36} x^2 - \frac{(1 - \alpha)^2}{2} x - \frac{(1 - \alpha)^2(4 - x)^2}{36} |\zeta|^2 + \frac{(1 - \alpha)^2(1 - 3\alpha)(4 - x)x}{36} \operatorname{Re} \zeta \\ &=: F(\alpha, x, |\zeta|, \operatorname{Re} \zeta). \end{aligned} \tag{2.6}$$

We now proceed further in the proof through several steps.

Case I(a): For $\alpha = 0$, from (2.6) we have $F(0, x, |\zeta|, \operatorname{Re} \zeta) \leq F(0, x, |\zeta|, |\zeta|) =: \Upsilon(x, y)$, where

$$\Upsilon(x, y) = 1 + \frac{x^2}{18} - \frac{x}{2} - \frac{(4 - x)^2}{36} y^2 + \frac{(4 - x)x}{36} y, \quad (x, y) \in [0, 1] \times [0, 1]. \tag{2.7}$$

On the boundary line segments of the rectangular region $[0, 4] \times [0, 1]$, we have $\Upsilon(0, y) = 1 - 4y/9 \leq 1$, $\Upsilon(4, y) = -1/9$, $y \in (0, 1)$. Since $\Upsilon'(x, 0) = (2x - 9)/18 < 0$, it follows that

$$\Upsilon(x, 0) = \frac{x^2}{18} - \frac{x}{2} + 1 \leq \Upsilon(0, 0) = 1, \quad \forall x \in (0, 4).$$

Further when $y = 1$, we find that $F(x, 1) = \frac{5}{9} - \frac{x}{6} \leq \frac{5}{9}$, $\forall x \in (0, 4)$. It is matter of simple calculation now to verify that the function Υ has no critical point in the domain $(0, 4) \times (0, 1)$. Therefore, keeping the conclusions of the above discussion together, we arrive at

$$H_{3,1}^T(f) \leq \max \left\{ 1, -\frac{1}{9}, \frac{5}{9} \right\} = 1. \tag{2.8}$$

Now we find the minimum in the case $\alpha = 0$. From (2.7), we have

$$\begin{aligned} H_{3,1}^T(f) &= F(0, x, |\zeta|, \operatorname{Re} \zeta) \\ &\geq F(x, 1, -1) = \frac{1}{18} (x^2 - 7x + 10) \\ &\geq -\frac{1}{8}. \end{aligned}$$

Therefore,

$$H_{3,1}^T(f) \geq -\frac{1}{8}. \tag{2.9}$$

Thus, for $\alpha = 0$, from (2.8) and (2.9), we have

$$-\frac{1}{8} \leq |H_{3,1}^T(f)| \leq 1. \tag{2.10}$$

The upper bound is sharp in case of the function \tilde{f}_0 given in (2.1), whereas the equality in lower bound holds for the function f_2 given by

$$\tilde{f}_2(z) = \int_0^z \frac{1-t^2}{1-2\sqrt{7/8}t+t^2} dt = z + \frac{1}{2}\sqrt{\frac{7}{2}}z^2 + \frac{1}{2}z^3 + \frac{1}{8}\sqrt{\frac{7}{2}}z^4 + \frac{1}{20}z^5 + \dots \quad (z \in \mathbb{D}). \tag{2.11}$$

Case I(b) For $\alpha = 1/3$, we have

$$\begin{aligned} F(1/3, x, |\zeta|, \operatorname{Re} \zeta) &= 1 + \frac{1}{81}x^2 - \frac{2}{9}x - \frac{1}{81}(4-x)^2|\zeta|^2 \\ &\leq \frac{1}{81}(x-9)^2 \\ &\leq 1. \end{aligned}$$

For minimum, consider

$$\begin{aligned} F(1/3, x, |\zeta|, \operatorname{Re} \zeta) &= 1 + \frac{1}{18}x^2 - \frac{2}{9}x - \frac{1}{81}(4-x)^2|\zeta|^2 \\ &\geq \frac{5}{81}(13-2x) \\ &\geq \frac{25}{81}. \end{aligned}$$

Thus, we have, for $\alpha = 1/3$:

$$\frac{25}{81} \leq H_{3,1}^T(f) \leq 1. \tag{2.12}$$

The lower bound equals in case of the function \tilde{f}_1 defined in (2.2). The upper bound is sharp in case of the function \tilde{f}_0 defined in (2.1).

Case (II): If $0 < \alpha < 1/3$, then $H_{3,1}^T(f) = F(\alpha, x, |\zeta|, \operatorname{Re} \zeta) \leq F(\alpha, x, |\zeta|, |\zeta|) = G(x, y)$, where

$$G(x, y) := 1 + \frac{1}{36}(2-3\alpha)(1-\alpha)^2x^2 + \frac{1}{36}(1-3\alpha)(1-\alpha)^2(4-x)xy - \frac{1}{36}(1-\alpha)^2(4-x)^2y^2 - \frac{1}{2}(1-\alpha)^2x.$$

On the boundary line segments of the rectangular region $[0, 4] \times [0, 1]$, we have

$$\begin{aligned} G(0, y) &= 1 - \frac{4}{9}(1-\alpha)^2y^2 \leq 1, \forall y \in [0, 1] \\ G(4, y) &= \frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1), \forall y \in [0, 1] \\ G(x, 0) &= \frac{1}{36}(2-3\alpha)(1-\alpha)^2x^2 - \frac{1}{2}(1-\alpha)^2x + 1. \end{aligned}$$

Now computation reveals that $G(x, 0)$ has no maximum in $(0, 4)$ and $G(4, 0) = (-12\alpha^3 + 14\alpha^2 + 8\alpha - 1) / 9$. Further, as before, it can be verified that the function

$$G(x, 1) = -\frac{1}{18}(2\alpha + 1) (3\alpha^2x + \alpha(4 - 6x) + 3x - 10)$$

has no critical point inside $(0, 4)$, and $G(0, 1) = -(2\alpha - 5)(2\alpha + 1)/9$. Consider the case when the function G is defined inside the rectangular region $(x, y) \in (0, 4) \times (0, 1)$. It is a matter of routine calculation to see that the function G has no maximum inside this region. For $\alpha \in (0, 1/3)$, on the basis of above discussion, we have

$$H_{3,1}^T(f) \leq \max \left\{ 1, \frac{1}{9} (-12\alpha^3 + 14\alpha^2 + 8\alpha - 1) \right\} = 1.$$

The equality attained for the function defined by (2.2).

We now find the minimum of $H_{3,1}^T(f)$ for the case $0 < \alpha < 1/3$. In this case, we have $H_{3,1}^T(f) = F(\alpha, x, |\zeta|, \operatorname{Re} \zeta) \geq F(\alpha, x, |\zeta|, -|\zeta|) \geq F(\alpha, x, 1, -1) =: h(x)$. Here we see that

$$h'(x) = \frac{1}{18}(a - 1)^2(6a - 7) - \frac{1}{9}(a - 1)^2(3a - 1)x = 0$$

if and only if

$$x = x_2 = \frac{6\alpha - 7}{2(3\alpha - 1)}.$$

We now consider two subcases, namely (a) $0 < \alpha < 1/18$; and (b) $1/18 \leq \alpha < 1/3$.

(a) It is easy to check that $x_2 \in (0, 4)$ for $0 < \alpha < 1/18$. Further, the values of h at the critical point is given by

$$h(x_2) = \frac{(2\alpha^2 - 7\alpha + 1)^2}{8(3\alpha - 1)}, \quad 0 < \alpha < \frac{1}{18}.$$

Also at the end points of the line segment $0 \leq x \leq 4$, we have

$$h(0) = 1 - \frac{4}{9}(1 - \alpha)^2 \quad \text{and} \quad h(4) = \frac{1}{9} (-12\alpha^3 + 14\alpha^2 + 8\alpha - 1).$$

From the above, for $0 < \alpha < 1/18$, we conclude that

$$H_{3,1}^T(f) \geq \min\{h(0), h(4), h(x_2)\} = h(x_2).$$

Thus, for $0 < \alpha < 1/18$, we have

$$\frac{(2\alpha^2 - 7\alpha + 1)^2}{8(3\alpha - 1)} \leq H_{3,1}^T(f) \leq 1. \tag{2.13}$$

The equality for the lower bound is attained for the function

$$\begin{aligned} \tilde{f}_5(z) &= \int_0^z \left((1 - \alpha) \frac{1 - t^2}{1 - u_0 t + t^2} + \alpha \right) dt, \quad \text{where } u_0 = \sqrt{\frac{6\alpha - 7}{2(3\alpha - 1)}} \\ &= z + \frac{1 - \alpha}{2\sqrt{2}} \sqrt{\frac{6\alpha - 7}{3\alpha - 1}} z^2 + \frac{(\alpha - 1)(2\alpha + 1)}{2(3\alpha - 1)} z^3 + \dots \end{aligned}$$

(b) For $1/18 \leq \alpha < 1/3$, we find that h has no critical point in $(0, 4)$, hence the minimum will be attained at the end points. From the above, for $1/18 \leq \alpha < 1/3$, we conclude that

$$\begin{aligned} H_{3,1}^T(f) &\geq \min \left\{ 1 - \frac{4}{9}(1 - \alpha)^2, \frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1) \right\} \\ &= \frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1). \end{aligned}$$

Thus, for $1/18 \leq \alpha < 1/3$, we have

$$\frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1) \leq H_{3,1}^T(f) \leq 1. \tag{2.14}$$

The equality in the lower bound occurs for the function \tilde{f}_1 and that of upper bound for the function \tilde{f}_0 .

Case (III): Now consider the case $1/3 < \alpha < 1$. As before, we have $H_{3,1}^T(f) = F(\alpha, x, |\zeta|, \operatorname{Re} \zeta) \leq F(\alpha, x, |\zeta|, -|\zeta|) = H(x, y)$, where

$$H(x, y) := \frac{1}{36}(2 - 3\alpha)(1 - \alpha)^2 x^2 - \frac{1}{36}(1 - \alpha)^2(4 - x)^2 y^2 - \frac{1}{36}(1 - 3\alpha)(1 - \alpha)^2(4 - x)xy - \frac{1}{2}(1 - \alpha)^2 x + 1.$$

On the boundary line segment of the rectangular region $[0, 4] \times [0, 1]$, we have

$$H(0, y) = 1 - \frac{4}{9}(1 - \alpha)^2 y^2 \leq 1$$

$$H(4, y) = \frac{4}{9}(2 - 3\alpha)(1 - \alpha)^2 - 2(1 - \alpha)^2 + 1$$

$$H(x, 0) = \frac{1}{36}(2 - 3\alpha)(1 - \alpha)^2 x^2 - \frac{1}{2}(1 - \alpha)^2 x + 1$$

$$H(x, 1) = \frac{1}{36}(2 - 3\alpha)(1 - \alpha)^2 x^2 - \frac{1}{36}(1 - \alpha)^2(4 - x)^2 - \frac{1}{2}(1 - \alpha)^2 x - \frac{1}{36}(1 - 3\alpha)(1 - \alpha)^2(4 - x)x + 1.$$

The function $H(x, 0)$ is decreasing and for all $x \in (0, 4)$ and $1/3 < \alpha < 1$ and so $G(x, 0) \leq G(0, 0) = 1$. Moreover the function $H(x, 1)$ has no critical point in $(0, 4)$. A similar computations show that the function H has no critical point in $(x, y) \in (0, 4) \times (0, 1)$. Therefore, on the basis of above discussion, we see that

$$H_{3,1}^T(f) \leq \max \left\{ 1, \frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1), 1 - \frac{4}{9}(1 - \alpha)^2 \right\} = 1.$$

For $1/3 < \alpha < 1$ equality occurs in case of the function \tilde{f}_0 defined in (2.1).

We now proceed to find the minimum and for this we consider $H_{3,1}^T(f) = F(x, |\zeta|, \operatorname{Re} \zeta) \geq F(x, |\zeta|, |\zeta|) \geq F(x, 1, 1) = h_1(x)$. Now since $h_1'(x) = -(2\alpha + 1)(3\alpha^2 - 6\alpha + 3)/18 < 0$ ($1/3 < \alpha < 1$), it follows that the minimum attained at the end point $x = 4$ and $h_1(4) = -(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1)/9$. Therefore, we have

$$H_{3,1}^T(f) \geq \frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1)$$

with equality for the function \tilde{f}_1 defined in (2.2). Thus, in this case, we have

$$\frac{1}{9}(-12\alpha^3 + 14\alpha^2 + 8\alpha - 1) \leq H_{3,1}^T(f) \leq 1. \tag{2.15}$$

The estimations given in (2.10), (2.12), (2.13), (2.14) and (2.15) end the proof. \square

Example 2.4 For the function $\tilde{f}_0(z) = z$, $z \in \mathbb{D}$, we have $a_2 = a_3 = a_4 = 0$ and clearly $f \in \mathcal{R}$. Thus we have $H_{3,1}^T(f) = 1$, which is the upper bound as investigated in Theorem 2.3. The function \tilde{f}_2 given by (2.11) is also an example in the class \mathcal{R} which gives $H_{3,1}^T(f) = -1/8$, which is the lower bound in case of Theorem 2.3 in the case when $\alpha = 0$.

Conclusion

It should be noted that the class $\mathcal{R}(\alpha)$ is a subclass of \mathcal{S} , so it is interesting to compare the lower and upper bounds on Hermitian–Toeplitz determinants of second and third orders. Obradović and Tuneski [27] proved that $-3 \leq H_{2,1}^T(f) \leq 1$ and $-1 \leq H_{3,1}^T(f) \leq 8$. Here we find that the upper bound on $T_{2,1}(f)$ for both the classes are same whereas the lower bound for the class $\mathcal{R}(\alpha)$ is less than that of the class \mathcal{S} . If we compare the upper and lower bounds on $H_{3,1}^T(f)$, we find that the corresponding bounds in case of the class $\mathcal{R}(\alpha)$ lies entirely in the range $[-1, 8]$ of the bound for the class \mathcal{S} . Thus, the bounds obtained in this paper very much follow the expected outcomes which confirm the correctness of technicality and procedures adopted in proving our results. It would be interesting to find the lower and upper bounds for $H_{4,1}^T(f)$ for the class $\mathcal{R}(\alpha)$.

Acknowledgment

The authors would like to express their thanks to the editor and the anonymous referee for their valuable suggestions. The second author was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (No. 2019R1I1A3A01050861).

References

- [1] Ali RM. Coefficients of the inverse of strongly starlike functions. Bulletin of the Malaysian Mathematical Sciences Society 2003; 26 (1): 63-71.
- [2] Ali MF, Thomas DK, Vasudevarao A. Toeplitz determinants whose elements are the coefficients of analytic and univalent functions. Bulletin of the Australian Mathematical Society 2018; 97: 253-264.
- [3] Altınkaya Ş, Yalçın S. Upper bound of second Hankel determinant for bi-Bazilevič functions. Mediterranean Journal of Mathematics 2016; 13 (6): 4081-4090.
- [4] Arif M, Ullah I, Raza M, Zaprawa P. Investigation of the fifth Hankel determinant for a family of functions with bounded turnings. Mathematica Slovaca 2020; 70 (2): 319-328.
- [5] Böttcher A, Grudsky SM. Infinite Toeplitz matrices. In: Toeplitz matrices, asymptotic linear algebra and functional analysis. Texts and readings in mathematics (2000), Hindustan Book Agency, Gurgaon.
- [6] Bogoya JM, Böttcher A, Grudsky SM, Maximenko EA. Eigenvalues of Hermitian Toeplitz matrices with smooth simple-loop symbols. Journal of Mathematical Analysis and Applications 2015; 422(2): 1308-1334.
- [7] Bouras B. Hankel determinants of a linear combination of three successive Catalan numbers. Mediterranean Journal of Mathematics 2013; 10 (2): 693-705.
- [8] Chan NN, Kwong MK. Hermitian matrix inequalities and a conjecture, The American Mathematical Monthly 1985; 92 (8): 533-541.

- [9] Cho NE, Kumar V, Kwon OS, Sim YJ. Coefficient bounds for certain subclasses of starlike functions. *Journal of Inequalities and Applications* 2019; Paper No. 276: 13 pp.
- [10] Cho NE, Kumar S., Kumar V. Coefficient functionals for starlike functions associated with the modified sigmoid function and the Bell numbers, *Asian-European Journal of Mathematics* 2021. doi:10.1142/S1793557122500425.
- [11] Cho NE, Kumar V, Kwon OS, Sim YJ. Sharp coefficient bounds for certain p -valent functions. *Bulletin of the Malaysian Mathematical Sciences Society* 2019; 42: 405-416.
- [12] Cudna K, Kwon OS, Lecko A, Sim YJ, Smiarowska B. The second and third-order Hermitian-Toeplitz determinants for starlike and convex functions of order α . *Boletán de la Sociedad Matemática Mexicana* 2020; 26 (2): 361-375.
- [13] Deniz E, Budak L. Second Hankel determinat for certain analytic functions satisfying subordinate condition. *Mathematica Slovaca* 2018; 68 (2): 463-471.
- [14] Goodman AW. *Univalent functions, vol. I : Mariner, Tampa* 1983.
- [15] Gray RM. Toeplitz and circulant matrices: A Review, *Foundations and Trends in Communications and Information Theory* 2006; 2 (3): 155–239.
- [16] Kanas S, Analouei Adegani E, Zireh A. An unified approach to second Hankel determinant of bi-subordinate functions, *Mediterranean Journal of Mathematics* 2017; 14 (6): 12pp.
- [17] Hayman WK. On the second Hankel determinant of mean univalent functions. *Proceedings of the London Mathematical Society* 1968; 18 (3): 77-94.
- [18] Krishna DV, Ramreddy T. Coefficient inequality for multivalent bounded turning functions of order α . *Problemy Analiza-Issues of Analysis* 2016; 5 (1): 45-54.
- [19] Krishna DV, Venkateswarlu B, Ramreddy T. Third Hankel determinant for certain subclass of p -valent functions. *Complex Variables and Elliptic Equations* 2015; 60 (9): 1301-1307.
- [20] Kumar V. Hermitian-Toeplitz determinants for certain classes of close-to-convex functions. *Bulletin of the Iranian Mathematical Society* 2021. doi:10.1007/s41980-021-00564-0.
- [21] Kumar V, Kumar S. Bounds on Hermitian-Toeplitz and Hankel determinants for strongly starlike functions. *Boletán de la Sociedad Matemática Mexicana* 2021; 27 (55): 16pp. doi:10.1007/s40590-021-00362-y.
- [22] Kumar V, Srivastava R, Cho NE. Sharp estimation of Hermitian-Toeplitz determinants for Janowski type starlike and convex functions. *Miskolc Mathematical Notes* 2020; 21 (2): 939-952.
- [23] Kurt V, Simsek Y. On the Hermitian matrix inequalities. *Pure and Applied Mathematika Sciences* 1992; 36: 29-32.
- [24] Lee SK, Ravichandran V, Supramaniam S. Bounds for the second Hankel determinant of certain univalent functions. *Journal of Inequalities and Applications* 2013; 281: 17pp.
- [25] Libera RJ, Zlotkiewicz EJ. Coefficient bounds for the inverse of a function with derivatives in \mathcal{P} . *Proceedings of the American Mathematical Society* 1983; 87: 251-257.
- [26] Mustafa N, Mrugusundaramoorthy G, Janani T. Second Hankel determinant for a certain subclass of bi-univalent functions. *Mediterranean Journal of Mathematics* 2018; 15 (3): 17 pp.
- [27] Obradović M, Tuneski N. Hermitian-Toeplitz determinants for the class of univalent functions. *Armenian Journal of Mathematics* 2021, 13 (4): 1-10. doi:10.52737/18291163-2021.13.4-1-10.
- [28] Pommerenke C. On the coefficients and Hankel determinants of univalent functions. *Journal of the London Mathematical Society* 1966; 41: 111-122.
- [29] Trench WF. Some spectral properties of Hermitian Toeplitz matrices. *SIAM Journal on Matrix Analysis and Applications* 1994; 15 (3): 938-942.
- [30] Wilkes DM, Morgera SD, Noor F, Hayes MH. A Hermitian Toeplitz matrix is unitarily similar to a real Toeplitz-plus-Hankel matrix. *IEEE Transactions on Signal Processing* 1991; 39(9): 2146-2148.

- [31] Zaprawa P. Third Hankel determinants for subclasses of univalent functions. *Mediterranean Journal of Mathematics* 2017; 14 (1): 10pp.