

1-1-2021

Physiological, biochemical, and metabolic responses of abiotic plant stress: salinity and drought

KIARASH JAMSHIDI GOHARRIZI

Michael R. Hamblin

Soraya Karami

Maryam Nazari

Follow this and additional works at: <https://dctubitak.researchcommons.org/botany>



Part of the [Botany Commons](#)

Recommended Citation

GOHARRIZI, KIARASH JAMSHIDI; Hamblin, Michael R.; Karami, Soraya; and Nazari, Maryam (2021) "Physiological, biochemical, and metabolic responses of abiotic plant stress: salinity and drought," *Turkish Journal of Botany*. Vol. 45: No. 7, Article 4. <https://doi.org/10.3906/bot-2108-30>
Available at: <https://dctubitak.researchcommons.org/botany/vol45/iss7/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals.

Physiological, biochemical, and metabolic responses of abiotic plant stress: salinity and drought

Kiarash Jamshidi GOHARRIZI^{1*}, Michael R HAMBLIN², Soraya KARAMI^{3*}, Maryam NAZARI⁴

¹Department of Plant Breeding, Yazd Branch, Islamic Azad University, Yazd, Iran

²Laser Research Centre, Faculty of Health Science, University of Johannesburg, Doornfontein, South Africa

³Department of Agriculture, Payame Noor University (PNU), Tehran, Iran

⁴Department of Agronomy and Plant Breeding, Faculty of Agriculture, Bu-Ali Sina University, Hamedan, Iran

Received: 16.08.2021 • Accepted/Published Online: 27.11.2021 • Final Version: 28.12.2021

Abstract: The most important types of abiotic stress that affect agricultural crops throughout the world are drought and salinity. These stresses will continue to worsen as the current climate crisis progresses. Plants have evolved a complex set of mechanisms in order to protect themselves from salt and drought. At the macrobiological level, these include alterations in growth rate, water balance, antioxidant defenses, and photosynthesis. Altered metabolites include proline, carbohydrates, glycine betaine, gamma-aminobutyric acid, and polyamines. It is not surprising, thus, that one of the most important research areas of plant biology is the study of plant responses to abiotic stress and stress tolerance mechanisms. The methods, used in this study, are diverse, including the study of physiological properties, biochemical research, and metabolomics approaches toward abiotic stress. This awareness should lead to the development of (near) future, sustainable and better-adapted agriculture in the sense of global warming and environmental emission scenarios. Also, different information presented in this overview can be regarded by the scientific community to produce tolerant cultivars in response to salinity and drought stresses.

Key words: Salinity and drought stresses, physiological characteristics, biochemical traits, metabolomics

1. Introduction

Salinity and drought are two of the main stressors that affect plants' life throughout the world (Nazari et al., 2019; Jamshidi Goharrizi et al., 2020a). These stresses are able to alter the water distribution within the plant and can disturb the ionic balance in the plant resulting in the generation of reactive oxygen species (ROS) (Wang et al., 2003; Jamshidi Goharrizi et al., 2020b). Moreover, the increased ROS can reduce the overall plant production yield in agriculture (Wang et al., 2003; Jamshidi Goharrizi et al., 2019a).

In the last 25 years, increasing salinity has reduced agricultural land productivity by 30%, and this is expected to reach more than 50% by the year 2050 (Yan et al., 2003; Wang et al., 2008). Moreover, more than 45% of agricultural land around the world suffers from constant or repeated droughts, and this is also increasing every year (Ashraf and Foolad, 2007). Osmotic stress, including salinity and drought, affects different plant pathways including physiology and metabolism, with a severity based on the intensity and time duration of the stress, and finally suppresses crop production yields and plant growth (Munns, 2005; Rahnama et al., 2010; James et al.,

2011; Nazari et al., 2020). Although the region affected by drought and salinity is still expanding, genetic resources with high drought/salinity tolerance must be determined (Ibrahim et al., 2019b; Nazari et al., 2019). Therefore, the task of current and potential agriculture is to increase the availability of food to an ever-growing human population under declining environmental conditions in many parts of the world. It is of general interest to mitigate the impact of different forms of abiotic stresses (Fita et al., 2015; Jamshidi Goharrizi et al., 2018). Given its undoubted academic importance and functional consequences in agriculture, the study of abiotic stress resistance mechanisms is one of the most productive lines of research in plant biology. The various types of abiotic stresses imposed by the environment are typically interconnected and often have an osmotic component, which affects the homeostasis of plant cells (Duque et al., 2013).

This paper presents a detailed summary of the present understanding of the physiological, biochemical, and metabolic mechanisms that govern the adaptation of plants to salinity and drought stresses.

* Correspondence: jamshidi_kiarash@yahoo.com

2. Physiological Mechanisms of Drought Stress

2.1. Effect on Plant Growth

Drought stress is considered a complex process, which can alter the normal process of plant growth at various stages throughout the life span (Yordanov et al., 2000; Jamshidi Goharrizi et al., 2019a; Pakzad et al., 2019; Pakzad et al., 2021). In response to drought stress and under water deficit conditions, more efficient usage of water by the plants is the most critical factor affecting survival and growth. Also, it is well known that water availability can change the growth pattern in plants. For example, the root system is increased in water deficit conditions (Hamblin et al., 1990); in contrast, under normal growth situations and without any stress, the size of the leaf area and the rate of photosynthesis in plants have a positive correlation together, and, therefore, a bigger leaf area is considered useful to accelerate the growth rate of plants (Poorter and Remkes, 1990). The main disadvantage of having a larger leaf area in plants is the loss of more water by transpiration, even in water deficit conditions. Therefore, although a bigger leaf area can increase the growth rate of plants, it also involves a higher usage of water. In plants, better growth and more efficient water usage together determine not only biomass production but also the loss and absorption of water. Moreover, drought stress reduces the accumulation of both fresh and dry biomasses in plants, but the fresh biomass is decreased more than the dry biomass under water deficit stress (Ramos et al., 1999).

2.2. Changes in Plant Water Distribution

Plant yields are considerably reduced because of drought stress, and overcoming this stressor can be extremely difficult (Jamshidi Goharrizi et al., 2020a). Nevertheless, the process of osmotic adjustment in plants is able to overcome the destructive effects of this stress to some extent (Turner, 1986). For this purpose, several different strategies are used by plants, such as tolerance, escape, and avoidance of cell dehydration in order to survive in drought conditions (Turner, 1986). Plants attempt to resist the damage caused by drought stress by preserving the cell turgor to maintain their metabolism. Resistance to drought stress includes two different mechanisms, which are changes in the elastic attributes of tissues, and osmotic adjustment (Munns, 1988).

The main response of plants in response to water shortage is osmotic adjustment, which can maintain the cell turgor (Ludlow and Muchow, 1990) and the metabolism of plants and therefore can allow continual plant growth and yield under water deficit stress (Shackel et al., 1982). Moreover, osmotic adjustment in plants leads to dehydration of bulk tissue and reduced cellular osmotic potential, so roots shoots can develop and grow relatively more under water deficit stress (Bray, 1997).

In plants, the balance of both the metabolic process and the water potential is disturbed under drought stress, but water potential is affected in this situation later than the metabolic process, and, therefore, water potential is more tolerant to water deficit stress compared to the metabolic process (Jones and Corlett, 1992). Several studies have confirmed that there is a relationship between the cell volume and the cellular turgor pressure in sunflower (Maury et al., 2000) and common bean (Zlatko Stoyanov, 2005), but, under drought stress conditions, the balance between these two parameters is significantly disturbed (Blake et al., 1991). Another study showed that the leaf water potential had a great effect on plant functions and determined the response of most plant cultivars in response to drought stress (White et al., 2000).

For many years, it was not clear whether the overall proline content could determine the tolerance of plants under drought stress or not, but recently several studies suggested that the content of this amino acid is much higher in drought-tolerant cultivars compared to drought-sensitive ones (Jamshidi Goharrizi et al., 2020d; Jamshidi Goharrizi et al., 2020f). Moreover, in 1992 the relationship of proline content and turgor pressure was discovered, and it was then considered to be an index for drought injury in plants (Irigoyen et al., 1992).

2.3. Effect on Photosynthesis

Under drought stress, several metabolic processes in plants undergo major changes; photosynthesis is one of the most important metabolic processes in plants, which is affected by drought. Under water deficit stress conditions, the fundamental metabolic processes and the basic structures of plants are damaged, which result in reduced carbon assimilation and damage to the photosynthetic apparatus (Ali and Ashraf, 2011; Gollmack et al., 2011). Several studies have shown that a reduction in leaf photosynthesis occurs due to stomatal limitations under moderate drought stress and non-stomatal limitations in response to intense drought stress (Degl'Innocenti et al., 2009; Misson et al., 2010).

The reduction in photosynthesis results in the absorption of more light energy by the plants than is strictly required for photosynthesis, and, therefore, this excess energy is able to produce reactive oxygen species, like hydrogen peroxide, H_2O_2 , and superoxide $O_2^{\cdot-}$. The reactive oxygen species (ROs) are able to prevent the synthesis of the photosystem II (PSII) core D1 (Takahashi and Murata, 2008), which can reduce the overall activity of the photosynthetic electron transport chain.

Moreover, CO_2 fixation in the Calvin cycle is highly susceptible to different environmental stresses such as salinity (Al-Taweel et al., 2007), low temperature, and high temperature (Greer et al., 1986). In response to these destructive stresses, the reduction of the photosynthesis

rate most likely occurs due to a decrease in the synthesis of D1 protein, which occurs as a result of an interruption in CO₂ fixation. However, the precise mechanism of how the photosynthetic electron transport chain is affected by drought

3. Physiological Mechanisms of Salinity Stress

3.1. Effect on Plant Growth

The productivity and yield of plants are greatly affected by soil salinity, which is considered to be an important environmental problem. Salinity stress is able to destroy plant growth through the followings: 1) nutritional imbalance, 2) water deficit stress, and 3) excessive uptake of ions such as Na⁺ and Cl⁻, which can themselves be cytotoxic (Tsugane et al., 1999; Hernández et al., 2001; Isayenkov, 2012).

Numerous studies have confirmed the relationship between the concentration of NaCl in the soil and reduced plant growth. With an increasing concentration of NaCl, the dimensions of the plants are significantly reduced (Beltagi et al., 2006; Mustard and Renault, 2006; Jamil et al., 2007). Moreover, other studies have shown a negative relationship between the total leaf area and high levels of NaCl (López-Aguilar et al., 2003; Netondo et al., 2004; Bohra and Vyas, 2006; Chen et al., 2007; Zhao et al., 2007a; Liu et al., 2009). Furthermore, the harmful effect of salinity stress on leaf number was shown in another study. This study also proved that leaf number was greatly affected by the concentration of salinity stress (López-Aguilar et al., 2003). The negative (and positive) effects of salt stress on the dry and fresh weights of shoots and biomass have also been reported in several studies. These researchers showed that negative or positive effects of salt stress depended on the level and type of salinity stress and on the type of plant species (Bayuelo-Jiménez et al., 2002; Niazi et al., 2005; Saqib et al., 2006; Turan et al., 2007; Taffouo et al., 2009; Taffouo et al., 2010).

3.2. Changes in Plant Water Balance

Under salinity stress, it is known that the osmotic potential and the uptake of water from the soil are decreased, resulting in a disturbance of the water balance in plants (Munns, 2005). The changes in plant water balance in response to salinity stress allow plants to attempt to overcome this stressful condition using osmotic regulation. This is carried out by increasing the negativity of the osmotic (pressure) potential of the leaf sap (Rodríguez et al., 1997; Gama et al., 2009). In salt-stressed plants, stomatal closure frequently occurs, which lessens tissue dehydration by the reduction of water loss (Fricke et al., 2004). Moreover, the toxic effects of ions also decrease because of reduced transport through the xylem as a result of lowered transpiration (Kerstiens et al., 2002). In a previous study, it was found that barley cultivars exposed to salinity stress

decreased their stomatal conductivity and transpiration rates, and the reduction in these parameters caused increased growth as well as reduced the concentration of toxic ions (Veselov et al., 2008). Most likely, the triggering of abscisic acid production in salt-treated plants is one reason for stomatal closure (Mulholland et al., 2003; Fricke et al., 2004; Veselov et al., 2008), but it has been shown that the maintenance of water balance in plants can also be affected by hydraulic parameters and chemical substances (Tardieu and Simonneau, 1998). The stomatal variation in water vapor, which confirms the relationship between stomatal and hydraulic conductivity in salt-treated plants, is extremely susceptible to dynamic disturbances in water transport (Cochard et al., 2002; Meinzer, 2002; Bunce, 2006).

The hydrostatic pressure gradient causes water to enter the root xylem from the soil when plants are actively transpiring, but this balance condition is considerably altered when transpiration is limited by salinity stress. In these situations, the cell-to-cell pathway is the main route of water movement (Steudle, 2000), and most of this water is transported by aquaporins or water channels (Morillon and Chrispeels, 2001). Plasma membrane aquaporins, which belong to the phosphatidylinositol 4,5-bisphosphate (PI4,5P₂) or PIP₂ subfamily, are relatively abundant phospholipids in the root system. An artificially increased expression of this subfamily increased the activity of water channels in plant cells (Chaumont et al., 2000; Javot et al., 2003). In one study, an increase in the expression of PIP₂ aquaporins was observed in maize plants subjected to salinity stress (Zhu et al., 2005).

3.3. Effects on Photosynthesis

Decreased productivity in response to salinity stress in plants is common, and this reduction is usually associated with a decreased photosynthesis rate. Although some factors that can restrict or change the photosynthesis rate in salt-treated plants have been identified, the fundamental mechanisms of decreased photosynthesis are still somewhat unclear (Steduto et al., 2000). Several studies have shown that salinity stress can change many biochemical processes in plants. In this regard, the photosynthesis rate, which is measured by the number of photosynthetic pigments, is one of the most important biochemical processes that undergo major alterations. Several studies have shown that salinity stress is able to reduce the quantity of photosynthetic pigments in salt-stressed plants (Sultana et al., 1999; Misra et al., 2006; Taffouo et al., 2010). Following exposure of plants to salt stress, carotenoids, chlorophylls, and various enzymes, which are part of the photosynthetic apparatus, are strongly affected (Chrysargyris et al., 2018).

Stomatal closure is the main reason for the reduced photosynthesis rate in salt-stressed plants. In addition, the toxicity of salt to the photosynthetic apparatus could

be another reason for the reduction of photosynthesis in plants subjected to salinity stress. However, there are conflicting reports on whether stomatal or non-stomatal factors are most responsible for reduced photosynthesis in salt-treated plants. For example, in sunflowers, beans, and cotton, stomatal limitation was the main reason for the reduction of the photosynthesis rate in response to salt stress, while other reports implicated non-stomatal limitation for the reduction of photosynthesis (Steduto et al., 2000).

4. Biochemical Mechanisms of Tolerance to Salinity and Drought

4.1. Production of substances that affect osmotic adjustment

Salinity and drought stress exert deleterious effects on plant metabolism via ion toxicity, osmotic stress, and oxidative stress (Liang et al., 2018). In contrast, plants have evolved a series of biochemical processes to survive in abiotic stress conditions. The initial response to osmotic stress involves osmotic adjustment. Osmotic adjustment is necessary to maintain the cell turgor and to allow plant survival and maintain productivity. In order to promote osmotic adjustment at the cellular level and provide stress tolerance, plants synthesize several osmotic-regulating compounds, such as soluble proline, carbohydrates, glycine betaine, gamma-aminobutyric acid (GABA), polyamines, and others (Ashraf and Foolad, 2007; Chen and Jiang, 2010; Gupta et al., 2014; Slama et al., 2015; Blum, 2017; Rady et al., 2018; Jamshidi Goharrizi et al., 2020f). At the beginning of salinity or drought stress, the accumulation of osmotic regulating substance in the cytoplasm leads to the protection of cellular functions. Also, these compounds increase the activity of antioxidant enzymes and protect the activity of enzymatic systems (Ashraf and Foolad, 2007; Wei et al., 2009; Hossain and Fujita, 2010; Hayat et al., 2012; Kaya et al., 2013; Theerakulpisut and Phongngarm, 2013; Filippou et al., 2014; Najjaa et al., 2018; Khalid et al., 2020). Several studies have looked at the effect of exogenous application of osmoprotectant substances on the growth and tolerance in drought and salinity stressed plants (Mäkelä et al., 2019; Nakhaie et al., 2020).

4.2. Proline

Proline is the main osmotic balancing compound, which exists in plant cells in its free form. Proline has a low molecular weight, high water solubility, and zero net charge at the physiological pH value range (Trovato et al., 2019). The proline content increases in conditions of biotic and abiotic stress (Goharrizi et al., 2021) and is considered to be one of the main physiological indices of plant resistance to stress (Trovato et al., 2008; Kaur and Asthir, 2015; Dar et al., 2016; Furlan et al., 2020; Jamshidi Goharrizi

et al., 2020f; Khalid et al., 2020). In a study from our laboratory, the proline content was significantly increased in response to drought, salinity, and combined drought plus salinity stresses in different pistachio rootstocks (Jamshidi Goharrizi et al., 2020f; Jamshidi Goharrizi et al., 2020c). Two important factors, which determine salt tolerance, are the balance between the K^+/Na^+ ratio and the accumulation of proline (Kim et al., 2002). Results of several studies showed that the content of proline was at its highest level, while the K^+/Na^+ ratio was at its lowest level in tolerant cultivars of different plants (Gharsallah et al., 2016; Jamshidi Goharrizi et al., 2020f).

Many reports have shown that the harmful effects of salt stress on plant growth could be reduced by proline (Khedr et al., 2003; Szabados and Savoure, 2010; Aqsa et al., 2013; Butt et al., 2016; Ami et al., 2020). Also, the accumulation of proline improved the root capacity for water uptake in drought conditions, protected chloroplast structures by suppressing ROS generation in drought-stressed plants, and is also involved in the tolerance towards photoinhibition (Chimenti et al., 2006; Moustakas et al., 2011; Semida et al., 2020). The findings of Semida et al. (2020) demonstrated that exogenous application of proline could rescue drought-stressed onion (*Allium cepa*) plants and increase plant growth. Supplementation with proline could reduce the irrigation frequency and increase the crop yield in drought-stressed fennel (*Foeniculum vulgare*) plants (Zali and Ehsanzadeh, 2018). Proline pretreatment protected *Aloe vera* plants against salt stress by modifying the K/Na ratio and water balance, and increasing the antioxidant systems (Nakhaie et al., 2020). Moreover, exogenous application of proline to drought-stressed *Arabidopsis thaliana* plants led to a notable increase in the accumulation of proline and soluble sugars (Moustakas et al., 2011).

4.3. Accumulations of carbohydrates

Under salt and drought stress, the accumulation of sugars and carbohydrates e.g., glucose, fructose, fructans, trehalose, and starch are changed in plants (Anjum et al., 2017; Bianco and Scalisi, 2017). These carbohydrates can reduce the effects of stress by osmotic regulation, improving carbon and energy storage, and scavenging reactive oxygen species (Chen and Jiang, 2010; Nguyen et al., 2010; Yin et al., 2010; Bianco and Scalisi, 2017; Woodrow et al., 2017; Traversari et al., 2020). The studies showed that sucrose accumulation in the leaves in response to drought stress provides the necessary energy for the protection of cells during high respiration rates (Burke, 2007) and promotes plant tolerance to osmotic stresses (salinity and drought) (Van den Ende and Valluru, 2009; Al Hassan et al., 2016; Sami et al., 2016; Kumar et al., 2017a). Indeed, soluble sugars (sucrose, glucose, and fructose) are main energy sources and carry out osmoregulation

in plant cells (Rosa et al., 2009; Bolouri-Moghaddam et al., 2010; Sami et al., 2016; Du et al., 2020b). The sugars are also involved in the regulation of biological processes (Rosa et al., 2009), stability of membranes and protoplasts (Keunen et al., 2013), and the protection of enzymes from high intracellular concentrations of inorganic ions in both normal and stress conditions. Many studies have reported an increased accumulation of sugars in response to drought and salinity stress (Singh et al., 2015). For instance, Tang et al. (2013) showed that the water-soluble carbohydrate concentration increased under salt stress in *Lolium perenne* L. Palma et al. (2013) reported that salt stress caused the content of total soluble sugars and proline to increase in *Medicago sativa*. Zaher-Ara et al. (2016) showed that the accumulation of soluble sugars was involved in drought tolerance in citrus plants. Najjaa et al. (2018) reported that soluble sugars increased in *Allium roseum* in both salt and drought stress conditions. On the other hand, Abdallah et al. (2016) demonstrated that the exogenous application of trehalose reduced the adverse effects of salinity stress in rice plants. Li et al. (2020) reported that the glucose contents increased in *Lilium davidii* var. *unicolor* under drought stress conditions. Fructan can also protect plants against ROS damage, as reducing oxidative stress can increase drought stress tolerance (Rigui et al., 2019). Fructan accumulation under salinity stress could protect a tolerant wheat cultivar and avoid any yield loss under salt stress (Sharbatkhari et al., 2016). Also, it has been reported that fructan could protect sprouting *Helianthus tuberosus* L plants subjected to salt stress (Luo et al., 2018).

4.4. Glycine betaine

Glycine betaine (GB, trimethylglycine) is an amphoteric quaternized amino acid, which is present in plants, animals, and some microorganisms (Rhodes and Hanson, 1993; Chen and Murata, 2008). Glycine betaine has an overall neutral and is stable at a wide range of pH values. Numerous studies have demonstrated that GB accumulation is involved in the tolerance to salinity and drought stress conditions in many plants such as potato (You et al., 2019), *Phasoulis vulgaris* (Desoky et al., 2019; Sofy et al., 2020), stevia (Rameeh et al., 2017), tomato (De la Torre-González et al., 2018), cucumber (Youssef et al., 2018), pigeonpea (Kumar et al., 2017b), sugar beet (Li et al., 2016a), cowpea (Manaf, 2016), wheat (Gupta and Thind, 2019), and lettuce (Shams et al., 2016). Glycine betaine mainly exists in the chloroplasts where it protects the thylakoid membrane and maintains the efficiency of photochemical reactions (Tian et al., 2017). In addition, it promotes plant defensive responses such as osmotic homeostasis, protection of enzyme activity, and the expression of genes related to resistance (Chen and Murata, 2011; Giri, 2011; Ahmad et al., 2013; He et al.,

2013; Karabudak et al., 2014; Xu et al., 2018; Annunziata et al., 2019; Kahraman et al., 2019; Mäkelä et al., 2019; Sun et al., 2020). Under drought or salinity stress, GB protects the biological membranes from oxidative damage (Gupta et al., 2014; Wutipraditkul et al., 2015; Kumar et al., 2017c; Wei et al., 2017; Rady et al., 2018; Ahmad et al., 2020), modifies the cell water balance through osmotic adjustment (Gupta et al., 2014; Lai et al., 2014; Yildiztugay et al., 2014), and improves photosynthesis (Zhao et al., 2007b; Wang et al., 2010; Nusrat et al., 2014; Athar et al., 2015; Tian et al., 2017; Wei et al., 2017; Wang et al., 2019b; You et al., 2019; Nawaz and Wang, 2020), resulting in increased plant tolerance.

4.5. Gamma-aminobutyric acid

Gamma-aminobutyric acid (GABA) is an important component of the free amino acid storage system in plants, which provides a substrate for carbon and nitrogen metabolism, and is necessary for plant growth (Ansari et al., 2014). The accumulation of GABA in plants is rapidly increased in various biotic and abiotic stress conditions (Kanwal et al., 2014; Malekzadeh et al., 2014; Nayyar et al., 2014; Wang et al., 2014; Boonburapong et al., 2016; Mei et al., 2016; Vijayakumari and Puthur, 2016; Xiang et al., 2016; Otto et al., 2017; Yoon et al., 2017; Rezaei-Chiyaneh et al., 2018). Also, under salinity stress, GABA acts as an active osmolyte and also has ROS scavenging activity, suggesting that GABA may be associated with plant tolerance (Sheteiwy et al., 2019; Jalil and Ansari, 2020). Under salinity stress, the administration of exogenous GABA protected seedling growth and maintained the normal function of plants by lowering the transport of salt ions into the leaves (Wang et al., 2017b). Moreover, GABA can mitigate the harmful effects of salinity stress by improving photosynthesis and promoting the activity of antioxidant enzymes (Li et al., 2016b; Xiang et al., 2016). GABA can prevent oxidative and osmotic damage (Al-Quraan and Al-Omari, 2017), maintain cell morphology, improve cell function, and stabilize the intracellular PH value (Wang et al., 2017a). Mekonnen et al. (2016) reported that, under drought stress, the accumulation of GABA regulated stomatal opening in *Arabidopsis thaliana*. Vijayakumari and Puthur (2016) reported that the accumulation of proline and total sugars occurred earlier in plants that had been primed with GABA compared with control plants, and the activity of the antioxidant enzymes was enhanced in response to drought stress.

4.6. Polyamines

Polyamines (PAs), which include spermidine, spermine, and the diamine obligate precursor putrescine, are organic nitrogen-containing polycations of low molecular weight. They are involved in the regulation of plant growth and development and govern the defensive responses to abiotic stresses, such as drought and salinity (Takahashi and Takechi, 2010; Hussain et al., 2011; Hu et al., 2014;

Sobieszczuk-Nowicka and Legocka, 2014; Sun et al., 2016). Hu et al. (2014) reported that exogenously applied spermidine could maintain the structure of chloroplasts and protect the photosynthetic capacity in tomato plants under salinity and alkalinity stress. Li et al. (2018a) showed that exogenous spermidine application could reduce the harmful effects of drought stress in maize plants and improved the growth, photosynthetic pigment content, photosynthesis rate, and photochemical quenching parameters. On the other hand, exogenous spermidine application altered the level of phytohormones, osmolytes, and photosynthetic pigments. It also regulated the redox balance, enhanced antioxidant enzymes, regulated endogenous PA production, and preserved the vacuole structure, which all improved the tolerance of the plants to drought and salinity (Baniyadi et al., 2018; Hassan et al., 2018; Li et al., 2018b; Li et al., 2018a; Mohammadi et al., 2018; Mostafaei et al., 2018; Zhang et al., 2018; Liu et al., 2019; Du et al., 2020a; Hassan et al., 2020; Semida et al., 2020; Zhong et al., 2020).

4.7. Antioxidant Defenses

One of the biochemical changes that occur under drought and salinity stress is the generation of ROS such as hydrogen peroxide (H_2O_2), hydroxyl radicals ($\bullet HO$), superoxide radical anion ($O_2\bullet-$), and singlet oxygen (1O_2). One of the target areas of ROS is the cell membrane, so that ROS with a negative effect on membrane integrity may impair its motility and morphology and possibly lead to cell death (Wang et al., 2019a). In addition to membrane integrity, ROS can damage important cellular components, such as nucleic acids, proteins, carbohydrates, lipids, and enzymes, if not correctly regulated (Miller et al., 2010; Jamshidi Goharrizi et al., 2019b; Jamshidi Goharrizi et al., 2020e). To restore the normal balance of ROS under drought stress, plants increase the expression of different enzymes that can scavenge ROS leading to increased tolerance of salinity and drought stress (Miller et al., 2010; Dar et al., 2017). In addition to antioxidant enzymes, various antioxidant compounds can chemically quench the ROS, thus protecting plants/tissues against oxidative stress. These enzymatic and non-enzymatic ROS scavenging systems can also protect plants against the adverse effects of drought stress (Zhang et al., 2016; Dar et al., 2017; Schneider et al., 2019). The enzymatic antioxidant systems include superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), glutathione peroxidase (GPX) and peroxiredoxin (PRDX). Non-enzymatic antioxidants include ascorbic acid (AsA) and glutathione (GS). Antioxidants are required for ROS detoxification during normal plant growth, and especially in stress conditions (Ahmad et al., 2014; Das and Roychoudhury, 2014; Ahmed et al., 2015). Kumar et al. (2017a) reported that the increased activity of SOD and CAT, as well as

soluble carbohydrates, increased the tolerance of *Nerium oleander* to drought and salt stress. Jamshidi Goharrizi et al. (2020f) reported that the activity of antioxidants such as ascorbate peroxidase, catalase, and guaiacol peroxidase (GP) was enhanced in pistachio rootstocks under drought or salinity stress or a combination of the two stresses. Numerous studies have shown that the increased activity of enzymatic and non-enzymatic antioxidant systems are associated with tolerance to drought and salinity in a variety of plants, including wheat (Ahanger and Agarwal, 2017; Ahmadi et al., 2018; Dugasa et al., 2019), maize (AbdElgawad et al., 2016), rice (Islam et al., 2016; Basu et al., 2017; Vighi et al., 2017), *Amaranthus tricolor* (Sarker et al., 2018), soybeans (Liu et al., 2017b), *Chenopodium quinoa* (Fischer et al., 2017), *Thymus vulgaris* & *T. daenensis* (Bistgani et al., 2019), citrus (Hussain et al., 2018), cotton (Ibrahim et al., 2019a), *Nicotiana tabacum* (Da Silva et al., 2017), *Fargesia rufa* (Liu et al., 2017a), cabbage (Sahin et al., 2018), *Anacardium occidentale* (Lima et al., 2018), *Cucumis sativus* (Ouzounidou et al., 2016), sweet basil (Jakovljević et al., 2017), *Carthamus tinctorius* (Golkar and Taghizadeh, 2018), and *Tagetes minuta* (Moghaddam et al., 2019). The results of a study by Sheikh-Mohamadi et al. (2017) showed that high levels of diamine oxidase (histaminase: DAO) and polyamine oxidase were associated with increased tolerance to drought and salinity.

5. Metabolomics profiling in the response to drought and salinity stress

Metabolomics is a fundamental technology to study aspects of functional genomics and systems biology. Metabolomics looks at the final products of gene expression and protein activity and provides a wide overview of the biochemical status of plant tissue under different stresses (Bowne et al., 2018). The regulation of metabolic pathways is a strategy used by plants to increase tolerance under drought and salt stresses. The metabolic profile of plants under drought stress is somewhat similar to plants exposed to salt stress (Table 1). In other words, the signaling pathways of drought and salt stress exhibit some degree of overlap because both of them can lead to cell dehydration, osmotic imbalance, and oxidative stress (Golldack et al., 2014). On the other hand, there are differences in how the metabolic pathways in different species respond to drought, and understanding these differences can help to improve crop productivity (Parida et al., 2018). The TCA (tricarboxylic acid) cycle and metabolism pathways of galactose, starch, sucrose, nitrogen, and antioxidant as well as oxylipin were all important pathways that responded to drought stress (Rangani et al., 2020; Catola et al., 2016).

5.1. Carbohydrates.

The results of metabolomics analysis have shown that the accumulation of several carbohydrates, such as sucrose,

Table 1. Important metabolites, which are altered in both drought and salinity stress.

Metabolite Class	Metabolites	Reference
Sugars	Glucose	(Shi et al., 2015; Nam et al., 2016; Li et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020)
	Sucrose	(Shi et al., 2015; Gupta and De, 2017; Li et al., 2017; Rangani et al., 2020; Xiong et al., 2021)
	Galactose	(Li et al., 2017; Kumari and Parida, 2018; 2019a; Rangani et al., 2020)
	Mannose	(Shi et al., 2015; Li et al., 2017; Mibei et al., 2018; Rangani et al., 2020)
	Myo-inositol	(Li et al., 2017; Mibei et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021)
	Glycerol	(Nam et al., 2016; Gupta and De, 2017; Guo et al., 2018; Kumari and Parida, 2018; Rangani et al., 2020)
	Fructose	(Shi et al., 2015; Gupta and De, 2017; Li et al., 2017; Guo et al., 2018; Kumari and Parida, 2018)
	Trehalose	(Shi et al., 2015; Nam et al., 2016; Mibei et al., 2018; Xiong et al., 2021)
Amino acids	Aspartic acid	(Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Guo et al., 2018; Kumari and Parida, 2018; Rangani et al., 2020)
	Glutamic acid	(Shi et al., 2015; Gupta and De, 2017; Li et al., 2017; Guo et al., 2018; Kumari and Parida, 2018; Mibei et al., 2018; Nawaz and Wang, 2020; Rangani et al., 2020)
	Serine	(Alla et al., 2012; Shi et al., 2015; Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021)
	Glycine	(Alla et al., 2012; Shi et al., 2015; Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020)
	Histidine	(Alla et al., 2012; Gupta and De, 2017; Kumari and Parida, 2018; Xiong et al., 2021)
	Arginine	(Kumari and Parida, 2018; Rangani et al., 2020)
	Threonine	(Shi et al., 2015; Nam et al., 2016; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2020)
	Alanine	(Alla et al., 2012; Shi et al., 2015; Nam et al., 2016; Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021)
	Proline	(Alla et al., 2012; Shi et al., 2015; Nam et al., 2016; Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020)
	Valine	(Alla et al., 2012; Shi et al., 2015; Nam et al., 2016; Li et al., 2017; Guo et al., 2018; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021)
	Methionine	(Alla et al., 2012; Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020)
	Isoleucine	(Shi et al., 2015; Nam et al., 2016; Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021)
	Leucine	(Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020)
	Phenylalanine	(Gupta and De, 2017; Li et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020)
	Lysine	(Shi et al., 2015; Nam et al., 2016; Kumari and Parida, 2018; Rangani et al., 2020)
	Asparagin	(Shi et al., 2015; Nam et al., 2016; Li et al., 2017)
Organic acids	Tryptophan	(Gupta and De, 2017; Mostafaei et al., 2018; Kang et al., 2019)
	Glyceric acid	(Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Guo et al., 2018; Rangani et al., 2020)
	GABA	(Guo et al., 2018; Xiong et al., 2021)
	Shikimic acid	(Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020)
TCA cycle	Oxalic acid	(Nam et al., 2016; Gupta and De, 2017; Guo et al., 2018; Kumari and Parida, 2018; Rangani et al., 2020)
	Malic acid	(Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Guo et al., 2018; Kumari and Parida, 2018; Kang et al., 2019; Rangani et al., 2020)
	Citric acid	(Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021)
	Fumaric acid	(Xiong et al., 2021)
	Succinic acid	(Nam et al., 2016; Gupta and De, 2017; Li et al., 2017; Guo et al., 2018)

glucose, mannose, galactose, erythrose, sorbose, glycerol, galactose, rhamnose, and xylose were significantly changed under water deficit and salinity stress (Kanani et al., 2010; Sanchez et al., 2012; Marček et al., 2019; Rangani et al., 2020; Kumari and Parida, 2018). Besides, Kanani et al. (2010) reported that the concentrations of glycine, N-acetylglutamate, and allantoin were increased in salt stress in *Arabidopsis thaliana*. On the other hand, metabolomics profiling showed a positive relationship between concentrations of sugars and resistance to drought stress in drought-tolerant varieties of *Brachypodium distachyon* (Shi et al., 2015). Marček et al. (2019) showed that the accumulation of sugars such as sucrose, glucose, fructose, and several disaccharides were enhanced in different genotypes of winter wheat under drought stress.

Sugar alcohols are one type of important osmotic adjustment substances. Metabolic profiling has shown that sugar alcohols and metabolites, such as mannitol, inositol, tagatose, d-arabitol, galactose, 1-kestose, myo-inositol, and galactinol were increased when plants were exposed to salt or drought stress (Li et al., 2017; Kumari and Parida, 2018; Rangani et al., 2020; Xiong et al., 2021).

5.2. Amino acids

Amino acid metabolism results in the accumulation of osmoprotectant amino acids, such as proline, glutamate, aspartate, isoleucine, glycine, GABA and phenylalanine, tryptophan, and alanine (Shi et al., 2015; Li et al., 2017; Marček et al., 2019; Xiong et al., 2021). The results showed that high concentrations of amino acids such as isoleucine and phenylalanine may increase glycolysis in order to reduce salt stress and can also act as an osmotic regulator (Kumari and Parida, 2018; Rangani et al., 2020). It has been reported that the levels of alanine, phenylalanine, and tyrosine were enhanced in drought stress, while levels of serine, methionine, glycine, glutamate, and arginine were decreased (Rangani et al., 2020). On the other hand, Xiong et al. (2021) reported that levels of valine, leucine, fumarate, citrulline, and isoleucine were increased in *Phlox subulata* under drought stress, while the levels of alanine and tryptophan were decreased. The results of a study by Marček et al. (2019) showed that amino acids such as proline, threonine, GABA, and glutamine were associated with tolerance to drought stress in wheat genotypes. It was reported that the concentrations of proline, methionine, arginine, lysine, were changed in drought-tolerant wheat genotypes (Michaletti et al., 2018).

5.3. Metabolites involved in ROS scavenging

Results of metabolomics profiling in different species of plants have shown how various metabolites involved in ROS scavenging are altered in response to drought and salinity stress. Some osmoprotectant substances play a simultaneous role as ROS scavenging agents. For example, the results of a metabolomic analysis by Jia et

al. (2020) showed that some soluble carbohydrates, such as glucose, sucrose, fructose, galactose, lactose, trehalose, and myo-inositol were significantly increased in tolerant and susceptible species of poplar (*Populus* spp.) trees in response to drought stress. The accumulation of these metabolites protects cells from damage through osmotic adjustment and by ROS scavenging in drought conditions. The results of this study showed that various non-enzymatic antioxidants, including ascorbate, α -tocopherol, zeaxanthin, and β -carotene, were increased in tolerant species. α -Tocopherol prevented lipid peroxidation by translocating to the lipid membranes and quenching ROS free radicals (Kim et al., 2017).

The carotenoids, β -carotene and zeaxanthin, are involved in photosynthesis and photoprotection, are precursors to phytohormones such as ABA and strigolactone, and are also antioxidants and robustly scavenge free radicals (Cazzonelli, 2011; Mibei et al., 2017). Flavonoids are also secondary metabolites that can quench free radicals and mediate the activation of ROS-induced signaling cascades. Metabolic profiling has shown that many flavonoids are increased in plants subjected to salinity and drought stress (Nakabayashi et al., 2014; Pandey et al., 2015; Goufo et al., 2017; Li et al., 2017; Biswas et al., 2018; Gai et al., 2020). Increased levels of several polyphenols, such as sorbitol (Biswas et al., 2018) and gallic acid (Li et al., 2017) were observed in various plant species subjected to salinity stress.

A raised concentration of isoleucine and phenylalanine can help to scavenge oxygen free radicals. Phenylalanine production via the shikimate pathway is increased under salinity and drought stress. On the other hand, phenylalanine is involved in the production of secondary metabolites that act as ROS scavengers (Kumari and Parida, 2018; Rangani et al., 2020). It was reported that antioxidants such as hydroxylamine and putrescine were increased under neutral salt stress in soybeans (Li et al., 2017).

5.4. Organic acids

Metabolic profiling has shown that various organic acids are involved in plant tolerance to drought and salinity stress. Rangani et al. (2020) reported that galactaric acid, oxalic acid, malic acid, and tartaric acid were all increased in *Salvadora persica* leaves in response to drought stress, while the levels of myristic acid, ribonic acid, tartaric acid, stearic acid, lactic acid, and 2-oxoglutaric acid were all decreased. The accumulation of organic acids, such as 3-hydroxybutyric acid, galacturonic acid, 15-alkyl oleic acid, linoleic acid, and glycuronic acid was also reported to occur after salt stress in soybean plants (Li et al., 2017). It was reported that the accumulation of malic and oxalic acids was increased in several wheat genotypes under drought stress (Marček et al., 2019). On the other hand,

fatty acids and their derivatives play an important role in plant resistance and the storage of energy under abiotic stress. Elevated levels of fatty acids, such as palmitic acid, lauric acid, myristic acid, and sterol acids have been reported to mediate resistance to salinity and drought stress (Kanani et al., 2010; Biswas et al., 2018). The increased accumulation of citric and benzoic acids was also reported in *Phlox subulata* under drought stress (Xiong et al., 2021). Moreover, other organic acids such as pyruvic acid, amino adipic acid (Michaletti et al., 2018), O-acetyl salicylic acid (Biswas et al., 2018) have been reported to be involved in plant tolerance to drought and salinity stress (Marček et al., 2019). The 16-hydroxypalmitic acid pathway, which is involved in cuticular wax biosynthesis was reported to increase tolerance to drought (Jia et al., 2020).

5.5. Tricarboxylic acid cycle

The tricarboxylic acid (TCA) cycle is one of the main pathways in energy metabolism in all living cells. This cycle involves the oxidation of respiratory substrates to finally produce ATP. Kanani et al. (2010) reported that the levels of all the TCA cycle intermediates ranging from citrate to fumarate were increased in salt stress. The role of the TCA cycle in different plant species has been investigated in both drought and salinity stress conditions (Nam et al., 2016; Gupta and De, 2017; Kumari and Parida, 2018; Rangani et al., 2020).

5.6. Phytohormones

Phytohormones are the key regulators in many plant responses to abiotic stresses (Wani et al., 2016). Plants must be able to modulate their growth and development in response to different internal and external stimuli (Wolters and Jürgens, 2009). Phytohormones are signaling molecules that exist in only small amounts in plant cells but are designed to regulate these responses. Their crucial role in plant adaptation to ever-changing conditions, by regulating growth, sink-source transition in leaves, overall development, and nutrient distribution is well understood (Fahad et al., 2015). Although the response of plants to abiotic stress depends on many different factors, phytohormones are the most significant endogenous agents that regulate physiological and molecular responses, which is important for the survival of plants as sessile organisms (Fahad et al., 2015). Phytohormones can function either at their site of biosynthesis or elsewhere in the plant after their transport (Peleg and Blumwald, 2011). Phytohormones are of major importance in the development of plants and in growth plasticity. Phytohormones are formed from auxins such as indole-3-acetic acid (IAA), cytokinins (CKs), abscisic acid (ABA), ethylene (ET), gibberellins (GAs), salicylic acid (SA), brassinosteroids (BRs), and jasmonates (JAs); it has recently been discovered that strigolactones (SL) are new phytohormones. According to the results obtained,

the levels and activities of phytohormones, such as ABA, SA, and JA increase under drought and salinity conditions (Rangani et al., 2020), so they can act as signaling molecules to promote plant survival under these stress (Kumari and Parida, 2018). Moreover, there is increasing evidence that IAA plays an integral role in the adaptation of plants to salinity stress (Iqbal et al., 2014; Fahad et al., 2015). In plants under water stress, CK levels decreases, and on the other hand, increasing the level of ABA leads to an increase in the ABA/CK ratio. Indeed, the decreased CK levels function to increase apical dominance, and along with ABA acts to modulate the stomatal aperture, which helps in the adaptation to drought stress (O'Brien and Benková, 2013).

6. Conclusion

In summary, as mentioned above, salinity (in water and soil) and drought are important factors that hinder the crop production in areas affected by water scarcity and salt. Regarding the effects of salinity stress, it was found that the metabolic activities of plants could be limited due to the accumulation of salt. Hence, the occurrence of salt stress can be the cause of severe changes in the development and growth of plants, which ultimately reduces plant survival. In contrast, against various ionic and osmotic components of the stress of salinity, several complicated physiological and biochemical reactions (single or combined) are activated at the whole, organ, or cell plant level to overcome or adapt to these harsh conditions. These reactions can occur in the form of various strategies in plants including avoidance, exclusion, ion compartmentalization, extrusion, and so on (single or combined). Therefore, a correct understanding of the mechanisms related to salt tolerance in tissue/organ/cell of plants can provide a new opportunity to integrate biochemical and physiological knowledge to amend and improve salinity tolerance in order to improve economic productivity and facilitate cultivation, especially in salt-affected areas. In addition to salinity, another usual event in many areas is drought. Hence, various mechanisms have been developed by many plant species to deal with this limitation (water supply). Briefly, plant species can prevent stress of drought via maximizing water uptake (e.g., tapping groundwater by longitudinal growth of roots) and/or minimizing water loss (e.g., reducing transpiration by small leaves and closing stomata), etc. However, these strategies can also affect other fundamental plant processes or systems such as the photosynthetic apparatus and related processes (RuBisCo activity, energy balance, and contents and components of chlorophyll) as well as metabolic activities. Despite significant advances in identifying and elucidating drought tolerance mechanisms, significant challenges remain.

In general, the points that should not be overlooked is the fact that, in selection and breeding programs, various indices related to drought and salinity resistance/tolerance/adaptation can be used; however, the interaction between the different indices in the environment and the plant complicates breeding programs for osmotic stresses. On the other hand, variable levels of multiple stresses affect plants under field conditions, but many researchers evaluate the response of plants to specific stress alone under controlled conditions (in laboratory or field conditions). Therefore, there occurs a significant gap between the knowledge obtained under controlled conditions and the knowledge needed to breed plants with the goal of enhancing tolerance/resistance/adaptation to osmotic stresses in field conditions. This is because; the plants' response to specific stress (alone) may not be a true and clear reflection of their behavior in the natural conditions, where several multiple stresses happen simultaneously. In addition, in many cases, the effects of salinity and drought stresses are indistinguishable from each other, because many of the physiological and biochemical mechanisms of osmotic stresses tolerance/resistance are common to many plant species when plants are exposed to the stress of salinity or drought, such as proline, which is an extremely important biochemical mechanism of tolerance to salinity and drought stresses in plants, whereas some mechanisms are specifically amplified

under a combination of stresses or vice versa. As a final point, since the response of plants to multiple stresses could not be deduced from the response to individual stress, one of the scopes of study that should be considered is the response of plants to a combination of stresses to develop stress-tolerant/resistant plants in field conditions. On the other hand, the plant's response to multiple stresses may not reflect the incremental impact of these restrictions, but rather, a specific response to a new situation in which many constraints operate simultaneously. Therefore, to accurately describe the actual response of plants to multiple stresses, the stresses must be imposed sequentially or simultaneously, and each set of environmental conditions must be considered as a completely new stress.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Declaration of funding

The present study was financially supported by Sahab Gostar Kerman Engineering Services Company, Kerman, Iran with Grant No. 1400-26 and Grant Code 1296.

Data availability statement

The data that support this study will be shared upon reasonable request to the corresponding authors.

References

- Abdallah M-S, Abdelgawad Z, El-Bassiouny H (2016). Alleviation of the adverse effects of salinity stress using trehalose in two rice varieties. *South African Journal of Botany* 103:275-282.
- AbdElgawad H, Zinta G, Hegab MM, Pandey R, Asard H et al. (2016). High salinity induces different oxidative stress and antioxidant responses in maize seedlings organs. *Frontiers in Plant Science* 7:276.
- Ahanger MA, Agarwal R (2017). Salinity stress induced alterations in antioxidant metabolism and nitrogen assimilation in wheat (*Triticum aestivum* L) as influenced by potassium supplementation. *Plant Physiology and Biochemistry* 115:449-460.
- Ahmad P, Jamsheed S, Hameed A, Rasool S, Sharma I et al. (2014). Drought stress induced oxidative damage and antioxidants in plants. In: *Oxidative Damage to Plants*. Elsevier, pp 345-367.
- Ahmad R, Lim CJ, Kwon SY (2013). Glycine betaine: a versatile compound with great potential for gene pyramiding to improve crop plant performance against environmental stresses. *Plant Biotechnology Reports* 7:49-57.
- Ahmad R, Ali S, Abid M, Rizwan M, Ali B et al. (2020). Glycinebetaine alleviates the chromium toxicity in *Brassica oleracea* L. by suppressing oxidative stress and modulating the plant morphology and photosynthetic attributes. *Environmental Science and Pollution Research* 27:1101-1111.
- Ahmadi J, Pour-Aboughadareh A, Fabriki-Ourang S, Mehrabi AA, Siddique KH (2018). Screening wild progenitors of wheat for salinity stress at early stages of plant growth: insight into potential sources of variability for salinity adaptation in wheat. *Crop and Pasture Science* 69:649-658.
- Ahmed IM, Nadira UA, Bibi N, Cao F, He X et al. (2015). Secondary metabolism and antioxidants are involved in the tolerance to drought and salinity, separately and combined, in Tibetan wild barley. *Environmental and Experimental Botany* 111:1-12.
- Al-Quraan N, Al-Omari H (2017). GABA accumulation and oxidative damage responses to salt, osmotic and H₂O₂ treatments in two lentil (*Lens culinaris* Medik) accessions. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology* 151:148-157.
- Al-Taweel K, Iwaki T, Yabuta Y, Shigeoka S, Murata N et al. (2007). A bacterial transgene for catalase protects translation of D1 protein during exposure of salt-stressed tobacco leaves to strong light. *Plant Physiology* 145:258.
- Al Hassan M, del Pilar López-Gresa M, Boscaiu M, Vicente O (2016). Stress tolerance mechanisms in *Juncus*: responses to salinity and drought in three *Juncus* species adapted to different natural environments. *Functional Plant Biology* 43:949-960.

- Ali Q, Ashraf M (2011). Induction of drought tolerance in maize (*Zea mays* L.) due to exogenous application of trehalose: growth, photosynthesis, water relations and oxidative defence mechanism. *Journal of Agronomy and Crop Science* 197:258-271.
- Alla MMN, Khedr AHA, Serag MM, Abu-Alnaga AZ, Nada RM (2012). Regulation of metabolomics in *Atriplex halimus* growth under salt and drought stress. *Plant Growth Regulation* 67:281-304.
- Ami K, Planchais S, Cabassa C, Guivarc'h A, Véry AA et al. (2020). Different proline responses of two Algerian durum wheat cultivars to in vitro salt stress. *Acta Physiologiae Plantarum* 42:21.
- Anjum SA, Ashraf U, Tanveer M, Khan I, Hussain S et al. (2017). Drought induced changes in growth, osmolyte accumulation and antioxidant metabolism of three maize hybrids. *Frontiers in Plant Science* 8:69.
- Annunziata MG, Ciarmiello LF, Woodrow P, Dell'Aversana E, Carillo P (2019). Spatial and temporal profile of glycine betaine accumulation in plants under abiotic stresses. *Frontiers in plant science* 10:230.
- Ansari MI, Hasan S, Jalil SU (2014). Leaf senescence and GABA shunt. *Bioinformation* 10:734.
- Aqsa T, Khalid N, Khalid H, Bhatti KH, Siddiqi EH et al. (2013). Foliar application of proline for salt tolerance of two wheat (*Triticum aestivum* L.) cultivars. *World Applied Sciences Journal* 22:547-554.
- Ashraf M, Foolad MR (2007). Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environmental and Experimental Botany* 59:206-216.
- Athar HuR, Zafar Z, Ashraf M (2015). Glycinebetaine improved photosynthesis in canola under salt stress: evaluation of chlorophyll fluorescence parameters as potential indicators. *Journal of Agronomy and Crop Science* 201:428-442.
- Baniasadi F, Saffari VR, Moud AAM (2018). Physiological and growth responses of *Calendula officinalis* L. plants to the interaction effects of polyamines and salt stress. *Scientia Horticulturae* 234:312-317.
- Basu S, Giri RK, Benazir I, Kumar S, Rajwanshi R et al. (2017). Comprehensive physiological analyses and reactive oxygen species profiling in drought tolerant rice genotypes under salinity stress. *Physiology and Molecular Biology of Plants* 23:837-850.
- Bayuelo-Jiménez JS, Debouck DG, Lynch JP (2002). Salinity tolerance in *Phaseolus* Species during early vegetative growth. *Crop Science* 42:2184-2192.
- Beltagi M, Ismail M, Mohamed F (2006). Induced salt tolerance in common bean (*Phaseolus vulgaris* L.) by gamma irradiation. *Pakistan Journal of Biological Sciences* 9:1143-1148.
- Bianco RL, Scalisi A (2017). Water relations and carbohydrate partitioning of four greenhouse-grown olive genotypes under long-term drought. *Trees* 31:717-727.
- Bistgani ZE, Hashemi M, DaCosta M, Craker L, Maggi F et al. (2019). Effect of salinity stress on the physiological characteristics, phenolic compounds and antioxidant activity of *Thymus vulgaris* L. and *Thymus daenensis* Celak. *Industrial Crops and Products* 135:311-320.
- Biswas S, Biswas AK, De B (2018). Metabolomics analysis of *Cajanus cajan* L. seedlings unravelled amelioration of stress induced responses to salinity after halopriming of seeds. *Plant Signaling & Behavior* 13:e1489670.
- Blake TJ, Bevilacqua E, Zwiazek JJ (1991). Effects of repeated stress on turgor pressure and cell elasticity changes in black spruce seedlings. *Canadian Journal of Forest Research* 21:1329-1333.
- Blum A (2017). Osmotic adjustment is a prime drought stress adaptive engine in support of plant production. *Plant, cell & environment* 40:4-10.
- Bolouri-Moghaddam MR, Le Roy K, Xiang L, Rolland F, Van den Ende W (2010). Sugar signalling and antioxidant network connections in plant cells. *The FEBS Journal* 277:2022-2037.
- Boonburapong B, Laloknam S, Incharoensakdi A (2016). Accumulation of gamma-aminobutyric acid in the halotolerant cyanobacterium *Aphanothece halophytica* under salt and acid stress. *Journal of applied phycology* 28:141-148.
- Bowne J, Bacic A, Tester M, Roessner U (2018). Abiotic stress and metabolomics. *Annual Plant Reviews online* 61-85.
- Bray EA (1997). Plant responses to water deficit. *Trends in Plant Science* 2:48-54.
- Bunce JA (2006). How do leaf hydraulics limit stomatal conductance at high water vapour pressure deficits? *Plant, Cell & Environment* 29:1644-1650.
- Burke JJ (2007). Evaluation of source leaf responses to water-deficit stresses in cotton using a novel stress bioassay. *Plant Physiology* 143:108-121.
- Butt M, Ayyub C, Amjad M, Ahmad R (2016). Proline application enhances growth of chilli by improving physiological and biochemical attributes under salt stress. *Pakistan Journal of Agricultural Sciences* 53:43-49.
- Catola S, Marino G, Emiliani G, Huseynova T, Musayev M et al. (2016). Physiological and metabolomic analysis of *Punica granatum* (L.) under drought stress. *Planta* 243:441-449.
- Chaumont F, Barrieu F, Jung R, Chrispeels MJ (2000). Plasma membrane intrinsic proteins from maize cluster in two sequence subgroups with differential aquaporin activity. *Plant Physiology* 122:1025-1034.
- Chen C, Tao C, Peng H, Ding Y (2007). Genetic analysis of salt stress responses in asparagus bean (*Vigna unguiculata* (L.) ssp. *sesquipedalis* Verdc.). *Journal of Heredity* 98:655-665.
- Chen H, Jiang JG (2010). Osmotic adjustment and plant adaptation to environmental changes related to drought and salinity. *Environmental Reviews* 18:309-319.
- Chen TH, Murata N (2011). Glycinebetaine protects plants against abiotic stress: mechanisms and biotechnological applications. *Plant, Cell & Environment* 34:1-20.

- Chen TH, Murata N (2008). Glycinebetaine: an effective protectant against abiotic stress in plants. *Trends in plant science* 13:499-505.
- Chimenti CA, Marcantonio M, Hall A (2006). Divergent selection for osmotic adjustment results in improved drought tolerance in maize (*Zea mays* L.) in both early growth and flowering phases. *Field Crops Research* 95:305-315.
- Chrysargyris A, Tzision A, Xylia P, Tzortzakis N (2018). Effects of salinity on tagetes growth, physiology, and shelf life of edible flowers stored in passive modified atmosphere packaging or treated with ethanol. *Frontiers in Plant Science* 9:1765-1765.
- Cochard H, Coll L, Le Roux X, Améglio T (2002). Unraveling the effects of plant hydraulics on stomatal closure during water stress in walnut. *Plant Physiology* 128:282.
- Da Silva CJ, Fontes EPB, Modolo LV (2017). Salinity-induced accumulation of endogenous H₂S and NO is associated with modulation of the antioxidant and redox defense systems in *Nicotiana tabacum* L. cv. Havana. *Plant Science* 256:148-159.
- Dar MI, Naikoo MI, Khan FA, Rehman F, Green ID et al. (2017). An introduction to reactive oxygen species metabolism under changing climate in plants. In: *Reactive oxygen species and antioxidant systems in plants: Role and Regulation under Abiotic Stress*. Springer, pp 25-52.
- Dar MI, Naikoo MI, Rehman F, Naushin F, Khan FA (2016). Proline accumulation in plants: roles in stress tolerance and plant development. In: *Osmolytes and Plants Acclimation to Changing Environment: Emerging Omics Technologies*. Springer, pp 155-166.
- Das K, Roychoudhury A (2014). Reactive oxygen species (ROS) and response of antioxidants as ROS-scavengers during environmental stress in plants. *Frontiers in Environmental Science* 2:53.
- De la Torre-González A, Montesinos-Pereira D, Blasco B, Ruiz J (2018). Influence of the proline metabolism and glycine betaine on tolerance to salt stress in tomato (*Solanum lycopersicum* L.) commercial genotypes. *Journal of Plant Physiology* 231:329-336.
- Degl'Innocenti E, Hafsi C, Guidi L, Navari-Izzo F (2009). The effect of salinity on photosynthetic activity in potassium-deficient barley species. *Journal of Plant Physiology* 166:1968-1981.
- Desoky E, Ibrahim SA, Merwad ARM (2019). Mitigation of salinity stress effects on growth, physio-chemical parameters and yield of snapbean (*Phaseolus vulgaris* L.) by exogenous application of glycine betaine. *International Letters of Natural Sciences* 76:60-71.
- Du H, Liu G, Liu D, Yu J, Liu H et al. (2020a). Polyamines conjugated to embryo tonoplast enhance the tolerance of maize to drought stress by maintaining the vacuole conformation. *Russian Journal of Plant Physiology* 67:724-732.
- Du Y, Zhao Q, Chen L, Yao X, Zhang W et al. (2020b). Effect of drought stress on sugar metabolism in leaves and roots of soybean seedlings. *Plant Physiology and Biochemistry* 146:1-12.
- Dugasa MT, Cao F, Ibrahim W, Wu F (2019). Differences in physiological and biochemical characteristics in response to single and combined drought and salinity stresses between wheat genotypes differing in salt tolerance. *Physiologia Plantarum* 165:134-143.
- Duque AS, de Almeida AM, da Silva AB, da Silva JM, Farinha AP et al. (2013). Abiotic stress responses in plants: unraveling the complexity of genes and networks to survive. *Abiotic stress-plant responses and applications in agriculture* 13:49-101.
- Fahad S, Hussain S, Bano A, Saud S, Hassan S et al. (2015). Potential role of phytohormones and plant growth-promoting rhizobacteria in abiotic stresses: consequences for changing environment. *Environmental Science and Pollution Research* 22:4907-4921.
- Filippou P, Bouchagier P, Skotti E, Fotopoulos V (2014). Proline and reactive oxygen/nitrogen species metabolism is involved in the tolerant response of the invasive plant species *Ailanthus altissima* to drought and salinity. *Environmental and Experimental Botany* 97:1-10.
- Fischer S, Wilckens R, Jara J, Aranda M, Valdivia W et al. (2017). Protein and antioxidant composition of quinoa (*Chenopodium quinoa* Willd.) sprout from seeds submitted to water stress, salinity and light conditions. *Industrial Crops and Products* 107:558-564.
- Fita A, Rodríguez-Burruezo A, Boscaiu M, Prohens J, Vicente O (2015). Breeding and domesticating crops adapted to drought and salinity: a new paradigm for increasing food production. *Frontiers in Plant Science* 6:978.
- Fricke W, Akhilarova G, Veselov D, Kudoyarova G (2004). Rapid and tissue-specific changes in ABA and in growth rate in response to salinity in barley leaves. *Journal of Experimental Botany* 55:1115-1123.
- Furlan AL, Bianucci E, Giordano W, Castro S, Becker DF (2020). Proline metabolic dynamics and implications in drought tolerance of peanut plants. *Plant Physiology and Biochemistry* 151:566-578.
- Gai Z, Wang Y, Ding Y, Qian W, Qiu C et al. (2020). Exogenous abscisic acid induces the lipid and flavonoid metabolism of tea plants under drought stress. *Scientific Reports* 10:1-13.
- Gama PBS, Tanaka K, Eneji AE, Eltayeb AE, Siddig KE (2009). Salt-induced stress effects on biomass, photosynthetic rate, and reactive oxygen species-scavenging enzyme accumulation in common bean. *Journal of Plant Nutrition* 32:837-854.
- Gharsallah C, Fakhfakh H, Grubb D, Gorsane F (2016). Effect of salt stress on ion concentration, proline content, antioxidant enzyme activities and gene expression in tomato cultivars. *AoB Plants* 8: plw055.
- Giri J (2011). Glycinebetaine and abiotic stress tolerance in plants. *Plant Signaling & Behavior* 6:1746-1751.
- Goharrizi KJ, Meru G, Kermani SG, Heidarinezhad A, Salehi F (2021). Short-term cold stress affects physiological and biochemical traits of pistachio rootstocks. *South African Journal of Botany* 141:90-98.

- Golkar P, Taghizadeh M (2018). In vitro evaluation of phenolic and osmolite compounds, ionic content, and antioxidant activity in safflower (*Carthamus tinctorius* L.) under salinity stress. *Plant Cell, Tissue and Organ Culture (PCTOC)* 134:357-368.
- Golldack D, Lüking I, Yang O (2011). Plant tolerance to drought and salinity: stress regulating transcription factors and their functional significance in the cellular transcriptional network. *Plant Cell Reports* 30:1383-1391.
- Golldack D, Li C, Mohan H, Probst N (2014). Tolerance to drought and salt stress in plants: unraveling the signaling networks. *Frontiers in Plant Science* 5:151-151.
- Goufo P, Moutinho-Pereira JM, Jorge TF, Correia CM, Oliveira MR et al. (2017). Cowpea (*Vigna unguiculata* L. Walp.) metabolomics: osmoprotection as a physiological strategy for drought stress resistance and improved yield. *Frontiers in Plant Science* 8:586.
- Greer DH, Berry JA, Björkman O (1986). Photoinhibition of photosynthesis in intact bean leaves: role of light and temperature, and requirement for chloroplast-protein synthesis during recovery. *Planta* 168:253-260.
- Guo S-H, Hu N, Li Q-S, Yang P, Wang L-L et al. (2018). Response of edible amaranth cultivar to salt stress led to Cd mobilization in rhizosphere soil: a metabolomic analysis. *Environmental Pollution* 241:422-431.
- Gupta N, Thind S (2019). Foliar application of glycine betaine alters sugar metabolism of wheat leaves under prolonged field drought stress. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences* 89:877-884.
- Gupta N, Thind SK, Bains NS (2014). Glycine betaine application modifies biochemical attributes of osmotic adjustment in drought stressed wheat. *Plant Growth Regulation* 72:221-228.
- Gupta P, De B (2017). Metabolomics analysis of rice responses to salinity stress revealed elevation of serotonin, and gentisic acid levels in leaves of tolerant varieties. *Plant Signaling & Behavior* 12:e1335845.
- Hamblin A, Tennant D, Perry MW (1990). The cost of stress: dry matter partitioning changes with seasonal supply of water and nitrogen to dryland wheat. *Plant and Soil* 122:47-58.
- Hassan F, Ali E, Alamer K (2018). Exogenous application of polyamines alleviates water stress-induced oxidative stress of *Rosa damascena* Miller var. *trigintipetala* Dieck. *South African Journal of Botany* 116:96-102.
- Hassan N, Ebeed H, Aljaarany A (2020). Exogenous application of spermine and putrescine mitigate adversities of drought stress in wheat by protecting membranes and chloroplast ultra-structure. *Physiology and Molecular Biology of Plants* 26:233-245.
- Hayat S, Khalique G, Irfan M, Wani AS, Tripathi BN et al. (2012). Physiological changes induced by chromium stress in plants: an overview. *Protoplasma* 249:599-611.
- He C, He Y, Liu Q, Liu T, Liu C et al. (2013). Co-expression of genes ApGSMT2 and ApDMT2 for glycinebetaine synthesis in maize enhances the drought tolerance of plants. *Molecular Breeding* 31:559-573.
- Hernández JA, Ferrer MA, Jiménez A, Barceló AR, Sevilla F (2001). Antioxidant systems and O₂⁻/H₂O₂ production in the apoplast of pea leaves. Its relation with salt-induced necrotic lesions in minor veins. *Plant Physiology* 127:817-831.
- Hossain MA, Fujita M (2010). Evidence for a role of exogenous glycinebetaine and proline in antioxidant defense and methylglyoxal detoxification systems in mung bean seedlings under salt stress. *Physiology and Molecular Biology of Plants* 16:19-29.
- Hu L, Xiang L, Zhang L, Zhou X, Zou Z et al. (2014). The photoprotective role of spermidine in tomato seedlings under salinity-alkalinity stress. *PLoS One* 9:e110855.
- Hussain S, Khalid MF, Saqib M, Ahmad S, Zafar W et al. (2018). Drought tolerance in citrus rootstocks is associated with better antioxidant defense mechanism. *Acta Physiologiae Plantarum* 40:135.
- Hussain SS, Ali M, Ahmad M, Siddique KH (2011). Polyamines: natural and engineered abiotic and biotic stress tolerance in plants. *Biotechnology Advances* 29:300-311.
- Ibrahim W, Zhu YM, Chen Y, Qiu CW, Zhu S et al. (2019a). Genotypic differences in leaf secondary metabolism, plant hormones and yield under alone and combined stress of drought and salinity in cotton genotypes. *Physiologia Plantarum* 165:343-355.
- Ibrahim W, Qiu CW, Zhang C, Cao F, Shuijin Z et al. (2019b). Comparative physiological analysis in the tolerance to salinity and drought individual and combination in two cotton genotypes with contrasting salt tolerance. *Physiologia Plantarum* 165:155-168.
- Bohra A, Vyas A (2006). Biomass production, productivity and physiological changes in moth bean genotypes at different salinity levels. *American Journal of Plant Physiology* 1:210-213.
- Iqbal N, Umar S, Khan NA, Khan MIR (2014). A new perspective of phytohormones in salinity tolerance: regulation of proline metabolism. *Environmental and Experimental Botany* 100:34-42.
- Irigoyen JJ, Einerich DW, Sánchez-Díaz M (1992). Water stress induced changes in concentrations of proline and total soluble sugars in nodulated alfalfa (*Medicago sativa*) plants. *Physiologia Plantarum* 84:55-60.
- Islam F, Ali B, Wang J, Farooq MA, Gill RA et al. (2016). Combined herbicide and saline stress differentially modulates hormonal regulation and antioxidant defense system in *Oryza sativa* cultivars. *Plant Physiology and Biochemistry* 107:82-95.
- Jakovljević DZ, Topuzović MD, Stanković MS, Bojović BM (2017). Changes in antioxidant enzyme activity in response to salinity-induced oxidative stress during early growth of sweet basil. *Horticulture, Environment, and Biotechnology* 58:240-246.
- Jalil SU, Ansari MI (2020). Physiological role of gamma-aminobutyric acid in salt stress tolerance. In: *Salt and Drought Stress Tolerance in Plants*. Springer, pp 337-350.

- James RA, Blake C, Byrt CS, Munns R (2011). Major genes for Na⁺ exclusion, Nax1 and Nax2 (wheat HKT1; 4 and HKT1; 5), decrease Na⁺ accumulation in bread wheat leaves under saline and waterlogged conditions. *Journal of Experimental Botany* 62:2939-2947.
- Jamil M, Lee KJ, Kim JM, Kim HS, Rha ES (2007). Salinity reduced growth PS2 photochemistry and chlorophyll content in radish. *Scientia Agricola* 64:111-118.
- Jamshidi Goharrizi K, Baghizadeh A, Afroushteh M, Amirmahani F, Kermani SG (2020a). Effects of salinity stress on proline content and expression of Δ 1-pyrroline-5-carboxylate synthase and vacuolar-type H⁺ subunit E genes in wheat. *Plant Genetic Resources* 18:334-342.
- Jamshidi Goharrizi K, Moosavi SS, Amirmahani F, Salehi F, Nazari M (2020b). Assessment of changes in growth traits, oxidative stress parameters, and enzymatic and non-enzymatic antioxidant defense mechanisms in *Lepidium draba* plant under osmotic stress induced by polyethylene glycol. *Protoplasma* 257:459-473.
- Jamshidi Goharrizi K, Baghizadeh A, Kalantar M, Fatehi F (2020c). Assessment of changes in some biochemical traits and proteomic profile of UCB-1 pistachio rootstock leaf under salinity stress. *Journal of Plant Growth Regulation* 39:608-630.
- Jamshidi Goharrizi K, Amirmahani F, Salehi F (2020d). Assessment of changes in physiological and biochemical traits in four pistachio rootstocks under drought, salinity and drought+salinity stresses. *Physiologia Plantarum* 168:973-989.
- Jamshidi Goharrizi K, Fatehi F, Nazari M, Salehi F, Maleki M (2020e). Assessment of changes in the content of sulforaphane and expression levels of CYP79F1 and myrosinase genes and proteomic profile of *Lepidium draba* plant under water-deficit stress induced by polyethylene glycol. *Acta Physiologiae Plantarum* 42:101.
- Jamshidi Goharrizi K, Baghizadeh A, Kalantar M, Fatehi F (2020f). Combined effects of salinity and drought on physiological and biochemical characteristics of pistachio rootstocks. *Scientia Horticulturae* 261:108970.
- Jamshidi Goharrizi K, Riahi-Madvar A, Rezaee F, Pakzad R, Bonyad FJ et al. (2019a). Effect of salinity stress on enzymes' activity, ions concentration, oxidative stress parameters, biochemical traits, content of sulforaphane, and CYP79F1 gene expression level in *Lepidium draba* plant. *Journal of Plant Growth Regulation* 39:1075-1094.
- Jamshidi Goharrizi K, Amirmahani F, Fatehi F, Nazari M, Moosavi SS (2019b). Functional annotation of two hypothetical proteins (LOC105974023 and LOC109219975), as proteins involved in response to salinity: an in silico approach. *Journal of Genetic Resources* 5:72-82.
- Jamshidi Goharrizi K, Wilde HD, Amirmahani F, Moemeni MM, Zaboli M et al. (2018). Selection and validation of reference genes for normalization of qRT-PCR gene expression in wheat (*Triticum durum* L.) under drought and salt stresses. *Journal of Genetics* 97:1433-1444.
- Javot H, Lauvergeat V, Santoni V, Martin-Laurent F, Güçlü J et al. (2003). Role of a single aquaporin isoform in root water uptake. *The Plant Cell* 15:509-522.
- Jia H, Wang L, Li J, Sun P, Lu M et al. (2020). Comparative metabolomics analysis reveals different metabolic responses to drought in tolerant and susceptible poplar species. *Physiologia Plantarum* 168:531-546.
- Jones HG, Corlett JE (1992). Current topics in drought physiology. *The Journal of Agricultural Science* 119:291-296.
- Kahraman M, Sevim G, Bor M (2019). The role of proline, glycinebetaine, and trehalose in stress-responsive gene expression. In: *Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants*. Springer, pp 241-256.
- Kanani H, Dutta B, Klapa MI (2010). Individual vs. combinatorial effect of elevated CO₂ conditions and salinity stress on *Arabidopsis thaliana* liquid cultures: comparing the early molecular response using time-series transcriptomic and metabolomic analyses. *BMC Systems Biology* 4:177.
- Kanwal S, Rastogi RP, Incharoensakdi A (2014). Glutamate decarboxylase activity and gamma-aminobutyric acid content in *Synechocystis* sp. PCC 6803 under osmotic stress and different carbon sources. *Journal of Applied Phycology* 26:2327-2333.
- Karbudak T, Bor M, Özdemir F, Türkan İ (2014). Glycine betaine protects tomato (*Solanum lycopersicum*) plants at low temperature by inducing fatty acid desaturase7 and lipoxygenase gene expression. *Molecular Biology Reports* 41:1401-1410.
- Kaur G, Asthir B (2015). Proline: a key player in plant abiotic stress tolerance. *Biologia Plantarum* 59:609-619.
- Kaya C, Sonmez O, Aydemir S, Ashraf M, Dikilitas M (2013). Exogenous application of mannitol and thiourea regulates plant growth and oxidative stress responses in salt-stressed maize (*Zea mays* L.). *Journal of Plant Interactions* 8:234-241.
- Kerstiens G, Tych W, Robinson MF, Mansfield TA (2002). Sodium-related partial stomatal closure and salt tolerance of *Aster tripolium*. *New Phytologist* 153:509-515.
- Keunen E, Peshev D, Vangronsveld J, Van Den Ende W, Cuypers A (2013). Plant sugars are crucial players in the oxidative challenge during abiotic stress: extending the traditional concept. *Plant, Cell & Environment* 36:1242-1255.
- Khalid MF, Hussain S, Anjum MA, Ahmad S, Ali MA et al. (2020). Better salinity tolerance in tetraploid vs diploid volkamer lemon seedlings is associated with robust antioxidant and osmotic adjustment mechanisms. *Journal of Plant Physiology* 244:153071.
- Khedr AHA, Abbas MA, Wahid AAA, Quick WP, Abogadallah GM (2003). Proline induces the expression of salt-stress-responsive proteins and may improve the adaptation of *Pancreaticum maritimum* L. to salt-stress. *Journal of Experimental Botany* 54:2553-2562.
- Kim JM, To TK, Matsui A, Tanoi K, Kobayashi NI et al. (2017). Acetate-mediated novel survival strategy against drought in plants. *Nature Plants* 3:1-7.

- Kim SJ, Matsuo T, Watanabe M, Watanabe Y (2002). Effect of nitrogen and sulphur application on the glucosinolate content in vegetable turnip rape (*Brassica rapa* L.). *Soil Science and Plant Nutrition* 48:43-49.
- Kumar D, Al Hassan M, Naranjo MA, Agrawal V, Boscaiu M et al. (2017a). Effects of salinity and drought on growth, ionic relations, compatible solutes and activation of antioxidant systems in oleander (*Nerium oleander* L.). *PloS One* 12:e0185017.
- Kumar P, Sharma V, Atmaram CK, Singh B (2017b). Regulated partitioning of fixed carbon (14C), sodium (Na⁺), potassium (K⁺) and glycine betaine determined salinity stress tolerance of gamma irradiated pigeonpea [*Cajanus cajan* (L.) Millsp]. *Environmental Science and Pollution Research* 24:7285-7297.
- Kumar V, Shriram V, Hoque TS, Hasan MM, Burritt DJ et al. (2017c). Glycinebetaine-mediated abiotic oxidative-stress tolerance in plants: physiological and biochemical mechanisms. In: *Stress Signaling in Plants: Genomics and Proteomics Perspective*, Volume 2. Springer, pp 111-133.
- Kumari A, Parida AK (2018). Metabolomics and network analysis reveal the potential metabolites and biological pathways involved in salinity tolerance of the halophyte *Salvadora persica*. *Environmental and Experimental Botany* 148:85-99.
- Lai SJ, Lai MC, Lee RJ, Chen YH, Yen HE (2014). Transgenic *Arabidopsis* expressing osmolyte glycine betaine synthesizing enzymes from halophilic methanogen promote tolerance to drought and salt stress. *Plant Molecular Biology* 85:429-441.
- Li G, Wu H, Sun Y, Zhang S (2016a). Betaine aldehyde dehydrogenase (BADH) expression and betaine production in sugarbeet cultivars with different tolerances to drought stress. *Sugar Tech* 18:420-423.
- Li L, Gu W, Li J, Li C, Xie T et al. (2018a). Exogenously applied spermidine alleviates photosynthetic inhibition under drought stress in maize (*Zea mays* L.) seedlings associated with changes in endogenous polyamines and phytohormones. *Plant Physiology and Biochemistry* 129:35-55.
- Li L, Gu W, Li C, Li W, Li C et al. (2018b). Exogenous spermidine improves drought tolerance in maize by enhancing the antioxidant defence system and regulating endogenous polyamine metabolism. *Crop and Pasture Science* 69:1076-1091.
- Li M, Guo S, Yang X, Meng Q, Wei X (2016b). Exogenous gamma-aminobutyric acid increases salt tolerance of wheat by improving photosynthesis and enhancing activities of antioxidant enzymes. *Biologia Plantarum* 60:123-131.
- Li M, Guo R, Jiao Y, Jin X, Zhang H et al. (2017). Comparison of salt tolerance in soja based on metabolomics of seedling roots. *Frontiers in Plant Science* 8:1101.
- Li W, Wang Y, Zhang Y, Wang R, Guo Z et al. (2020). Impacts of drought stress on the morphology, physiology, and sugar content of Lanzhou lily (*Lilium davidii* var. unicolor). *Acta Physiologiae Plantarum* 42:1-11.
- Liang W, Ma X, Wan P, Liu L (2018). Plant salt-tolerance mechanism: a review. *Biochemical and Biophysical Research Communications* 495:286-291.
- Lima CS, Ferreira-Silva SL, Carvalho FEL, Neto MCL, Aragão RM et al. (2018). Antioxidant protection and PSII regulation mitigate photo-oxidative stress induced by drought followed by high light in cashew plants. *Environmental and Experimental Botany* 149:59-69.
- Liu CG, Wang QW, Jin YQ, Pan KW, Wang YJ (2017a). Photoprotective and antioxidative mechanisms against oxidative damage in *Fargesia rufa* subjected to drought and salinity. *Functional Plant Biology* 44:302-311.
- Liu H, Sun G, Dong L, Yang L, Yu S et al. (2017b). Physiological and molecular responses to drought and salinity in soybean. *Biologia Plantarum* 61:557-564.
- Liu Q, Arnold R, Yang S, Wu J, Li Z et al. (2019). Foliar application of exogenous polyamines to ameliorate drought-induced oxidative damage and physiological inhibition in *Toona ciliata* seedlings. *Australian Forestry* 82:139-150.
- Liu R, Sun W, Chao M, Ji C, Wang M et al. (2009). Leaf anatomical changes of *Bruguiera gymnorhiza* seedlings under salt stress. *Journal of Tropical and Subtropical Botany* 17:169-175.
- López-Aguilar R, Orduño-Cruz A, Lucero-Arce A, Murillo-Amador B, Troyo-Diéguez E (2003). Response to salinity of three grain legumes for potential cultivation in arid areas. *Soil Science and Plant Nutrition* 49:329-336.
- Ludlow MM, Muchow RC (1990) A critical evaluation of traits for improving crop yields in water-limited environments. In: Brady NC (ed) *Advances in Agronomy*. Academic Press, pp 107-153.
- Luo R, Song X, Li Z, Zhang A, Yan X et al. (2018). Effect of soil salinity on fructan content and polymerization degree in the sprouting tubers of Jerusalem artichoke (*Helianthus tuberosus* L.). *Plant Physiology and Biochemistry* 125:27-34.
- Mäkelä PS, Jokinen K, Himanen K (2019). Roles of endogenous glycinebetaine in plant abiotic stress responses. In: *Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants*. Springer, pp 153-173.
- Malekzadeh P, Khara J, Heydari R (2014). Alleviating effects of exogenous gamma-aminobutyric acid on tomato seedling under chilling stress. *Physiology and Molecular Biology of Plants* 20:133-137.
- Marček T, Hamow KÁ, Véghe B, Janda T, Darko E (2019). Metabolic response to drought in six winter wheat genotypes. *PloS One* 14:e0212411.
- Maury P, Berger M, Mojayad F, Planchon C (2000). Leaf water characteristics and drought acclimation in sunflower genotypes. *Plant and Soil* 223:155-162.
- Mei X, Chen Y, Zhang L, Fu X, Wei Q et al. (2016). Dual mechanisms regulating glutamate decarboxylases and accumulation of gamma-aminobutyric acid in tea (*Camellia sinensis*) leaves exposed to multiple stresses. *Scientific Reports* 6:23685.

- Mekonnen DW, Flüggé UI, Ludewig F (2016). Gamma-aminobutyric acid depletion affects stomata closure and drought tolerance of *Arabidopsis thaliana*. *Plant Science* 245:25-34.
- Mibei EK, Ambuko J, Giovannoni JJ, Onyango AN, Owino WO (2017). Carotenoid profiling of the leaves of selected African eggplant accessions subjected to drought stress. *Food Science & Nutrition* 5:113-122.
- Michaletti A, Naghavi MR, Toorchi M, Zolla L, Rinalducci S (2018). Metabolomics and proteomics reveal drought-stress responses of leaf tissues from spring-wheat. *Scientific Reports* 8:1-18.
- Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R (2010). Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant, Cell & Environment* 33:453-467.
- Misra AN, Latowski D, Strzalka K (2006). The xanthophyll cycle activity in kidney bean and cabbage leaves under salinity stress. *Russian Journal of Plant Physiology* 53:102-109.
- Misson L, Limousin JM, Rodriguez R, Letts MG (2010). Leaf physiological responses to extreme droughts in Mediterranean *Quercus ilex* forest. *Plant, Cell & Environment* 33:1898-1910.
- Moghaddam M, Farhadi N, Panjtandoust M, Ghanati F (2019). Seed germination, antioxidant enzymes activity and proline content in medicinal plant *Tagetes minuta* under salinity stress. *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology* 154:1-8.
- Mohammadi H, Ghorbanpour M, Brestic M (2018). Exogenous putrescine changes redox regulations and essential oil constituents in field-grown *Thymus vulgaris* L. under well-watered and drought stress conditions. *Industrial Crops and Products* 122:119-132.
- Morillon R, Chrispeels MJ (2001). The role of ABA and the transpiration stream in the regulation of the osmotic water permeability of leaf cells. *Proceedings of the National Academy of Sciences* 98:14138-14143.
- Mostafaei E, Zehtab-Salmasi S, Salehi-Lisar Y, Ghassemi-Golezani K (2018). Changes in photosynthetic pigments, osmolytes and antioxidants of Indian mustard by drought and exogenous polyamines. *Acta Biologica Hungarica* 69:313-324.
- Moustakas M, Sperdouli I, Kouna T, Antonopoulou CI, Therios I (2011). Exogenous proline induces soluble sugar accumulation and alleviates drought stress effects on photosystem II functioning of *Arabidopsis thaliana* leaves. *Plant Growth Regulation* 65:315.
- Mulholland BJ, Taylor IB, Jackson AC, Thompson AJ (2003). Can ABA mediate responses of salinity stressed tomato. *Environmental and Experimental Botany* 50:17-28.
- Munns R (1988). Why measure osmotic adjustment? *Functional Plant Biology* 15:717-726.
- Munns R (2005). Genes and salt tolerance: bringing them together. *New phytologist* 167:645-663.
- Mustard J, Renault S (2006). Response of red-osier dogwood (*Cornus sericea*) seedlings to NaCl during the onset of bud break. *Canadian Journal of Botany* 84:844-851.
- Najjaa H, Boubakri A, Arfa AB, Zouari N, Neffati M (2018). Salinity and drought stresses improve antioxidant potential of *Allium roseum* L., an edible medicinal plant. *Acta Physiologiae Plantarum* 40:195.
- Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y et al. (2014). Enhancement of oxidative and drought tolerance in *Arabidopsis* by overaccumulation of antioxidant flavonoids. *The Plant Journal* 77:367-379.
- Nakhaie A, Habibi G, Vaziri A (2020). Exogenous proline enhances salt tolerance in acclimated *Aloe vera* by modulating photosystem II efficiency and antioxidant defense. *South African Journal of Botany* 00:1-10.
- Nam KH, Shin HJ, Pack IS, Park JH, Kim HB et al. (2016). Metabolomic changes in grains of well-watered and drought-stressed transgenic rice. *Journal of the Science of Food and Agriculture* 96:807-814.
- Nawaz M, Wang Z (2020). Abscisic acid and glycine betaine mediated tolerance mechanisms under drought stress and recovery in *axonopus compressus*: a new insight. *Scientific Reports* 10:1-10.
- Nayyar H, Kaur R, Kaur S, Singh R (2014). γ -Aminobutyric acid (GABA) imparts partial protection from heat stress injury to rice seedlings by improving leaf turgor and upregulating osmoprotectants and antioxidants. *Journal of Plant Growth Regulation* 33:408-419.
- Nazari M, Goharrizi KJ, Moosavi SS, Maleki M (2019). Expression changes in the TaNAC2 and TaNAC69-1 transcription factors in drought stress tolerant and susceptible accessions of *Triticum boeoticum*. *Plant Genetic Resources* 17:471-479.
- Nazari M, Moosavi SS, Maleki M, Jamshidi Goharrizi K (2020). Chloroplastic acyl carrier protein synthase I and chloroplastic 20 kDa chaperonin proteins are involved in wheat (*Triticum aestivum*) in response to moisture stress. *Journal of Plant Interactions* 15:180-187.
- Netondo GW, Onyango JC, Beck E (2004). Sorghum and salinity: II. gas exchange and chlorophyll fluorescence of sorghum under salt stress. *Crop Science* 44:806-811.
- Nguyen G, Hailstones D, Wilkes M, Sutton B (2010). Drought stress: role of carbohydrate metabolism in drought-induced male sterility in rice anthers. *Journal of Agronomy and Crop Science* 196:346-357.
- Niazi B, Athar M, Salim M, Rozema J (2005). Growth and ionic relations of fodderbeet and seabet under saline environments. *International Journal of Environmental Science & Technology* 2:113-120.
- Nusrat N, Shahbaz M, Perveen S (2014). Modulation in growth, photosynthetic efficiency, activity of antioxidants and mineral ions by foliar application of glycinebetaine on pea (*Pisum sativum* L.) under salt stress. *Acta Physiologiae Plantarum* 36:2985-2998.
- O'Brien JA, Benková E (2013). Cytokinin cross-talking during biotic and abiotic stress responses. *Frontiers in Plant Science* 4:451.

- Otto MSG, Francisco JG, Gonzalez BT, de Almeida Calvo L, de Mattos EM et al. (2017). Changes in γ -aminobutyric acid concentration, gas exchange, and leaf anatomy in *Eucalyptus* clones under drought stress and rewatering. *Acta Physiologiae Plantarum* 39:208.
- Ouzounidou G, Giannakoula A, Ilias I, Zamanidis P (2016). Alleviation of drought and salinity stresses on growth, physiology, biochemistry and quality of two *Cucumis sativus* L. cultivars by Si application. *Brazilian Journal of Botany* 39:531-539.
- Pakzad R, Goharrizi KJ, Riahi-Madvar A, Amirmahani F, Mortazavi M et al. (2021). Identification of *Lepidium draba* Δ 1-pyrroline-5-carboxylate synthetase (P5CS) and assessment of its expression under NaCl stress: P5CS identification in *L. draba* plant. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences* 91:195-203.
- Pakzad R, Fatehi F, Kalantar M, Maleki M (2019). Evaluating the antioxidant enzymes activities, lipid peroxidation and proteomic profile changing in UCB-1 pistachio rootstock leaf under drought stress. *Scientia Horticulturae* 256:108617.
- Palma F, Tejera NA, Lluch C (2013). Nodule carbohydrate metabolism and polyols involvement in the response of *Medicago sativa* to salt stress. *Environmental and Experimental Botany* 85:43-49.
- Pandey S, Patel MK, Mishra A, Jha B (2015). Physio-biochemical composition and untargeted metabolomics of cumin (*Cuminum cyminum* L.) make it promising functional food and help in mitigating salinity stress. *PLoS One* 10:e0144469.
- Parida AK, Panda A, Rangani J (2018). Metabolomics-guided elucidation of abiotic stress tolerance mechanisms in plants. In: *Plant Metabolites and Regulation Under Environmental Stress*. Elsevier, pp 89-131.
- Peleg Z, Blumwald E (2011). Hormone balance and abiotic stress tolerance in crop plants. *Current Opinion in Plant Biology* 14:290-295.
- Poorter H, Remkes C (1990). Leaf area ratio and net assimilation rate of 24 wild species differing in relative growth rate. *Oecologia* 83:553-559.
- Rady MO, Semida WM, Abd El-Mageed TA, Hemida KA, Rady MM (2018). Up-regulation of antioxidative defense systems by glycine betaine foliar application in onion plants confer tolerance to salinity stress. *Scientia Horticulturae* 240:614-622.
- Rahnama A, James RA, Poustini K, Munns R (2010). Stomatal conductance as a screen for osmotic stress tolerance in durum wheat growing in saline soil. *Functional Plant Biology* 37:255-263.
- Rameeh V, Gerami M, Omran VG, Ghavampour S (2017). Impact of glycine betaine on salinity tolerance of stevia (*Stevia rebaudiana bertonii*) under in vitro condition. *Cercetari Agronomice in Moldova* 50:95-105.
- Ramos MLG, Gordon AJ, Minchin FR, Sprent JI, Parsons R (1999). Effect of water stress on nodule physiology and biochemistry of a drought tolerant cultivar of common bean (*Phaseolus vulgaris* L.). *Annals of Botany* 83:57-63.
- Rangani J, Panda A, Parida AK (2020). Metabolomic study reveals key metabolic adjustments in the xerohalophyte *Salvadora persica* L. during adaptation to water deficit and subsequent recovery conditions. *Plant Physiology and Biochemistry* 150:180-195.
- Rezaei-Chiyaneh E, Seyyedi SM, Ebrahimian E, Moghaddam SS, Damalas CA (2018). Exogenous application of gamma-aminobutyric acid (GABA) alleviates the effect of water deficit stress in black cumin (*Nigella sativa* L.). *Industrial Crops and Products* 112:741-748.
- Rhodes D, Hanson A (1993). Quaternary ammonium and tertiary sulfonium compounds in higher plants. *Annual Review of Plant Biology* 44:357-384.
- Rigui AB, Carvalho V, dos Santos ALW, Morvan-Bertrand A, Prud'Homme MP et al. (2019). Fructan and antioxidant metabolisms in plants of *Lolium perenne* under drought are modulated by exogenous nitric oxide. *Plant Physiology and Biochemistry* 145:205-215.
- Rodriguez P, Dell'Amico J, Morales D, Sanchez Blanco MdJ, Alarcón J (1997). Effect of salinity on growth, shoot water relations and root hydraulic conductivity in tomato plants. *The Journal of Agricultural Science* 128:439-444.
- Rosa M, Prado C, Podazza G, Interdonato R, González JA et al. (2009). Soluble sugars: metabolism, sensing and abiotic stress: a complex network in the life of plants. *Plant Signaling & Behavior* 4:388-393.
- Sahin U, Ekinci M, Ors S, Turan M, Yildiz S et al. (2018). Effects of individual and combined effects of salinity and drought on physiological, nutritional and biochemical properties of cabbage (*Brassica oleracea* var. capitata). *Scientia Horticulturae* 240:196-204.
- Sami F, Yusuf M, Faizan M, Faraz A, Hayat S (2016). Role of sugars under abiotic stress. *Plant Physiology and Biochemistry* 109:54-61.
- Sanchez DH, Schwabe F, Erban A, Udvardi MK, Kopka J (2012). Comparative metabolomics of drought acclimation in model and forage legumes. *Plant, Cell & Environment* 35:136-149.
- Saqib M, Zörb C, Schubert S (2006). Salt-resistant and salt-sensitive wheat genotypes show similar biochemical reaction at protein level in the first phase of salt stress. *Journal of Plant Nutrition and Soil Science* 169:542-548.
- Sarker U, Islam MT, Oba S (2018). Salinity stress accelerates nutrients, dietary fiber, minerals, phytochemicals and antioxidant activity in *Amaranthus tricolor* leaves. *PLoS One* 13:e0206388.
- Schneider JR, Caverzan A, Chavarria G (2019). Water deficit stress, ROS involvement, and plant performance. *Archives of Agronomy and Soil Science* 65:1160-1181.
- Semida WM, Abdelkhalik A, Rady MO, Marey RA, Abd El-Mageed TA (2020). Exogenously applied proline enhances growth and productivity of drought stressed onion by improving photosynthetic efficiency, water use efficiency and up-regulating osmoprotectants. *Scientia Horticulturae* 272:109580.

- Shackel KA, Foster KW, Hall AE (1982). Genotypic differences in leaf osmotic potential among grain sorghum cultivars grown under irrigation and drought. *Crop Science* 22:1121-1125.
- Shams M, Yildirim E, Ekinçi M, Turan M, Dursun A et al. (2016). Exogenously applied glycine betaine regulates some chemical characteristics and antioxidative defence systems in lettuce under salt stress. *Horticulture, Environment, and Biotechnology* 57:225-231.
- Sharbatkhari M, Shobbar ZS, Galeshi S, Nakhoda B (2016). Wheat stem reserves and salinity tolerance: molecular dissection of fructan biosynthesis and remobilization to grains. *Planta* 244:191-202.
- Sheikh-Mohamadi MH, Etemadi N, Nikbakht A, Arab M, Majidi MM et al. (2017). Antioxidant defence system and physiological responses of Iranian crested wheatgrass (*Agropyron cristatum* L.) to drought and salinity stress. *Acta Physiologiae Plantarum* 39:245.
- Sheteiwy MS, Shao H, Qi W, Hamoud YA, Shaghaleh H et al. (2019). GABA-alleviated oxidative injury induced by salinity, osmotic stress and their combination by regulating cellular and molecular signals in rice. *International Journal of Molecular Sciences* 20:5709.
- Shi H, Ye T, Song B, Qi X, Chan Z (2015). Comparative physiological and metabolomic responses of four *Brachypodium distachyon* varieties contrasting in drought stress resistance. *Acta Physiologiae Plantarum* 37:122.
- Singh M, Kumar J, Singh S, Singh VP, Prasad SM (2015). Roles of osmoprotectants in improving salinity and drought tolerance in plants: a review. *Reviews in Environmental Science and Bio/Technology* 14:407-426.
- Slama I, Abdelly C, Bouchereau A, Flowers T, Savoure A (2015). Diversity, distribution and roles of osmoprotective compounds accumulated in halophytes under abiotic stress. *Annals of Botany* 115:433-447.
- Sobieszczuk-Nowicka E, Legocka J (2014). Plastid-associated polyamines: their role in differentiation, structure, functioning, stress response and senescence. *Plant Biology* 16:297-305.
- Sofy MR, Elhawat N, Alshaal T (2020). Glycine betaine counters salinity stress by maintaining high K^+/Na^+ ratio and antioxidant defense via limiting Na^+ uptake in common bean (*Phaseolus vulgaris* L.). *Ecotoxicology and Environmental Safety* 200:110732.
- Steduto P, Albrizio R, Giorio P, Sorrentino G (2000). Gas-exchange response and stomatal and non-stomatal limitations to carbon assimilation of sunflower under salinity. *Environmental and Experimental Botany* 44:243-255.
- Sultana N, Ikeda T, Itoh R (1999). Effect of NaCl salinity on photosynthesis and dry matter accumulation in developing rice grains. *Environmental and Experimental Botany* 42:211-220.
- Sun H, Luo M, Zhou X, Zhou Q, Sun Y et al. (2020). Exogenous glycine betaine treatment alleviates low temperature-induced pericarp browning of 'Nanguo' pears by regulating antioxidant enzymes and proline metabolism. *Food Chemistry* 306:125626.
- Sun J, Lu N, Xu H, Maruo T, Guo S (2016). Root zone cooling and exogenous spermidine root-pretreatment promoting *Lactuca sativa* L. growth and photosynthesis in the high-temperature season. *Frontiers in Plant Science* 7:368.
- Szabados L, Savoure A (2010). Proline: a multifunctional amino acid. *Trends in Plant Science* 15:89-97.
- Taffouo V, Wamba O, Youmbi E, Nono G, Akoa A (2010). Growth, yield, water status and ionic distribution response of three bambara groundnut (*Vigna subterranea* (L.) Verdc.) landraces grown under saline conditions. *International Journal of Botany* 6:53-58.
- Taffouo VD, Kouamou JK, Ngalangue LMT, Ndjedji BAN, Akoa A (2009). Effects of salinity stress on growth, ions partitioning and yield of some cowpea (*Vigna unguiculata* L. Walp.) cultivars. *International Journal of Botany* 5:135-143.
- Takahashi S, Murata N (2008). How do environmental stresses accelerate photoinhibition? *Trends in Plant Science* 13:178-182.
- Takahashi T, Kakeji J-I (2010). Polyamines: ubiquitous polycations with unique roles in growth and stress responses. *Annals of Botany* 105:1-6.
- Tang J, Camberato JJ, Yu X, Luo N, Bian S et al. (2013). Growth response, carbohydrate and ion accumulation of diverse perennial ryegrass accessions to increasing salinity. *Scientia Horticulturae* 154:73-81.
- Tardieu F, Simonneau T (1998). Variability among species of stomatal control under fluctuating soil water status and evaporative demand: modelling isohydric and anisohydric behaviours. *Journal of Experimental Botany* 49:419-432.
- Theerakulpisut P, Phongngarm S (2013). Alleviation of adverse effects of salt stress on rice seedlings by exogenous trehalose. *Asian Journal of Crop Science* 5:405-415.
- Tian F, Wang W, Liang C, Wang X, Wang G et al. (2017). Over accumulation of glycine betaine makes the function of the thylakoid membrane better in wheat under salt stress. *The Crop Journal* 5:73-82.
- Traversari S, De Carlo A, Traversi ML, Minnocci A, Francini A et al. (2020). Osmotic adjustments support growth of poplar cultured cells under high concentrations of carbohydrates. *Plant Cell Reports* 39:1-12.
- Trovato M, Forlani G, Signorelli S, Funck D (2019). Proline metabolism and its functions in development and stress tolerance. In: *Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants*. Springer, pp 41-72.
- Trovato M, Mattioli R, Costantino P (2008). Multiple roles of proline in plant stress tolerance and development. *Rendiconti Lincei* 19:325-346.
- Tsugane K, Kobayashi K, Niwa Y, Ohba Y, Wada K et al. (1999). A recessive *Arabidopsis* mutant that grows photoautotrophically under salt stress shows enhanced active oxygen detoxification. *The Plant Cell* 11:1195-1206.
- Turan MA, Katkat V, Taban S (2007). Salinity-induced stomatal resistance, proline, chlorophyll and. *International Journal of Agricultural Research* 2:483-488.

- Turner NC (1986). Adaptation to water deficits: a changing perspective. *Functional Plant Biology* 13:175-190.
- Van den Ende W, Valluru R (2009). Sucrose, sucrosyl oligosaccharides, and oxidative stress: scavenging and salvaging? *Journal of Experimental Botany* 60:9-18.
- Veselov DS, Sharipova GV, Veselov SU, Kudoyarova GR (2008). The effects of NaCl treatment on water relations, growth, and aba content in barley cultivars differing in drought tolerance. *Journal of Plant Growth Regulation* 27:380-386.
- Vighi I, Benitez L, Amaral M, Moraes G, Auler P et al. (2017). Functional characterization of the antioxidant enzymes in rice plants exposed to salinity stress. *Biologia Plantarum* 61:540-550.
- Vijayakumari K, Puthur JT (2016). γ -Aminobutyric acid (GABA) priming enhances the osmotic stress tolerance in *Piper nigrum* Linn. plants subjected to PEG-induced stress. *Plant Growth Regulation* 78:57-67.
- Wang C, Fan L, Gao H, Wu X, Li J et al. (2014). Polyamine biosynthesis and degradation are modulated by exogenous gamma-aminobutyric acid in root-zone hypoxia-stressed melon roots. *Plant Physiology and Biochemistry* 82:17-26.
- Wang G-P, Hui Z, Li F, Zhao M-R, Zhang J et al. (2010). Improvement of heat and drought photosynthetic tolerance in wheat by overaccumulation of glycinebetaine. *Plant Biotechnology Reports* 4:213-222.
- Wang L, Huang X, Lim DJ, Laserna AKC, Li SFY (2019a). Uptake and toxic effects of triphenyl phosphate on freshwater microalgae *Chlorella vulgaris* and *Scenedesmus obliquus*: insights from untargeted metabolomics. *Science of the Total Environment* 650:1239-1249.
- Wang N, Cao F, Richmond MEA, Qiu C, Wu F (2019b). Foliar application of betaine improves water-deficit stress tolerance in barley (*Hordeum vulgare* L.). *Plant Growth Regulation* 89:109-118.
- Wang W, Vinocur B, Altman A (2003). Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta* 218:1-14.
- Wang X, Yang P, Gao Q, Liu X, Kuang T et al. (2008). Proteomic analysis of the response to high-salinity stress in *Physcomitrella patens*. *Planta* 228:167-177.
- Wang Y, Gu W, Meng Y, Xie T, Li L et al. (2017a). γ -Aminobutyric acid imparts partial protection from salt stress injury to maize seedlings by improving photosynthesis and upregulating osmoprotectants and antioxidants. *Scientific Reports* 7:1-13.
- Wang Y, Huang Y-Y, Wang Y, Lyu P, Hamblin MR (2017b). Red (660 nm) or near-infrared (810 nm) photobiomodulation stimulates, while blue (415 nm), green (540 nm) light inhibits proliferation in human adipose-derived stem cells. *Scientific Reports* 7:1-10.
- Wani SH, Kumar V, Shriram V, Sah SK (2016). Phytohormones and their metabolic engineering for abiotic stress tolerance in crop plants. *The Crop Journal* 4:162-176.
- Wei D, Zhang W, Wang C, Meng Q, Li G et al. (2017). Genetic engineering of the biosynthesis of glycinebetaine leads to alleviate salt-induced potassium efflux and enhances salt tolerance in tomato plants. *Plant Science* 257:74-83.
- Wei GP, Yang LF, Zhu YL, Chen G (2009). Changes in oxidative damage, antioxidant enzyme activities and polyamine contents in leaves of grafted and non-grafted eggplant seedlings under stress by excess of calcium nitrate. *Scientia Horticulturae* 120:443-451.
- White DA, Turner NC, Galbraith JH (2000). Leaf water relations and stomatal behavior of four allopatric *Eucalyptus* species planted in Mediterranean southwestern Australia. *Tree Physiology* 20:1157-1165.
- Wolters H, Jürgens G (2009). Survival of the flexible: hormonal growth control and adaptation in plant development. *Nature Reviews Genetics* 10:305-317.
- Woodrow P, Ciarmiello LF, Annunziata MG, Pacifico S, Iannuzzi F et al. (2017). Durum wheat seedling responses to simultaneous high light and salinity involve a fine reconfiguration of amino acids and carbohydrate metabolism. *Physiologia Plantarum* 159:290-312.
- Wutipraditkul N, Wongwan P, Buaboocha T (2015). Alleviation of salt-induced oxidative stress in rice seedlings by proline and/or glycinebetaine. *Biologia Plantarum* 59:547-553.
- Xiang L, Hu L, Xu W, Zhen A, Zhang L et al. (2016). Exogenous γ -aminobutyric acid improves the structure and function of photosystem II in muskmelon seedlings exposed to salinity-alkalinity stress. *PLoS One* 11:e0164847.
- Xiong Y, Qu Y, Han H, Chen F, Li L et al. (2021). Unraveling physiological and metabolomic responses involved in *Phlox subulata* L. tolerance to drought stress. *Plant Molecular Biology Reporter* 39:98-111.
- Xu Z, Sun M, Jiang X, Sun H, Dang X et al. (2018). Glycinebetaine biosynthesis in response to osmotic stress depends on jasmonate signaling in watermelon suspension cells. *Frontiers in Plant Science* 9:1469.
- Yan J, Tang H, Huang Y, Shi Y, Li J et al. (2003). Dynamic analysis of QTL for plant height at different developmental stages in maize (*Zea mays* L.). *Chinese Science Bulletin* 48:2601-2607.
- Yildiztugay E, Ozfidan-Konakci C, Kucukoduk M, Duran Y (2014). Variations in osmotic adjustment and water relations of *Sphaerophysa kotschyana*: glycine betaine, proline and choline accumulation in response to salinity. *Botanical Studies* 55:1-9.
- Yin YG, Kobayashi Y, Sanuki A, Kondo S, Fukuda N et al. (2010). Salinity induces carbohydrate accumulation and sugar-regulated starch biosynthetic genes in tomato (*Solanum lycopersicum* L. cv. 'Micro-Tom') fruits in an ABA-and osmotic stress-independent manner. *Journal of Experimental Botany* 61:563-574.
- Yoon YE, Kuppasamy S, Cho KM, Kim PJ, Kwack YB et al. (2017). Influence of cold stress on contents of soluble sugars, vitamin C and free amino acids including gamma-aminobutyric acid (GABA) in spinach (*Spinacia oleracea*). *Food Chemistry* 215:185-192.

- Yordanov I, Velikova V, Tsonev T (2000). Plant responses to drought, acclimation, and stress tolerance. *Photosynthetica* 38:171-186.
- You L, Song Q, Wu Y, Li S, Jiang C et al. (2019). Accumulation of glycine betaine in transplastomic potato plants expressing choline oxidase confers improved drought tolerance. *Planta* 249:1963-1975.
- Youssef SM, Abd Elhady SA, Aref RM, Riad GS (2018). Salicylic acid attenuates the adverse effects of salinity on growth and yield and enhances peroxidase isozymes expression more competently than proline and glycine betaine in cucumber plants. *Gesunde Pflanzen* 70:75-90.
- Zaher-Ara T, Boroomand N, Sadat-Hosseini M (2016). Physiological and morphological response to drought stress in seedlings of ten citrus. *Trees* 30:985-993.
- Zali AG, Ehsanzadeh P (2018). Exogenously applied proline as a tool to enhance water use efficiency: case of fennel. *Agricultural Water Management* 197:138-146.
- Zhang M, Smith JAC, Harberd NP, Jiang C (2016). The regulatory roles of ethylene and reactive oxygen species (ROS) in plant salt stress responses. *Plant Molecular Biology* 91:651-659.
- Zhang Y, Li Z, Li YP, Zhang XQ, Ma X et al. (2018). Chitosan and spermine enhance drought resistance in white clover, associated with changes in endogenous phytohormones and polyamines, and antioxidant metabolism. *Functional Plant Biology* 45:1205-1222.
- Zhao GQ, Ma BL, Ren CZ (2007a). Growth, gas exchange, chlorophyll fluorescence, and ion content of naked oat in response to salinity. *Crop Science* 47:123-131.
- Zhao XX, Liang C, Fang Y, Wang YQ, Wang W (2007b). Effect of glycinebetaine on function of thylakoid membranes in wheat flag leaves under drought stress. *Biologia Plantarum* 51:584-588.
- Zhong M, Song R, Wang Y, Shu S, Sun J et al. (2020). TGase regulates salt stress tolerance through enhancing bound polyamines-mediated antioxidant enzymes activity in tomato. *Environmental and Experimental Botany* 179:104191.
- Zhu C, Schraut D, Hartung W, Schäffner AR (2005). Differential responses of maize MIP genes to salt stress and ABA. *Journal of Experimental Botany* 56:2971-2981.
- Zlatko Stoyanov Z (2005). Effects of water stress on leaf water relations of young bean plants. *Journal of Central European Agriculture* 6:5-14.