

1-1-2021

The curative effects of dietary yeast beta-1,3/1,6-glucan on oxidative stress and apoptosis in laying quails (*Coturnix coturnix japonica*) exposed to lead

İSMAİL SEVEN

PINAR TATLI SEVEN


SEDA İFLAZOĞLU MUTLU

GÖZDE ARKALI

NURGÜL BİRBEN

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>

 Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

Recommended Citation

SEVEN, İSMAİL; SEVEN, PINAR TATLI; MUTLU, SEDA İFLAZOĞLU; ARKALI, GÖZDE; BİRBEN, NURGÜL; ARSLAN, ASLIHAN SUR; and AKSAKAL, MESUT (2021) "The curative effects of dietary yeast beta-1,3/1,6-glucan on oxidative stress and apoptosis in laying quails (*Coturnix coturnix japonica*) exposed to lead," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 45: No. 1, Article 4. <https://doi.org/10.3906/vet-2007-25>

Available at: <https://journals.tubitak.gov.tr/veterinary/vol45/iss1/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The curative effects of dietary yeast beta-1,3/1,6-glucan on oxidative stress and apoptosis in laying quails (*Coturnix coturnix japonica*) exposed to lead

Authors

İSMAİL SEVEN, PINAR TATLI SEVEN, SEDA İFLAZOĞLU MUTLU, GÖZDE ARKALI, NURGÜL BİRBEN, ASLIHAN SUR ARSLAN, and MESUT AKSAKAL

The curative effects of dietary yeast beta-1,3/1,6-glucan on oxidative stress and apoptosis in laying quails (*Coturnix coturnix japonica*) exposed to lead

İsmail SEVEN^{1*}, Pinar TATLI SEVEN², Seda İFLAZOĞLU MUTLU²,
Gözde ARKALI³, Nurgül BİRBEN⁴, Aslıhan SUR ARSLAN⁵, Mesut AKSAKAL³

¹Department of Plant and Animal Production, Vocational School of Sivrice, Firat University, Elazığ, Turkey

²Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, Firat University, Elazığ, Turkey

³Department of Physiology, Faculty of Veterinary Medicine, Firat University, Elazığ, Turkey

⁴The Ministry of Food, Agriculture and Livestock, Veterinary Control Institute Office, Elazığ, Turkey

⁵Department of Veterinary Medicine, Vocational School of Kepsut, Balıkesir, Turkey

Received: 06.07.2020 • Accepted/Published Online: 03.10.2020 • Final Version: 23.02.2021

Abstract: The objective of this study was to investigate the effects of supplementation of beta-glucan to the basal diet of laying quails exposed to lead (Pb) toxicity. A total of 112 birds were randomly divided into 4 groups. The first group was fed a corn-soybean based basal diet without any supplementation (control). The second (Pb) and third (beta-glucan) groups were supplemented with 100 mg/kg of Pb (as Pb acetate) and 100 mg/kg of beta-glucan, respectively. The fourth group (Pb + beta-glucan) was fed a basal diet supplemented with 100 mg/kg of Pb and 100 mg/kg of beta-glucan. It was determined that serum total protein, albumin, and creatinine values of the Pb + beta-glucan group partially improved ($P < 0.05$) with the addition of beta-glucan, but globulin and blood urea nitrogen (BUN) values were similar to the Pb group ($P < 0.001$). Globulin and albumin/globulin ratio were not affected by the supplements. Aspartate aminotransferase (AST) ($P < 0.01$) and alanine transaminase (ALT) ($P < 0.001$) enzyme activities in the beta-glucan and Pb + beta-glucan groups were similar to the control group, unlike the Pb group. However, ALP enzyme activity was similar in all supplemented groups, unlike the control group ($P < 0.01$). Malondialdehyde ($P < 0.001$) was reduced in the liver, heart, and kidney tissues of beta-glucan supplemented groups compared to the Pb group. Glutathione levels of liver and heart tissues in the Pb + beta-glucan group were significantly higher than in the Pb group ($P < 0.001$). Catalase and glutathione peroxidase activities were also significantly increased in the liver tissue of the beta-glucan group compared to the Pb group. The caspase-3 and caspase-9 protein expression levels in the liver tissue of the beta-glucan group decreased ($P < 0.001$) compared to the Pb group. As a result, we conclude that beta-glucan helps reduce the harmful effects of Pb toxicity in laying quails.

Key words: Beta-glucan, lead toxicity, oxidative damage, apoptosis, quail

1. Introduction

Lead (Pb), which is found naturally in the earth, is the most important toxic metal and also one of the most prevalent heavy metal contaminants in the world [1,2]. Pb threatens human and animal life in many ways, especially during the development stages [3]. Hematopoietic, kidney, reproductive, and central nervous systems are parts of the human body that are vulnerable to exposure to high levels of Pb [4]. Pb poisoning is a real threat to public health, especially in developing countries. Therefore, great efforts are being made to reduce the dangers of the metal in terms of occupational and public health. Animals can easily be exposed to Pb because it is a global environmental pollutant, primarily in the industrial regions [4].

Beta-glucans are carbohydrates consisting of glucose chains in the cell wall of yeast, fungi, bacteria, and cereals [5]. Beta-1,3/1,6-glucan is formed as a result of the formation of glucose monomers by chaining with beta-1,3 glycosidic bonds and by the binding of glucose monomers to these chains by beta-1,6 linkages. Beta-glucans can be found in many organisms, from bacteria to trees and from seeds to fungi. Beta-1,3/1,6-glucan from polysaccharides, which has different functions, helps the mechanical wall hardness and integrity of the cell wall [6–8]. Beta-glucan has several beneficial health outcomes such as antibacterial, antitumor, antiviral, enhancing immunity, and wound healing activities [9]. Some features of beta-glucans, such as molecular weight, degree of branching, solubility, primary structure, and polymer charge can

* Correspondence: iseven@firat.edu.tr

change its effect on the immune system and biological activity [10].

Pb poisoning in animals can usually occur from a large number of environmental sources and can be monitored via contamination of soil and feed because of industrial pollution and agricultural applications. A wide range of toxic effects of Pb in animal systems is observed. Poisoning of wild birds following ingestion of Pb from ammunition in relation to hunting activities has been recognized. Pb taken in this way endangers the growth and survival of nestlings, causes adverse effects on reproduction by reducing egg production and plasma calcium level, causes hemolytic anemia and behavioral impairments [11,12]. Exposure to high doses of Pb can result in reproductive disorders, poor performance, oxidative stress, apoptosis, poisoning, or even death in animals [10,13–19]. Pb can cause oxidative damage to the reproductive organs, kidney, heart, and liver. The mechanism of oxidative stress induced by Pb involves the effects of Pb on the antioxidant defense systems, membranes, and DNA of cells [3,20]. While new cells are formed in living organisms, some of the existing cells are eliminated by cell death, thereby maintaining a constant balance. These existing cells are destroyed by various types of cell death, such as physiological, programmed cell death (apoptosis), and pathological cell death (necrosis). Reactive oxygen species (ROS) can react with macromolecules due to oxidation and cause them to undergo apoptosis or necrosis [21,22]. Beta-glucans show their antioxidant effects as a strong intracellular free radical scavenger [23]. Free oxygen radicals cause lipid peroxidation and consequently membrane damage and destruction [24]. Beta-1,3-glucan has a protective effect against oxidative damage by suppressing lipid peroxidation. Also, it controls lipid peroxidation due to inhibiting the increased levels of malondialdehyde (MDA), which is an indication of lipid peroxidation [25]. It directs the immune system to recognize and destroy macrophages in mutated cells. Beta-glucans show their effects in tissue regeneration and repair by accelerating the repair of damaged tissue. They show their adjuvant effects by increasing antibacterial, antifungal, antiviral, and antiparasitic effects [6].

Nowadays, many researchers focus on the effects of natural dietary supplements on oxidative stress and apoptosis, and the relationship between oxidative stress and apoptosis. The aim of this study was to examine the effects of beta-glucan, which has various natural biological functions, on laying quails exposed to Pb toxicity, which has become increasingly widespread during today's industrialization period.

2. Materials and methods

2.1. Supplements

Beta-1,3/1,6-glucan originating from *Saccharomyces cerevisiae* containing at least 70% beta-1,3/1,6-glucan was

purchased from Pure Bulk Inc. (Roseburg, OR, USA). Pb (II) acetate trihydrate $[(CH_3COO)_2Pb \cdot 3H_2O]$ was purchased from Merck (CAS #: 6080-56-4).

2.2. Animal, diet, and experimental design

A total of 112 laying quails (*Coturnix coturnix japonica*) were obtained from a local commercial company. Each group consisted of 28 female animals aged 5 weeks (each with four replicates consisting of 7 birds). The quails were randomly divided into 4 groups, which had 4 replicates including 7 quails in each group. The research was carried out using standard cages (34 × 62.5 × 23 cm from the CIMUKA company) designed for laying quails with the approval of the Local Ethics Committee for Animal Experiments of the Ministry of Food, Agriculture, and Livestock (05.29.2018/2018-1). Water and feed were provided ad libitum. The photoperiod was 16/8 h (light/dark). The birds were kept in quail layer cages in a temperature-controlled room. A basal diet prepared according to National Research Center [26] was supplied by a commercial company (Table 1). The experimental groups were arranged as follows: fed a corn-soybean basal diet and no supplementation (group I; control), supplementation with 100 mg/kg of Pb (as Pb (II) acetate trihydrate) to the diet (group II; Pb), supplementation with 100 mg/kg of beta-glucan to basal diet (group III; beta-glucan), and supplementation with 100 mg/kg of Pb and 100 mg/kg of beta-glucan to basal diet (group IV; Pb + beta-glucan). At the end of the experiment (after 56 days), 6 quails from each group were slaughtered for analysis by decapitation.

Research studies on Pb toxicity in quail were used for determining the dosage of Pb [27,28]. The dose of beta-glucan was determined according to Moon et al. [29] and Zhu and Wu [30].

2.3. Oxidative stress

2.3.1. Sample collection and homogenate preparation

After the quails were slaughtered, liver, heart, and kidney tissues were washed with phosphate buffer (PBS) and each was wrapped in aluminum foil, placed in polyethylene bags, and labeled. The tissues were kept at -20 °C until analysis. For the analysis, the tissues were weighed and transferred to glass tubes while maintaining the coldness. Tris buffer (pH 7.4) was added to the tissues in a 1/10 ratio. While maintaining coldness, the tissue was homogenized in the homogenizer. The tissue total protein was determined by Lowry et al. [31] method, which used bovine serum albumin as the standard.

2.3.2. Determination of malondialdehyde (MDA) level

The amount of MDA produced in the liver, heart, and kidney tissues was used as an indicator of lipid peroxidation level. The colored complex of MDA formed by thiobarbituric acid (TBA) was measured at 532 nm. The

Table 1. Ingredients and nutrient composition of basal ration.

Ingredients	%	Nutritional composition	%
Maize	66.27	Dry matter	89.9
Soybean meal (44% CP)	24.44	Crude protein	17.0
Wheat bran	1.31	Crude cellulose	3.31
Salt	0.25	Ether extract	1.89
L-Lysine hydrochloride	0.21	Crude ash	9.78
L-Threonine	0.13	Calcium ^b	2.50
Sodium bicarbonate	0.10	Available phosphorus ^b	0.35
DL-Methionine	0.12	Sodium ^b	0.16
Vitamin-Mineral premix ^a	0.32	Lysine ^b	1.00
Limestone	5.46	Threonine ^b	0.75
Calcium phosphate	1.39	ME, kcal/kg ^{b, c}	2800
Total	100		

^aVitamin-mineral premix (per 1 kg): vitamin A, 8000 IU; vitamin D3, 3000 IU; vitamin E, 25 IU; menadione, 1.5 mg; vitamin B12, 0.02 mg; biotin, 0.1 mg; folacin, 1 mg; niacin, 50 mg; pantothenic acid, 15 mg; pyridoxine, 4 mg; riboflavin, 10 mg; thiamin, 3 mg; copper (copper sulphate), 10 mg; iodine (ethylenediamine dihydriodide), 1.0 mg; iron (ferrous sulphate monohydrate), 50 mg; manganese (manganese sulphate monohydrate), 60 mg; zinc (zinc sulphate monohydrate), 60 mg; selenium (sodium selenite), 0.42 mg.

^bCalculated.

^cMetabolizable energy (ME) = 53 + 38 [(Crude protein, %) + (2.25 x Ether extract, %) + (1.1 x Starch, %) + (Sugar, %)]

MDA levels were determined by the spectrophotometric method defined by Placer et al. [32].

2.3.3. Tissue glutathione (GSH) level

The GSH level was determined by the spectrophotometric method of Sedlak and Lindsay [33]. Since the color intensity of the colored complex formed by the 5,5'-dithio-bis-(2-nitrobenzoic acid) (DTNB) was directly proportional to the concentration of GSH in the environment, the samples were measured at 412 nm.

2.3.4. Measurement of glutathione peroxidase (GSH-Px) levels in tissue

The level of GSH-Px activity in the liver, heart, and kidney tissues was determined by the spectrophotometric method described by Lawrence and Burk [34]. The yellow color complex that was formed as a result of mixing the samples with DTNB solution was measured at 412 nm.

2.3.5. Measurement of catalase (CAT) enzyme level in tissue

The CAT activity in tissue was determined by the spectrophotometric method described by Goth [35]. When the tissue is incubated with the substrate containing hydrogen peroxide (H₂O₂), H₂O₂ is cleaved to H₂O and O₂ by catalase activity. The ammonium molybdate added to the medium combines with H₂O₂ to terminate the reaction.

During this period, the color change was measured at 405 nm.

2.4. Western blotting

Tissues were homogenized with cold RIPA lysis buffer (Santa Cruz Biotechnology, Sc24948A), centrifuged at 14,000 rpm at 4 °C for 45 min, and the supernatants were separated. Total protein content was determined spectrophotometrically according to the BCA (Thermo Fisher, 23227) method [36]. Samples were electrophoresed by loading 60 µg of protein in each well, and the first well with a protein marker, to the polyacrylamide gel (stacking gel 4%; separating gel 12%) [37]. After sodium dodecyl sulphate-polyacrylamide gel electrophoresis, specific proteins were transferred to polyvinylidene difluoride (PVDF) membrane by western blotting [38]. The membranes of PVDF were blocked by 5% bovine serum albumin (BSA), and then washed 3 times with the Tris buffered saline Tween 20 (TBST) to prevent nonspecific binding for 5 min. After blocking, the membranes were incubated with primary antibodies caspase-9 (ab69514; 1/500), caspase-3 (ab90437; 1/500), and beta-actin (ab8226; 1/5000) overnight. After incubation, the membranes were washed 3 times with TBST for 5 min. Subsequently, the bands that were obtained by chemiluminescent conjugate (ECL; Bio-Rad, 1705060) on membranes treated with the appropriate

secondary antibody (ab97240; 1/10,000, ab6721; 1/10,000) were visualized in the chemiluminescence imaging system (Bio-Rad ChemiDoc™ XRS+) [39,40]. The band intensities in images were measured with the appropriate analysis system (Bio-Rad Image Lab™ Software, version 5.2.1, Bio-Rad Laboratories, Inc., USA). Relative expression levels of protein were normalized to beta-actin, which was used as an internal control.

2.5. Statistical analyses

All values were presented as mean \pm SE. Whether the values obtained as a result of the study show a normal distribution was determined by the Shapiro–Wilk test. As a result of Shapiro–Wilk normality analysis, it was determined that the data showed normal distribution. The differences between the mean of groups were detected using one-way ANOVA and the posthoc Duncan test in IBM SPSS Statistics 22 computer program [41]. The results were noted as significant at $P < 0.05$.

3. Results

When the effects of beta-glucans on the serum biochemical parameters of the experimental groups were examined, it was seen that serum total protein, albumin, and creatinine values of Pb + beta-glucan group were partially improved ($P < 0.05$) by the addition of beta-glucan, but globulin and BUN values were similar to Pb group ($P < 0.001$). Globulin and albumin/globulin ratio were not affected by supplements (Table 2). Serum AST ($P < 0.01$) and ALT ($P < 0.001$) activities of beta-glucan and Pb + beta-glucan groups were similar to the control and the group different than the Pb group, but ALP activities of the control group were higher than in the other groups ($P < 0.01$; Table 2).

The MDA, GSH, GSH-Px, and CAT values of kidney, liver, and heart tissues are presented in Table 3. The highest MDA levels in all tissues were in the Pb group ($P < 0.001$). The MDA levels of heart, kidney, and liver tissues of the beta-glucan group were similar to the control, and also the supplementation of beta-glucan in the Pb + beta-glucan group significantly reduced the MDA level ($P < 0.001$). The GSH, GSH-Px, and CAT values of kidney, heart, and liver tissues in the Pb group were significantly lower than in the other groups ($P < 0.05$). Similar to the control group, caspase-3 (Figure 1) and caspase-9 (Figure 2) protein expression levels of the beta-glucan group decreased according to the Pb group in liver tissue ($P < 0.001$).

4. Discussion

Birds can reduce the amount of metal residue in their eggs by reducing the accumulation of the mineral. This type of protection may be enough to prevent the accumulation of some metals such as Mn and Cr but is inadequate for Pb [42]. Therefore, birds are very sensitive to Pb exposure. Oxidative stress has been reported to be the main mechanism of toxicity induced by Pb [16,17,27,43]. Oxidative stress formed by the effect of Pb occurs due to two different mechanisms working at the same time. The first one is the formation of ROS, like singlet oxygen, hydroperoxides, and H_2O_2 , and the second one is the depletion of the antioxidant reserves [44,45]. Apoptosis is a type of programmed cell death regulated by Bcl-2 and caspase protein families [46].

Recent studies show that Pb can cause oxidative stress by increasing oxidation production, suppressing the activities of antioxidant enzymes, and eventually activating

Table 2. Effects of beta-glucan on serum biochemical parameters of experimental groups.

Parameters	Control	Pb	Beta-glucan	Pb + Beta-glucan	P
Glu (mg/dL)	139.50 \pm 5.90 ^b	190.50 \pm 12.89 ^a	149.00 \pm 10.96 ^b	177.83 \pm 5.96 ^a	**
TP (g/dL)	3.08 \pm 0.10 ^a	2.42 \pm 0.09 ^b	2.82 \pm 0.18 ^{ab}	2.87 \pm 0.20 ^{ab}	*
Albumin (g/dL)	1.52 \pm 0.09 ^a	1.15 \pm 0.06 ^b	1.30 \pm 0.10 ^{ab}	1.33 \pm 0.08 ^{ab}	*
Globulin (g/dL)	1.57 \pm 0.09	1.27 \pm 0.04	1.52 \pm 0.09	1.53 \pm 0.14	NS
A/G ratio	1.00 \pm 0.12	0.90 \pm 0.04	0.86 \pm 0.02	0.90 \pm 0.06	NS
Cre (mg/dL)	0.02 \pm 0.00 ^b	0.04 \pm 0.01 ^a	0.02 \pm 0.00 ^b	0.03 \pm 0.00 ^{ab}	*
BUN (mg/dL)	11.67 \pm 1.09 ^b	19.70 \pm 1.58 ^a	12.20 \pm 1.24 ^b	16.25 \pm 1.08 ^a	**
Enzymes					
AST (U/L)	294.17 \pm 10.27 ^b	382.60 \pm 14.98 ^a	296.33 \pm 18.89 ^b	300.40 \pm 20.24 ^b	**
ALT (U/L)	4.25 \pm 0.16 ^b	7.75 \pm 0.70 ^a	3.40 \pm 0.24 ^b	3.20 \pm 0.48 ^b	***
ALP (U/L)	562.40 \pm 19.95 ^a	400.17 \pm 25.48 ^b	465.75 \pm 28.92 ^b	445.17 \pm 17.30 ^b	**

Pb: lead; Glu: glucose; TP: total protein; Cre: creatinine; A/G: albumin/globulin ratio; BUN: blood urea nitrogen; AST: aspartate aminotransferase; ALT: alanine transaminase; ALP: alkaline phosphatase; ^{a,b}: mean \pm SE values with different superscripts within a row differ significantly; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; NS: nonsignificant

Table 3. Malondialdehyde (MDA, nmol/g), glutathione (GSH, nmol/g), glutathione peroxidase (GSH-Px, IU/g protein) and catalase (CAT, kU/g protein) values of kidney, heart, and liver tissues of experimental groups.

Parameters		Control	Pb	Beta-glucan	Pb + Beta-glucan	P
Kidney	MDA	30.90 ± 1.33 ^c	39.40 ± 1.77 ^a	29.71 ± 1.22 ^c	35.25 ± 1.12 ^b	***
	GSH	2.47 ± 0.11 ^a	2.12 ± 0.29 ^b	2.54 ± 0.26 ^a	2.33 ± 0.12 ^{ab}	*
	GSH-Px	22.32 ± 1.83 ^a	16.80 ± 1.74 ^b	21.95 ± 0.96 ^a	20.19 ± 1.18 ^a	*
	CAT	5.32 ± 0.18 ^a	3.41 ± 0.55 ^b	5.94 ± 0.50 ^a	5.24 ± 0.61 ^a	*
Heart	MDA	15.67 ± 1.47 ^c	24.31 ± 0.80 ^a	15.18 ± 0.92 ^c	20.88 ± 1.30 ^b	***
	GSH	3.69 ± 0.02 ^a	2.78 ± 0.12 ^c	3.78 ± 0.15 ^a	3.08 ± 0.03 ^b	***
	GSH-Px	24.90 ± 1.63 ^a	18.91 ± 1.24 ^b	25.02 ± 2.19 ^a	23.81 ± 1.15 ^a	*
	CAT	10.53 ± 0.88 ^a	4.98 ± 0.38 ^c	10.50 ± 0.46 ^a	7.62 ± 0.17 ^b	***
Liver	MDA	28.04 ± 1.22 ^c	48.82 ± 1.69 ^a	27.60 ± 1.75 ^c	37.59 ± 1.71 ^b	***
	GSH	5.06 ± 0.26 ^a	3.02 ± 0.20 ^c	5.68 ± 0.38 ^a	3.98 ± 0.24 ^b	***
	GSH-Px	33.00 ± 2.83 ^a	22.77 ± 0.78 ^b	33.20 ± 1.03 ^a	30.38 ± 2.01 ^a	**
	CAT	9.22 ± 0.53 ^a	2.35 ± 0.06 ^c	9.81 ± 0.50 ^a	4.79 ± 0.59 ^b	***

Pb: lead; ^{a,b,c}: mean ± SE values with different superscripts within a row differ significantly; * P < 0.05; ** P < 0.01; *** P < 0.001




Figure 1. Liver tissue caspase-3 protein expression levels (P < 0.001; mean ± SE).

apoptosis [18,47]. Kofuji et al. [48] reported that beta-glucan from barley has higher antioxidant activity than gelatin, pectin from apple and citrus, curdlan, dextrin, chitosan, pullulan, gellan gum, and sodium alginate.

Bahakaim et al. [49] used Glycomoss (foreign product) or Mox (local product) as a source of mannan oligosaccharides (MOS) and beta-glucans (28% MOS and 34% beta-glucans) 0.5 and 0.75 g/kg in quail diet from 7 to 13 weeks of age. In MOS and beta-glucans supplemented

groups, when compared with the control, they reported that the plasma total protein and globulin were significantly increased, but the A/G ratio was significantly decreased (P < 0.05). Also, the plasma concentration of albumin and the activities of AST and ALT enzymes were not affected in this research. Similarly, Mousa et al. [50] reported that dietary MOS plus beta-glucans supplementation (0.5 and 0.75 g/kg) increased the plasma total protein, globulin, and glucose, and decreased the A/G ratio (P < 0.05). However,




Figure 2. Liver tissue caspase-9 protein expression levels ($P < 0.001$; mean \pm SE).

plasma albumin, AST, and ALT values were not affected by dietary beta-glucan. Unlike our study, Bahakaim et al. [49] and Mousa et al. [50] reported that supplementation with beta-glucan and MOS did not change the plasma albumin, AST, and ALT values, increased the globulin and total protein levels and decreased the A/G ratio. This may be attributed to exposure to Pb toxicity or to the effect of MOS and/or MOS and glucan interaction. The organism naturally exhibits defense behavior against Pb-induced oxidative stress and apoptosis [18,47]. Since one of the organs most affected by this condition is the liver, there is an increase in liver enzymes AST and ALT activities. Similar to our study, Alagawany et al. [27] reported that Pb (100 mg/kg) supplementation to the diet of quails for 8 weeks has significantly decreased total protein, albumin, and globulin, and increased ALT and AST enzyme activities ($P < 0.001$). Also, Seven et al. [16] reported that the addition of 200 mg/kg Pb to broiler diet for 42 days as compared to the control group has significantly increased AST activity ($P < 0.05$), but there was no effect on glucose, ALP, and albumin. Similarly, Yuan et al. [19], in their study, in which the basal diet (corn-soybean meal) of Hy-Line Brown hens was supplemented with 15, 30, and 60 mg/kg of Pb, reported that the supplementation with Pb at 60 mg/kg resulted in increased levels of BUN, urea acid, and AST activity ($P < 0.05$).

Seven et al. [16] reported that the 200 mg/kg of Pb supplementation to broiler diet significantly increased the MDA level in plasma ($P < 0.01$) and liver tissue ($P < 0.05$), and the GSH values in plasma and heart tissue ($P < 0.01$). In addition to the above effects, they reported that

Pb supplementation increased the superoxide dismutase (SOD) activity of plasma and CAT activities of hemolysate, liver, and heart tissues. The water-soluble yeast beta-1,3-glucan, which can participate in the body's natural defense mechanisms, has a perfect radical scavenging effect [10]. By means of exhibiting an action positively correlated with its dose, water-soluble yeast beta-1,3-glucan may increase SOD activity in serum [10,51]. Yeast beta-glucan has a powerful antioxidative effect that can increase antioxidant activity by removing free radicals, preventing the activity of ROS and antioxidants, and increasing the body's antioxidant capacity [10,52,53]. In a study investigating the protective role of beta-glucan against oxidative damage caused by sepsis in rats, Şener et al. [25] gave 50 mg/kg beta-glucan to rats once a day by intragastric gavage for 10 days. They reported that the application of beta-glucan according to the saline-treated group significantly decreased the MDA levels in the brain ($P < 0.05$), liver, heart, lung, diaphragm ($P < 0.01$), and kidney tissues ($P < 0.001$). Moreover, beta-glucan use prevented the decline of GSH levels in the brain, heart, lung, diaphragm, kidney ($P < 0.05$), and liver tissues ($P < 0.001$). In a study that examined the effects of 0, 50, 250, 500, and 1000 mg/kg Pb solution on liver histology and oxidative stress for 49 days, Kou et al. [43] reported that the activities of GSH-Px, SOD (only 500 mg/kg), CAT enzymes, and total antioxidant capacity significantly decreased in liver tissues of Pb groups, but MDA levels increased. They also noted that Pb exposure caused damages to bird health by inducing ultrastructural and microstructural injury, lipid metabolism disorder, and oxidative damage in liver tissue.

Alagawany et al. [27] reported that supplementation of Pb (100 mg/kg) to the diet of quails for 8 weeks increased serum MDA level compared to the control ($P < 0.001$), decreased the SOD and CAT ($P < 0.001$). However, they reported that the GSH level of Pb and control groups was similar ($P < 0.001$). In a study that explored whether or not the mannan/beta-glucans from the yeast cell wall (BYCW) (0, 0.2, 1, and 5 mg/mL) have a protective role in suppressing the cellular ROS in intestinal porcine epithelial cells from the jejunum (IPEC-J2) cells damaged by deoxynivalenol (DON; 1 μ M), Guo et al. [54] reported that treatment with BYCW in the attack of DON, which was a *Fusarium* mycotoxin, caused a drop in MDA and ROS and a rise of GSH in IPEC-J2 cells. Moreover, they reported that there were similar improvements in MDA, ROS, and GSH levels in the cells after treatment with CPDT (25 μ M) and BYCW (0.2, 1, 5 mg/mL) for 12 and 24 h in the IPEC-J2 cells under DON attack. All these research reports are compatible with our study. Kofuji et al. [48] reported that beta-glucan from barley has higher antioxidant activity than oat and black yeast-derived beta-glucan and they also reported that this effect is directly proportional to beta-glucan density. In addition, they reported that the hydroxyl radical scavenging activity of beta-glucans extracted under alkali or acid conditions was slightly higher than extracted in neutral conditions with warm water, and this activity was reduced with a drop in their molecular size.

Caspase-3 and caspase-9 have been noted to play a significant role in the formation phase of apoptosis caused by different stimuli [18,55]. Xu et al. [18] reported that Pb could induce an increase in caspase-3 activity, and the apoptosis induced by Pb occurs via the activation of apoptosis practitioners such as caspase-3. In a study that investigated the cytotoxic effect of Pb in human

leukemia (HL-60) cells, which were cultured with lead nitrate for 24 h, Yedjou et al. [47] reported that Pb at very low concentration has a similar effect with control on the viability of HL-60 cells, but significantly decreased between 25 and 50 μ g/mL of lead nitrate ($P < 0.05$). Also, they noted that the apoptosis of HL-60 cells induced by Pb was associated with the activation of caspase-3, and the numbers of caspase-3 positive cells (apoptotic cells) were significantly increased in all doses (10, 20, 30, and 40 μ g/mL) of lead nitrate ($P < 0.05$). These research results are in line with ours.

5. Conclusion

Research on beta-glucan added to poultry diets focuses on its effects on the immune system, but there is no research on its effects on oxidative stress and/or apoptosis in the literature. Here, we demonstrated that the addition of Pb, which is known as the inducer of oxidative stress and apoptosis, to the diet of laying quails had a negative effect on some blood parameters (glucose, total protein, albumin, creatinine, and BUN) and liver enzymes (AST, ALT, and ALP). In addition, Pb caused oxidative stress in the kidney, heart, and liver tissues. Beta-glucan was able to ameliorate these adverse effects. Moreover, Pb sensitized liver cells to apoptosis through the activation of caspase-3 and caspase-9, but this effect was suppressed by beta-glucan. To increase the profitability in poultry farming, the effects of beta-glucan, a natural product, on oxidative stress and apoptosis need to be investigated in future research.

Acknowledgments/disclaimers/ conflict of interest

This research was funded by the Firat University Scientific Research Projects Unit (project number: SMYO.19.01).

The authors of this research declare that there is no conflict of interest in the publication of this article.

References

1. Tchounwou PB, Yedjou CG, Patlolla AK, Sutton DJ. Heavy Metals toxicity and the environment. In: Luch A (editor). *Molecular, Clinical and Environmental Toxicology* (Volume 3: Environmental Toxicology). Berlin, Germany: Springer, Basel; 2012. pp. 133-164.
2. Wani AL, Ara A, Usmani JA. Lead toxicity: a review. *Interdisciplinary Toxicology* 2015; 8 (2): 55-64. doi: 10.1515/intox-2015-0009
3. Sharma R, Panwar K, Mogra S. Effects of prenatal and neonatal exposure to lead on white blood cells in Swiss mice. *Journal of Cell and Molecular Biology* 2012; 10 (1): 33-40.
4. Assi MA, Hezmee MN, Haron AW, Sabri MY, Rajion MA. The detrimental effects of lead on human and animal health. *Veterinary World* 2016; 9 (6): 660-671. doi: 10.14202/vetworld.2016.660-671
5. Volman JJ, Ramakers D, Plat J. Dietary modulation of immune function by beta-glucans. *Physiology & Behavior* 2009; 94 (2): 276-284. doi: 10.1016/j.physbeh.2007.11.045
6. Bashir KMI, Choi JS. Clinical and physiological perspectives of β -glucans: the past, present, and future. *International Journal of Molecular Sciences* 2017; 18 (9):1906. doi: 10.3390/ijms18091906
7. Kaur R, Sharma M, Ji D, Xu M, Agyei D. Structural features, modification, and functionalities of beta-glucan. *Fibers* 2020; 8 (1): 1. doi: 10.3390/fib8010001
8. Zhu F, Du B, Xu B. A critical review on production and industrial applications of beta-glucans. *Food Hydrocolloid* 2016; 52: 275-288. doi: 10.1016/j.foodhyd.2015.07.003

9. Wang Q, Sheng X, Shi A, Hu H, Yang Y et al. β -Glucans: Relationships between modification, conformation and functional activities. *Molecules* 2017; 22 (2): 257. doi:10.3390/molecules22020257
10. Yuan H, Lan P, He Y, Li C, Ma X. Effect of the modifications on the physicochemical and biological properties of β -glucan—a critical review. *Molecules* 2020; 25 (1): 57. doi: 10.3390/molecules25010057.
11. Lucia M, Andre JM, Gontier K, Diot N, Veiga J et al. Trace element concentrations (mercury, cadmium, copper, zinc, lead, aluminium, nickel, arsenic, and selenium) in some aquatic birds of the Southwest Atlantic Coast of France. *Archives of Environmental Contamination and Toxicology* 2010; 58 (3): 844-853. doi: 10.1007/s00244-009-9393-9
12. Pain DJ, Mateo R, Green RE. Effects of lead from ammunition on birds and other wildlife: A review and update. *Ambio* 2019; 48 (9): 935-953. doi: 10.1007/s13280-019-01159-0.
13. Ahmed WM, Abdel-Hameed AR, Moghazy FME. Some reproductive and health aspects of female buffaloes in relation to blood lead concentration. *International Journal of Dairy Science* 2008; 3 (2): 63-70. doi:10.3923/ijds.2008.63.70
14. Burki TK. Nigeria's lead poisoning crisis could leave a long legacy. *Lancet* 2012; 379 (9818): 792. doi: 10.1016/s0140-6736(12)60332-8.
15. McDowell LR. *Minerals in Animal and Human Nutrition*. 2nd ed. Amsterdam, Netherlands: Elsevier Science BV; 2003.
16. Seven I, Aksu T, Tatli Seven P. The effects of propolis on biochemical parameters and activity of antioxidant enzymes in broilers exposed to lead-induced oxidative stress. *Asian Australasian Journal of Animal Sciences* 2010; 23 (11): 1482-1489. doi: 10.5713/ajas.2010.10009
17. Seven I, Aksu T, Tatli Seven P. The effects of propolis and vitamin C supplemented feed on performance, nutrient utilization and carcass characteristics in broilers exposed to lead. *Livestock Science* 2012; 148 (1-2): 10-15. doi:10.1016/j.livsci.2012.05.001
18. Xu LH, Mu FF, Zhao JH, He Q, Cao CL et al. Lead induces apoptosis and histone hyperacetylation in rat cardiovascular tissues. *PLoS ONE* 2015; 10 (6): e0129091. doi:10.1371/journal.pone.0129091
19. Yuan C, Song HH, Jiang YJ, Azzam MMM, Zhu S et al. Effects of lead contamination in feed on laying performance, lead retention of organs and eggs, protein metabolism, and hormone levels of laying hens. *The Journal of Applied Poultry Research* 2013; 22 (4): 878-884. doi: 10.3382/japr.2013-00801
20. Ahamed M, Siddiqui MKJ. Low level lead exposure and oxidative stress: Current opinions. *Clinica Chimica Acta* 2007; 383 (1-2): 57-64. doi: 10.1016/j.cca.2007.04.024
21. Kannan K, Jain SK. Oxidative stress and apoptosis. *Pathophysiology* 2000; 7 (3): 153-163. doi:10.1016/S0928-4680(00)00053-5
22. Loh KP, Huang SH, De Silva R, Tan BKH, Zhu YZ. Oxidative stress: apoptosis in neuronal injury. *Current Alzheimer Research* 2006; 3 (4): 327-337. doi: 10.2174/156720506778249515
23. Kayali H, Ozdag MF, Kahraman S, Aydin A, Gonul E et al. The antioxidant effect of beta-glucan on oxidative stress status in experimental spinal cord injury in rats. *Neurosurgical Review*. 2005; 28 (4): 298-302. doi: 10.1007/s10143-005-0389-2
24. Tatli Seven P, Yilmaz S, Seven I, Tuna Kelestemur G. Effects of propolis in animals exposed oxidative stress. In: Lushchak VI (editor). *Oxidative Stress - Environmental Induction and Dietary Antioxidants*. 1st ed. Rijeka, Croatia: InTECH; 2012. pp. 267-288.
25. Şener G, Toklu H, Ercan F, Erkanlı G. Protective effect of β -glucan against oxidative organ injury in rat model of sepsis. *International Immunopharmacology* 2005; 5 (9): 1387-1396. doi: 10.1016/j.intimp.2005.03.007
26. NRC (National Research Council). *Nutrient Requirement of Poultry*. 9th ed. Washington DC, USA: National Academy Press, 1994.
27. Alagawany M, Abd El-Hack ME, Farag MR, Elnesr SS, El-Kholy MS et al. Dietary supplementation of *yucca schidigera* extract enhances productive and reproductive performances, blood profile, immune function, and antioxidant status in laying Japanese quails exposed to lead in the diet. *Poultry Science* 2018; 97 (9): 3126-3137. doi: 10.3382/ps/pey186
28. Farag MR, Alagawany M, Abd El-Hack ME, El-Sayed SAA, Ahmed SYA et al. *Yucca schidigera* extract modulates the lead-induced oxidative damage, nephropathy and altered inflammatory response and glucose homeostasis in Japanese quails. *Ecotoxicology and Environmental Safety* 2018; 156: 311-321. doi: 10.1016/j.ecoenv.2018.03.010
29. Moon SH, Lee I, Feng X, Lee HY, Kim J et al. Effect of dietary Beta-glucan on the performance of broilers and the quality of broiler breast meat. *Asian-Australasian Journal of Animal Sciences* 2016; 29 (3): 384-389. doi: 10.5713/ajas.15.0141
30. Zhu M, Wu S. The growth performance and nonspecific immunity of loach *paramisgurnus dabryanus* as affected by dietary β -1,3-glucan. *Fish & Shellfish Immunology* 2018; 83: 368-372. doi: 10.1016/j.fsi.2018.09.049
31. Lowry OH, Rosenbrough NJ, Farr AL, Randal RJ. Protein measurement with the folin phenol reagent. *The Journal of Biological Chemistry* 1951; 193 (1): 265-275.
32. Placer ZA, Cushman LL, Johnson BC. Estimation of product of lipid peroxidation (malonyldialdehyde) in biochemical systems. *Analytical Biochemistry* 1966; 16 (2): 359-364. doi:10.1016/0003-2697(66)90167-9
33. Sedlak J, Lindsay RH. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Analytical Biochemistry* 1968; 25 (1): 192-205. doi:10.1016/0003-2697(68)90092-4
34. Lawrence RA, Burk RF. Glutathione peroxidase activity in selenium-deficient rat liver. *Biochemical and Biophysical Research Communications* 1976; 71 (1): 952-958. doi: 10.1016/0006-291x(76)90747-6
35. Goth LA. Simple method for determination of serum catalase activity and revision of reference range. *Clinica Chimica Acta* 1991; 196 (2-3): 143-151. doi: 10.1016/0009-8981(91)90067-M

36. Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH et al. Measurement of protein using bicinchoninic acid. *Analytical Biochemistry* 1985; 150 (1): 76-85. doi: 10.1016/0003-2697(85)90442-7
37. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227 (5259): 680-685. doi:10.1038/227680a0
38. Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences of the United States of America* 1979; 76 (9): 4350-4354. doi: 10.1073/pnas.76.9.4350
39. Kielkopf CL, Bauer W, Urbatsch IL. Expressing cloned genes for protein production, purification and analysis. In: Green MR, Sambrook J (editors). *Molecular Cloning: A Laboratory Manual*. 4thed. New York, USA: Cold Spring Harbor Laboratory Press; 2012. pp. 1599-1625.
40. Bass JJ, Wilkinson DJ, Rankin D, Phillips BE, Szewczyk NJ et al. An overview of technical considerations for Western blotting applications to physiological research. *Scandinavian Journal of Medicine and Science in Sports* 2017; 27 (1): 4-25. doi: 10.1111/sms.12702
41. SPSS. IBM SPSS Statistics for Windows, version 22.0. NY, Armonk, USA: IBM; 2013.
42. Hui CA. Concentrations of chromium, manganese, and lead in air and in avian eggs. *Environmental Pollution* 2002; 120 (2): 201-206. doi:10.1016/S0269-7491(02)00158-6
43. Kou H, Ya J, Gao X, Zhao H. The effects of chronic lead exposure on the liver of female Japanese quail (*Coturnix japonica*): Histopathological damages, oxidative stress and AMP-activated protein kinase based lipid metabolism disorder. *Ecotoxicology and Environmental Safety* 2020; 190:110055. doi: 10.1016/j.ecoenv.2019.110055
44. Flora SJS. Nutritional components modify metal absorption, toxic response and chelation therapy. *Journal of Nutritional & Environmental Medicine* 2009; 12 (1): 53-67. doi:10.1080/13590840220123361
45. Flora G, Gupta D, Tiwari A. Toxicity of lead: A review with recent updates. *Interdisciplinary Toxicology* 2012; 5 (2): 47-58. doi: 10.2478/v10102-012-0009-2
46. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH. Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biology* 2013; 14 (1): 32. doi: 10.1186/1471-2121-14-32.
47. Yedjou CG, Milner JN, Howard CB, Tchounwou BP. Basic apoptotic mechanisms of lead toxicity in human leukemia (HL-60) cells. *International Journal of Environmental Research and Public Health* 2010; 7 (5): 2008-2017. doi:10.3390/ijerph7052008
48. Kofuji K, Aoki A, Tsubaki K, Konishi M, Isobe Y et al. Antioxidant Activity of β -Glucan. *ISRN Pharmaceutics* 2012; 2012:125864. doi: 10.5402/2012/125864
49. Bahakaim ASA, Mousa SMM, Soliman MM. Effect of mannan oligosaccharides and β -glucans on productive performance, egg quality and blood biochemical and hematological parameters of laying Japanese quail. *Egyptian Poultry Science Journal*, 2015; 35 (4): 1109-1122.
50. Mousa SMM, Soliman MM, Bahakaim ASA. Effect of mannan oligosaccharides and β -glucans on productive performance and some physiological and immunological parameters of growing Japanese quail chicks. *Egyptian Poultry Science Journal*, 2014; 34 (2): 433-451.
51. Gu K, Zhao L, Wang L, Bu D, Liu N et al. Effects of dietary supplementation of yeast beta-glucan on performance, serum biochemical indices and antioxidant capacity of transition dairy cows. *Chinese Journal of Animal Nutrition* 2018; 30 (6): 2164-2171.
52. Shan-bai DHX, Xiao-bo HU. Study on antitumor and antioxidant activities of yeast beta-1,3-glucan hydrolysates in vivo. *Journal of Huazhong Agricultural University* 2007; 26 (6): 871-874.
53. Tang Q, Huang G. Progress in polysaccharide derivatization and properties. *Mini Reviews in Medicinal Chemistry* 2016; 16 (15): 1244-1257. doi: 10.2174/1389557516999160612164003
54. Guo W, Gu X, Tong Y, Wang X, Wu J et al. Protective effects of mannan/ β -glucans from yeast cell wall on the deoxyriyalenol-induced oxidative stress and autophagy in IPEC-J2 cells. *International Journal of Biological Macromolecules* 2019; 135: 619-629. doi:10.1016/j.ijbiomac.2019.05.180
55. Lockshin RA. Programmed cell death: history and future of a concept. *Journal de la Société de Biologie* 2005; 199 (63): 169-173. doi: 10.1051/jbio:2005017