

1-1-2021

A regularized trace formula for "weighted" Sturm-Liouville equation with point delta-interaction

MANAF MANAFI

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

MANAFI, MANAF (2021) "A regularized trace formula for "weighted" Sturm-Liouville equation with point delta-interaction," *Turkish Journal of Mathematics*: Vol. 45: No. 4, Article 22. <https://doi.org/10.3906/mat-2101-34>

Available at: <https://journals.tubitak.gov.tr/math/vol45/iss4/22>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A regularized trace formula for "weighted" Sturm-Liouville equation with point δ - interaction

Manaf Dzh. MANAFOV^{1,2,*†} 

¹Department of Mathematics, Faculty of Arts and Sciences, Adiyaman University, Adiyaman, Turkey

²Institute of Mathematics and Mechanics, Azerbaijan National Academy of Science, Baku, Azerbaijan

Received: 13.01.2021

Accepted/Published Online: 30.05.2021

Final Version: 27.07.2021

Abstract: In this study, we obtain a formula for the regularized trace formula for "weighted" Sturm–Liouville equation with point δ - interaction. At the beginning, for the correct determination of solutions of analyzed equation at the point of discontinuity, the matching conditions are required. As a result, an equation is derived for the eigenvalues of the differential operator given in this study.

Key words: "weighted" Sturm–Liouville equation, regularized trace formula, point δ -interaction

1. Introduction

The theory of regularized trace of ordinary differential operators has a long history. Gelfand and Levitan [6] first obtained the trace formula for the Sturm–Liouville differential equation. After this study, several mathematicians were interested in developing trace formulas for different differential operators. The current situation of this subject and studies related to it are presented in the comprehensive survey paper [13]. The regularized trace for differential equations are found in [4, 5, 7–9]. However, there is a small number of words on the regularized trace for Sturm–Liouville operators with singular potentials (see [14–16]). Note that, the trace formulas have applications in the approximate calculation of the eigenvalues of the related operator [4, 13].

We consider the boundary value problem (BVP) for the differential equation

$$ly := -y'' + q(x)y = \lambda y, \quad x \in \left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right), \quad (1.1)$$

with the boundary conditions

$$U(y) := y'(0) = 0, \quad V(y) := y(\pi) = 0, \quad (1.2)$$

and conditions at the point $x = \frac{\pi}{2}$,

$$I(y) := \begin{cases} y\left(\frac{\pi}{2} + 0\right) = y\left(\frac{\pi}{2} - 0\right) \equiv y\left(\frac{\pi}{2}\right), \\ y'\left(\frac{\pi}{2} + 0\right) - y'\left(\frac{\pi}{2} - 0\right) = -\alpha \lambda y\left(\frac{\pi}{2}\right), \end{cases} \quad (1.3)$$

where $q(x)$ is real-valued function in $W_2^1(0, \pi)$ and $\alpha > 0$; λ is spectral parameter.

*Correspondence: mmanafov@adiyaman.edu.tr

†In memory of M. G. Gasymov (1939–2008)

2010 AMS Mathematics Subject Classification: 34A55, 34B24, 47E05

Notice that, we can understand problem (1.1),(1.3) as studying the equation

$$-y'' + q(x)y = \lambda\rho(x)y, \quad x \in (0, \pi), \tag{1.4}$$

when $\rho(x) = 1 + \alpha\delta(x - \frac{\pi}{2})$, where $\delta(x)$ is the Dirac function (see [1]). In this aspect, various inverse spectral problems for the equation (1.4) have been investigated in [12].

In the present paper, after construction of the Hilbert space related to (1.4), we obtain the formula of the first order regularized trace for "weighted" Sturm–Liouville equation with point δ - interaction.

2. Construction of the Hilbert space related to the problem and some properties of its spectral characteristics

In the Hilbert space $\aleph := L_2[0, \pi] \oplus \mathbb{C}$ of two component vectors, we define an inner product by

$$\langle f, g \rangle_{\aleph} := \int_0^{\pi} f_1(x)\bar{g}_1(x) dx + \frac{1}{\alpha}f_2\bar{g}_2 \tag{2.1}$$

for

$$f = \begin{pmatrix} f_1(x) \\ f_2 \end{pmatrix}, \quad g = \begin{pmatrix} g_1(x) \\ g_2 \end{pmatrix},$$

where $f_1(x), g_1(x) \in L_2(0, \pi)$, $f_2, g_2 \in \mathbb{C}$. In the Hilbert space \aleph we define the operator L

$$L : \aleph \rightarrow \aleph$$

with domain

$$D(L) : \left\{ f \in \aleph \mid \begin{array}{l} f_1, f_1' \in AC\left(\left(0, \frac{\pi}{2}\right) \cup \left(\frac{\pi}{2}, \pi\right)\right), \quad lf_1 \in L_2\left[\left(0, \pi\right) \setminus \left\{\frac{\pi}{2}\right\}\right], \\ f_2 = \alpha f_1\left(\frac{\pi}{2}\right), \quad U(f_1) = V(f_1) = 0 \end{array} \right\}$$

and operator rule

$$L(f) := \begin{pmatrix} lf_1 \\ f_1'(\frac{\pi}{2} - 0) - f_1'(\frac{\pi}{2} + 0) \end{pmatrix}.$$

Here $AC(\cdot)$ denotes the set of all absolutely continuous functions on related interval. In particular, those functions will have limits at the point $\frac{\pi}{2}$.

It is clear that the eigenvalues of the operator L and the BVP (1.1)-(1.3) are same and eigenfunctions of (1.1)-(1.3) coincide with the first component of corresponding eigenelements of the operator L .

Theorem 2.1 *The operator L is symmetric.*

Proof Let $f, g \in D(L)$. From the inner product defined in (2.1), we obtain

$$\begin{aligned} \langle Lf, g \rangle_{\aleph} - \langle f, Lg \rangle_{\aleph} &= \left[W\left(f, \bar{g}; \frac{\pi}{2} - 0\right) - W\left(f, \bar{g}; 0\right) \right] \\ &\quad + \left[W\left(f, \bar{g}; \pi\right) - W\left(f, \bar{g}; \frac{\pi}{2} + 0\right) \right] \\ &\quad + \frac{1}{\alpha} \left[\alpha W\left(f, \bar{g}; \frac{\pi}{2} + 0\right) - \alpha W\left(f, \bar{g}; \frac{\pi}{2} - 0\right) \right], \end{aligned}$$

where $W(f, g; x) = f(x)g'(x) - f'(x)g(x)$ is the wronskian of the functions f and g . Since f and \bar{g} satisfy the same boundary conditions (1.2) and from the conditions at the point $x = \frac{\pi}{2}$, we obtain $\langle Lf, g \rangle_{\mathbb{R}} = \langle f, Lg \rangle_{\mathbb{R}}$ for $f, g \in D(L)$. So L is symmetric. \square

Corollary 2.2 *The function $W(f, g; x)$ is continuous on $(0, \pi)$.*

Corollary 2.3 *All eigenvalues of the problem (1.1)-(1.3) are real and if λ_1 and λ_2 are two different eigenvalues of the problem (1.1)-(1.3), the corresponding eigenfunctions $y_1(x)$ and $y_2(x)$ are orthogonal in the sense of*

$$\int_0^{\pi} y_1(x)y_2(x) dx + \frac{1}{\alpha}y_1\left(\frac{\pi}{2}\right)y_2\left(\frac{\pi}{2}\right) = 0.$$

Let $\varphi(x, \lambda), \psi(x, \lambda), C(x, \lambda), S(x, \lambda)$ be solutions of (1.1) under the initial conditions

$$\begin{aligned} C(0, \lambda) &= S'(0, \lambda) = \varphi(0, \lambda) = \psi'(0, \lambda) = 1, \\ C'(0, \lambda) &= S(0, \lambda) = \varphi'(0, \lambda) = \psi(\pi, \lambda) = 0, \end{aligned}$$

and under the conditions (1.3).

For each fixed x , the functions $\varphi(x, \lambda), \psi(x, \lambda), C(x, \lambda), S(x, \lambda)$ are entire in λ . Clearly,

$$U(\varphi) = \varphi'(0, \lambda) = 0, \quad V(\psi) = \psi(\pi, \lambda) = 0.$$

Denote

$$\Delta(\lambda) = W(\varphi, \psi; x). \tag{2.2}$$

By virtue of Corollary 2.2 and the Ostrogradskii–Liouville theorem (see [3], $\Delta(\lambda)$ does not depend on x . The function $\Delta(\lambda)$ is called *characteristic function* of L . Substituting $x = 0$ and $x = \pi$ into (2.2), we get

$$\Delta(\lambda) = V(\varphi) = U(\psi). \tag{2.3}$$

The function $\Delta(\lambda)$ is entire in λ , and it has an at most countable set of zeros $\{\lambda_n\}_{n=1,2,3,\dots}$.

Now, consider the solution $\varphi(x, \lambda)$. Let $C_0(x, \lambda)$ and $S_0(x, \lambda)$ be smooth solutions of (1.1) on the interval $(0, \pi)$ under the initial conditions $C_0(0, \lambda) = S'_0(0, \lambda) = 1, C'_0(0, \lambda) = S_0(0, \lambda) = 0$. Then,

$$C(x, \lambda) = C_0(x, \lambda), \quad S(x, \lambda) = S_0(x, \lambda), \quad x < \frac{\pi}{2}, \tag{2.4}$$

$$\begin{aligned} C(x, \lambda) &= A_1C_0(x, \lambda) + B_1S_0(x, \lambda), \\ S(x, \lambda) &= A_2C_0(x, \lambda) + B_2S_0(x, \lambda), \end{aligned} \quad x > \frac{\pi}{2}, \tag{2.5}$$

where

$$\begin{aligned} A_1 &= 1 + \alpha\lambda C_0\left(\frac{\pi}{2}, \lambda\right) S_0\left(\frac{\pi}{2}, \lambda\right), \quad B_1 = -\alpha\lambda [C_0\left(\frac{\pi}{2}, \lambda\right)]^2, \\ A_2 &= \alpha\lambda [S_0\left(\frac{\pi}{2}, \lambda\right)]^2, \quad B_2 = 1 - \alpha\lambda C_0\left(\frac{\pi}{2}, \lambda\right) S_0\left(\frac{\pi}{2}, \lambda\right). \end{aligned} \tag{2.6}$$

Let $\lambda = k^2$. It is easy to verify that the function $C_0(x, \lambda)$ satisfies the following integral equation:

$$C_0(x, \lambda) = \cos kx + \int_0^x \frac{\sin k(x-t)}{k} q(t) C_0(t, \lambda) dt. \tag{2.7}$$

Solving the equation (2.7) by the method of successive approximations, we obtain

$$C_0(x, \lambda) = \cos kx + \frac{\sin kx}{2k} \int_0^x q(t) dt + \frac{\cos kx}{4k^2} \left\{ q(x) - q(0) - \frac{1}{2} \left[\int_0^x q(t) dt \right]^2 \right\} + O\left(\frac{1}{k^3} \exp(|Imk|x)\right), \tag{2.8}$$

Analogously,

$$S_0(x, \lambda) = \frac{\sin kx}{k} - \frac{\cos kx}{2k^2} \int_0^x q(t) dt + \frac{\sin kx}{4k^2} \left\{ q(x) + q(0) - \frac{1}{2} \left[\int_0^x q(t) dt \right]^2 \right\} + O\left(\frac{1}{k^4} \exp(|Imk|x)\right), \tag{2.9}$$

By virtue of (2.6), (2.8) and (2.9),

$$A_1 = \frac{\alpha}{2} k \sin k\pi + 1 - \frac{\alpha}{2} \cos k\pi \int_0^{\frac{\pi}{2}} q(t) dt + \frac{\alpha}{4k} \sin k\pi \left\{ q\left(\frac{\pi}{2}\right) - \left[\int_0^{\frac{\pi}{2}} q(t) dt \right]^2 \right\} + O\left(\frac{1}{k^2}\right),$$

$$B_1 = -\frac{\alpha}{2} k^2 (1 + \cos k\pi) - \frac{\alpha}{2} k \sin k\pi \int_0^{\frac{\pi}{2}} q(t) dt - \frac{\alpha}{4} \left[q\left(\frac{\pi}{2}\right) + q(0) - \frac{\alpha}{4} \cos k\pi \left\{ q\left(\frac{\pi}{2}\right) - q(0) - \left[\int_0^{\frac{\pi}{2}} q(t) dt \right]^2 \right\} \right] + O\left(\frac{1}{k}\right),$$

$$A_2 = \frac{\alpha}{2} (1 - \cos k\pi) - \frac{\alpha}{2k} \sin k\pi \int_0^{\frac{\pi}{2}} q(t) dt + O\left(\frac{1}{k^2}\right),$$

$$B_2 = -\frac{\alpha}{2} k \sin k\pi + 1 + \frac{\alpha}{2} \cos k\pi \int_0^{\frac{\pi}{2}} q(t) dt + O\left(\frac{1}{k}\right).$$

Since $\varphi(x, \lambda) = C(x, \lambda)$, we calculate using (2.4)–(2.9)

$$\varphi(x, \lambda) = \cos kx + \frac{\sin kx}{2k} \int_0^x q(t) dt + \frac{\cos kx}{4k^2} \left\{ q(x) - q(0) - \frac{1}{2} \left[\int_0^x q(t) dt \right]^2 \right\} + O\left(\frac{1}{k^3} \exp(|Imk|x)\right), \quad x < \frac{\pi}{2}, \tag{2.10}$$

$$\begin{aligned} \varphi(x, \lambda) &= -\frac{\alpha}{2} k \sin kx + \frac{\alpha}{2} k \sin k(\pi - x) + \cos kx \left[1 + \frac{\alpha}{4} \int_0^x q(t) dt \right] \\ &+ \frac{\alpha}{4} \cos k(\pi - x) \left[\int_0^x q(t) dt - \int_0^{\frac{\pi}{2}} q(t) dt \right] - \frac{\alpha}{8k} \sin kx \left\{ q(x) + 2q\left(\frac{\pi}{2}\right) - q(0) - \frac{1}{2} \left[\int_0^x q(t) dt \right]^2 \right\} \\ &- \frac{\alpha}{8k} \sin k(\pi - x) \left\{ q(x) + 2q\left(\frac{\pi}{2}\right) - q(0) - \frac{1}{2} \left[\int_0^x q(t) dt - \int_0^{\frac{\pi}{2}} q(t) dt \right]^2 \right\} + O\left(\frac{1}{k^2} \exp(|\tau|x)\right), \quad x > \frac{\pi}{2}. \end{aligned} \tag{2.11}$$

It follows from (2.3) and (2.11) that

$$\Delta(\lambda) = -\frac{\alpha}{2} \left(k \sin k\pi - w_1 \cos k\pi + w_2 - w_3 \frac{\sin k\pi}{k} \right) + O\left(\frac{1}{|k|^2}\right), \tag{2.12}$$

where

$$\begin{aligned} w_1 &= \frac{2}{\alpha} + \frac{1}{2} \int_0^\pi q(t) dt, & w_2 &= \frac{1}{2} \left[\int_0^{\frac{\pi}{2}} q(t) dt - \int_{\frac{\pi}{2}}^\pi q(t) dt \right], \\ w_3 &= -\frac{1}{4} \left\{ q(\pi) + 2q\left(\frac{\pi}{2}\right) - q(0) - \frac{1}{2} \left[\int_0^\pi q(t) dt \right]^2 \right\}. \end{aligned} \tag{2.13}$$

Using (2.12) and Rouché’s theorem, by the well-known method (see [2]) for $n \rightarrow \infty$,

$$k_n = n + o(1).$$

Analogously, by using Rouché’s theorem, one can prove that, for sufficiently large values of n , every circle $\sigma_n(\delta) = \{k : |k - n| \leq \delta\}$ contains exactly one zero of $\Delta(\lambda)$. Since $\delta > 0$ is arbitrary, we must have

$$k_n = n + \varepsilon_n, \quad \varepsilon_n = o(1), \quad n \rightarrow \infty. \tag{2.14}$$

Since k_n are zeros of $\Delta(\lambda)$, from (2.12), we get

$$n \cdot \sin \varepsilon_n \pi - \left(w_1 \cos \varepsilon_n \pi + (-1)^{n-1} w_2 \right) + v_n = 0, \tag{2.15}$$

where $v_n = n \cdot \sin \varepsilon_n \pi + o(\exp(|\tau_n|))$, $\tau_n = \text{Im} k_n$. Hence, $\sin \varepsilon_n \pi = o\left(\frac{1}{n}\right)$, that is $\varepsilon_n = o\left(\frac{1}{n}\right)$. Using (2.15), we get more precisely

$$\varepsilon_n = \frac{1}{\pi n} \left(w_1 + (-1)^{n-1} w_2 \right) + o\left(\frac{1}{n}\right). \tag{2.16}$$

Substituting (2.16) into (2.14), we get

$$k_n = n + \frac{1}{\pi n} \left(w_1 + (-1)^{n-1} w_2 \right) + \frac{\xi_n}{n}, \quad (\xi_n) \in l_2. \tag{2.17}$$

3. Trace of the problem

The series

$$s_\lambda := \sum_{n=1}^\infty \left[\lambda_n - n^2 - \frac{2}{\pi} \left(w_1 + (-1)^{n-1} w_2 \right) \right] \tag{3.1}$$

converges and is called *the regularized trace of first order* for the problem (1.1)-(1.3). The goal of this paper is to find its sum.

Theorem 3.1 *Suppose that $q(x) \in W_2^1(0, \pi)$, then the following first order regularized trace formula holds*

$$s_\lambda = w_3 - \frac{2w_1}{\pi} - \frac{w_1^2}{2},$$

where w_i ($i = 1, 3$) satisfy the relations (2.13).

Proof Since $\Delta(\lambda)$ is an entire function of order $\frac{1}{2}$, from Hadamard's theorem (see, [10], Section 4.2), using (2.12) we obtain:

$$\Delta(\lambda) = A \prod_{n=1}^{\infty} \left(1 - \frac{\lambda}{\lambda_n}\right), \tag{3.2}$$

where A is a certain constant to be determined below.

Let $\lambda = -\mu^2$. We calculate the sum s_λ of the series (3.1) by comparing the asymptotic expressions from formulas (2.12) and (3.2) as $\mu \rightarrow \infty$. From the (3.2), we have:

$$\Delta(-\mu^2) = \frac{(\lambda_1 + \mu^2) \sinh \mu\pi}{\mu\pi} C\Phi(\mu), \tag{3.3}$$

where

$$C = \frac{A}{\lambda_1} \prod_{n=2}^{\infty} \frac{n^2}{\lambda_n}, \quad \Phi(\mu) = \prod_{n=2}^{\infty} \left(1 - \frac{n^2 - \lambda_n}{\mu^2 + n^2}\right).$$

We study the asymptotic behaviour of the function $\Phi(\mu)$ for large positive μ . For this, we need the following formulas (see, [11]):

$$\sum_{j=2}^{\infty} \frac{1}{j} \sum_{n=1}^{\infty} \frac{|n^2 - \lambda_n|^j}{(\mu^2 + n^2)^j} = O\left(\frac{1}{\mu^3}\right), \tag{3.4}$$

$$\sum_{n=1}^{\infty} \frac{1}{\mu^2 + n^2} = \frac{\pi \coth \pi\mu}{2\mu} - \frac{1}{2\mu^2} = \frac{\pi}{2\mu} - \frac{1}{2\mu^2} + O(\exp(-2\pi\mu)), \tag{3.5}$$

and since $\sup_n \left| \lambda_n - n^2 - \frac{2}{\pi} (w_1 + (-1)^{n-1} w_2) \right| n^2 < \infty$,

$$\frac{1}{\mu^2} \sum_{n=1}^{\infty} \left(\lambda_n - n^2 - \frac{2}{\pi} (w_1 + (-1)^{n-1} w_2) \right) \frac{n^2}{\mu^2 + n^2} \leq \frac{1}{\mu^2} \left(\sum_{n=1}^{\infty} \zeta_n^2 \right)^{\frac{1}{2}} \left(\sum_{n=1}^{\infty} \frac{1}{\mu^2 + n^2} \right)^{\frac{1}{2}} = O\left(\frac{1}{\mu^3}\right). \tag{3.6}$$

From (3.4), (3.5) and (3.6) we calculate:

$$\begin{aligned} \ln \Phi(\mu) &= \sum_{n=2}^{\infty} \ln \left(1 - \frac{n^2 - \lambda_n}{\mu^2 + n^2}\right) = - \sum_{n=2}^{\infty} \sum_{j=1}^{\infty} \frac{1}{j} \left(\frac{n^2 - \lambda_n}{\mu^2 + n^2}\right)^j \\ &= \frac{w_1}{\mu} + \frac{1}{\mu^2} \left(s_\lambda + \frac{2w_1}{\pi}\right) + O\left(\frac{1}{\mu^3}\right). \end{aligned}$$

Therefore, we get

$$\Phi(\mu) = 1 + \frac{w_1}{\mu} + \frac{1}{\mu^2} \left(s_\lambda + \frac{2w_1}{\pi} + \frac{w_1^2}{2}\right) + O\left(\frac{1}{\mu^3}\right),$$

and from (3.3)

$$\Delta(-\mu^2) = \frac{1}{2} ce^{\mu\pi} \left\{ \mu + w_1 + \frac{1}{\mu} \left(s_\lambda + \frac{2w_1}{\pi} + \frac{w_1^2}{2}\right) \right\} + O\left(\frac{1}{\mu^2}\right). \tag{3.7}$$

We now study the asymptotic behaviour of the function $\Delta(-\mu^2) = \varphi(\pi, -\mu^2)$ for large negative $\lambda = -\mu^2$. Then, according to formula (2.11), we have

$$\Delta(-\mu^2) = \frac{1}{4}\alpha e^{\mu\pi} \left\{ \mu + w_1 + \frac{1}{\mu}w_3 \right\} + O\left(\frac{1}{\mu^2}\right). \tag{3.8}$$

It follows from the equalities (3.3), (3.7), (3.8) and comparing the coefficients of μ , we obtain

$$c = \frac{\alpha}{2}, \quad s_\lambda = w_3 - \frac{2w_1}{\pi} - \frac{w_1^2}{2},$$

completing the proof of Theorem 3.1. □

4. Numerical example

The boundary value problem

$$-y'' = \lambda \left(1 + \alpha \delta \left(x - \frac{\pi}{2} \right) \right), \quad x \in (0, \pi), \tag{4.1}$$

with the boundary conditions

$$y'(0) = y(\pi) = 0, \tag{4.2}$$

is a special case of and conditions at the point $x = \frac{\pi}{2}$,

$$\begin{cases} y\left(\frac{\pi}{2} + 0\right) = y\left(\frac{\pi}{2} - 0\right) \equiv y\left(\frac{\pi}{2}\right), \\ y'\left(\frac{\pi}{2} + 0\right) - y'\left(\frac{\pi}{2} - 0\right) = -\alpha\lambda y\left(\frac{\pi}{2}\right), \end{cases} \tag{4.3}$$

is a special case of problem (1.1)-(1.3) when $q(x) = 0$.

The first order regularized trace formula for the problem (4.1), (4.2) may be written by

$$s_\lambda := \sum_{n=1}^{\infty} \left(\lambda_n - n^2 - \frac{4}{\pi\alpha} \right) = -\frac{4}{\pi\alpha} - \frac{2}{\alpha^2}.$$

References

- [1] Albeverio S, Gesztesy F, Hoegh-Krohn R, Holden H. (with an appendix by Exner P.) Solvable Models in Quantum Mechanics (second edition). American Mathematical Society Chelsea Publishing 2005.
- [2] Bellman R, Cooke K. Differential-Difference Equations. Academic Press 1963.
- [3] Coddington EA, Levinson N. Theory of Ordinary Differential Equations. McGraw-Hill 1955.
- [4] Dikii LA. Trace formulas for Sturm-Liouville differential equations. Uspekhi Matematicheskikh Nauk 1958; 13 (3): 111-143 (in Russian).
- [5] Gasymov MG. On the sum of differences of eigenvalues of two self-adjoint operators. Soviet Mathematics Doklady 1963; 150: 1202-1205 (in Russian).
- [6] Gelfand IM, Levitan BM. On a formula for eigenvalues of a differential operator of second order. Soviet Mathematics Doklady 1953; 88: 593-596 (in Russian).

- [7] Gesztesy F, Holden H, Simon B, Zhao Z. A trace formula for multidimensional Schrödinger operators. *Journal of Functional Analysis* 1996; 141: 449-465.
- [8] Guseinov GS, Levitan BM. On trace formulas for Sturm-Liouville operators. *Vestnik Moskovskogo Universiteta Seriya 1 Matematika Mekhanika* 1978; 1: 40-49 (in Russian).
- [9] Lax PD. Trace formulas for the Schrödinger operator. *Communications on Pure and Applied Mathematics* 1994; 47: 503-512.
- [10] Levin BY. *Lectures on entire functions*. Providence, RI: American Mathematical Society 1996.
- [11] Levitan BM. Calculation of the regularized trace for the Sturm-Liouville operator. *Uspekhi Matematicheskikh Nauk* 1964; 19 (1): 161-165 (in Russian).
- [12] Manafov MD. On the spectrum of a class of non-self-adjoint "weighted" operator with point δ -interactions. *Proceedings of the Institute of Mathematics and Mechanics Azerbaijan National Academy of Sciences* 2014; 40: 283-289.
- [13] Sadovnichii VA, Podolskii VE. Traces of operators. *Uspekhi Matematicheskikh Nauk* 2006; 61 (5): 89-156; English translation: *Russian Mathematical Surveys* 2006; 61 (5): 885-953.
- [14] Savchuk AM. First order regularized trace of the Sturm-Liouville operator with δ -potential. *Uspekhi Matematicheskikh Nauk* 2000; 55 (6): 155-156; English translation: *Russian Mathematical Surveys* 2000; 55 (6): 1168-1169.
- [15] Savchuk AM, Shkalikov AA. The trace formula for Sturm-Liouville operator with singular potentials. *Matematicheskie Zametki* 2001; 69 (3): 427-442; English translation: *Mathematical Notes* 2001; 69 (3): 387-400.
- [16] Vinokurov VA, Sadovnichii VA. The asymptotics of eigenvalues and eigenfunctions and a trace formula for a potential with delta functions. *Differentsial'nye Uravneniya* 2002; 38 (6): 735-751; English translation: *Differential Equations* 2002; 38 (6): 772-789.