

1-1-2021

Liftings and covering morphisms of crossed modules in group-groupoids

SERAP DEMİR KARAKAŞ

OSMAN MUCUK

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

KARAKAŞ, SERAP DEMİR and MUCUK, OSMAN (2021) "Liftings and covering morphisms of crossed modules in group-groupoids," *Turkish Journal of Mathematics*: Vol. 45: No. 3, Article 20. <https://doi.org/10.3906/mat-2102-56>

Available at: <https://journals.tubitak.gov.tr/math/vol45/iss3/20>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Liftings and covering morphisms of crossed modules in group-groupoids

Serap DEMİR KARAKAŞ* , Osman MUCUK 

Department of Mathematics, Faculty of Arts and Sciences, Erciyes University, Kayseri, Turkey

Received: 11.02.2021

Accepted/Published Online: 16.04.2021

Final Version: 20.05.2021

Abstract: In this work we introduce lifting and covering of a crossed module in the category of group-groupoids; and then we prove the categorical equivalence of horizontal actions of a double group-groupoid and lifting crossed modules of corresponding crossed module in group-groupoids. These allow us to produce more examples of double group-groupoids.

Key words: Crossed module, group-groupoid, double group-groupoid, action, covering morphism

1. Introduction

The concept of covering groupoid has a significant role in the utilizations of groupoids (see [4, 16]). It is well known that the groupoid actions on sets and the covering morphisms of a certain groupoid G are categorically equivalent (see [6] for topological version). An analogous equivalence was given in [9, Proposition 3.1] for a group-groupoid G which is used under the name 2-group in [2] and G -groupoids or group object in the category of groupoids in [11]. In [1], this result was generalized by assuming G is an internal groupoid in the category of groups with operations appeared in [27, 28]. That result is adapted to Leibniz algebras setting in [29], to categorical groups in [25] and to categorical ring in [22].

Double groupoids which are useful for Seifert-van Kampen Theorem to determine the fundamental groupoids of topological spaces [7] were defined by Ehresmann in [13, 14] to be internal groupoids in the category of groupoids. It was proved in [10] that double groupoids are categorically equivalent to crossed modules in the sense of Whitehead [31, 32]. Due to this equivalence some algebraic structures such as normality and quotient of double groupoids were characterized in [21] (see [23] for similar structures in group-groupoids).

By Loday [18] cat^1 -groups and crossed modules in groups; cat^2 -groups and crossed squares are categorically equivalent. More generally by Ellis and Steiner [15] cat^n -groups are equivalent to crossed n -cubes. The readers are also referred to [3] for algebraic structures on groupoids and algebraic descriptions of homotopy n -types. Due to [30] crossed modules in group-groupoids are equivalent to double group-groupoids and to crossed squares; and therefore to cat^2 -groups.

It was proved in [8, Theorem1.7] that the categories of horizontal actions and horizontal action morphisms for a double Lie groupoid are equivalent. Recently this result is extended to double group-groupoids in [12]; and action and covering notions of double group-groupoids are characterized.

In this paper, by means of the latter equivalence, we aim to introduce the notions of lifting and covering

*Correspondence: srpdmr89@gmail.com

2010 AMS Mathematics Subject Classification: 20L05, 57M10, 18D35, 18D05

of a crossed module in group-groupoids and prove the categorical equivalence of them. For the convenience of the reader in the second section we give preliminaries on groupoids, actions, coverings and group-groupoids. Section 3 contains a brief summary of double groupoids, double group-groupoids together with actions, coverings and crossed module in the category of group-groupoids. In Section 4, we characterize a crossed module in the category of group-groupoids corresponding to a double group-groupoid horizontally acting on a group-groupoid, define the lifting notion of crossed modules in group-groupoids and we give some examples. We prove that the category of horizontal actions of a double group-groupoid on group-groupoids and the category of lifting crossed modules in the category of group-groupoids are equivalent; and obtain a categorical equivalence between liftings and coverings of a crossed module in group-groupoids. These results extend [24, Theorem 4.6] and [24, Theorem 5.1] respectively. The results of the paper enable us to produce more examples of double group-groupoids.

2. Groupoids and group-groupoids

A groupoid is a small category whose morphisms are isomorphisms (see [4, 19] for more details). Indeed a groupoid G includes a set G of morphisms or arrows and G_0 of objects with source and target point maps $d_0, d_1: G \rightarrow G_0$ and object inclusion map $\epsilon: G_0 \rightarrow G$ with the property that $d_0\epsilon = d_1\epsilon = 1_{G_0}$. An associative partial composition $G_{d_1} \times_{d_0} G \rightarrow G, (g, h) \mapsto g \circ h$, where $G_{d_1} \times_{d_0} G$ is the pullback of d_0 and d_1 is defined. Here if $g, h \in G$ and $d_1(g) = d_0(h)$, then the composite $g \circ h$ is well defined such that $d_0(g \circ h) = d_0(g)$ and $d_1(g \circ h) = d_1(h)$. Moreover, for $x \in G_0$ the morphism $\epsilon(x)$ acts as the identity and it is denoted by 1_x . There is a map $G \rightarrow G$ called inversion which assigns to every element g its inverse g^{-1} such that $d_0(g^{-1}) = d_1(g)$, $d_1(g^{-1}) = d_0(g)$, $g \circ g^{-1} = \epsilon(d_0(g))$, $g^{-1} \circ g = \epsilon(d_1(g))$. In a groupoid G , all maps defined above are called structural maps. If $x \in G_0$, the star $St_G x$ of x is defined by the set $\{g \in G; d_0(g) = x\}$. The fundamental groupoid πX of a topological space X is an example of groupoid whose objects are the points of X and morphisms are the homotopy classes of the paths relative to the end points.

A morphism $f: G \rightarrow H$ of groupoids includes the maps $f_1: G \rightarrow H$ and $f_0: G_0 \rightarrow H_0$ satisfying $d_0 f_1 = f_0 d_0$, $d_1 f_1 = f_0 d_1$, $f_1 \epsilon = \epsilon f_0$ and preserving the composite $f(g \circ h) = f(g) \circ f(h)$, for $g, h \in G$.

A groupoid G whose sets of objects and morphisms are equipped with group structures is called a group-groupoid whenever the group operation written additively $G \times G \rightarrow G, (g, h) \mapsto g + h$, the inverse $G \rightarrow G, g \mapsto -g$ and the unit map $\{0\} \rightarrow G$, where $\{0\}$ is singleton, are groupoid morphisms. Here note that the additive map is a morphism of groupoids if and only if the interchange rule

$$(g + h) \circ (k + l) = (g \circ k) + (h \circ l)$$

is satisfied for $g, h, k, l \in G$ whenever the composites are well defined. A group-groupoid morphism is a group structure preserving morphism of underlying groupoids. We thus obtain a category GpGd of group-groupoids.

A groupoid G whose morphism and object sets have topologies and the structural maps are continuous is called topological groupoid (see [5, 19]). A topological group-groupoid is defined in [17] to be a topological groupoid which has topological group structures on the sets of objects and morphisms.

A covering morphism of groupoids, $p: \tilde{G} \rightarrow G$, is a groupoid morphism with the property that for every $\tilde{x} \in \tilde{G}_0$ the restriction $St_{\tilde{G}} \tilde{x} \rightarrow St_G p(\tilde{x})$ is bijective. A covering morphism of topological groupoid is a covering morphism of groupoids in which each restriction to the star is a homeomorphism. A covering morphism of topological group-groupoids is defined in [20, Definition 4.1] as a covering morphism of topological groupoid.

The following construction on action appears in [4, p.373].

An action of a groupoid G on a set X includes a function $\omega: X \rightarrow G_0$ and a partial function $\varphi: X_\omega \times_{d_0} G \rightarrow X$, $(x, g) \mapsto x \bullet g$ where $X_\omega \times_{d_0} G$ is pullback of ω and d_0 with the following properties.

- (i) $\omega(x \bullet g) = d_1(g)$ for $(x, g) \in X_\omega \times_{d_0} G$;
- (ii) $x \bullet (g \circ h) = (x \bullet g) \bullet h$ for $(g, h) \in G_{d_1} \times_{d_0} G$ and $(x, g) \in X_\omega \times_{d_0} G$;
- (iii) $x \bullet \epsilon(\omega(x)) = x$ for $x \in X$.

Such an action is denoted by (X, ω) . A morphism of these actions from (X, ω) to (X', ω') is a function $f: X \rightarrow X'$ with the properties $\omega'f = \omega$ and $f(x \bullet g) = f(x) \bullet g$. So for a given groupoid G , we have a category denoted by $\text{GpdAct}(G)$.

Following [4], for such an action there is a groupoid $G \ltimes X$, called semidirect product groupoid. Here the object set is X . The elements of $(G \ltimes X)(x, y)$ are the pairs (g, x) in which $g \in G(\omega(x), \omega(y))$ and $x \bullet g = y$. The groupoid composition is as follows.

$$(g, x) \circ (h, y) = (g \circ h, x)$$

The projection map $p: G \ltimes X \rightarrow G$ is a covering morphism of groupoids. This assignment determines a categorical equivalence between actions and coverings of G [6].

An action of a group-groupoid G on a group X by a group morphism $\omega: X \rightarrow G_0$ (See [9, Section 3] for more details) is a groupoid action of G on the underlying set of X by ω , satisfying the interchange rule

$$(x \bullet g) + (y \bullet h) = (x + y) \bullet (g + h)$$

for $g, h \in G$ and $x, y \in X$.

A morphism from a group-groupoid action (X, ω) to (X, ω') include $f: X \rightarrow X'$ as a morphism of group and of underlying operations of G . Therefore there is a category $\text{GpGpdAct}(G)$ of group-groupoid actions and morphisms of them.

Besides, the categories of group-groupoid coverings and actions are equivalent for a fixed group-groupoid [9, Proposition 3.1].


3. Double group-groupoids and crossed modules

A double groupoid is a groupoid object in the category of groupoids. In other words it consists of a quadruple of sets $(S; H, V; P)$ such that there are two groupoid structures on H and V with object set P and two groupoid structures on S which are vertical one based on H denoted by S_V and horizontal one based on V denoted as S_H . Therefore a double groupoid has four related groupoid structures and compatible structural maps.

In a double groupoid we write multiplication for groupoid compositions in H and V ; and $1_b^H \in H$ and $1_b^V \in V$ for the identity elements for $b \in P$. The source, target, object inclusion, composition for H are $d_0^H, d_1^H: H \rightarrow P$, $\epsilon^H: P \rightarrow H$ and $m^H: H * H \rightarrow H$ respectively and similar notations are used for V .

The horizontal groupoid S_H has source and target $d_0^h, d_1^h: S \rightarrow V$, object inclusion $\epsilon^h: V \rightarrow S$ and partial composition $\circ_h: S * S \rightarrow S$, $(s_1, s_2) \rightarrow s_1 \circ_h s_2$. The vertical groupoid S_V has source and target $d_0^v, d_1^v: S \rightarrow H$, object inclusion $\epsilon^v: H \rightarrow S$ and partial composition $\circ_v: S * S \rightarrow S$, $(s_1, s_2) \rightarrow s_1 \circ_v s_2$. For a square s write s^{-h} and s^{-v} for the inverses of s in S_H and S_V , respectively.

Elements of S are squares with boundaries as follows:



A double groupoid has the following interchange rule for $s_1, s_2, s_3, s_4 \in S$

$$(s_1 \circ_h s_2) \circ_v (s_3 \circ_h s_4) = (s_1 \circ_v s_3) \circ_h (s_2 \circ_v s_4)$$

A morphism $\varphi = (\varphi_s, \varphi_h, \varphi_v, \varphi_p): (S'; H', V'; P') \rightarrow (S; H, V; P)$ of double groupoids consists of four maps that commute with structural maps. These form a category DGpd of double groupoids.

A double group-groupoid is defined in [30] to be an internal groupoid in the category GpGd . Hence it consists of four related group-groupoids S_H, S_V, H and V provided with the following interchange rules

$$(s_1 \circ_h s_2) + (s_3 \circ_h s_4) = (s_1 + s_3) \circ_h (s_2 + s_4)$$

,

$$(s_1 \circ_v s_2) + (s_3 \circ_v s_4) = (s_1 + s_3) \circ_v (s_2 + s_4)$$

.

By the interchange rule in double group-groupoid, horizontal and vertical groupoid compositions can be written in terms of group operation for $d_1^h(s_1) = d_0^h(s)$ and $d_1^v(\alpha_1) = d_0^v(\alpha)$ as follows:

$$s_1 \circ_h s = s_1 - \varepsilon^h(d_1^h)^h(s_1) + s = s - \varepsilon^h(d_1^h)^h(s) + s_1 \tag{3.1}$$

$$\alpha_1 \circ_v \alpha = \alpha_1 - \varepsilon^v(d_1^v)^v(\alpha_1) + \alpha = \alpha - \varepsilon^v(d_1^v)^v(\alpha) + \alpha_1 \tag{3.2}$$

whenever the necessary operations are defined; and for the squares $s, s_1 \in \text{Ker } d_0^h$ and $\alpha, \alpha_1 \in \text{Ker } d_0^v$, we have

$$s + s_1 - s = \varepsilon^h d_1^h(s) + s_1 - \varepsilon^h d_1^h(s)$$

and

$$\alpha + \alpha_1 - \alpha = \varepsilon^v d_1^v(\alpha) + \alpha_1 - \varepsilon^v d_1^v(\alpha).$$

There is a category of double group-groupoids denoted by DGpGpd .

Action of a double group-groupoid on a group-groupoid below comes from [12].

A horizontal action of double group-groupoid $\mathcal{S} = (S; H, V; P)$ on a group-groupoid G via a morphism $\omega: G \rightarrow V$ of group-groupoids includes an action of horizontal group-groupoid S_H on V via $\omega: G \rightarrow V$ and an action of H on P via $\omega_0: G_0 \rightarrow P$ with the following properties.

(i) $d_1^G(g \bullet s) = d_1^G(g) \bullet d_1^v(s)$ and $d_0^G(g \bullet s) = d_0^G(g) \bullet d_0^v(s)$ for each $s \in S, g \in G$ with $d_0^h(s) = \omega(g)$.

(ii) For $s_1, s_2 \in S$ and $g_1, g_2 \in G$ we have

$$(g_1 g_2) \bullet (s_1 \circ_v s_2) = (g_1 \bullet s_1) \circ_v (g_2 \bullet s_2) \tag{3.3}$$

(iii) For all $a \in H$ and $x \in G_0$ with $d_0^H(a) = \omega_0(x)$ we have $1_x^G \bullet \varepsilon^v(a) = 1_{ax}^G$ and for $x, x_1 \in G_0$ and $a, a_1 \in H$ we have

$$(x_1 + x) \bullet (a_1 + a) = (x_1 \bullet a_1) + (x \bullet a)$$

We write (G, ω) for such an action. Due to structure of group-groupoid we have an interchange rule

$$(g_1 + g_2) \bullet (s_1 + s_2) = (g_1 \bullet s_1) + (g_2 \bullet s_2). \tag{3.4}$$

Similarly vertical action of double group-groupoids can be restated. See [8] for the study about horizontal action of Lie double groupoid and related examples.

A morphism $f: (G, \omega) \rightarrow (G', \omega')$ of such actions consists of group homomorphisms $f: G \rightarrow G'$ and $f_0: G_0 \rightarrow G'_0$ provided that $f(g \bullet s) = f(g) \bullet s$ and $f_0(x \bullet h) = f_0(x) \bullet h$ such that $\omega'f = \omega$ and $\omega'_0 f_0 = \omega_0$. Thus for a fixed double group-groupoid S we have a category $\text{DGpGpdAct}_H(S)$ of horizontal actions of double group-groupoids.

A morphism $\varphi = (\varphi_s, \varphi_h, \varphi_v, \varphi_p): (S'; H', V'; P') \rightarrow (S; H, V; P)$ of double group-groupoids is called covering morphism associated with the horizontal action if (φ_s, φ_v) and (φ_h, φ_p) are covering morphisms of ordinary group-groupoids [12, Definition 2.2]. Then we have a category $\text{Cov}_H \text{DGpGpd}/S$ of coverings of S .

A crossed module which is due to Whitehead in [31, 32] is defined to be group homomorphism $\partial: A \rightarrow B$ with a right action $(a, b) \mapsto a.b$ of B on A with the following rules.

$$[\text{CM1}] \quad \partial(a.b) = -b + \partial(a) + b, \text{ and}$$

$$[\text{CM2}] \quad a_1.\partial(a) = -a + a_1 + a.$$

We know by [30, Proposition 3.9] that (G, H, ∂) is a crossed module in group-groupoids if (G, H, ∂_1) is a crossed module in groups. A morphism (f, g) from (G', H', ∂) to (G, H, ∂) is defined to be two group-groupoid morphisms $f: G' \rightarrow G$ and $g: H' \rightarrow H$ with the property that $(f, g): (G', H', \partial) \rightarrow (G, H, \partial)$ is a morphism of crossed module in groups. Therefore there is a category XModGpGd of crossed modules in group-groupoids.

We need some techniques of the proof for the following result in later parts and hence we only state the main ideas.

Theorem 3.1 [30, Theorem 4.7] Crossed modules in group-groupoids and double group-groupoids are categorically equivalent.

$$\text{XModGpGd} \simeq \text{DGpGpd}$$

Proof For a crossed module in group-groupoid (G, H, ∂) , one has a corresponding double group-groupoid $(H \times G, H_0 \times G_0, H, H_0)$ in which the compositions are defined as follows:

$$(h', g') \circ_h (h, g) = (h' \circ h, g' \circ g)$$

,

$$(h', g') \circ_v (h, g) = (h, g' + g')$$

. By [30, Lemma 3.4], group operation of $G \times H$ is

$$(h_1, g_1) + (h, g) = (h_1 + h, g_1 + h_1.g). \tag{3.5}$$

Conversely for a double group-groupoid $(S; H, V; P)$ one has a crossed module in group-groupoid (K, V, ∂) where

$$K = (\text{Ker } d_0^h, \text{Ker } d_0^H, d_0^v, d_1^v, \varepsilon^v, n^v, m^v)$$

and

$$V = (V, P, d_0^V, d_1^V, \varepsilon^V, n^V, m^V)$$

the boundary map is $\partial = (\partial_1 = d_1^h, \partial_0 = d_1^H)$ and the action of V on $\text{Ker } d_0^h$ is given by

$$a.b = -\varepsilon^h(b) + a + \varepsilon^h(b) \tag{3.6}$$

for $a \in \text{Ker } d_0^h$ and $b \in V$. We refer to the cited reference for more details.

□

We now state how to construct a double group-groupoid from a topological group-groupoid and obtain the corresponding crossed module in group-groupoids.

Example 3.2 We know that for a topological group X , the fundamental groupoid πX is a group-groupoid. Hence if G is a topological group-groupoid, then G and G_0 are topological groups and then we have related four group-groupoids $(\pi G, \pi G_0)$, $(\pi G, G)$, $(\pi G_0, G_0)$ and (G, G_0) . So we have a quadruple $(\pi G; \pi G_0, G; G_0)$ which becomes a double group-groupoid. Therefore by Theorem 3.1 we have a corresponding crossed module $d_1: St_{\pi G}0 \rightarrow G$ in group-groupoids.

4. Liftings and coverings of crossed modules in group-groupoids

In this section evaluating the equivalence of the categories in Theorem 3.1 we obtain the lifting notion for a crossed module in the category of group-groupoids associated with a horizontal action of a double group-groupoid on a group-groupoid. We first have the following preparation.

For given a double group-groupoid $\mathcal{S} = (S; H, V; P)$ horizontally acting on a group-groupoid G via a morphism $\omega: G \rightarrow V$ of group-groupoids suppose that (A, B, ∂) is the crossed module of group-groupoids associated with \mathcal{S} by Theorem 3.1. We then actually have the following.

(i) a morphism of group-groupoids $\omega: G \rightarrow B$

(ii) an action of G on $A = \text{Ker } d_0^h$ via ω defined by

$$A \times G \rightarrow A; (a, g) \mapsto a.g = -\varepsilon^h(\omega(g)) + a + \varepsilon^h(\omega(g)) \tag{4.1}$$

(iii) an action of G_0 on $A_0 = \text{Ker } d_0^H$ via ω_0 defined by

$$A_0 \times G_0 \rightarrow A_0; (x, y) \mapsto x.y = -\varepsilon^H(\omega_0(y)) + x + \varepsilon^H(\omega_0(y))$$

(iv) a group-groupoid morphism

$$\varphi: A \rightarrow G \quad \varphi_1(a) = 0_G \bullet a, \quad \varphi_0(x) = 0_{G_0} \bullet x \tag{4.2}$$

such that $\omega\varphi = \partial$.

We now state the following theorem.

Theorem 4.1 (A, G, φ) is a crossed module in group-groupoids.

Proof By [30, Proposition 3.9], we need to prove that (A, G, φ_1) satisfies the axioms of a crossed module of groups.

[CM1]

$$\begin{aligned}
 \varphi_1(a.g) &= \varphi_1(-\varepsilon^h(\omega(g)) + a + \varepsilon^h(\omega(g))) && \text{(by Eq.4.1)} \\
 &= 0_G \bullet (-\varepsilon^h(\omega(g)) + a + \varepsilon^h(\omega(g))) && \text{(by Eq.4.2)} \\
 &= (-g + g) \bullet (-\varepsilon^h(\omega(g)) + a + \varepsilon^h(\omega(g))) \\
 &= ((-g) \bullet \varepsilon^h(\omega(-g))) + g \bullet (a + \varepsilon^h(\omega(g))) && \text{(by Eq. 3.4)} \\
 &= (-g) + (0_G + g) \bullet (a + \varepsilon^h(\omega(g))) && \text{(by } g \bullet \varepsilon^h(\omega(g)) = g) \\
 &= (-g) + 0_G \bullet a + g \bullet \varepsilon^h(\omega(g)) && \text{(by Eq. 3.4)} \\
 &= -g + \varphi_1(a) + g && \text{(by Eq.4.2)}
 \end{aligned}$$

[CM2]

$$\begin{aligned}
 a_1\varphi_1(a) &= a_1(0_G \bullet a) && \text{(by Eq.4.2)} \\
 &= -\varepsilon^h(\omega(0_G \bullet a)) + a_1 + \varepsilon^h(\omega(0_G \bullet a)) && \text{(by Eq. 4.1)} \\
 &= -\varepsilon^h(d_1^h(a)) + a_1 + \varepsilon^h(d_1^h(a)) && \text{(by } \omega(0_G \bullet a) = d_1^h(a)) \\
 &= a_1 d_1^h(a) && \text{(by Eq.3.6)} \\
 &= -a + a_1 + a
 \end{aligned}$$

Therefore (A, G, φ) becomes a crossed module of group-groupoids as required. □

Therefore we can state definition below.

Definition 4.2 Suppose that (A, B, ∂) is a crossed module in group-groupoids and $\omega: G \rightarrow B$ is a morphism of group-groupoids. A crossed module (A, G, φ) in which G acts on A via ω is called a lifting of (A, B, ∂) if the following diagram commutes, i.e. $\omega\varphi = \partial$

$$\begin{array}{ccc}
 & & G \\
 & \nearrow \varphi & \downarrow \omega \\
 A & \xrightarrow{\quad} & B
 \end{array}$$

We will denote such a lifting by (φ, G, ω) .

A morphism $\rho: (\varphi, G, \omega) \rightarrow (\varphi', G', \omega')$ of such liftings is a morphism $\rho: G \rightarrow G'$ of group-groupoids satisfying $\rho\varphi = \varphi'$ and $\omega'\rho = \omega$. Therefore we have a category $\text{LXModGpGd}/(A, B, \partial)$ of liftings and morphisms of them.

Example 4.3 For every crossed module (A, B, ∂) of group-groupoids, $(\partial, B, 1_B)$ becomes a lifting of (A, B, ∂) .

Example 4.4 If N is a normal subgroup-groupoid of G as defined in [23, Definition 2.10], there exists a morphism $\partial: G \rightarrow H$ of group-groupoids with $\text{Ker } \partial = N$ by [23, Theorem 2.19]. Hence for a crossed module in group-groupoid $\partial: G \rightarrow H$ with $\text{Ker } \partial = N$, there is a unique morphism $\tilde{\partial}: G/N \rightarrow H$ with the commutative diagram below.

$$\begin{array}{ccc} & & G/N \\ & \nearrow \eta & \downarrow \tilde{\partial} \\ G & \xrightarrow{\partial} & H \end{array}$$

This means that $(G, G/N, \eta)$ is a lifting of (G, H, ∂) by group-groupoid morphism $\tilde{\partial}$.

The following theorem extends [24, Example 4.8].

Theorem 4.5 If $p: \tilde{G} \rightarrow G$ is a covering morphism of topological group-groupoids, then there exists a crossed module in group-groupoids $d_1: \text{St}_{\pi_G}0 \rightarrow G$ with a lifting $\tilde{d}_1: \text{St}_{\pi_G}0 \rightarrow \tilde{G}$.

Proof We observe from Example 3.2 that for a topological group-groupoid G , $d_1: \text{St}_{\pi_G}0 \rightarrow G$ is a crossed module in group-groupoids. Since $p: \tilde{G} \rightarrow G$ is a covering morphism of groupoids, for a path α in G with initial point 0 , identity, there exists a path $\tilde{\alpha}$ in \tilde{G} with initial point $\tilde{0}$ such that $p(\tilde{\alpha}) = \alpha$. Thus we obtain a function $\tilde{d}_1: \text{St}_{\pi_G}0 \rightarrow \tilde{G}$ which assigns the homotopy class $[\alpha]$ of α to the final point of $\tilde{\alpha}$. So we have the commutative diagram below.

$$\begin{array}{ccc} & & \tilde{G} \\ & \nearrow \tilde{d}_1 & \downarrow p \\ \text{St}_{\pi_G}0 & \xrightarrow{d_1} & G \end{array}$$

Hence $(\tilde{d}_1, \tilde{G}, p)$ becomes a lifting of $(\text{St}_{\pi_G}0, G, d_1)$. □

As a result we can give the following categorical equivalence.

Theorem 4.6 Let S be a double group-groupoid and (A, B, ∂) be the crossed module in group-groupoid which corresponds to S . Then the following categories are equivalent.

$$\text{DGpGpdAct}_H(S) \simeq \text{LXModGpGd}/(A, B, \partial)$$

Proof Let us begin with defining a functor $\theta: \text{DGpGpdAct}_H(S) \rightarrow \text{LXModGpGd}/(A, B, \partial)$ which assigns each horizontal action (G, ω) of the double group-groupoid S to a lifting (φ, G, ω) of (A, B, ∂) in which φ is defined by

$$\varphi_1: A \rightarrow G, \varphi_1(a) = 0_G \bullet a$$

$$\varphi_0: A_0 \rightarrow G_0, \varphi_0(b) = 0_{G_0} \bullet b$$

such that $\omega\varphi = \partial$.

Let us consider the functor $\delta: \text{LXModGpGd}/(A, B, \partial) \rightarrow \text{DGpGpdAct}_H(\mathbb{S})$ which assigns every lifting (φ, G, ω) to a horizontal action of double group-groupoid (G, ω) of S on G with action

$$S \times G \rightarrow G, (s, g) \mapsto g \bullet s = \varphi_1(s - \varepsilon^h(d_0^h(s))) + g$$

$$H \times G_0 \rightarrow G_0, (h, x) \mapsto x \bullet h = \varphi_0(h - \varepsilon^H(d_0^H(h))) + x$$

We now proceed to show that $\theta \circ \delta$ and $\delta \circ \theta$ are naturally isomorphic to $1_{\text{LXModGpGd}/(A, B, \partial)}$ and $1_{\text{DGpGpdAct}_H(\mathbb{S})}$, respectively. If $(\varphi, G, \omega) \in \text{LXModGpGd}/(A, B, \partial)$ then $(\theta \circ \delta)(\varphi, G, \omega) = (\varphi', G, \omega)$ where

$$\varphi'_1(a) = 0_G \bullet a = \varphi_1(a - \varepsilon^h(d_0^h(a))) + 0_G = \varphi_1(a)$$

and

$$\varphi'_0(b) = x \bullet b = \varphi_0(b - \varepsilon^H(d_0^H(b))) + x = \varphi_0(b)$$

. Therefore $\theta \circ \delta = 1$.

Conversely if $(G, \omega) \in \text{DGpGpdAct}_H(\mathbb{S})$ with an action of S on G by

$$S \times G \rightarrow G, (s, g) \mapsto g \bullet s; \quad H \times G_0 \rightarrow G_0, (h, x) \mapsto x \bullet h$$

then we have an induced action defined by

$$\begin{aligned} g \bullet' s &= \varphi_1(s - \varepsilon^h(d_0^h(s))) + g \\ &= 0_G \bullet (s - \varepsilon^h(d_0^h(s))) + (g \bullet \varepsilon^h(\omega(g))) \\ &= (0_G + g) \bullet (s - \varepsilon^h(d_0^h(s))) + \varepsilon^h(\omega(g)) && \text{(by } d_0^h(s) = \omega(g)) \\ &= g \bullet s. \end{aligned}$$

The action of H on G_0 can be checked in a similar way. Hence $\delta \circ \theta = 1$ which completes the proof. \square

The definition of covering of double group-groupoid was given in [12]. Evaluating the detailed proof of Theorem 3.1, we can characterize the morphism of crossed modules in group-groupoids corresponding to covering of double group-groupoid as follows.

Definition 4.7 A morphism (f, g) of crossed modules in group-groupoids from (A', B', ∂') to (A, B, ∂) is called covering morphism if $f: A' \rightarrow A$ and $g: B' \rightarrow B$ are isomorphisms.

Therefore we obtain a category of $\text{XModCov}/(A, B, \partial)$ of coverings for a given crossed module (A, B, ∂) in group-groupoids.

Example 4.8 The morphism $(1_A, 1_B): (A, B, \partial) \rightarrow (A, B, \partial)$ of crossed module in group-groupoids is a covering morphism.

By using the categorical equivalence given in Theorem 3.1, we introduce the below corollary.

Corollary 4.9 The category $\text{Cov}_{\mathbb{H}}\text{DGpGpd}/S$ of coverings of double group-groupoid S and the category $\text{XModCov}/(A, B, \partial)$ of coverings of corresponding crossed module in group-groupoids are equivalent.

The proof of the following corollary follows from [24, Theorem 5.2].

Corollary 4.10 Let (A, B, ∂) be a crossed module in group-groupoids. Then the category $\text{LXModGpGd}/(A, B, \partial)$ of liftings and the category $\text{XModCov}/(A, B, \partial)$ of coverings are equivalent.

Acknowledgement

We would like to thank the anonymous referees for their useful comments to improve the paper; and also the editors for editorial process.

References

- [1] Akız HF, Alemdar N, Mucuk O, Şahan T. Coverings of internal groupoids and crossed modules in the category of groups with operations. *Georgian Mathematical Journal* 2013; 20 (2): 223-238.
- [2] Baez JC, Lauda AD. Higher-dimensional algebra V: 2-groups. *Theory and Applications of Categories* 2004; 12: 423-491.
- [3] Brown R. Computing homotopy types using crossed n -cubes of groups. *Adams Memorial Symposium on Algebraic Topology, 1*. London Mathematical Society Lecture Note Series, 175. Cambridge, UK: Cambridge University Press, 1992, pp. 187-210.
- [4] Brown R. *Topology and Groupoids*. Deganwy, UK: BookSurge LLC, 2006.
- [5] Brown R, Hardy JPL. Topological groupoids I: Universal constructions. *Mathematische Nachrichten* 1976; 71: 273-286.
- [6] Brown R, Danesh-Naruie G, Hardy JPL. Topological groupoids II: Covering morphisms and G -spaces. *Mathematische Nachrichten* 1976; 74: 143-156.
- [7] Brown R, Higgins PJ, Sivera R. Nonabelian Algebraic Topology: filtered spaces, crossed complexes, cubical homotopy groupoids. *European Mathematical Society Tracts in Mathematics* 2011; 15.
- [8] Brown R, Mackenzie KCH. Determination of a double Lie groupoid by its core diagram. *Journal of Pure and Applied Algebra* 1992; 80: 237-272.
- [9] Brown R, Mucuk O. Covering groups of non-connected topological groups revisited. *Mathematical Proceedings of the Cambridge Philosophical Society* 1994; 115: 97-110.
- [10] Brown R, Spencer CB. Double groupoids and crossed modules. *Cahiers de Topologie et Géométrie Différentielle Catégoriques* 1976; 17: 343-362.
- [11] Brown R, Spencer CB. G -groupoids, crossed modules and the fundamental groupoid of a topological group. *Koninklijke Nederlandse Akademie van Wetenschappen A* 1976; 38: 296-302.
- [12] Demir S, Mucuk O. Actions of double group-groupoids and covering morphism. *Gazi University Journal of Science* 2020; 33 (4): 810-819. doi: 10.35378/gujs. 604849
- [13] Ehresmann C. Categories doubles et categories structurees. *Comptes Rendus Mathématique Académie des Sciences*. Paris 1963; 256: 1198-1201. (in French)
- [14] Ehresmann C. Categories structures. *Annales Scientifiques de l'École Normale Supérieure* 1963; 80: 349-426. (in French)
- [15] Ellis G, Steiner R. Higher-dimensional crossed modules and the homotopy groups of $(n + 1)$ -ads. *Journal of Pure and Applied Algebra* 1987; 46: 117-136.

- [16] Higgins PJ. Notes on categories and groupoids. Van Nostrand Reinhold Mathematical Studies, No. 32. New York, NY, USA: Van Nostrand Reinhold Co., 1971.
- [17] Icen I, Özcan AF. Topological crossed modules and G -groupoids. *Algebras, Groups and Geometries* 2001; 18 (4): 401-410.
- [18] Loday JL. Spaces with finitely many nontrivial homotopy groups. *Journal of Pure and Applied Algebra* 1982; 24 (2): 179-202.
- [19] Mackenzie KCH. Lie groupoids and Lie algebroids in differential geometry. London Mathematical Society Lecture Note Series, 124. Cambridge, UK: Cambridge University Press, 1987.
- [20] Mucuk O, Akız HF. Covering morphisms of topological internal groupoids. *Hacettepe Journal of Mathematics and Statistics* 2020; 49 (3): 1020-1029.
- [21] Mucuk O, Demir S. Normality and quotient in crossed modules over groupoids and double groupoids. *Turkish Journal of Mathematics* 2018; 42: 2336-2347.
- [22] Mucuk O, Demir S. Covering groupoids of categorical rings. *Filomat* 2015; 29 (1): 39-49.
- [23] Mucuk O, Şahan T, Alemdar N. Normality and quotients in crossed modules and group-groupoids. *Applied Categorical Structures* 2015; 23 (3): 415-428.
- [24] Mucuk O, Şahan T. Group-groupoid actions and liftings of crossed modules. *Georgian Mathematical Journal* 2019; 26 (3): 437-447.
- [25] Mucuk O, Şahan T. Covering groupoids of categorical groups. *Hacettepe Journal of Mathematics and Statistics* 2013; 42 (4): 419-430.
- [26] Norrie K. Actions and automorphisms of crossed modules. *Bulletin de la Société Mathématique de France* 1990; 118 (2): 129-146.
- [27] Orzech G. Obstruction theory in algebraic categories I. *Journal of Pure and Applied Algebra* 1972; 2: 287-314.
- [28] Orzech G. Obstruction theory in algebraic categories II. *Journal of Pure and Applied Algebra* 1972; 2: 315-340.
- [29] Şahan T, Erciyes A. Actions of internal groupoids in the category of Leibniz Algebras. *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics* 2019; 68 (1): 619-632.
- [30] Temel S, Şahan T, Mucuk O. Crossed modules, double group-groupoids and crossed squares. *Filomat* 2020; 34 (6): 1755-1769. doi: 10.2298/FIL2006755T
- [31] Whitehead JHC. Note on a previous paper entitled on adding relations to homotopy groups. *Annals of Mathematics* 1946; 47 (2): 806-810.
- [32] Whitehead JHC. Combinatorial homotopy II. *Bulletin of the American Mathematical Society* 1949; 55: 453-496.