

1-1-2020

## Some results on a system of multiterm fractional integro-differential equations

SHAHRAM REZAPOUR

FETHİYE MÜGE SAKAR

SEHER MELİKE AYDOĞAN

ELAHE RAVASH

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

REZAPOUR, SHAHRAM; SAKAR, FETHİYE MÜGE; AYDOĞAN, SEHER MELİKE; and RAVASH, ELAHE (2020) "Some results on a system of multiterm fractional integro-differential equations," *Turkish Journal of Mathematics*: Vol. 44: No. 6, Article 4. <https://doi.org/10.3906/mat-1903-51>  
Available at: <https://dctubitak.researchcommons.org/math/vol44/iss6/4>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

## Some results on a system of multiterm fractional integro-differential equations

Shahram REZAPOUR<sup>1,2</sup> , Fethiye Müge SAKAR<sup>3,\*</sup> , Seher Melike AYDOĞAN<sup>4</sup> ,  
Elahe RAVASH<sup>1</sup> 

<sup>1</sup>Department of Mathematics, Azarbaijan Shahid Madani University, Tabriz, Iran

<sup>2</sup>Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

<sup>3</sup>Department of Business and Administration, Dicle University, Diyarbakır, Turkey

<sup>4</sup>Department of Mathematics, İstanbul Technical University, İstanbul, Turkey

Received: 14.03.2019

Accepted/Published Online: 10.08.2020

Final Version: 16.11.2020

**Abstract:** In present study, we investigate the existence of solution for a multiterm fractional integro-differential system with special boundary conditions under some different conditions. In this way, we provide different results for the existence of solutions for the system and also for obtaining unique solution for the system under different conditions. We also present three numerical examples in which by using the Legendre multiwavelet method, we approximate solutions of the system. These examples illustrate our main results.

**Key words:** Caputo derivation, Legendre multiwavelet, system of multiterm fractional integro-differential equations

### 1. Introduction

It is known that fractional calculus can describe most natural phenomena and has been published a lot of works in this field from analytical view up to applied one (see for example, [1], [4], [5], [7]-[27], [29]-[37], [39], [40], [42]-[46], [49] and [51]). Some researchers have been considered multiterm fractional differential equations (see for example, [28] and [38]). It is known that the Caputo fractional derivative is defined by  ${}^c D^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t \frac{f^{(n)}(t)}{(t-s)^{\alpha-n+1}} dt$ , where  $n-1 < \alpha \leq n$  ([41]). Also, the fractional integral of order  $\alpha$  is defined by  $I^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(s)}{(t-s)^{1-\alpha}} ds$ , when  $\alpha > 0$  ([41]). We need the following results.

**Lemma 1.1** ([41]) *Let  $\alpha > 0$  and  $n = \lceil \alpha \rceil + 1$ . Then,  $I^{\alpha c} D^\alpha f(t) = f(t) + c_0 + c_1 t + c_2 t^2 + \dots + c_{n-1} t^{n-1}$ , where  $c_0, c_1, \dots, c_{n-1}$  are some real numbers.*

**Theorem 1.2** ([48]) *Let  $M$  be a closed convex and nonempty subset of a Banach space  $X$ . Let  $A, B$  be the Operators such that (i)  $Ax + By \in M$  whenever  $x, y \in M$ , (ii)  $A$  is compact and continuous, (iii)  $B$  is a contraction mapping. Then there exists  $z \in M$  such that  $z = Az + Bz$ .*

In 2011, Ahmed and Nieto studied the existence of solution for the fractional equation  ${}^c D^q x(t) = f(t, x(t))$  with boundary conditions  $x(0) = -x(T)$ ,  ${}^c D^p x(0) = -{}^c D^p x(T)$ , where  $t \in [0, T]$ ,  $T > 0$ ,  $1 < q \leq 2$ ,  $0 < p < 1$  and  $f$  is a continuous function ([2]). In 2013, Wang, Guo and Tang reviewed the antiperiodic fractional

\*Correspondence: mugesakar@hotmail.com

2010 AMS Mathematics Subject Classification: 26A33, 34A08, 65T60

boundary value problem  ${}^c D^\alpha x(t) = f(t, x(t))$  with boundary condition  $x(0) = -x(T)$ ,  ${}^c D^p x(0) = -{}^c D^p x(T)$  and  ${}^c D^q x(0) = -{}^c D^q x(T)$ , where  $t \in J = [0, T]$ ,  $T > 0$ ,  $2 < \alpha \leq 3$ ,  $0 < p < 1 < q < 2$  and  $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function ([50]). By use and mixing main idea of [2], [28], [38] and [50]), we investigate the existence of solutions for the  $k$ -dimensional system of fractional integro-differential equations

$$\left\{ \begin{array}{l} {}^c D^{\alpha_1} x_1(t) = f_1\left(t, x_1(t), \dots, x_k(t), I^{\beta_{11}} x_1(t), \dots, I^{\beta_{1k}} x_k(t), {}^c D^{\gamma_{11}} x_1(t), \dots, {}^c D^{\gamma_{1k}} x_k(t)\right), \\ {}^c D^{\alpha_2} x_2(t) = f_2\left(t, x_1(t), \dots, x_k(t), I^{\beta_{21}} x_1(t), \dots, I^{\beta_{2k}} x_k(t), {}^c D^{\gamma_{21}} x_1(t), \dots, {}^c D^{\gamma_{2k}} x_k(t)\right), \\ \vdots \\ {}^c D^{\alpha_k} x_k(t) = f_k\left(t, x_1(t), \dots, x_k(t), I^{\beta_{k1}} x_1(t), \dots, I^{\beta_{kk}} x_k(t), {}^c D^{\gamma_{k1}} x_1(t), \dots, {}^c D^{\gamma_{kk}} x_k(t)\right), \end{array} \right.$$

with boundary condition  $x_i(0) + x_i(1) = \sum_{j=1}^k a_i x_i(\xi_j)$  and

$$\sum_{j=1}^k {}^c D^{\gamma_{ij}} x_i(0) + \sum_{j=1}^k {}^c D^{\gamma_{ij}} x_i(1) = \sum_{j=1}^k b_i x'_i(\eta_j),$$

where  $a_i, b_i \in \mathbb{R}$ ,  $1 \neq b_i \Gamma(2 - \gamma_{ij})$ ,  $a_i \neq 2$ ,  $0 < \xi_1 < \xi_2 < \dots < \xi_k < 1$ ,  $0 < \eta_1 < \eta_2 < \dots < \eta_k < 1$ ,  $1 < \alpha_i \leq 2$ ,  $0 < \gamma_{ij} \leq 1$ ,  $\beta_{ij} > 0$  for  $i = 1, \dots, k$ ,  $t \in I = [0, 1]$  and  $f_1, \dots, f_k \in C(I \times \mathbb{R}^{3k}, \mathbb{R})$ . Consider the Banach spaces

$$X = \{x : x \in C(I) \text{ and } {}^c D^{\gamma_{ij}} x \in C(I) \text{ for } i, j = 1, 2, \dots, k\}$$

endowed with the norm  $\|x\| = \max_{t \in I} |x(t)| + \max_{t \in I} |{}^c D^\gamma x(t)|$  and  $X^k = X \times X \times \dots \times X$  endowed with the norm  $\|(x_1, x_2, \dots, x_k)\|_* = \|x_1\| + \|x_2\| + \dots + \|x_k\|$ .

## 2. Main results

Now, we are ready to state and prove our main results.

**Lemma 2.1** *Let  $y \in C(I)$ ,  $1 \neq b \Gamma(2 - \gamma_j)$ ,  $a \neq 2$ ,  $0 < \xi_1 < \xi_2 < \dots < \xi_k < 1$ ,  $0 < \eta_1 < \eta_2 < \dots < \eta_k < 1$ ,  $1 < \alpha \leq 2$  and  $0 < \gamma_j \leq 1$  for  $j = 1, \dots, k$ . Then the problem  ${}^c D^\alpha x(t) = y(t)$  with boundary conditions  $x(0) + x(1) = a \sum_{j=1}^k x(\xi_j)$  and  $\sum_{j=1}^k {}^c D^{\gamma_j} x(0) + \sum_{j=1}^k {}^c D^{\gamma_j} x(1) = b \sum_{j=1}^k x'(\eta_j)$  has the unique solution*

$$x(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + \frac{1}{(a-2)\Gamma(\alpha)} \left( \int_0^1 (1-s)^{\alpha-1} y(s) ds - a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha-1} y(s) ds \right) - \sum_{j=1}^k \frac{\Gamma(2-\gamma_j)(1-a\xi_j+t(a-2))}{(a-2)(1-b\Gamma(2-\gamma_j))} \left( \frac{1}{\Gamma(\alpha-\gamma_j)} \int_0^1 (1-s)^{\alpha-\gamma_j-1} y(s) ds - \frac{b}{\Gamma(\alpha-1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha-2} y(s) ds \right).$$

**Proof** By using some calculations, one can check that the given  $x$  is a solution for the problem satisfying the boundary conditions. On the other hand by using Lemma 1.1, there exist  $b_1, b_2 \in \mathbb{R}$  such that

$$x(t) = I^\alpha y(t) - b_1 - b_2 t = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds - b_1 - b_2 t$$

for all  $t \in I$ . Thus,  ${}^c D^{\gamma_j} x(t) = \frac{1}{\Gamma(\alpha-\gamma_j)} \int_0^t (t-s)^{\alpha-\gamma_j-1} y(s) ds - b_2 \frac{t^{1-\gamma_j}}{\Gamma(2-\gamma_j)}$  and

$$x'(\eta_j) = \frac{1}{\Gamma(\alpha-1)} \int_0^{\eta_j} (\eta_j-s)^{\alpha-2} y(s) ds - b_2.$$

Hence,  $x(0) + x(1) = a \sum_{j=1}^k x(\xi_j)$  and  $\sum_{j=1}^k {}^c D^{\gamma_j} x(0) + \sum_{j=1}^k {}^c D^{\gamma_j} x(1) = b \sum_{j=1}^k x'(\eta_j)$  and so

$$b_2 = - \sum_{j=1}^k \frac{\Gamma(2-\gamma_j)}{(1-b\Gamma(2-\gamma_j))} \left( \frac{1}{\Gamma(\alpha-\gamma_j)} \int_0^1 (1-s)^{\alpha-\gamma_j-1} y(s) ds - \frac{b}{\Gamma(\alpha-1)} \int_0^{\eta_j} (\eta_j-s)^{\alpha-2} y(s) ds \right)$$

and

$$b_1 = \sum_{j=1}^k \frac{a}{(a-2)\Gamma(\alpha)} \int_0^{\xi_j} (\xi_j-s)^{\alpha-1} y(s) ds - \frac{1}{(a-2)\Gamma(\alpha)} \int_0^1 (1-s)^{\alpha-1} y(s) ds + \sum_{j=1}^k \frac{(1-a\xi)\Gamma(2-\gamma_j)}{(1-b\Gamma(2-\gamma_j))(a-2)} \left( \frac{1}{\Gamma(\alpha-\gamma_j)} \int_0^1 (1-s)^{\alpha-\gamma_j-1} y(s) ds - \frac{b}{\Gamma(\alpha-1)} \int_0^{\eta_j} (\eta_j-s)^{\alpha-2} y(s) ds \right).$$

Therefore, we obtain

$$x(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds + \frac{1}{(a-2)\Gamma(\alpha)} \left( \int_0^1 (1-s)^{\alpha-1} y(s) ds - a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j-s)^{\alpha-1} y(s) ds \right) - \sum_{j=1}^k \frac{\Gamma(2-\gamma_j)(1-a\xi_j+t(a-2))}{(a-2)(1-b\Gamma(2-\gamma_j))} \left( \frac{1}{\Gamma(\alpha-\gamma_j)} \int_0^1 (1-s)^{\alpha-\gamma_j-1} y(s) ds - \frac{b}{\Gamma(\alpha-1)} \int_0^{\eta_j} (\eta_j-s)^{\alpha-2} y(s) ds \right).$$

This completes the proof. □

Define the operator  $E : X^k \rightarrow X^k$  by  $Ex(t) = \begin{pmatrix} E_1 x(t) \\ E_2 x(t) \\ \vdots \\ E_k x(t) \end{pmatrix}$ , where  $x = (x_1, x_2, \dots, x_k)$  and

$$E_i x(t) = \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i-1} y(s) ds + \frac{1}{(a_i-2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i-1} y(s) ds - a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j-s)^{\alpha_i-1} y(s) ds \right) - \sum_{j=1}^k \frac{\Gamma(2-\gamma_{ij})(1-a_i\xi_j+t(a-2))}{(a-2)(1-b_i\Gamma(2-\gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i-\gamma_{ij})} \int_0^1 (1-s)^{\alpha_i-\gamma_{ij}-1} y(s) ds - \frac{b_i}{\Gamma(\alpha_i-1)} \int_0^{\eta_j} (\eta_j-s)^{\alpha_i-2} y(s) ds \right).$$

Observe that the main problem has a solution  $u_0$  if and only if  $u_0$  is a fixed point of the operator  $E$ . Put

$$\begin{aligned} \Delta_{i1} &= \sum_{j=1}^k (a_i - 1 + a_i \xi_j^{\alpha_i}) (1 - b_i \Gamma(2 - \gamma_{ij})) \Gamma(\alpha_i - \gamma_{ij} + 1), \\ \Delta_{i2} &= \sum_{j=1}^k \Gamma(2 - \gamma_{ij}) (3 + a_i \xi_j + a_i) (\Gamma(\alpha_i + 1) + b_i \eta_j^{\alpha_i} \Gamma(\alpha_i - \gamma_{ij} + 1)), \\ \Omega_{i1} &= \sum_{j=1}^k (a_i - 1 + a_i \xi_j^{\alpha_i - \gamma_{ij}}) (1 - b_i \Gamma(2 - \gamma_{ij})) \Gamma(\alpha_i - 2\gamma_{ij} + 1) \end{aligned}$$

and  $\Omega_{i2} = \sum_{j=1}^k \Gamma(2 - \gamma_{ij}) (3 + a_i \xi_j + a_i) (\Gamma(\alpha_i - \gamma_{ij} + 1) + b_i \eta_j^{\alpha_i - \gamma_{ij}} \Gamma(\alpha_i - 2\gamma_{ij} + 1))$  for  $i = 1, \dots, k$ .

**Theorem 2.2** *Suppose that  $f_1, \dots, f_k : I \times \mathbb{R}^{3k} \rightarrow \mathbb{R}$  are continuous and there exist  $l_1, \dots, l_k \in C([0, 1], (0, \infty))$ , nondecreasing maps  $\lambda_1, \dots, \lambda_k \in C([0, \infty), (0, \infty))$  and a positive constant  $K$  such that*

$$\begin{aligned} |f_i(t, x_1(t), \dots, x_k(t), z_1(t), \dots, z_k(t), y_1(t), \dots, y_k(t))| &\leq \sum_{j=1}^k l_i(t) (\lambda_i(|x_j(t)| + |y_j(t)|)), \\ |f_i(t, x_1(t), \dots, x_{3k}(t)) - f_i(t, y_1(t), \dots, y_{3k}(t))| &\leq K \sum_{j=1}^{2k} |x_j(t) - y_j(t)| \end{aligned}$$

for all  $t \in I$  and

$$\begin{aligned} K &\leq \sum_{j=1}^k \left( \frac{\Delta_{i1} + \Delta_{i2}}{2(a_i - 2)\Gamma(\alpha_i + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)} \right. \\ &\quad \left. + \frac{\Omega_{i1} + \Omega_{i2}}{2(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - 2\gamma_{ij} + 1)} \right)^{-1}. \end{aligned}$$

Then the main problem has a solution.

**Proof** Choose a real number

$$\begin{aligned} r &\geq \sum_{j=1}^k 2M \left( \frac{\Delta_{i1} + \Delta_{i2}}{(a_i - 2)\Gamma(\alpha_i + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)} \right. \\ &\quad \left. + \frac{\Omega_{i1} + \Omega_{i2}}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - 2\gamma_{ij} + 1)} \right). \end{aligned}$$

Let  $B_r = \{x = (x_1, x_2, \dots, x_k) \in X^k : \|x\| \leq r\}$  and  $M = \sup_{t \in [0, 1]} \|f(t, 0, \dots, 0)\|$ . We show that  $E(B_r) \subset B_r$ . For each  $x = (x_1, x_2, \dots, x_k) \in B_r$ , we have

$$|E_i x(t)| = \left| \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i-1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right|$$

$$\begin{aligned}
 & + \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i-1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \\
 & \quad - a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i-1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \\
 & \quad \left. - \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \right. \right. \\
 & \quad \left. \left. \int_0^1 (1-s)^{\alpha_i - \gamma_{ij} - 1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \right. \\
 & \quad \left. \left. - \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right) \right) \\
 & \leq \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i-1} (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, 0, \dots, 0)| + \\
 & |f_i(s, 0, \dots, 0)|) ds + \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i-1} (|f_i(x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) \right. \\
 & - f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds + a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i-1} (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1 \\
 & \quad \dots, {}^c D^{\gamma_{ik}} x_k) + f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds + \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \\
 & \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^1 (1-s)^{\alpha_i - \gamma_{ij} - 1} (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, 0, \dots, 0)| \right. \\
 & + |f_i(s, 0, \dots, 0)|) ds + \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) \\
 & - f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds \leq \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i-1} (K\|x\| + M) ds + \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i-1} \right. \\
 & (K\|x\| + M) ds + a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i-1} (K\|x\| + M) ds + \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \right. \\
 & \left. \int_0^1 (1-s)^{\alpha_i - \gamma_{ij} - 1} (K\|x\| + M) ds + \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} (K\|x\| + M) ds \leq (K\|x\| + M) \right. \\
 & \left. \left[ \frac{1}{\Gamma(\alpha_i + 1)} + \sum_{j=1}^k \left( \frac{1 + a_i \xi_j}{(a_i - 2)\Gamma(\alpha_i + 1)} + \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij}) + 1} + \frac{b_i \eta_j^{\alpha_i}}{\Gamma(\alpha_i + 1)} \right) \right) \right].
 \end{aligned}$$

Also, we have

$$\begin{aligned}
 |{}^c D^{\gamma_{ij}} T_i x(t)| &= \left| \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^t (t-s)^{\alpha_i - \gamma_{ij} - 1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \\
 &+ \frac{1}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij})} \left( \int_0^1 (1-s)^{\alpha_i - 1 - \gamma_{ij}} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \\
 &\quad \left. - a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i - 1 - \gamma_{ij}} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right) \\
 &- \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - 2\gamma_{ij})} \int_0^1 (1-s)^{\alpha_i - 2\gamma_{ij} - 1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, \right. \\
 &\quad \left. {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds - \frac{b_i}{\Gamma(\alpha_i - 1 - \gamma_{ij})} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2 - \gamma_{ij}} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, \right. \\
 &\quad \left. {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right) \leq \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^t (t-s)^{\alpha_i - 1} (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, \\
 &\quad {}^c D^{\gamma_{ik}} x_k) - f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds + \frac{1}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij})} \left( \int_0^1 (1-s)^{\alpha_i - \gamma_{ij} - 1} (|f_i(s, x_1, \dots, x_k, \right. \\
 &\quad \left. I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds + a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i - \gamma_{ij} - 1} \right. \\
 &\quad \left. (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) + f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds \right) \\
 &+ \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - 2\gamma_{ij})} \int_0^1 (1-s)^{\alpha_i - 2\gamma_{ij} - 1} (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, \right. \\
 &\quad \left. I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds + \frac{b_i}{\Gamma(\alpha_i - \gamma_{ij} - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - \gamma_{ij} - 2} \right. \\
 &\quad \left. (|f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, 0, \dots, 0)| + |f_i(s, 0, \dots, 0)|) ds \right) \\
 &\leq (K\|x\| + M) \left[ \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^t (t-s)^{\alpha_i - \gamma_{ij} - 1} ds + \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i - 1 - \gamma_{ij}} ds \right. \right. \\
 &\quad \left. \left. - a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i - 1 - \gamma_{ij}} ds \right) - \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - 2\gamma_{ij})} \int_0^1 (1-s)^{\alpha_i - 2\gamma_{ij} - 1} ds \right. \right. \\
 &\quad \left. \left. - \frac{b_i}{\Gamma(\alpha_i - 1 - \gamma_{ij})} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2 - \gamma_{ij}} ds \right) \right] \leq (Kr + M) \left[ \frac{1}{\Gamma(\alpha_i - \gamma_{ij} + 1)} + \sum_{j=1}^k \left( \frac{1 + a_i \xi_j^{\alpha_i - \gamma_{ij}}}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)} \right. \right. \\
 &\quad \left. \left. + \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - 2\gamma_{ij} + 1)} + \frac{b_i \eta_j^{\alpha_i - \gamma_{ij}}}{\Gamma(\alpha_i - \gamma_{ij} + 1)} \right) \right].
 \end{aligned}$$

Thus,

$$\|Ex(t)\|_* = \|E(x_1, x_2, x_k)(t)\|_* \leq \sum_{j=1}^k (Kr + M) \left( \frac{\Delta_{i1} + \Delta_{i2}}{(a_i - 2)\Gamma(\alpha_i + 1)(1 - b_i\Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)} + \frac{\Omega_{i1} + \Omega_{i2}}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)(1 - b_i\Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - 2\gamma_{ij} + 1)} \right) \leq r\left(\frac{1}{2}\right) + \frac{r}{2} = r$$

and so  $E(B_r) \subseteq B_r$ . Now, define the operators  $F_i$  and  $G_i$  on  $B_r$  by

$$F_i x(t) = \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i-1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds$$

and

$$\begin{aligned} G_i x(t) &= \frac{1}{(a-2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i-1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \\ &\quad \left. - a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i-1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \\ &\quad \left. - \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a\xi_j + t(a-2))}{(a-2)(1 - b\Gamma(2 - \gamma_{ij}))} \right. \\ &\quad \left. \times \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^1 (1-s)^{\alpha_i - \gamma_{ij} - 1} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right. \right. \\ &\quad \left. \left. - \frac{b}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i-2} f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) ds \right) \right). \end{aligned}$$

Now, we prove that  $G_i$  is a contraction with the constant

$$\begin{aligned} \Lambda' &= \sum_{j=1}^k K \left[ \frac{(1 + a_i \xi^{\alpha_i})\Gamma(\alpha_i - \gamma_{ij} + 1) + (1 + a_i \xi^{\alpha_i - \gamma_{ij}})\Gamma(\alpha_i + 1)}{(a_i - 2)\Gamma(\alpha_i + 1)\Gamma(\alpha_i - \gamma_{ij} + 1)} + \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i\Gamma(2 - \gamma_{ij}))} \right. \\ &\quad \left. \left( \frac{\Gamma(\alpha_i - 2\gamma_{ij} + 1) + \Gamma(\alpha_i - \gamma_{ij} + 1)}{\Gamma(\alpha_i - \gamma_{ij} + 1)\Gamma(\alpha_i - 2\gamma_{ij} + 1)} + \frac{b_i \eta^{\alpha_i} \Gamma(\alpha_i - \gamma_{ij} + 1) + b \eta^{\alpha_i - \gamma_{ij}} \Gamma(\alpha_i + 1)}{\Gamma(\alpha_i + 1)\Gamma(\alpha_i - \gamma_{ij} + 1)} \right) \right] < 1. \end{aligned}$$

Note that,

$$\begin{aligned} |G_i x(t) - G_i y(t)| &\leq \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \left( \int_0^1 (1-s)^{\alpha_i-1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) \right. \\ &\quad \left. - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k) | ds + a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i-1} |f_i(s, x_1, \dots, x_k, \right. \\ &\quad \left. I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k) | ds \right) \end{aligned}$$



$$\begin{aligned}
 & + \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^1 (1 - s)^{\alpha_i - \gamma_{ij} - 1} |f_i(s, x_1, \dots, x_k, \right. \\
 & I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k) | ds \\
 & \left. + \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - \right. \\
 & \left. f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k) | ds \right) \leq K \|x - y\| \left( \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \left( \int_0^1 (1 - s)^{\alpha_i - 1} ds + \right. \right. \\
 & \left. \left. a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i - 1} ds \right) + \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^1 (1 - s)^{\alpha_i - \gamma_{ij} - 1} ds \right. \right. \\
 & \left. \left. + \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} ds \right) \right) \leq \sum_{j=1}^k K \|x - y\| \left( \frac{1 + a_i \xi_j^{\alpha_i}}{(a_i - 2)\Gamma(\alpha_i + 1)} + \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \right. \\
 & \left. \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij} + 1)} + \frac{b_i \eta_j^{\alpha_i}}{\Gamma(\alpha_i + 1)} \right) \right).
 \end{aligned}$$

Hence,

$$\begin{aligned}
 |{}^c D^{\gamma_{ij}} G_i x(t) - {}^c D^{\gamma_{ij}} G_i y(t)| & \leq \sum_{j=1}^k K \|x - y\| \left( \frac{1 + a_i \xi_j^{\alpha_i - \gamma_{ij}}}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)} + \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \right. \\
 & \left. \left( \frac{1}{\Gamma(\alpha_i - 2\gamma_{ij} + 1)} + \frac{b_i \eta_j^{\alpha_i - \gamma_{ij}}}{\Gamma(\alpha_i + 1 - \gamma_{ij})} \right) \right)
 \end{aligned}$$

and so

$$\begin{aligned}
 \|Gx(t) - Gy(t)\|_* & \leq \sum_{j=1}^k K \|x - y\|_* \left[ \frac{(1 + a_i \xi_j^{\alpha_i})\Gamma(\alpha_i - \gamma_{ij} + 1) + (1 + a_i \xi_j^{\alpha_i - \gamma_{ij}})\Gamma(\alpha_i + 1)}{(a_i - 2)\Gamma(\alpha_i + 1)\Gamma(\alpha_i - \gamma_{ij} + 1)} + \frac{\Gamma(2 - \gamma_{ij})(3 + a_i \xi_j + a_i)}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \right. \\
 & \left. \left( \frac{\Gamma(\alpha_i - 2\gamma_{ij} + 1) + \Gamma(\alpha_i - \gamma_{ij} + 1)}{\Gamma(\alpha_i - \gamma_{ij} + 1)\Gamma(\alpha_i - 2\gamma_{ij} + 1)} + \frac{b_i \eta_j^{\alpha_i} \Gamma(\alpha_i - \gamma_{ij} + 1) + b \eta_j^{\alpha_i - \gamma_{ij}} \Gamma(\alpha_i + 1)}{\Gamma(\alpha_i + 1)\Gamma(\alpha_i - \gamma_{ij} + 1)} \right) \right] = \Lambda' \|x - y\|_*.
 \end{aligned}$$

Thus,  $G$  is a contraction. Since  $f_1, \dots, f_k$  are continuous, It is easy to check that  $F_1, \dots, F_k$  are so. We show that  $F_1, \dots, F_k$  are uniformly bounded on  $B_r$ . Let  $x \in B_r$ . Then, we have

$$\begin{aligned}
 |F_i x(t)| & \leq \frac{1}{\Gamma(\alpha_i)} \int_0^t (t - s)^{\alpha_i - 1} |f_i(s, x_1(s), \dots, x_k(s), I^{\beta_{i1}} x(s), \dots, I^{\beta_{ik}} x(s), {}^c D^{\gamma_{i1}} x_1(s), \dots, {}^c D^{\gamma_{ik}} x_k(s))| ds \\
 & \leq \frac{l_i(t)}{\Gamma(\alpha)} \sum_{j=1}^k \lambda_i (|x_j| + |{}^c D^{\gamma_{ij}} x_j|) \int_0^t (t - s)^{\alpha_i - 1} ds, \\
 |{}^c D^{\gamma_{ij}} F_i x(t)| & \leq \frac{1}{\Gamma(\alpha_i + \beta_{ij})} \int_0^t (t - s)^{\alpha_i - \gamma_{ij} - 1} |f_i(s, x_1(s), \dots, x_k(s), I^{\beta_{i1}} x_1(s), \dots, I^{\beta_{ik}} x_k(s),
 \end{aligned}$$

$${}^c D^{\gamma_{i1}} x_1(s), \dots, {}^c D^{\gamma_{ik}} x_k(s) | ds \leq \frac{l_i(t)}{\Gamma(\alpha_i - \gamma_{ij})} \sum_{j=1}^k \lambda_i (|x_j| + |{}^c D^{\gamma_{ij}} x_j|) \int_0^t (t-s)^{\alpha_i - \gamma_{ij} - 1} ds.$$

Hence,  $\|Fx\|_* \leq \sum_{i=1}^k (\frac{1}{\Gamma(\alpha_i)} + \frac{1}{\Gamma(\alpha_i - \gamma_{ij})}) \|l_i\| (\sum_{j=1}^k \lambda_i(r))$ . By using the Arzela-Ascoli Theorem, we conclude that the map  $F$  is compact. For  $0 \leq t_1 < t_2 \leq 1$ , we have

$$\|Fx(t_1) - Fx(t_2)\|_* \leq \sum_{i=1}^k \|l_i\| \lambda_i(r) \left| \frac{2(t_2 - t_1)^{\alpha_i} + t_1^{\alpha_i} - t_2^{\alpha_i}}{\Gamma(\alpha_i + 1)} - \sum_{j=1}^k \frac{2(t_2 - t_1)^{\alpha_i - \gamma_{ij}} + t_1^{\alpha_i - \gamma_{ij}} - t_2^{\alpha_i - \gamma_{ij}}}{\Gamma(\alpha_i - \gamma_{ij} + 1)} \right|$$

and so right side of the inequality tends to zero whenever  $t_2 \rightarrow t_1$ . Now by using Theorem 1.2, the operator  $T$  has at least one fixed point which is a solution for the problem.  $\square$

**Theorem 2.3** Suppose that  $f_1, \dots, f_k : I \times \mathbb{R}^{3k} \rightarrow \mathbb{R}$  are continuous functions and there exist a positive constant  $K$  such that  $|f_i(t, x_1(t), \dots, x_{3k}(t)) - f_i(t, y_1(t), \dots, y_{3k}(t))| \leq K \sum_{j=1}^{2k} |x_j(t) - y_j(t)|$  for all  $t \in I$  and  $i = 1, \dots, k$ . If  $\Lambda = \sum_{i=1}^k K [\sum_{j=1}^k \frac{\Gamma(\alpha_i - \gamma_{ij} + 1)(\Delta_{i1} + \Delta_{i2})\Gamma(\alpha_i - 2\gamma_{ij} + 1) + \Gamma(\alpha_i + 1)(\Omega_{i1} + \Omega_{i2})\Gamma(\alpha_i - \gamma_{ij} + 1)}{(a_i - 2)\Gamma(\alpha_i + 1)\Gamma(\alpha_i - \gamma_{ij} + 1)(1 - b\Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)\Gamma(\alpha_i - 2\gamma_{ij} + 1)}] < 1$ , then the main problem has a unique solution.

**Proof** Let  $x = (x_1, \dots, x_k), y = (y_1, \dots, y_k) \in X^k$ . Then, we have

$$\begin{aligned} |E_i(x_1, x_2, \dots, x_k)(t) - E_i(y_1, y_2, \dots, y_k)(t)| &\leq \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i - 1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, \\ &{}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds + \frac{1}{(a_i - 2)\Gamma(\alpha_i)} \\ &(\int_0^1 (1-s)^{\alpha_i - 1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, \\ &{}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds + a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha_i - 1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) \\ &- f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds) + \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \\ &(\frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^1 (1-s)^{\alpha_i - \gamma_{ij} - 1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, \\ &I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds - \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, \\ &{}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds) \leq \frac{1}{\Gamma(\alpha_i)} \int_0^t (t-s)^{\alpha_i - 1} \\ &(K \sum_{j=1}^k (|x_j - y_j| + |{}^c D^{\gamma_{ij}} x_j - {}^c D^{\gamma_{ij}} y_j|)) ds + \frac{1}{(a_i - 2)\Gamma(\alpha_i)} (\int_0^1 (1-s)^{\alpha_i - 1} (K \sum_{j=1}^k (|x_j - y_j| + |{}^c D^{\gamma_{ij}} x_j - {}^c D^{\gamma_{ij}} y_j|)) ds + \end{aligned}$$

$$\begin{aligned}
 & a_i \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha-1} (K \sum_{j=1}^k (|x_j - y_j| + |{}^c D^{\gamma_{ij}} x_j - {}^c D^{\gamma_{ij}} y_j|)) ds + \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \right. \\
 & \int_0^1 (1 - s)^{\alpha_i - \gamma_{ij} - 1} (K \sum_{j=1}^k (|x_j - y_j| + |{}^c D^{\gamma_{ij}} x_j - {}^c D^{\gamma_{ij}} y_j|)) ds + \frac{b_i}{\Gamma(\alpha_i - 1)} \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - 2} (K \sum_{j=1}^k (|x_j - y_j| + \\
 & \left. |{}^c D^{\gamma_{ij}} x_j - {}^c D^{\gamma_{ij}} y_j|)) ds \leq \sum_{j=1}^k K \|x_j - y_j\| \left( \frac{\Delta_{i1} + \Delta_{i2}}{(a_i - 2)\Gamma(\alpha_i + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)} \right)
 \end{aligned}$$

for all  $t \in I$ . Also, we have

$$\begin{aligned}
 & |{}^c D^{\gamma_{ij}} E_i(x_1, x_2, \dots, x_k)(t) - {}^c D^{\gamma_{ij}} E_i(y_1, y_2, \dots, y_k)(t)| \leq \frac{1}{\Gamma(\alpha_i - \gamma_{ij})} \int_0^t (t - s)^{\alpha_i - \gamma_{ij} - 1} |f_i(s, x_1, \dots, x_k, \\
 & I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds + \\
 & \frac{1}{(a - 2)\Gamma(\alpha_i - \gamma_{ij})} \left( \int_0^1 (1 - s)^{\alpha_i - \gamma_{ij} - 1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) \right. \\
 & \left. - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k) \right) ds + a \sum_{j=1}^k \int_0^{\xi_j} (\xi_j - s)^{\alpha - \gamma_{ij} - 1} |f_i(s, x_1, \dots, x_k, \\
 & I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds + \\
 & \sum_{j=1}^k \frac{\Gamma(2 - \gamma_{ij})(1 - a_i \xi_j + t(a_i - 2))}{(a_i - 2)(1 - b_i \Gamma(2 - \gamma_{ij}))} \left( \frac{1}{\Gamma(\alpha_i - 2\gamma_{ij})} \int_0^1 (1 - s)^{\alpha_i - 2\gamma_{ij} - 1} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, \right. \\
 & \left. {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k) \right) ds - \frac{b_i}{\Gamma(\alpha_i - \gamma_{ij} - 1)} \\
 & \int_0^{\eta_j} (\eta_j - s)^{\alpha_i - \gamma_{ij} - 2} |f_i(s, x_1, \dots, x_k, I^{\beta_{i1}} x_1, \dots, I^{\beta_{ik}} x_k, {}^c D^{\gamma_{i1}} x_1, \dots, {}^c D^{\gamma_{ik}} x_k) - f_i(s, y_1, \dots, y_k, I^{\beta_{i1}} y_1, \dots, \\
 & I^{\beta_{ik}} y_k, {}^c D^{\gamma_{i1}} y_1, \dots, {}^c D^{\gamma_{ik}} y_k)| ds \leq \sum_{j=1}^k K \|x_j - y_j\| \left( \frac{\Omega_{i1} + \Omega_{i2}}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - 2\gamma_{ij} + 1)} \right)
 \end{aligned}$$

and so

$$\begin{aligned}
 & \|E_i(x_1, x_2, \dots, x_k) - E_i(y_1, y_2, \dots, y_k)\| \leq \sum_{j=1}^k K \|x_j - y_j\| \left[ \frac{\Delta_{i1} + \Delta_{i2}}{(a_i - 2)\Gamma(\alpha_i + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)} \right. \\
 & \left. + \frac{\Omega_{i1} + \Omega_{i2}}{(a_i - 2)\Gamma(\alpha_i - \gamma_{ij} + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - 2\gamma_{ij} + 1)} \right]
 \end{aligned}$$

for  $i = 1, \dots, k$ . Thus, we obtain

$$\|E(x_1, x_2, \dots, x_k) - E(y_1, y_2, \dots, y_k)\|_* \leq K \|x - y\|_* \sum_{i=1}^k \left[ \sum_{j=1}^k \frac{\Delta_{i1} + \Delta_{i2}}{(a_i - 2)\Gamma(\alpha_i + 1)(1 - b_i \Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i - \gamma_{ij} + 1)} \right]$$

$$+ \sum_{j=1}^k \frac{\Omega_{i1} + \Omega_{i2}}{(a_i - 2)\Gamma(\alpha_i + \beta_{ij} + 1)(1 - b_i\Gamma(2 - \gamma_{ij}))\Gamma(\alpha_i + \beta_{ij} - \gamma_{ij} + 1)}] = \Lambda \|x - y\|_*.$$

Since  $\Lambda < 1$ ,  $E$  has a unique fixed point which is unique solution for the main problem. □

Now, we introduced a class of wavelet basis constructed by Alpert for  $L^2[0, 1]$  ([3]). First, we review Legendre multiwavelets briefly ([6]). For functions  $\varphi^m \in L^2(\mathbb{R})$  ( $m = 0, 1, \dots, r$ ), consider a reference subspace  $V_0 = \overline{\langle \varphi^m(\cdot - k) : k \in \mathbb{Z}, m = 0, 1, \dots, r \rangle}$  be generated as the  $L^2$ -closure of the linear span of the integer translation of the functions. Also, consider other subspaces  $V_j = \overline{\langle \varphi_{j,k}^m = \varphi^m(2^j x - k) : k \in \mathbb{Z}, m = 0, 1, \dots, r, j \in \mathbb{Z} \rangle}$ . We say that the functions  $\varphi^m \in L^2(\mathbb{R})$  ( $m = 0, 1, \dots, r$ ) generate a multiresolution analysis (MRA) whenever generate a nested sequence of closed subspaces  $\dots \subset V_{-2} \subset V_{-1} \subset V_0 \subset V_1 \subset V_2 \subset \dots$  such that  $\cup_{j \in \mathbb{Z}} V_j$  is dense in  $L^2(\mathbb{R})$ ,  $\cap_{j \in \mathbb{Z}} V_j = \{0\}$ ,  $f(x) \in V_j \iff f(x + 2^{-j}) \in V_j \iff f(2x) \in V_{j+1}$  and  $\{\varphi^m(\cdot - k)\}_{k \in \mathbb{Z}}$  is an Riesz basis of  $V_0$  ([6]). If  $\varphi^m$  generate an MRA, then  $\varphi^m$  are nominated scaling functions ([6]). The scaling functions are called orthogonal whenever  $\varphi^m(\cdot - k) \perp \varphi^{m'}(\cdot - k')$  for  $m \neq m'$  and  $k \neq k'$  ([6]). Since the subspaces  $V_j$  are nested, there exist complementary orthogonal subspaces  $W_j$  such that  $V_{j+1} = V_j \oplus W_j$  for all  $j$  ([6]). This gives an orthogonal decomposition of  $L^2(\mathbb{R})$  as  $L^2(\mathbb{R}) = \bigoplus_{j \in \mathbb{Z}} V_j = \bigoplus_{j \in \mathbb{Z}} W_j$  ([6]). We say that  $\psi^0, \psi^1, \dots, \psi^r \in L^2(\mathbb{R})$  are wavelets whenever those supply the complementary orthogonal subspaces  $W_j$  of an MRA, that is,  $W_j = \overline{\langle \psi_{j,k}^m = \psi^m(2^j x - k) : k \in \mathbb{Z}, m = 0, 1, \dots, r, j \in \mathbb{Z} \rangle}$  ([6]). If  $\psi_{j,k}^m \perp \psi_{j',k'}^{m'}$  for  $j \neq j'$ ,  $k \neq k'$  and  $m \neq m'$ , then  $\psi^0, \psi^1, \dots, \psi^r$  are called orthonormal wavelets ([6]). It is known that Alpert multiwavelets systems with multiplicity  $r$  consist of  $r + 1$  scaling functions and  $r + 1$  wavelets ([6]). The  $r$ th order scaling functions are the  $r + 1$  functions  $\varphi^0(x), \dots, \varphi^r(x)$ , where  $\varphi^i(x)$  is a polynomial of  $i$ th order and all  $\varphi$ 's form orthonormal basis (see [3], [6]), that is,  $\varphi^i(x) = \sum_{k=0}^i a_{ik} x^k$  for some  $a_{ik} \geq 0$  and  $\int_0^1 \varphi^i(x)\varphi^k(x) = \delta_{i,k}$  for all  $i$  and  $k$  ([6]). The two-scale relation for scaling functions of order  $r$  are in the form  $\varphi_i(x) = \sum_{k=0}^r c_{i,j} \varphi_j^i(2x) + \sum_{k=0}^r c_{i,r+j+1} \varphi_j^i(2x - 1)$  ([6]). The coefficients  $\{c_{i,j}\}$  could be obtained uniquely ([6]). Suppose that  $Q_m^k$  are the orthonormal projections from  $L^2[0, 1]$  onto  $S_m^k$  ([6]). If  $f \in C^k[0, 1]$ , then  $\|Q_m^k f - f\| \leq 2^{-mk} \frac{2}{4^k k!} \sup_{x \in [0,1]} |f^{(k)}(x)|$  ([6]). It is known that each function  $x(t)$  which is square integrable on the interval  $[0, 1]$  can be expanded by the scaling functions, that is,

$$x(t) \approx \sum_{k=0}^{2^J-1} \sum_{m=0}^r c_{J,k} \varphi_{J,k}^m(t) = C^T \Phi_J(t)$$

and the corresponding wavelet functions,  $x(t) \approx \sum_{m=0}^r \{c_{0,0}^m \varphi_{0,0}^m(t) + \sum_{j=0}^{J-1} \sum_{k=0}^{2^j-1} d_{j,k}^m \psi_{j,k}^m(t)\} = D^T \Psi_J(t)$ , where  $c_{J,k}^m = \int_0^1 x(t) \varphi_{J,k}^m(t) dt$ ,  $d_{j,k}^m = \int_0^1 x(t) \psi_{j,k}^m(t) dt$  and  $C$  and  $D$  are  $n \times 1$  ( $n = (r + 1)2^J$ ) matrices given by  $C = [c_{J,0}^0, \dots, c_{J,0}^r, \dots, c_{J,2^J-1}^0, \dots, c_{J,2^J-1}^r]^T$  and

$$D = [c_{0,0}^0, \dots, c_{0,0}^r, d_{0,0}^0, \dots, d_{0,0}^r, \dots, d_{J-1,0}^0, \dots, d_{J-1,0}^r, \dots, d_{J-1,2^{J-1}-1}^0, \dots, d_{J-1,2^{J-1}-1}^r]^T$$

(see [3], [6] and [47]). By approximating  $x_i(t) \approx \sum_{k'=0}^r d_{ik'} \psi_{k'}(t) = D^T \Psi(t)$  and replacing in the system, we

obtain

$$\left\{ \begin{array}{l} \sum_{k'=\lceil \alpha_1 \rceil}^r d_{1k'} {}^c D^{\alpha_1}(\psi_{k'}(t)) = f_1\left(t, \sum_{k'=0}^r d_{1k'} \psi'_{k'}(t), \dots, \sum_{k'=0}^r d_{kk'} \psi_{k'}(t), \sum_{k'=0}^r d_{1k'} I^{\beta_{11}}(\psi_{k'}(t)), \right. \\ \dots, \sum_{k'=0}^r d_{kk'} I^{\beta_{1k}}(\psi_{k'}(t)), \sum_{k'=\lceil \gamma_{11} \rceil}^r d_{1k'} {}^c D^{\gamma_{11}}(\psi_{k'}(t)), \dots, \left. \sum_{k'=\lceil \gamma_{1k} \rceil}^r d_{kk'} {}^c D^{\gamma_{1k}}(\psi_{k'}(t)) \right), \\ \\ \sum_{k'=\lceil \alpha_2 \rceil}^r d_{2k'} {}^c D^{\alpha_2}(\psi_{k'}(t)) = f_2\left(t, \sum_{k'=0}^r d_{1k'} \psi_{k'}(t), \dots, \sum_{k'=0}^r d_{kk'} \psi_{k'}(t), \sum_{k'=0}^r d_{1k'} I^{\beta_{21}}(\psi_{k'}(t)), \right. \\ \dots, \sum_{k'=0}^r d_{kk'} I^{\beta_{2k}}(\psi_{k'}(t)), \sum_{k'=\lceil \gamma_{21} \rceil}^r d_{1k'} {}^c D^{\gamma_{21}}(\psi_{k'}(t)), \dots, \left. \sum_{k'=\lceil \gamma_{2k} \rceil}^r d_{kk'} {}^c D^{\gamma_{2k}}(\psi_{k'}(t)) \right), \\ \\ \vdots \\ \\ \sum_{k'=\lceil \alpha_k \rceil}^r d_{kk'} {}^c D^{\alpha_k}(\psi_{k'}(t)) = f_k\left(t, \sum_{k'=0}^r d_{1k'} \psi'_{k'}(t), \dots, \sum_{k'=0}^r d_{kk'} \psi'_{k'}(t), \sum_{k'=0}^r d_{1k'} I^{\beta_{k1}}(\psi_{k'}(t)), \right. \\ \dots, \sum_{k'=0}^r d_{kk'} I^{\beta_{kk}}(\psi_{k'}(t)), \sum_{k'=\lceil \gamma_{k1} \rceil}^r d_{1k'} {}^c D^{\gamma_{k1}}(\psi_{k'}(t)), \dots, \left. \sum_{k'=\lceil \gamma_{kk} \rceil}^r d_{kk'} {}^c D^{\gamma_{kk}}(\psi_{k'}(t)) \right). \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \sum_{k'=0}^r d_{1k'} \psi_{k'}(0) + \sum_{k'=0}^r d_{1k'} \psi_{k'}(1) = \sum_{j=1}^k \sum_{k'=0}^r d_{1k'} \psi_{k'}(\xi_j), \\ \\ \sum_{j=1}^k \sum_{k'=\lceil \gamma_{ij} \rceil}^r d_{1k'} {}^c D^{\gamma_{ij}} \psi_{k'}(0) + \sum_{j=1}^k \sum_{k'=\lceil \gamma_{ij} \rceil}^r d_{1k'} {}^c D^{\gamma_{ij}} \psi_{k'}(1) = \sum_{j=1}^k \sum_{k'=0}^r b_j d_{ik'} \psi'_{k'}(\eta_j). \end{array} \right.$$

for all  $j = 1, 2, \dots, k$ . By applying the Newton iterated method, we can calculate the coefficients  $d_{ij}$ . Here, we give three examples to illustrate our main results by using the method.

**Example 2.4** Consider the system of fractional integro-differential equations

$$\begin{cases} {}^c D^{\frac{4}{3}} x_1(t) = f(t) + 0.0078(x_1(t) + x_2(t) + I^{(2)} x_1(t)), \\ {}^c D^{\frac{5}{4}} x_2(t) = g(t) + 0.0078(x_1(t) + x_2(t) + I^{(3)} x_2(t)), \end{cases}$$

with boundary conditions  ${}^c D^{\frac{1}{4}} x_1(0) + {}^c D^{\frac{3}{4}} x_1(0) + {}^c D^{\frac{1}{4}} x_1(1) + {}^c D^{\frac{3}{4}} x_1(1) = 1.82955(x'_1(\frac{1}{3}) + x'_1(\frac{2}{3}))$ ,  $x_1(0) + x_1(1) = 8.39757(x_1(\frac{1}{5}) + x_1(\frac{2}{5}))$ ,  ${}^c D^{\frac{1}{5}} x_2(0) + {}^c D^{\frac{1}{3}} x_2(0) + {}^c D^{\frac{1}{5}} x_2(1) + {}^c D^{\frac{1}{3}} x_2(1) = 1.0907(x'_2(\frac{1}{3}) + x'_2(\frac{2}{3}))$  and  $x_2(0) + x_2(1) = 1.66667(x_2(\frac{1}{5}) + x_2(\frac{2}{5}))$ . Put  $f(t) = 0.0078(-t - t^{\frac{5}{2}} - \frac{\Gamma(\frac{7}{2})t^{\frac{9}{2}}}{\Gamma(\frac{11}{2})}) + \frac{\Gamma(\frac{7}{2})t^{\frac{7}{6}}}{\Gamma(\frac{13}{6})}$ ,  $g(t) = 0.0078(-t - t^{\frac{5}{2}} - \frac{t^4}{24}) + \frac{t^{-\frac{1}{4}}}{\Gamma(\frac{3}{4})}$ ,  $\alpha_1 = \frac{4}{3}$ ,  $\alpha_2 = \frac{5}{4}$ ,  $\beta_{11} = 2$ ,  $\beta_{22} = 3$ ,  $\gamma_{11} = \frac{1}{4}$ ,  $\gamma_{12} = \frac{3}{4}$ ,  $\gamma_{21} = \frac{1}{5}$ ,  $\gamma_{22} = \frac{1}{3}$ ,  $\xi_1 = 0.01$ ,  $\xi_2 = 0.03$ ,  $\eta_1 = 0.02$ ,  $\eta_2 = 0.04$ ,  $a_1 = 8.39757$ ,  $a_2 = 1.66667$ ,  $b_1 = 1.82955$ ,  $b_2 = 1.0907$ ,  $f_1(t, x_1, x_2, x_3, x_4) = f_1(t) + (x_1(t) + x_2(t) + I^{(2)} x_1(t))$  and  $f_2(t, y_1, y_2, y_3, y_4) = f_2(t) + (y_1(t) + y_2(t) + I^{(3)} y_2(t))$ . If  $l_1(t) = 2t + 2t^{\frac{5}{2}} + \frac{8t^{9/2}}{63} + \frac{15\sqrt{\pi}}{4\Gamma(13/6)}t^{7/6}$ ,  $l_2(t) = 2t + 2t^{5/2} + \frac{t^4}{12} + 0.816049t^{-1/4}$ ,  $\lambda_1 = \lambda_2 = 1$ ,  $\Delta_{11} = -11.8103$ ,  $\Delta_{12} = 25.4806$ ,  $\Delta_{21} = -0.00394216$ ,  $\Delta_{22} = 9.88335$ ,  $\Omega_{11} = -38.0068$ ,  $\Omega_{12} = 110.528$ ,  $\Omega_{21} = 0.206273$ ,  $\Omega_{22} = 154.908$  and  $K = 0.0078$ , then by using Theorem 2.2 the system has a solution. Note that, the exact solutions for the system are  $x_1(t) = t^{\frac{5}{2}}$  and  $x_2(t) = t$ . Check the following table for numerical errors.

$t_i$	The coefficient value of $x_1(t)$	Absolute error with Alpert's multiwavelets
0	0.284359	6.960027e-06
0.2	0.275145	4.153067e-05
0.4	0.096792	4.922978e-05
0.6	0.008913	6.0522151e-05
0.8	0.000627	5.932623e-05
1	0.000174	1.516906e-04

$t_i$	The coefficient value of $x_2(t)$	Absolute error with Alpert's multiwavelets
0	0.500000	3.9350240e-08
0.2	0.288675	2.146224e-07
0.4	-2.311520e-08	2.696232e-07
0.6	2.287753e-08	3.150615e-07
0.8	-6.240553e-09	3.4434256e-07
1	9.0859893e-09	4.463759e-07

**Example 2.5** Consider the system of fractional integro-differential equations

$$\begin{cases} {}^c D^{\frac{7}{4}} x_1(t) = f(t) + 0.1(x_1(t) + x_2(t) + I^{(2)} x_1(t)), \\ {}^c D^{\frac{3}{2}} x_2(t) = g(t) + 0.1(x_1(t) + x_2(t) + I^{(1)} x_2(t)), \end{cases}$$

with boundary conditions  ${}^c D^{\frac{1}{2}} x_1(0) + {}^c D^{\frac{1}{3}} x_1(0) + {}^c D^{\frac{1}{2}} x_1(1) + {}^c D^{\frac{1}{3}} x_1(1) = -1.19536(x'_1(\frac{1}{4}) + x'_1(\frac{1}{2}))$ ,  $x_1(0) + x_1(1) = 0(x_1(\frac{1}{5}) + x_1(\frac{2}{5}))$ ,  ${}^c D^{\frac{2}{3}} x_2(0) + {}^c D^{\frac{4}{5}} x_2(0) + {}^c D^{\frac{2}{3}} x_2(1) + {}^c D^{\frac{4}{5}} x_2(1) = 4.94398(x'_2(\frac{1}{3}) + x'_2(\frac{2}{3}))$  and  $x_2(0) + x_2(1) = 6.17143(x_2(\frac{1}{3}) + x_2(\frac{1}{2}))$ . Put  $f(t) = 0.1(t - t^2 - \frac{5t^3}{6} - \frac{t^4}{12}) - \frac{t^{-\frac{3}{4}}}{\Gamma(\frac{1}{4})} + \frac{2t^{\frac{1}{4}}}{\Gamma(\frac{5}{4})}$ ,  $g(t) = 0.1(t - t^2 - t^3 - \frac{t^4}{4}) + \frac{6t^{\frac{3}{2}}}{\Gamma(\frac{5}{2})}$ ,  $\alpha_1 = \frac{7}{4}$ ,  $\alpha_2 = \frac{3}{2}$ ,  $\beta_{11} = 2$ ,  $\beta_{22} = 1$ ,  $\gamma_{11} = \frac{1}{2}$ ,  $\gamma_{12} = \frac{1}{3}$ ,  $\gamma_{21} = \frac{2}{3}$ ,  $\gamma_{22} = \frac{4}{5}$ ,  $\xi_1 = \frac{1}{5}$ ,  $\xi_2 = \frac{2}{5}$ ,  $\eta_1 = \frac{1}{4}$ ,  $\eta_2 = \frac{1}{2}$ ,  $a_1 = 0$ ,  $a_2 = 6.17143$ ,  $b_1 = -1.19536$ ,  $b_2 = 4.94398$ ,  $f_1(t, x_1, x_2, x_3, x_4) = f_1(t) + (x_1(t) + x_2(t) + I^{(2)} x_1(t))$  and  $f_2(t, y_1, y_2, y_3, y_4) = f_2(t) + (y_1(t) + y_2(t) + I^{(1)} y_2(t))$ . If  $\Delta_{11} = -4.94431$ ,  $\Delta_{21} = -40.0375$ ,  $\Delta_{12} = 7.10496$ ,  $\Delta_{22} = 48.9359$ ,  $\Omega_{11} = -4.05109$ ,  $\Omega_{21} = -53.3475$ ,  $\Omega_{12} = 8.17966$ ,  $\Omega_{22} = 91.2142$  and  $K = 0.1$ , then by using Theorem 2.3, the system has a unique solution. In fact, the exact solutions of the system are  $x_1(t) = t^2 - t$  and  $x_2(t) = t^3$ . Check the following table for numerical errors.

$t_i$	The coefficient value of $x_1(t)$	Absolute error with Alpert's multiwavelets
0	-0.166667	8.1740945e-07
0.2	4.32993e-07	3.495237e-07
0.4	0.0745356	3.768863e-08
0.6	2.263278e-08	2.332886e-07
0.8	-4.69e-09	5.050237e-07
1	2.27e-09	8.1740945e-07

$t_i$	The coefficient value of $x_2(t)$	Absolute error with Alpert's multiwavelets
0	0.25	7.9843424e-07
0.2	0.2598079	2.466749e-07
0.4	0.1118033	3.70490706e-08
0.6	0.01889827	1.40109070e-07
0.8	-1.117e-08	3.5657498e-07
1	9.29385839e-09	7.4183424e-07

**Example 2.6** Consider the system of fractional integro-differential equations

$$\begin{cases} {}^c D^{\frac{6}{5}} x_1(t) = f(t) + 68(x_1(t) + x_2(t) + I^{(1)}x_1(t)), \\ {}^c D^{\frac{5}{4}} x_2(t) = g(t) + 68(x_1(t) + x_2(t) + I^{(2)}x_2(t)), \end{cases}$$

with boundary condition  ${}^c D^{\frac{2}{3}}x_1(0) + {}^c D^{\frac{1}{5}}x_1(0) + {}^c D^{\frac{2}{3}}x_1(1) + {}^c D^{\frac{1}{5}}x_1(1) = 1.2361(x'_1(\frac{1}{3}) + x'_1(\frac{2}{3}))$ ,  $x_1(0) + x_1(1) = 1.09373(x_1(\frac{2}{5}) + x_1(\frac{3}{5}))$ ,  ${}^c D^{\frac{5}{6}}x_2(0) + {}^c D^{\frac{4}{6}}x_2(0) + {}^c D^{\frac{5}{6}}x_2(1) + {}^c D^{\frac{4}{6}}x_2(1) = 2.0758(x'_2(\frac{1}{3}) + x'_2(\frac{2}{3}))$  and  $x_2(0) + x_2(1) = 1.5479(x_2(\frac{2}{5}) + x_2(\frac{3}{5}))$ . Put  $f(t) = \sum_{k=0}^{\infty} \frac{t^{k+\frac{4}{5}}}{\Gamma(k+\frac{9}{5})} + \frac{t^{-\frac{1}{5}}}{\Gamma(\frac{4}{5})} - 68(-1 + 2e^t + t + \frac{3t^2}{2} + e^{2t})$ ,  $g(t) = \frac{2^{\frac{5}{4}}e^{2t}(\Gamma(\frac{3}{4})-\Gamma(\frac{3}{4},2t))}{\Gamma(\frac{3}{4})} + \frac{8t^{\frac{3}{4}}}{3\Gamma(\frac{3}{4})} - 68(\frac{1}{4} + \frac{t}{2} + t^2 + \frac{t^4}{12} + e^t + \frac{5e^{2t}}{4})$ ,  $\alpha_1 = \frac{6}{5}$ ,  $\alpha_2 = \frac{5}{4}$ ,  $\beta_{11} = 1$ ,  $\beta_{22} = 2$ ,  $\gamma_{11} = \frac{2}{3}$ ,  $\gamma_{12} = \frac{1}{5}$ ,  $\gamma_{21} = \frac{5}{6}$ ,  $\gamma_{22} = \frac{4}{6}$ ,  $\xi_1 = \frac{2}{5}$ ,  $\xi_2 = \frac{3}{5}$ ,  $\eta_1 = \frac{1}{3}$ ,  $\eta_2 = \frac{2}{3}$ ,  $a_1 = 1.09373$ ,  $a_2 = 1.5479$ ,  $b_1 = 1.2361$ ,  $b_2 = 2.0757$ ,  $f_1(t, x_1, x_2, x_3, x_4) = f_1(t) + (x_1(t) + x_2(t) + I^{(1)}x_1(t))$  and  $f_2(t, y_1, y_2, y_3, y_4) = f_2(t) + (y_1(t) + y_2(t) + I^{(2)}y_2(t))$ . If  $K = 68$ , then by using Theorem 2.3, the system has a unique solution. In fact, the exact solutions of the system are  $x_1(t) = e^t + t$  and  $x_2(t) = e^{2t} + t^2$ . Check the following table for numerical errors.

$t_i$	The coefficient value of $x_1(t)$	Absolute error with Alpert's multiwavelets
0	2.2153067	3.8596012e-03
0.2	0.77649721	3.3994125e-04
0.4	0.063107775	2.30481104e-03
0.6	0.00562384	2.67062026e-03
0.8	-0.000549966	7.349461374e-04
1	0.00014101506	1.58216779e-03

$t_i$	The coefficient value of $x_2(t)$	Absolute error with Alpert's multiwavelets
0	3.561401498	2.203544165e-02
0.2	2.01399322	2.98524144e-03
0.4	0.50864556	4.11330544e-03
0.6	0.070565292	1.364426553e-03
0.8	0.0110211123	7.15643916e-03
1	0.0005111546	1.35564386e-02

**Acknowledgments**

Research of the first and fourth authors were supported by Azarbaijan Shahid Madani University (Tabriz, Iran). Also, the authors wish to thank the referees for their valuable suggestions. The authors declare that the study was realized in collaboration with equal responsibility. All authors read and approved the final manuscript.

**References**

[1] Agarwal RP, Baleanu D, Hedayati V, Rezapour S. Two fractional derivative inclusion problems via integral boundary condition. Applied Mathematics and Computation 2015; 257: 205-212.

[2] Ahmad B, Nieto JJ. Anti-periodic fractional boundary value problems. Computers and Mathematics with Applications 2011; 62: 1150-1156.

[3] Alpert BK. A class of bases in  $L^2$  for the sparse representation of integral operators. SIAM Journal of Mathematics 1993; 24: 246-262.

- [4] Kojabad EA, Rezapour S. Approximate solutions of a sum-type fractional integro-differential equation by using Chebyshev and Legendre polynomials. *Advances in Difference Equations* 2017; 2017: 351.
- [5] Alizadeh S, Baleanu D, Rezapour S. Analyzing transient response of the parallel RCL circuit by using the Caputo–Fabrizio fractional derivative. *Advances in Difference Equations* 2020; 2020: 55.
- [6] Alpert B, Beylkin G, Gines D, Vozovoi L. Adaptive solution of partial differential equations in multi-wavelet bases. *Journal of Computational Physics* 2002; 182: 149-190.
- [7] Alsaedi A, Baleanu D, Etemad S, Rezapour Sh. On coupled systems of time-fractional differential problems by using a new fractional derivative. *Journal of Function Spaces* 2016; 4626940.
- [8] Amara A, Etemad S, Rezapour S. Topological degree theory and Caputo–Hadamard fractional boundary value problems. *Advances in Difference Equations* 2020; 2020: 369.
- [9] Aydogan SM, Baleanu D, Mousalou A, Rezapour S. On high order fractional integro-differential equations including the Caputo–Fabrizio derivative. *Boundary Value Problems* 2018; 2018: 90.
- [10] Aydogan SM, Gomez GF, Baleanu D, Rezapour S, Samei ME. Approximate endpoint solutions for a class of fractional q-differential inclusions by computational results. *Fractals* 2020; 28 (8): 2040029.
- [11] Aydogan SM, Baleanu D, Mousalou A, Rezapour Sh. On approximate solutions for two higher-order Caputo-Fabrizio fractional integro-differential equations. *Advances in Difference Equations* 2017; 2017: 221.
- [12] Aydogan SM, Baleanu D, Mohammadi H, Rezapour S. On the mathematical model of Rabies by using the fractional Caputo–Fabrizio derivative. *Advances in Difference Equations* 2020; 2020: 382.
- [13] Baleanu D, Agarwal RP, Mohammadi H, Rezapour S. Some existence results for a nonlinear fractional differential equation on partially ordered Banach spaces. *Boundary Value Problems* 2013; 2013: 112.
- [14] Baleanu D, Etemad S, Rezapour S. A hybrid Caputo fractional modeling for thermostat with hybrid boundary value conditions. *Boundary Value Problems* 2020; 2020: 64.
- [15] Baleanu D, Etemad S, Rezapour S. On a fractional hybrid multi-term integro-differential inclusion with four-point sum and integral boundary conditions. *Advances in Difference Equations* 2020; 2020: 250.
- [16] Baleanu D, Etemad S, Pourrazi S, Rezapour S. On the new fractional hybrid boundary value problems with three-point integral hybrid conditions. *Advances in Difference Equations* 2019; 2019: 473.
- [17] Baleanu D, Ghafarnezhad K, Rezapour S. On the existence of solutions of a three step crisis integro-differential equation. *Advances in Difference Equations* 2018; 2018: 135.
- [18] Baleanu D, Ghafarnezhad K, Rezapour S, Shabibi M. On a strong-singular fractional differential equation. *Advances in Difference Equations* 2020; 2020: 350.
- [19] Baleanu D, Hedayati V, Rezapour S, Al Qurashi MM. On two fractional differential inclusions. *SpringerPlus* 2016; 5: 882.
- [20] Baleanu D, Mohammadi H, Rezapour S. Some existence results on nonlinear fractional differential equations. *Philos. Philosophical Transactions of the Royal Society A* 2013; 371: 20120144.
- [21] Baleanu D, Mohammadi H, Rezapour S. On a nonlinear fractional differential equation on partially ordered metric spaces. *Advances in Difference Equations* 2013; 2013: 83.
- [22] Baleanu D, Mohammadi H, Rezapour S. The existence of solutions for a nonlinear mixed problem of singular fractional differential equations. *Advances in Difference Equations* 2013; 2013: 359.
- [23] Baleanu D, Mohammadi H, Rezapour S. A fractional differential equation model for the COVID-19 transmission by using the Caputo–Fabrizio derivative. *Advances in Difference Equations* 2020; 2020: 299.
- [24] Baleanu D, Mohammadi H, Rezapour S. A mathematical theoretical study of a particular system of Caputo–Fabrizio fractional differential equations for the Rubella disease model. *Advances in Difference Equations* 2020; 2020: 184.
- [25] Baleanu D, Mohammadi H, Rezapour S. Analysis of the model of HIV-1 infection of CD4+ T-cell with a new approach of fractional derivative. *Advances in Difference Equations* 2020; 2020: 71.



- [26] Baleanu D, Mousalou A, Rezapour S. A new method for investigating approximate solutions of some fractional integro-differential equations involving the Caputo-Fabrizio derivative. *Advances in Difference Equations* 2017; 2017: 51.
- [27] Baleanu D, Mousalou A, Rezapour S. On fractional integro-differential inclusions via the extended fractional Caputo-Fabrizio derivation. *Boundary Value Problems* 2019; 2019: 79.
- [28] Baleanu D, Nazemi Z, Rezapour S. Existence and uniqueness of solutions for multi-term nonlinear fractional integro-differential equations. *Advances in Difference Equations* 2013; 2013: 368.
- [29] Baleanu D, Nazemi Z, Rezapour S. Attractivity for a k-dimensional system of fractional functional differential equations and global attractivity for a k-dimensional system of nonlinear fractional differential equations. *Journal of Inequalities and Applications* 2014; 2014: 31.
- [30] Baleanu D, Rezapour S, Saberpour Z. On the existence of solutions for some infinite coefficient-symmetric Caputo-Fabrizio fractional integro-differential equations. *Boundary Value Problems* 2017; 2017: 145.
- [31] Etemad S, Rezapour S. On a two-variables fractional partial differential inclusion via Riemann-Liouville derivative. *Novi Sad Journal Mathematics* 2016; 46 (2): 45-53.
- [32] Etemad S, Rezapour S. On a system of hyperbolic partial fractional differential inclusions. *Novi Sad Journal Mathematics* 2016; 46 (2): 145-161.
- [33] Etemad S, Rezapour S, Samei ME. On fractional hybrid and non-hybrid multi-term integro-differential inclusions with three-point integral hybrid boundary conditions. *Advances in Difference Equations* 2020; 2020: 161.
- [34] Etemad S, Rezapour S, Samei ME.  $\alpha$ - $\psi$ -contractions and solutions of a q-fractional differential inclusion with three-point boundary value conditions via computational results. *Advances in Difference Equations* 2020; 2020: 218.
- [35] Etemad S, Pourrazi S, Rezapour S. On a hybrid inclusion problem via hybrid boundary value conditions. *Advances in Difference Equations* 2020; 2020: 302.
- [36] Etemad S, Rezapour S. On the existence of solutions for fractional boundary value problems on the ethane graph. *Advances in Difference Equations* 2020; 2020: 276.
- [37] Etemad S, Rezapour S, Sakar FM. On a fractional Caputo-Hadamard problem with boundary value conditions via different orders of the Hadamard fractional operators. *Advances in Difference Equations* 2020; 2020: 272.
- [38] Ford NJ, Connolly A. Systems-based decomposition schemes for the approximate solution of multi-term fractional differential equations. *Journal of Computational and Applied Mathematics* 2009; 229: 382-391.
- [39] Ghorbanian V, Rezapour S. On a system of fractional finite difference inclusions. *Advances in Difference Equations* 2017; 2017: 325.
- [40] Mohammadi H, Etemad S, Rezapour S, Baleanu D. Two sequential fractional hybrid differential inclusions. *Advances in Difference Equations* 2020; 2020: 385.
- [41] Podlubny I. *Fractional differential equations*. San Diego, CA, USA: Academic Press, Inc., 1999.
- [42] Rezapour S, Hedayati V. On a Caputo fractional differential inclusion with integral boundary condition for convex-compact and nonconvex-compact valued multifunctions. *Kragujevac Journal Mathematics* 2017; 41 (1): 143-158.
- [43] Rezapour S, Samei ME. On the existence of solutions for a multi-singular pointwise defined fractional q-integro-differential equation. *Boundary Value Problems* 2020; 2020: 38.
- [44] Samei ME, Rezapour S. On a system of fractional q-differential inclusions via sum of two multi-term functions on a time scale. *Boundary Value Problems* 2020; 2020: 135.
- [45] Shabibi M, Rezapour S, Vaezpour SM. A singular fractional integro-differential equation. *Scientific Bulletin-University Politehnica of Bucharest Series A* 2017; 79 (1): 109-118.
- [46] Shabibi M, Postolache M, Rezapour S. Positive solutions for a singular sum fractional differential system. *International Journal of Analysis and Applications* 2017; 13 (1): 108-118.

- [47] Shamsi M, Razzaghi M. Solution of Hallen's integral equation using multiwavelets. *Computer Physics Communications* 2005; 168: 187-197.
- [48] Smart DR. *Fixed point theorems*. Cambridge, UK: Cambridge University Press, 1980.
- [49] Talae M, Shabibi M, Gilani A, Rezapour S. On the existence of solutions for a pointwise defined multi-singular integro-differential equation with integral boundary condition. *Advances in Difference Equations* 2020; 2020: 41.
- [50] Wang X, Guo X, Tang G. Anti-periodic fractional boundary value problems for nonlinear differential equations of fractional order. *Journal of Applied Mathematics and Computing* 2013; 41: 367-375.
- [51] Charandabi ZZ, Rezapour S, Eftefagh M. On a fractional hybrid version of the Sturm–Liouville equation. *Advances in Difference Equations* 2020; 2020: 301.