

1-1-2018

Effects of wheel traffic and farmyard manure applications on soilCO₂ emission and soil oxygen content

SEFA ALTIKAT

HASAN KAAAN KÜÇÜKERDEM

AYSUN ALTIKAT

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

Recommended Citation

ALTIKAT, SEFA; KÜÇÜKERDEM, HASAN KAAAN; and ALTIKAT, AYSUN (2018) "Effects of wheel traffic and farmyard manure applications on soilCO₂ emission and soil oxygen content," *Turkish Journal of Agriculture and Forestry*. Vol. 42: No. 4, Article 7. <https://doi.org/10.3906/tar-1709-79>
Available at: <https://journals.tubitak.gov.tr/agriculture/vol42/iss4/7>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Effects of wheel traffic and farmyard manure applications on soil CO₂ emission and soil oxygen content

Sefa ALTİKAT^{1*}, H. Kaan KÜÇÜKERDEM¹, Aysun ALTİKAT²

¹Department of Biosystem Engineering, Faculty of Agriculture, Iğdır University, Iğdır, Turkey

²Department of Environmental Engineering, Faculty of Engineering, Iğdır University, Iğdır, Turkey

Received: 20.09.2017 • Accepted/Published Online: 17.04.2018 • Final Version: 07.08.2018

Abstract: This 2-year field study investigated the effects of different wheel traffic passes, manure amounts, and manure application methods on soil temperature, soil moisture, CO₂ emission, and soil O₂ content. To achieve this purpose, three different wheel traffic applications (no traffic, one pass, and two passes) were used. In the experiments, two different methods of manure applications (surface and subsurface) and three different farmyard manure amounts were used with a control plot (N0), 40 Mg ha⁻¹ (N40), and 80 Mg ha⁻¹ (N80). Manure was applied in both years of the experiment in the first week of April. For the subsurface application, the manure was mixed in at approximately 10 cm of soil depth with a rotary tiller. According to the results, soil temperature, soil moisture, penetration resistance, and bulk density increased with increasing wheel traffic except CO₂ emission for 2014 and 2015. CO₂ emission values decreased with traffic. Subsurface manure application caused more CO₂ emission compared to surface application. The increase in manure amounts led to an increase in CO₂ emission and soil moisture content. The effects on soil O₂ content were observed only during 2015. Maximum oxygen values were obtained in the plots where compaction was not applied. In addition, surface manure application caused more soil O₂ content compared to subsurface application.

Key words: Compaction, CO₂ emission, soil temperature, O₂ content, soil moisture

1. Introduction

One of the most important greenhouse gas is carbon dioxide (CO₂) because it makes up to 60% of the total greenhouse gases. CO₂ is released from soil by respiration. These respirations can be classified as microbial respiration, root respiration, and faunal respiration. Soil respiration occurs at the soil surface or within the upper layer of soil.

The level of agricultural mechanization and the increase in field traffic increase soil compaction (Altikat, 2013). As a result of soil compaction, soil aggregate size distribution and soil porosity change. This situation affects the soil microbial population and functions because of the decrease in C mineralization (Grigal, 2000) and C-N ratio (Li et al., 2004). The soil pore system in compacted soil is generally unfavorable for oxidic microbes because this situation generally restricts the gas-water ratio (Beylich et al., 2010) and causes a lower O₂ diffusion rate (Bilen et al., 2010).

Compaction not only prevents O₂ from being transported to the root surface, but also prevents CO₂ and toxic gases (both evolved and resident) from being removed from around the roots and vented to the

atmosphere (Altikat, 2013; Akbolat, 2009). Poor gas exchange causes the anaerobic layer to move closer to the surface and reduces rooting volume.

Application of farmyard manure in the soil generally increases CO₂ emissions (Fangueiro et al., 2008). Farmyard manure application to the soil can usually be performed with two methods: surface and subsurface applications. In subsurface application, farmyard manure is generally spread on the soil and then mixed with tillage machinery such as a plow and rotary tiller. Liquid manure can also be injected into the soil. Some researchers have reported that injection of liquid manure can reduce the nutrient transport by run-off (Daverede et al., 2004) and reduce NH₃ emissions compared to surface application (Misselbrook et al., 1996).

The purpose of this study is to determine the effect of wheel traffic, farmyard manure application technique, and manure amount on soil CO₂ emissions, O₂ content, soil temperature, and moisture content. Our specific hypotheses were as follows: 1) an increase in wheel traffic will result in a decrease in CO₂ emissions from the soil to the atmosphere, and a similar effect will be seen in the

* Correspondence: sefa.altikat@igdir.edu.tr

soil oxygen capacity; 2) the CO₂ emission from the soil to the atmosphere will be proportional to the amount of fertilizer; 3) maximum CO₂ emission will be observed in subsurface manure application.

2. Materials and methods

2.1. Study site

The field experiments were set up in Iğdır (39°48'06.69"N, 44°34'58.30"E), which is located in the east of Turkey. The experimental field's altitude is 800 m. In the experimental area the mean annual precipitation, minimum average temperature, and maximum average temperature are 256 mm, -3.3 °C, and 25.9 °C, respectively (www.mgm.gov.tr). In addition, meteorological data for Julian days are reported in Table 1.

Topsoil samples were collected from depths of 0–30 cm before farmyard manure and wheel traffic applications. The soil classification is Aridisol, the soil texture is clay loam, and the other properties of the soil are as follows: CaCO₃ 6.53%, EC 1228 µS cm⁻¹, pH 8.0, P 27.24 ppm, K 0.037 mEq 100 g⁻¹, and soil organic matter 1.06%. The experiments started in May 2014 and June 2015.

2.2. Experimental design

Three different experimental factors, provided to different levels, were studied as follows: i) wheel traffic (C0 = no traffic, C1 = one pass, and C2 = two passes), ii) manure application rates (N0 = control plot, N40 = 40 Mg ha⁻¹, and N80 = 80 Mg ha⁻¹); and iii) manure application methods (A1 = surface application and A2 = subsurface application).

The soil was tilled using conventional tillage systems before the experiments so that the plants and roots in the soil were removed from the experimental area. For this purpose, a moldboard plow was set to a depth of 300 mm, followed immediately by two passes with a tandem disk harrow and one pass of a shaped spring tooth harrow.

A tractor (New Holland TD85D) was driven for soil compaction (one pass and two passes) on the plots (20 × 35 m) in three blocks to account for spatial variability. The tires of the tractor were 47 cm wide, with a diameter of 61 cm, and they were inflated to a pressure of 120 kPa. The total weight of the tractor was 3.4 Mg, applying a mean vertical contact pressure of approximately 71 kPa. The mean volumetric water content of the soil surface was 31% and 30% for the 2014 and 2015 experimental periods, respectively. Fermented solid farmyard manure was used in this study. Before the application of manure, farmyard manure was stored for fermentation for about 1 year. In all plots tilled with the moldboard plow it was followed immediately by two passes with a tandem disk harrow and one pass of a shaped spring tooth harrow for avoiding plant respiration, which may otherwise affect CO₂ emissions. Manure was applied both years of the experiment in the first week of April. For the subsurface application, the manure was mixed in at approximately 10 cm of soil depth with a rotary tiller. The experimental scheme for the field analysis is given in Table 2. The properties of the manure used in the experiment are: organic matter 352 g kg⁻¹, pH 7.2, electrical conductivity 3.4 dS m⁻¹, total N 16 g kg⁻¹, P 8.2 g kg⁻¹, K 6.9 g kg⁻¹, Ca 65 g kg⁻¹, and Mg 5.8 g kg⁻¹.

2.3. Measurement of CO₂ emissions, PAR, O₂ content, moisture, temperature, and penetration resistance

In the study, an automated ACE and Soil CO₂ Exchange System (ADC BioScientific Ltd., Hoddesdon, UK) was used for determining the CO₂ emission (Figure 1). Volumetric soil moisture percentage (%) and temperature (°C) were simultaneously measured via CO₂ device sensors during the CO₂ measurement period. Soil O₂ content levels were determined with an ICT O₂ measurement instrument (ICT International Pty. Ltd., Armidale, Australia) (Figure 1).

The soil CO₂ and O₂ measurements were taken approximately 2 to 10 days from 9 May to 30 June in 2014 and from 12 May to 30 June in 2015, between 1000 and

Table 1. Meteorological data through the Julian days.

First year of the experiment (2014)											
Julian days	129	131	135	138	141	144	149	159	166	173	180
Max. temperature (°C)	25.4	27.1	25.2	30.7	23.1	26.4	27.9	26.9	32.3	35.2	31.7
Precipitation (mm)	0	0	1	0	0	2.2	10.3	0	0	0	1.1
Moisture (%)	65.9	59.5	59	48.1	60	65.8	57	55.4	31.1	23.5	41.1
Second year of the experiment (2015)											
Julian days	132	134	138	140	142	146	152	160	166	174	180
Max. temperature (°C)	19.8	20.7	27.4	30.3	26.9	28.8	31.7	30.5	31.1	36.5	35.4
Precipitation (mm)	3.9	0	0.3	0	0.2	7.1	0	0	0	0	0
Moisture (%)	71.2	55	31.9	27.8	46.2	42.4	23.2	32.8	28.5	23.2	26.8

Table 2. Experimental scheme for the field analysis.

Wheel traffic (C)		Manure amount (N)					Method of manure application (A)			
C0: No traffic		N0: Control plot (no manure)					A1: Surface application			
C1: One pass		N40: 40 Mg ha ⁻¹ manure amount					A2: Subsurface application			
C2: Two passes		N80: 80 Mg ha ⁻¹ manure amount								
2014										
CO ₂ emission, soil temperature, and soil moisture measurements in Julian days										
129	131	135	138	141	144	149	159	166	173	180
Oxygen measurements in Julian days										
129	131	135	138	141	144	149	159	166	173	180
2015										
CO ₂ emission, soil temperature, and soil moisture measurements in Julian days										
132	134	138	140	142	146	152	160	166	174	180
Oxygen measurements in Julian days										
132	134	138	140	142	146	152	160	166	174	180

Shaded days are days of measurement.



Soil CO₂ exchange system



ICT oxygen measurement system

Figure 1. Soil CO₂ and O₂ measurement systems.

1200 hours with 7-min intervals. Every measurement took at least 10 min for CO₂ and O₂. Soil temperature and soil moisture at the depth of 10 cm were measured concurrently with CO₂ and O₂ measurements.

Cumulative CO₂ emissions (μmol m⁻² s⁻¹) were calculated as follows:

$$\sum_i^n \frac{(X_i + X_{i+1})}{2} X(t_{i+1} - t_i), \quad (1)$$

where X_i is the first week's CO₂ reading and X_{i+1} is the following week's CO₂ reading at times t_i and t_{i+1}, respectively; n is the CO₂ measurement of the final week's

events during the study period. Cumulative soil CO₂ emissions were then converted to Mg ha⁻¹ day⁻¹ of CO₂-C (Grote and Al-Kaisi, 2007).

A single photosynthetically active radiation (PAR) sensor (silicon photocell) was supplied as standard on every ACE station. The PAR reading has a range of 0–3000 μmol m⁻² s⁻¹ and was factory calibrated. The sensor was mounted on the ACE station close to the chamber at the end of the arm.

A penetrometer was used for determination of penetration resistance for soil compaction (FieldScout SC 900 Soil Compaction Meter, Spectrum Technologies,

Aurora, IL, USA). Penetration values were determined in 0–30 cm with 5-cm intervals. Soil bulk density was determined for the depth ranges of 0–20 cm with 5-cm intervals using stainless steel rings having dimensions of 50 mm in diameter by 50 mm in height (Çelik and Altıkat, 2010). Each plot was sampled three times.

2.4. Statistical analysis

JMP 7 and SPSS statistical programs were used to determine the effects of both the main factors and their interactions. Analysis of variance (balanced ANOVA) was used to assess the significance of each treatment for soil properties and CO₂ emissions and O₂ content. Means were compared when the F-test for treatment was significant at the 5% level by using Duncan’s multiple range tests.

3. Results and discussion

3.1. Penetration resistance and soil bulk density

Soil compaction primarily affects soil properties such as soil bulk density and penetration resistance. In our experiments, penetration resistance and soil bulk density were affected by wheel traffic in both 2014 and 2015 (Tables 3 and 4). Soil penetration resistance increased depending on wheel traffic and soil depth. In the 2014 experimental year, average penetration resistance (0–30 cm soil depth) in the control plot (C0) was observed as 0.907 MPa and this value was increased to 1.458 MPa and 2.166 MPa with one pass (C1) and two passes (C2), respectively. In 2015 these values were observed as 0.877 MPa, 1.43 MPa, and 1.913 MPa for C0, C1, and C2, respectively (Table 4). In

Table 3. Effects of wheel traffic on penetration resistance.

Penetration resistance (MPa) (2014, Julian days 129–149)							
Treatments	0–5 cm	5–10 cm	10–15 cm	15–20 cm	20–25 cm	25–30 cm	Average
C0	0.53 c [√]	0.483 c	0.581 c	0.818 c	1.519 c	1.5135 c	0.907
C1	0.895 b	1.029 b	1.257 b	1.940 b	1.904 b	1.724 b	1.458
C2	1.129 a	1.522 a	1.783 a	2.518 a	3.073 a	2.968 a	2.166
P < 0.001							
Penetration resistance (MPa) (2015, Julian days 132–152)							
Treatments	0–5 cm	5–10 cm	10–15 cm	15–20 cm	20–25 cm	25–30 cm	Average
C0	0.427 b	0.518 c	0.605 c	0.770 c	1.264 c	1.675 b	0.877
C1	0.562 b	0.921 b	1.198 b	1.650 b	1.977 b	2.270 a	1.430
C2	0.95 a	1.384 a	1.689 a	2.098 a	2.558 a	2.80 a	1.913
P < 0.001							

[√]: Means within the same column followed by the same letter are not significantly different.

C0: Control plot (no pass); C1: one pass; C2: two passes

Table 4. Effects of wheel traffic on soil bulk density.

Soil bulk density (g cm ⁻³) (2014, Julian days 129–149)					
Treatment	0–5 cm	5–10 cm	10–15 cm	15–20 cm	Average
C0	1.015 c [√]	1.064 c	1.125 c	1.146 c	1.088
C1	1.08 b	1.156 b	1.189 b	1.202 b	1.157
C2	1.26 a	1.310 a	1.220 a	1.349 a	1.285
P < 0.001					
Soil bulk density (g cm ⁻³) (2015, Julian days 132–152)					
Treatment	0–5 cm	5–10 cm	10–15 cm	15–20 cm	Average
C0	1.11 c	1.14 c	1.17 c	1.205 c	1.156
C1	1.16 b	1.17 b	1.225 b	1.255 b	1.203
C2	1.20 a	1.23 a	1.275 a	1.305 a	1.253
P < 0.001					

[√]: Means within the same column followed by the same letter are not significantly different.

C0: Control plot (no pass); C1: one pass; C2: two passes.

addition, increasing tractor passes had a similar effect on soil bulk density and for the first year of the experiment in the control plot (C0) 1.088 g cm⁻³, one pass (C1) 1.157 g cm⁻³, and two passes (C2) 1.285 g cm⁻³ were observed. In the second year of the experiment these values were 1.156, 1.203, and 1.253 g cm⁻³ for C0, C1, and C2, respectively.

Increasing field traffic is one of the most important parameters in soil compaction. Soil compaction caused by traffic of heavy vehicles and machinery results in soil structure deterioration, both in the topsoil and in the subsoil. By means of dynamic loading, soil physical properties such as pore size distribution and pore continuity are negatively affected, which entails a decrease in air and water permeability and results in increased soil strength or, in the presence of excess soil water, decreased soil strength due to kneading. Soil deformations were increased with the number of passes (Bakker and Davis, 1995; Şeker and Işıldar, 2000). While some studies stated that the critical limit of the number of passes was ten (Jorajuria and Laura, 2000), others claimed that the first pass of the wheel causes the soil compaction (Bakker and Davis, 1995). However, Balbuena et al. (2000) reported that the soil surface layer was significantly affected in 10 passes compared to 1 pass compared to the control plot at 50 cm in soil depth. In addition, axle load in excess of 9 Mg can cause increases in penetration resistance in a depth of >30 cm (Rengasamy, 2000).

Manure applications generally improve soil aggregation and produce an increase in soil organic matter. Manure applications alter soil pH and electrolyte concentrations, which can have adverse effects on soil structure (Haynes and Naidu, 1998).

3.2. Soil temperature and moisture

According to the obtained results, soil temperature and moisture contents increased with increasing compaction levels for both the 2014 and 2015 experimental periods. In the first year of the experiments average soil temperature and soil moisture content was determined as 34.25 °C and 46.19%. In the second year these values were 32.78 °C and 42.33%, respectively.

Soil compaction increased contact among soil particles and thus soil thermal conductivity increased (Guérif et al., 2001). Usowicz et al. (1996) reported that soil thermal conductivity was higher in compacted soil compared to the others. However, increasing soil compaction decreased evaporation and thus soil moisture content increased. Similar results were found in many other studies (Otten et al., 2000; Altikat et al., 2006).

Soil temperature and moisture values were also affected by wheel traffic in both the 2014 and 2015 experimental periods (Table 5). Maximum values were observed for C2 (two passes) for both temperature and moisture for all the experimental periods (Table 6). Effects of wheel traffic on

the soil moisture and temperature in terms of Julian days are given in Figure 2.

Soil temperature values were higher with the surface manure application and N0 manure amount values compared to subsurface application and other manure amounts. In the first year of the experiments, soil temperature was 34.49 °C and 34.57 °C for surface manure application and N0 manure amount, respectively (Table 6). Similar results were obtained in the second year of the experiment. In this year, soil temperature values were 32.65 °C and 33.21 °C for surface manure application and N0 manure amount, respectively (Table 6).

Values of soil moisture content in the subsurface manure application were higher compared to surface manure applications; however, the increase in the amount of manure increased soil moisture content in all of the experiments. In 2014 soil moisture content values were determined as 47.10% and 49.52% for subsurface manure applications and N80 manure amount, respectively (Table 6). Similar results were obtained in 2015. In this year, maximum soil moisture values were observed with subsurface manure applications and N80 manure amount at 43.01% and 42.98%, respectively (Table 6).

In all of the experimental periods maximum soil moisture content values were observed for C2 × N80, A2 × N80, and C1 × A1. In 3-way interactions, maximum values were determined for C2 × A2 × N80. Minimum soil temperature values were observed for A2 × N80, C0 × A2, and C0 × A2 × N80 interactions in both 2014 and 2015.

3.3. CO₂ emissions and O₂ level

CO₂ emissions were statistically significantly affected by field traffic in both years of the experiment (Table 5). It can be seen from Table 6 that the increase in field traffic caused a decrease in CO₂ emission, though the results of the second year of the experiment showed a very small decrease compared to the first year of the experiment (Table 6).

Minimum CO₂ emissions generally occurred at 129–131 Julian days in 2014 and 132–180 Julian days in 2015. However, peak volumes of CO₂ emission were observed at 159–166 Julian days and 138–180 Julian days in 2014 and 2015, respectively (Figure 3).

Effects of the method of application of manure on CO₂ emission can be seen in Figure 4. Subsurface application had higher CO₂ emissions compared to the surface application in both years of the experiment. CO₂ emission rates were also statistically significantly affected by manure amount in both experimental years (Table 5). We have found a clear positive relationship between manure amount and CO₂ emission rate and increasing the manure amount caused higher soil CO₂ emissions in both 2014 and 2015 (Table 6). The relationship between the amount of manure and CO₂ emissions can be seen in Figure 5.

Table 5. Analyses of variance for soil moisture, soil temperature, CO₂ emission, O₂ content, and cumulative CO₂ emission values.

2014					
Treatment	df	CO ₂ emission	O ₂ content	Soil temperature	Soil moisture
		P	P	P	P
C	2	0.001**	0.163 ns	0.001**	0.0012*
N	2	0.001**	0.940 ns	0.001**	0.001**
A	1	0.001**	0.259 ns	0.001**	0.0019*
C × N	4	0.001**	0.773 ns	0.001**	0.025*
C × A	2	0.001**	0.642 ns	0.001**	0.014*
A × N	2	0.001**	0.286 ns	0.001**	0.001**
C × A × N	4	0.001**	0.836 ns	0.001**	0.001**
2015					
Treatment	df	CO ₂ emission	O ₂ content	Soil temperature	Soil moisture
		P	P	P	P
C	2	0.0001**	0.0001**	0.001**	0.0012*
N	2	0.0001**	0.0003**	0.001**	0.001**
A	1	0.0005**	0.0001**	0.001**	0.0019*
C × N	4	0.479 ns	0.773 ns	0.001**	0.025*
C × A	2	0.045 ns	0.001**	0.001**	0.014*
A × N	2	0.289 ns	0.0001**	0.001**	0.001**
C × A × N	4	0.896 ns	0.0001**	0.001**	0.001**
Variance analyses for cumulative CO ₂ emission					
Treatment	df	2014	2015		
		P	P		
Wheel traffic (C)	2	0.001**	0.001**		
Manure amount (N)	2	0.958 ns	0.001**		
Methods of application (A)	1	0.0587 ns	0.001**		
C × N	4	0.882 ns	0.479 ns		
C × A	2	0.352 ns	0.045*		
A × N	2	0.568 ns	0.289 ns		
C × A × N	4	0.993 ns	0.896 ns		

C: Wheel traffic, N: manure amount, A: methods of manure application, df: degrees of freedom; ns: nonsignificant, *: statistically significant (P < 0.05), **: statistically highly significant (P < 0.01).

Cumulative CO₂ emission values are given in Table 6. By increasing the amount of manure, higher CO₂ emissions could be obtained. Furthermore, the application methods of manure were important. Subsurface manure application had higher CO₂ emissions in comparison to surface manure application.

One of the key factors influencing production of CO₂ is fertilization of arable soils (Bouwman et al., 2002). The effects of fertilization on CO₂ at the soil surface tend to occur within the initial 8 to 10 weeks following N application (Phillips, 2007), but the magnitude of these effects likely vary with temperature at the time of application. Manure

application in soil can increase CO₂ emission because organic C in the soil is an immediate source of C for soil microorganisms, which in turn emit CO₂ (Rao and Pathak, 1996). Several studies have indicated that microorganism activities happening in the sublayer of soil increase the CO₂ gas emission (Bilen et al., 2010; Altikat et al., 2012; Altikat, 2013). Rochette and Gregorich (1998) reported that manure doubled CO₂ emissions compared to artificial fertilizer and the increase in the amount of CO₂ emission was highly related (r² = 0.75) to soil temperature. Gregorich et al. (1998), in another study conducted to determine the effects of manure on CO₂ emission, found that an

Table 6. Effects of factors on the soil moisture, soil temperature, CO₂ emission, and O₂ concentration values.

Treatments			Soil temp. (°C)	Soil moist. (%)	CO ₂ emission (μmol m ⁻² s ⁻¹)	Cumulative CO ₂ (Mg ha ⁻¹ day ⁻¹)	O ₂ content (%)
2014	Wheel traffic (C)	C0	34.08 b [√]	44.76 b	5.26 a	0.206 a	19.97 ns
		C1	34.34 a	46.37 a	4.40 b	0.170 b	19.94 ns
		C2	34.39 a	47.44 a	3.81 c	0.137 c	19.87 ns
	Manure Amount (N)	N0	34.57 a	40.78 b	3.84 c	0.172 ns	19.94 ns
		N40	34.27 ab	48.26 a	4.36 b	0.171 ns	19.94 ns
		N80	33.98 b	49.52 a	5.28 a	0.170 ns	19.92 ns
	Application (A)	A1	34.49 a	45.28 b	4.14 b	0.166 ns	19.95 ns
		A2	34.05 b	47.10 a	4.84 a	0.176 ns	19.91 ns
	2015	Wheel traffic (C)	C0	31.08 c	38.25 c	5.20 a	0.197 a
C1			32.59 b	41.67 b	4.39 b	0.166 b	19.79 b
C2			34.48 a	45.40 a	3.75 c	0.142 c	19.59 c
Manure Amount (N)		N0	33.21 a	40.73 b	3.57 c	0.135 c	19.83 a
		N40	32.41 b	41.61 b	4.33 b	0.165 b	19.86 a
		N80	31.53 c b	42.98 a	5.44 a	0.207 a	19.67 b
Application (A)		A1	32.65 a	40.53 b	4.26 b	0.176 b	19.91 a
		A2	32.12 b	43.01 a	4.64 a	0.162 a	19.67 b

[√]: Means within the same column followed by the same letter are not significantly different.

C0: Control plot (no pass); C1: one pass; C2: two passes, N0: control plot (0 Mg ha⁻¹ manure amount), N40: 40 Mg ha⁻¹ manure amount, N80: 80 Mg ha⁻¹ manure amount, A1: surface manure application, A2: subsurface manure application, ns: nonsignificant.




Figure 2. Effect of wheel traffic on soil moisture and soil temperature. C0: Control plot (no pass); C1: one pass; C2: two passes.




Figure 3. Effect of wheel traffic on soil CO₂ emissions on Julian days. C0: Control plot (no pass); C1: one pass; C2: two passes. PAR: Photosynthetically active radiation.




Figure 4. The changes of CO₂ emission and PAR depending on manure application form on Julian days. PAR: Photosynthetically active radiation.

increase in the amount of manure resulted in higher CO₂ emission. Kimble et al. (1995), using manure at 1 t C ha⁻¹, could allocate more or less 2–5 t CO₂ ha⁻¹ year⁻¹ and the net benefits of manure application ranged from 5.3 to 7.2 t CO₂ ha⁻¹ year⁻¹ equivalent, with higher gains for SOC-poor than SOC-rich sites. Dong and Ouyang (2005) investigated

effects of chicken manure, swine waste, and cattle manure on the CO₂ and CH₄ emissions of a farmland planted with summer maize. Their results showed that CO₂ emissions had the same trends under different organic manure applications, which were influenced by soil temperature and soil water content.




Figure 5. Effects of manure amount on CO₂ emissions on Julian days.

N0: Control plot (no manure), N40: 40 Mg ha⁻¹ manure amount; N80: 80 Mg ha⁻¹ manure amount, PAR: photosynthetically active radiation.

Manure amounts and their interactions with the O₂ content were highly significant ($P < 0.01$) but this effect was not observed in the first year of the experiment (Table 5). Generally, in the experiments, increasing wheel traffic level decreased soil O₂ content (Table 6). Similar results were found in many other studies. In these studies, microbial processes took place in more anaerobic conditions in compacted soils, documented by lower O₂ diffusion rate (Torbert and Wood, 1992; Whalley et al., 1995). In the second year of the experiment increased manure amount decreased soil O₂ content. However, subsurface application also decreased the soil O₂ content (Table 6). When the results of interactions for the 2014 experimental year were examined, the lowest O₂ content was determined in the plot with C2 × A1 × N80.

In the compacted soils, physical changes in pore systems commonly led to less favorable conditions for toxic microbes. This situation was mainly attributed to restriction of gas fluxes due to reduced porosity (Otten et al., 2000; Beylich et al., 2010). Decreased soil microbial activity reduced soil respiration levels. Changes in CO₂ emissions from structured soils were shown to be highly variable and highly affected by structural changes

influenced by mechanical loading and soil management (Mordhorst et al., 2014). Soil compaction mostly leads to reduced CO₂ emissions. Goutal et al. (2012) and Akbolat (2009) stated that lower cumulative soil CO₂ fluxes were recorded plots exposed to traffic compared to the control plots.

3.4. Conclusions

In agricultural practices, soil compaction, manure amount, and manure application techniques directly affect agricultural production. In this study, increased wheel traffic levels caused a reduction in CO₂ emission and O₂ content. In general, due to increase of wheel traffic, the penetration resistance and soil bulk density increases. In this case, CO₂ emissions and soil O₂ capacity are reduced. Increased manure amount increased both CO₂ emissions and soil moisture but decreased soil temperature. As an expected result, the increased carbon level in the soil led to an increase in CO₂ emission. In addition to these results, the subsurface manure application increased CO₂ emissions and soil moisture. Subsurface manure application is advisable for fertilization in terms of CO₂ emissions, but in this situation soil organic matter may be more slowly soluble compared to surface manure application.

References

- Akbolat D (2009). The effect of scrapper usage on soil CO₂ emission during the seedbed preparation. Süleyman Demirel Üniversitesi Ziraat Fakültesi Dergisi 4: 23-30 (in Turkish with abstract in English).
- Altikat S (2013). Effects of aggregate size and compaction level on CO₂-C fluxes and microbial populations. Journal of the Agricultural Faculty of Gaziosmanpaşa University 30: 55-61.
- Altikat S, Celik A, Bilen S (2012). Effects of various tillage systems on soil CO₂-C fluxes and on bacteria and fungi populations in Zea mays. Afr J Agr Res 7: 2926-2934.
- Altikat S, Çelik A, Turgut N (2006). The effects of different soil particle size distribution and seed row compaction levels on the germination and yield of spring wheat. Atatürk Üniversitesi Ziraat Fakültesi Dergisi 37: 197-205 (in Turkish with abstract in English).
- Bakker DM, Davis RJ (1995). Soil deformation observations in a Vertisol under field traffic. Soil Res 33: 817-832.
- Balbuena RH, Terminiello AM, Claverie JA, Casado JP, Marlats R (2000). Soil compaction by forestry harvester operation: evolution of physical properties. Rev Bras Eng Agr Amb 4: 453-459.
- Beylich A, Oberholzer HR, Schrader S, Höper H, Wilke BM (2010). Evaluation of soil compaction effects on soil biota and soil biological processes in soils. Soil Tillage Res 109: 133-143.
- Bilen S, Çelik A, Altikat S (2010). Effects of strip and full-width tillage on soil carbon IV oxide-carbon (CO₂-C) fluxes and on bacterial and fungal populations in sunflower. Afr J Biotechnol 9: 6312-6319.
- Bouwman AF, Boumans LJM, Batjes NH (2002). Emissions of N₂O and NO from fertilized fields: summary of available measurement data. Glob. Biogeochem Cycles 16: 1058-1071.
- Çelik A, Altikat S (2010). Effects of various strip widths and tractor forward speeds in strip tillage on soil physical properties and yield of silage corn. Journal of Agricultural Sciences 16: 169-179.
- Daverede IC, Kravchenko AN, Hoefl RG, Nafziger ED, Bullock DG, Warren JJ, Gonzini LC (2004). Phosphorus runoff from incorporated and surface applied liquid swine manure and phosphorus manure. J Environ Qual 33: 1535-1544.
- Dong Y, Ouyang Z (2005). Effects of organic manures on CO₂ and CH₄ fluxes of farmland. Ying Yong Sheng Tai Xue Bao 16: 1303-1307 (in Chinese with abstract in English).
- Fangueiro D, Senbayran M, Trindade H, Chadwick D (2008). Cattle slurry treatment by screw press separation and chemically enhanced settling: effect on greenhouse gas emissions after land spreading and grass yield. Bioresource Tech 99: 7132-7142.
- Goutal F, Parent B, Bonnaud J, Demaison G, Nourrisso D, Epron, J (2012). Soil CO₂ content and efflux as affected by heavy traffic in forest in northeast France. Eur J Soil Sci 63: 261-271.
- Gregorich EG, Rochette P, McGuire S, Liang BC, Lessard R (1998). Soluble organic carbon and carbon dioxide fluxes in maize fields receiving spring-applied manure. J Environ Qual 27 : 209-214.
- Grigal DF (2000). Effects of extensive forest management on soil productivity. For Ecol Manage 138: 167-185.
- Grote JB, Al-Kaisi MM (2007). Topsoil placement effect on soil carbon stock improvement of exposed subsoil in Iowa. J Soil Water Conserv 62 : 86-93.
- Guérif J, Richard G, Dürr C, Machet JM, Recous S, Roger-Estrade J (2001). A review of tillage effects on crop residue management, seedbed conditions and seedling establishment. Soil Tillage Res 61: 13-32.
- Haynes RJ, Naidu R (1998). Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: a review. Nutr Cycl Agroecosyst 51: 123-137.
- Jorajuria D, Laura D (2000). Sobrecompactación del suelo agrícola parte I: influencia diferencial del peso y del número de pasadas. Rev Bras Eng Agr Amb 4: 445-452 (in Portuguese).
- Kimble MJ, Levine RE, Stewart BA (1995). Soil Management and Greenhouse Effect (Advances in Soil Science). Boca Raton, FL, USA: CRC Press.
- Li Q, Allen HL, Arthur G, Wollum AG (2004). Microbial biomass and bacterial functional diversity in forest soils: effects of organic matter removal, compaction, and vegetation control. Soil Tillage Res 36: 571-579.
- Misselbrook TH, Laws JA, Pain BF (1996). Surface application and shallow injection of cattle slurry on grassland: nitrogen losses, herbage, yield and nitrogen recoveries. Grass Forage Sci 51: 270-277.
- Mordhorst A, Peth S, Horn R (2014). Influence of mechanical loading on static and dynamic CO₂ efflux on differently textured and managed Luvisols. Geoderma 220: 1-13.
- Otten W, Longstaff D, Watts CW (2000). Method to quantify short-term dynamics in carbon dioxide emission following controlled soil deformation. Soil Sci Soc Am J 64: 1740-1748.
- Phillips RL (2007). Organic agriculture and nitrous oxide emissions at subzero soil temperatures. J Environ Qual 36: 23-30.
- Rao DLN, Pathak H (1996). Ameliorative influence of organic matter on biological activity of salt affected soils. Arid Land Res Manag 10: 311-319.
- Rengasamy P (2000). Subsoil constraints and agricultural productivity. J Indian Soc Soil Sci 48: 674-682.
- Rochette P, Gregorich EG (1998). Dynamics of soil microbial biomass C, soluble organic C and CO₂ evolution after three years of manure application. Can J Soil Sci 78: 283-290
- Şeker C, Işıldar A (2000). Effects of wheel traffic porosity and compaction of soil profile. Turk J Agric For 24: 71-77 (in Turkish with abstract in English).
- Torbert HA, Wood CW (1992). Effects of soil compaction and water-filled pore space on soil microbial activity and N losses. Commun Soil Sci Plant Anal 23: 1321-1331.
- Usowicz B, Kossowski J, Baranowski P (1996). Spatial variability of soil thermal properties in cultivated fields. Soil Tillage Res 39: 85-100.
- Whalley WR, Dumitru E, Dexter AR (1995). Biological effects of soil compaction. Soil Tillage Res 35: 53-68.