

1-1-2020

## Dynamics of a fluid equation with Neumann boundary conditions

Limin Chen

SHENGJUN LI

YUMEI ZOU

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---




### Recommended Citation

Chen, Limin; LI, SHENGJUN; and ZOU, YUMEI (2020) "Dynamics of a fluid equation with Neumann boundary conditions," *Turkish Journal of Mathematics*: Vol. 44: No. 5, Article 27. <https://doi.org/10.3906/mat-2002-115>

Available at: <https://journals.tubitak.gov.tr/math/vol44/iss5/27>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Dynamics of a fluid equation with Neumann boundary conditions

Limin CHEN<sup>1</sup> , Shengjun LI<sup>2,\*</sup> , Yumei ZOU<sup>3</sup> 

<sup>1</sup>Department of Basic Course, Changzhou Vocational Institute of Engineering, Changzhou, China

<sup>2</sup>School of Science, Hainan University, Haikou, China

<sup>3</sup>Department of Statistics and Finance, Shandong University of Science and Technology, Qingdao, China

Received: 27.02.2020

Accepted/Published Online: 28.07.2020

Final Version: 21.09.2020

**Abstract:** We study the dynamics of a Neumann boundary value problem arising in fluid dynamics. We prove the nonexistence, existence and uniqueness of positive solutions under suitable conditions. At the same time, under stricter conditions, we also obtain the dynamic properties of the Neumann boundary value problem, such as the stability and instability of positive solutions. The methods of proof mainly involve the upper and lower solutions method, eigenvalue theory and some analysis techniques.

**Key words:** Dynamic analysis, Neumann conditions, fluid dynamics

### 1. Introduction

In this paper, we study the dynamics of positive solutions of the Neumann boundary value problem

$$\begin{cases} -u'' + a(x)u = b(x)u^p, & x \in (0, 1), \\ u'(0) = u'(1) = 0, \end{cases} \quad (1.1)$$

where  $p > 0$ ,  $p \neq 1$ ,  $a$  and  $b$  are continuous positive functions on  $[0, 1]$ . By the upper and lower solutions method, eigenvalue theory and some analysis techniques, we will establish the existence and dynamic properties for (1.1).

Many papers, such as [9, 18, 19] and the references therein, have been devoted to the study of the existence and multiplicity of positive solutions for Neumann boundary value problem using degree theory, fixed point theorem and upper and lower solutions methods.

We study problem (1.1) for three motivations. Firstly, (1.1) has proved to be important in applications. For example, fluid dynamics [1, 10] can be described by

$$\begin{cases} -u'' + q^2u = (1 + \sin \pi x)u^2, & x \in (0, 1), \quad q > 0, \\ u'(0) = u'(1) = 0. \end{cases} \quad (1.2)$$

Subsequently, Bensedik and Boucekif [2] and Torres [13] discussed the existence and symmetry of positive solutions of such problems. Secondly, positivity of Green's function plays an important role in finding solutions to problem (1.2) in the literature [2, 13]. However, in this paper, we do not need to depend on the properties of

\*Correspondence: shjli626@126.com

2010 AMS Mathematics Subject Classification: 34C25, 34B18, 34D20, 34B60

Green’s function, and make full use of the characteristics of problem (1.1) to study. Finally, previous results on Neumann boundary value problem are concerned with existence, but few on dynamical properties [12, 19], such as stability and instability. In this paper, inspired by the stability study of periodic boundary value problems [3–5, 15–17], we will continue to discuss this topic and study the existence and dynamic analysis of positive solutions for a fluid dynamics equation with Neumann boundary value problem (1.1).

Throughout the paper, it is assumed that  $u$  is a continuous function and denote

$$u_M = \max_{x \in [0,1]} u(x), \quad u_m = \min_{x \in [0,1]} u(x),$$

and  $\Delta_u = \frac{u_M}{u_m}$  when  $u_m > 0$ . The remaining part of the paper is organized as follows. In section 2, some preliminary results are given and in section 3, the existence results are proved. Finally, section 4 studies the dynamic properties of problem (1.1).

## 2. Preliminaries

Let  $\lambda_1(s) < \lambda_2(s) \leq \lambda_3(s) \leq \dots$  all be eigenvalues of the linear equation

$$\begin{cases} u'' + [s(x) + \lambda]u = 0, & x \in (0, 1), \\ u'(0) = u'(1) = 0. \end{cases} \tag{2.1}$$

It is well known that the first eigenvalue  $\lambda_1(s)$  is simple and the corresponding eigenfunction does not change sign. When  $s(x) \equiv 0$ , the first eigenvalue is equal to 0 and the second eigenvalue is equal to  $\pi^2$ . Further properties of eigenvalues concerning (2.1) can be found in [7].

Consider the following Neumann boundary value problem

$$\begin{cases} u'' + g(x, u) = f(x), & x \in (0, 1), \\ u'(0) = u'(1) = 0. \end{cases} \tag{2.2}$$

**Definition 2.1** [19] *A function  $\alpha \in C^2[0, 1]$  is called a lower solution of (2.2) if  $\alpha$  satisfies*

$$\begin{cases} \alpha'' + g(x, \alpha) \geq f(x), & x \in (0, 1), \\ \alpha'(0) \geq 0 \geq \alpha'(1). \end{cases} \tag{2.3}$$

*A function  $\beta \in C^2[0, 1]$  is called an upper solution of (2.2) if  $\beta$  satisfies*

$$\begin{cases} \beta'' + g(x, \beta) \leq f(x), & x \in (0, 1), \\ \beta'(0) \leq 0 \leq \beta'(1). \end{cases} \tag{2.4}$$

**Lemma 2.2** [12] *Suppose that  $u$  is the solution of (2.2), then  $u$  is stable if the first eigenvalue  $\lambda_1(g'_u(x, u))$  of the equation*

$$\begin{cases} \varphi'' + [g'_u(x, u) + \lambda]\varphi = 0, & x \in (0, 1), \\ \varphi'(0) = \varphi'(1) = 0, \end{cases} \tag{2.5}$$

*is strictly positive. The solution  $u$  is unstable if the first eigenvalue  $\lambda_1(g'_u(x, u))$  is strictly negative.*

The following two lemmas can be deduced directly from the differential mean value theorem combined with Lemmas 2.3 and 2.4 in [19].

**Lemma 2.3** *Let  $\alpha, \beta \in C^2[0, 1]$  be the lower and upper solutions of (2.2) for all  $x \in [0, 1]$  with  $\alpha(x) \leq \beta(x)$ . Suppose that there exists sufficiently large  $L > 0$  such that  $g'_u(x, u) > -L$ , for all  $x \in [0, 1]$  with  $\alpha(x) \leq u \leq \beta(x)$ . Then the problem (2.2) has a solution  $u \in [\alpha_m, \beta_M]$ .*

**Lemma 2.4** *Let  $\alpha, \beta \in C^2[0, 1]$  be the lower and upper solutions of (2.2) for all  $x \in [0, 1]$  with  $\beta(x) \leq \alpha(x)$ . Suppose that*

$$g'_u(x, u) < \frac{\pi^2}{4}, \tag{2.6}$$

for all  $x \in [0, 1]$  with  $\beta(x) \leq u \leq \alpha(x)$ . Then the problem (2.2) has a solution  $u \in [\beta_m, \alpha_M]$ .

**Lemma 2.5** [19] *Let  $[w, v] = \{u \in C^2[0, 1] | w \leq u \leq v\}$ . Assume that for all  $x \in [0, 1]$  and  $u \in [w, v]$ , (2.6) holds. Then*

- (1) (2.2) has a unique solution on  $[w, v]$  if a solution exists and  $g'_u(x, u) < 0$  on  $[w, v]$ .
- (2) (2.2) has a unique solution on  $[w, v]$  if a solution exists and  $g'_u(x, u) > 0$  on  $[w, v]$ .

### 3. Existence analysis for problem (1.1)

#### 3.1. Nonexistence

Integrating the equation in the problem (1.1) on  $[0, 1]$ , and taking into account the boundary conditions, we get a necessary condition for the existence of positive solutions, namely

$$\int_0^1 a(x)u(x)dx = \int_0^1 b(x)u^p(x)dx.$$

So it is easy to obtain the following nonexistence result.

**Theorem 3.1** *If one of the following conditions holds*

- (1)  $a_m \geq 0$  and  $b_M < 0$ ,
- (2)  $a_m > 0$  and  $b_M \leq 0$ ,
- (3)  $a_M \leq 0$  and  $b_m > 0$ ,
- (4)  $a_M < 0$  and  $b_m \geq 0$ ,

then problem (1.1) does not have positive solutions.

#### 3.2. Existence

The method of lower and upper solutions is a quite useful and flexible tool for investigating not only the existence, localization and multiplicity of periodic solutions, but also qualitative properties, such as stability and instability. The monograph [6] presents a nice and complete historical review of the subject. We also refer to [8, 11, 14] for details and further references related to the method of lower and upper solutions.

**Theorem 3.2** Let  $0 < p < 1$ , constants  $c_1, c_2, d_1, d_2 \in (0, +\infty)$ , functions  $\omega, \sigma \in C^2[0, 1]$  be such that the equalities

$$\ddot{\omega} = c_1 b(x) - c_2 a(x), \quad x \in (0, 1), \quad \omega'(0) = \omega'(1) = 0, \tag{3.1}$$

and

$$\ddot{\sigma} = d_1 b(x) - d_2 a(x), \quad x \in [0, 1], \quad \sigma'(0) = \sigma'(1) = 0 \tag{3.2}$$

are fulfilled and let there exist  $u_0 \in (0, +\infty)$  such that

$$u_0(\omega(x) - m_\omega) + \sigma(x) - m_\sigma \leq c_2 u_0 + d_2 - (c_1 u_0 + d_1)^{1/p} \text{ for } x \in [0, 1]. \tag{3.3}$$

Then problem (1.1) has at least one positive solution.

**Proof** In order to prove Theorem 3.2 from Lemma 2.3, only a pair of positive well-ordered lower and upper solutions need to be constructed.

*Step 1. Construction of a lower solution.*

Put

$$\alpha(x) = c_2 u_0 + d_2 - [u_0(\omega(x) - m_\omega) + \sigma(x) - m_\sigma] \text{ for } x \in [0, 1].$$

Then, obviously  $\alpha \in C^2[0, 1]$  and in view of (3.1) and (3.2) we have

$$\alpha'' = (c_2 u_0 + d_2)a(x) - (c_1 u_0 + d_1)b(x) \text{ for } x \in [0, 1]. \tag{3.4}$$

Moreover, according to (3.3),

$$(c_1 u_0 + d_1)^{1/p} \leq \alpha(x) \leq c_2 u_0 + d_2 \text{ for } x \in [0, 1]. \tag{3.5}$$

Now (3.4) and (3.5) imply

$$\alpha'' + b(x)\alpha^p - a(x)\alpha \geq 0 \text{ for } x \in [0, 1].$$

Consequently,  $\alpha$  is a lower function to (1.1).

*Step 2. Construction of an upper solution.*

Note that the inequality  $0 < p < 1$  implies

$$\lim_{u \rightarrow +\infty} [(c_1 u + d_1)^{1/p} - (c_2 u + d_2)] = +\infty.$$

Therefore, we can choose  $u_1 > u_0$  such that

$$u_1(\omega(x) - m_\omega) + \sigma(x) - m_\sigma \leq (c_1 u_1 + d_1)^{1/p} - (c_2 u_1 + d_2) \text{ for } x \in [0, 1] \tag{3.6}$$

and put

$$\beta(x) = (c_1 u_1 + d_1)^{\frac{1}{p}} - [u_1(\omega(x) - m_\omega) + \sigma(x) - m_\sigma] \text{ for } x \in [0, 1]. \tag{3.7}$$

Then,  $\beta \in C^2[0, 1]$  and in view of (3.1) and (3.2) we have

$$\beta'' = (c_2 u_1 + d_2)a(x) - (c_1 u_1 + d_1)b(x) \text{ for } x \in [0, 1]. \tag{3.8}$$

Moreover, according to (3.6) and (3.7),

$$c_2u_1 + d_2 \leq \beta(x) \leq (c_1u_1 + d_1)^{1/p}. \tag{3.9}$$

Now (3.8) and (3.9) imply

$$\beta'' + b(x)\beta^p - a(x)\beta \leq 0 \text{ for } x \in [0, 1].$$

Consequently,  $\beta$  is an upper function to (1.1).

Moreover, (3.5) and (3.9) imply

$$0 < \alpha(x) \leq \beta(x) \text{ for } x \in [0, 1].$$

Thus  $\alpha$  and  $\beta$  are a pair of well-ordered lower and upper solutions. □

**Theorem 3.3** *Let  $p > 1$ , constants  $c_1, c_2, d_1, d_2 \in (0, +\infty)$ , functions  $\omega, \sigma \in C^2[0, 1]$  be such that the equalities (3.1) and (3.2) are fulfilled and let there exist  $u_0 \in (0, +\infty)$  such that*

$$u_0(\omega(x) - m_\omega) + \sigma(x) - m_\sigma \leq (c_1u_0 + d_1)^{1/p} - (c_2u_0 + d_2) \text{ for } x \in [0, 1]. \tag{3.10}$$

*Then problem (1.1) has at least one positive solution provided that*

$$pb_M\alpha_M^{p-1} - a_m < \frac{\pi^2}{4}, \tag{3.11}$$

where  $\alpha$  is defined by (3.16).

**Proof** The existence is proved using Lemma 2.4. To do so, let us construct a pair of unwell-ordered lower and upper solutions.

Let

$$\beta(x) = (c_1u_0 + d_1)^{\frac{1}{p}} - [u_0(\omega(x) - m_\omega) + \sigma(x) - m_\sigma] \text{ for } x \in [0, 1]. \tag{3.12}$$

Then,  $\beta \in C^2[0, 1]$  and in view of (3.1) and (3.2) we have

$$\beta'' = (c_2u_0 + d_2)a(x) - (c_1u_0 + d_1)b(x) \text{ for } x \in [0, 1]. \tag{3.13}$$

Moreover, according to (3.10) and (3.12),

$$c_2u_0 + d_2 \leq \beta(x) \leq (c_1u_0 + d_1)^{1/p}. \tag{3.14}$$

Now (3.13) and (3.14) imply

$$\beta'' + b(x)\beta^p - a(x)\beta \leq 0 \text{ for } x \in [0, 1].$$

Consequently,  $\beta$  is an upper function to (1.1).

Note that the inequality  $p > 1$  implies

$$\lim_{u \rightarrow +\infty} [c_2u + d_2 - (c_1u + d_1)^{1/p}] = +\infty.$$

Therefore, we can choose  $u_1 > u_0$  such that

$$u_1(\omega(x) - m_\omega) + \sigma(x) - m_\sigma \leq c_2u_1 + d_2 - (c_1u_1 + d_1)^{1/p} \text{ for } x \in [0, 1] \tag{3.15}$$

and put

$$\alpha(x) = c_2u_1 + d_2 - [u_1(\omega(x) - m_\omega) + \sigma(x) - m_\sigma] \text{ for } x \in [0, 1]. \tag{3.16}$$

Then, obviously  $\alpha \in C^2[0, 1]$  and in view of (3.1) and (3.2) we have

$$\alpha'' = (c_2u_1 + d_2)a(x) - (c_1u_1 + d_1)b(x) \text{ for } x \in [0, 1]. \tag{3.17}$$

Moreover, according to (3.15),

$$(c_1u_1 + d_1)^{1/p} \leq \alpha(x) \leq c_2u_1 + d_2 \text{ for } x \in [0, 1]. \tag{3.18}$$

Now (3.17) and (3.18) imply

$$\alpha'' + b(x)\alpha^p - a(x)\alpha \geq 0 \text{ for } x \in [0, 1].$$

Consequently,  $\alpha$  is a lower function to (1.1).

Moreover, (3.14) and (3.18) imply

$$0 < \beta(x) \leq \alpha(x) \text{ for } x \in [0, 1].$$

Thus  $\alpha$  and  $\beta$  are a pair of unwell-ordered lower and upper solutions.

Let us fix

$$g(x, u) = b(x)u^p - a(x)u.$$

Then by  $p > 1$ , for  $u \in [\beta_m, \alpha_M]$ , we have

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\leq pb_M\alpha_M^{p-1} - a_m. \end{aligned}$$

The above inequality and (3.11) imply that condition (2.6) of Lemma 2.4 is satisfied, so the result is guaranteed. □

**Theorem 3.4** *Assume that  $0 < p < 1, a_m > 0$ . Then problem (1.1) has at least one solution such that*

$$\left(\frac{b_m}{a_M}\right)^{\frac{1}{1-p}} \leq u(x) \leq \left(\frac{b_M}{a_m}\right)^{\frac{1}{1-p}}.$$

**Proof** Note that

$$\beta \equiv \left(\frac{b_M}{a_m}\right)^{\frac{1}{1-p}}$$

is a constant upper function and

$$\alpha \equiv \left(\frac{b_m}{a_M}\right)^{\frac{1}{1-p}}$$

is a constant lower function on the well order  $\alpha < \beta$ .

Note that

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\geq pb(x)\beta^{p-1} - a(x) \\ &\geq pb_m \frac{a_m}{b_M} - a_M. \end{aligned}$$

Take  $L = a_M - pb_m \frac{a_m}{b_M}$ . Then  $L > 0$  because  $p < 1$ . By Lemma 2.3, problem (1.1) has at least one solution. □

**Theorem 3.5** Assume that  $p > 1, b_m > 0$  and the following inequality holds

$$4pb_M \cdot \frac{a_M}{b_m} - 4a_m < \pi^2. \tag{3.19}$$

Then problem (1.1) has at least one solution such that

$$\left(\frac{a_m}{b_M}\right)^{\frac{1}{p-1}} \leq u(x) \leq \left(\frac{a_M}{b_m}\right)^{\frac{1}{p-1}}.$$

**Proof** Note that

$$\beta \equiv \left(\frac{a_m}{b_M}\right)^{\frac{1}{p-1}}$$

is a constant upper function and

$$\alpha \equiv \left(\frac{a_M}{b_m}\right)^{\frac{1}{p-1}}$$

is a constant lower function on the reserved order  $\alpha > \beta$ .

Note that

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\leq pb(x)\alpha^{p-1} - a(x) \\ &\leq pb_M \cdot \frac{a_M}{b_m} - a_m. \end{aligned}$$

By Lemma 2.4, a sufficient condition for the existence of solution of (1.1) is

$$pb_M \cdot \frac{a_M}{b_m} - a_m < \frac{\pi^2}{4},$$

which is equivalent to condition (3.19). □

#### 4. Dynamic analysis for problem (1.1)

In this section, based on the existence of positive solutions obtained in the previous section, the dynamic behaviors of equation (1.1) are further studied, including stability and instability.



#### 4.1. Stability

**Theorem 4.1** *Under the conditions of Theorem 3.2, if  $a_m - pb_M \cdot (c_1u_0 + d_1)^{\frac{p-1}{p}} > 0$ , then the solution  $u$  provided by Theorem 3.2 is unique and stable.*

**Proof** From  $a_m - pb_M \cdot (c_1u_0 + d_1)^{\frac{p-1}{p}} > 0$ , for  $u \in [\alpha_m, \beta_M]$ , we have

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\leq pb(x)\alpha_m^{p-1} - a(x) \\ &\leq pb_M \cdot (c_1u_0 + d_1)^{\frac{p-1}{p}} - a_m \\ &< 0, \end{aligned}$$

it follows from the second conclusion of Lemma 2.5 that (1.1) has a unique solution  $u$ . By the comparison theorem of eigenvalue [7] we have  $\lambda_1(g'_u(x, u)) > 0$ , so the unique solution  $u$  is stable. □

**Theorem 4.2** *Under the conditions of Theorem 3.4, if moreover  $b_m > 0$  and  $p\Delta < 1$ , then the solution  $u$  provided by Theorem 3.4 is unique and stable.*

**Proof** From  $0 < p\Delta < 1$ , for  $u \in [\alpha, \beta]$ , we have

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\leq pb(x)\alpha^{p-1} - a(x) \\ &\leq pb_M \cdot \frac{a_M}{b_m} - a_m \\ &< 0. \end{aligned}$$

The rest of the proof is similar to that in the proof of Theorem 4.1, so we omit the details. □

#### 4.2. Instability

**Theorem 4.3** *Under the conditions of Theorem 3.3, if  $pb_m \cdot (c_2u_0 + d_2)^{p-1} - a_M > 0$ , then the solution  $u$  provided by Theorem 3.3 is unique and unstable.*

**Proof** From  $pb_m \cdot (c_2u_0 + d_2)^{p-1} - a_M > 0$ , for  $u \in [\beta_m, \alpha_M]$ , we have

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\geq pb(x)\beta_m^{p-1} - a(x) \\ &\geq pb_m \cdot (c_2u_0 + d_2)^{p-1} - a_M \\ &> 0. \end{aligned}$$

Therefore, it follows from the third conclusion of Lemma 2.5 that (1.1) has a unique solution  $u$ . By the comparison theorem of eigenvalue we have  $\lambda_1(g'_u(x, u)) < 0$ , so the unique solution  $u$  is unstable. □

**Theorem 4.4** *Under the conditions of Theorem 3.5, if moreover  $a_m > 0$  and  $p > \Delta \geq 1$ , then the solution  $u$  provided by Theorem 3.5 is unique and unstable.*

**Proof** From  $p > \Delta$ , for  $u \in [\beta_m, \alpha_M]$ , we have

$$\begin{aligned} g'_u(x, u) &= pb(x)u^{p-1} - a(x) \\ &\geq pb(x)\beta_m^{p-1} - a(x) \\ &\geq pb_m \cdot \frac{a_m}{b_M} - a_M \\ &> 0. \end{aligned}$$

We follow the same strategy as in the proof of Theorem 4.3. □

**Example 4.5** *Let the coefficients in (1.1) be*

$$a(x) \equiv q^2, \text{ and } b(x) = 1 + \sin \pi x, \quad 0 \leq x \leq 1, \quad q > 0.$$

(1) (1.1) has at least one positive solution if  $0 < p < 1$ .

(2) (1.1) has a unique stable solution if  $0 < p < \frac{1}{2}$ .

**Proof** First we prove (1). It is easy to see that  $a_m > 0$ . The result follows from Theorem 3.4. Next we prove (2). Notice that  $\Delta = 2$ , so the uniqueness and stability are obtained from Theorem 4.2. □

**Remark 4.6** *According to Theorem 3.5, it is easy to obtain the existence of positive solutions of the problem 1.2, but its further dynamic properties are still an open problem for us, because the condition of Theorem 4.4 is not satisfied.*

## Acknowledgment

This work was supported by the National Natural Science Foundation of China (Grant No. 11861028,11501055).

## References

- [1] Benjamin TB. A new kind of solitary wave. *Journal of Fluid Mechanics* 1992; 245 (1): 401-411. doi: 10.1017/S002211209200051X
- [2] Bensedik A, Bouchekif M. Symmetry and uniqueness of positive solutions for a Neumann boundary value problem. *Applied Mathematics Letters* 2007; 20 (4): 419-426. doi: 10.1016/j.aml.2006.05.008
- [3] Chu J, Torres PJ, Wang F. Radial stability of periodic solutions of the Gylden-Meshcherskii-type problem. *Discrete and Continuous Dynamical Systems - Series A* 2015; 35 (5): 1921-1932. doi: 10.3934/dcds.2015.35.1921
- [4] Chu J, Torres PJ, Wang F. Twist periodic solutions for differential equations with a combined attractive-repulsive singularity. *Journal of Mathematical Analysis and Applications* 2016; 437 (2): 1070-1083. doi: 10.1016/j.jmaa.2016.01.057
- [5] Chu J, Wang F. Prevalence of stable periodic solutions for Duffing equations. *Journal of Differential Equations* 2016; 260 (11): 7800-7820. doi: 10.1016/j.jde.2016.02.003

- [6] De Coster C, Habets P. Two-point boundary value problems: lower and upper solutions. In: Chui CK (editor). *Mathematics in Science and Engineering*, Vol. 205. Amsterdam, Netherlands: Elsevier, 2006.
- [7] Hale JK. *Ordinary differential equations*. Malabar, FL, USA: Krieger Publishing Company, 1969.
- [8] Hakl R, Torres PJ. On periodic solutions of second-order differential equations with attractive-repulsive singularities. *Journal of Differential Equations* 2010; 248 (1): 111–126. doi: 10.1016/j.jde.2009.07.008
- [9] Li S, Xiu Z, Liang L. Positive solutions for nonlinear singular second order Neumann boundary value problems. *Journal of Mathematical and Computational Science* 2012; 5: 1353-1362.
- [10] Mays L, Norbury J. Bifurcation of positive solutions for a Neumann boundary value problem. *The ANZIAM Journal* 2001; 42: 327-340. doi: 10.1017/S1446181100011974
- [11] Njoku FI and Omari P. Stability properties of periodic solutions of a Duffing equation in the presence of lower and upper solutions. *Applied Mathematics and Computation* 2003; 135 (2-3): 471-490. doi: 10.1016/s0096-3003(02)00062-0
- [12] Oruganti S, Shi J, Shivaji R. Diffusive logistic equation with constant yield harvesting, I: Steady States. *Transactions of the American Mathematical Society* 2002; 354: 3601-3619. doi: 10.2307/3072927
- [13] Torres PJ. Some remarks on a Neumann boundary value problem arising in fluid dynamics. *The ANZIAM Journal* 2004; 45 (3): 327-332. doi: 10.1017/S1446181100013390
- [14] Torres PJ. Existence and stability of periodic solutions of a Duffing equation by using a new maximum principle. *Mediterranean Journal of Mathematics* 2004; 1 (4): 479-486. doi: 10.1007/s00009-004-0025-3.
- [15] Wang F, Cid JA, Zima M. Lyapunov stability for regular equations and applications to the Liebau phenomenon, *Discrete and Continuous Dynamical Systems* 2018; 38 (9): 4657-4674. doi: 10.3934/dcds.2018204
- [16] Wang F, Li S, Zou Y. A relationship between linear stability and index of oscillations for Lagrangian equation at higher order zones, *Nonlinear Analysis: Real World Applications* 2019; 47: 178-187. doi: 10.1016/j.nonrwa.2018.10.014
- [17] Wang F, Zhu H. Existence, uniqueness and stability of periodic solutions of a Duffing equation under periodic and anti-periodic eigenvalues conditions. *Taiwanese Journal of Mathematics* 2015; 19 (5): 1457-1468. doi: 10.11650/tjm.19.2015.3992
- [18] Wang F, Zhang F, Yu Y. Existence of positive solutions of Neumann boundary value problem via a convex functional compression-expansion fixed point theorem. *Fixed Point Theory* 2010; 11: 395-400.
- [19] Xing H, Chen H, He X. Exact multiplicity and stability of solutions of second-order Neumann boundary value problem. *Applied Mathematics and Computation* 2014; 232: 1104-1111. doi: 10.1016/j.amc.2014.01.119