

1-1-2020

## On Ulam's type stability criteria for fractional integral equations including Hadamard type singular kernel

YASEMİN BAŞCI

SÜLEYMAN ÖĞREKÇİ

ADİL MISIR

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)




---

### Recommended Citation

BAŞCI, YASEMİN; ÖĞREKÇİ, SÜLEYMAN; and MISIR, ADİL (2020) "On Ulam's type stability criteria for fractional integral equations including Hadamard type singular kernel," *Turkish Journal of Mathematics*: Vol. 44: No. 4, Article 30. <https://doi.org/10.3906/mat-1910-70>  
Available at: <https://journals.tubitak.gov.tr/math/vol44/iss4/30>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## On Ulam's type stability criteria for fractional integral equations including Hadamard type singular kernel

Yasemin BAŞCI<sup>1,\*</sup>, Süleyman ÖĞREKÇİ<sup>2</sup>, Adil MISİR<sup>3</sup>

<sup>1</sup>Department of Mathematics, Faculty of Arts and Sciences, Bolu Abant İzzet Baysal University, Bolu, Turkey

<sup>2</sup>Department of Mathematics, Faculty of Arts and Sciences, Amasya University, Amasya, Turkey

<sup>3</sup>Department of Mathematics, Faculty of Sciences, Gazi University, Ankara, Turkey

Received: 18.10.2019

Accepted/Published Online: 10.06.2020

Final Version: 08.07.2020

**Abstract:** In this paper, we deal with the Hyers-Ulam-Rassias (HUR) and Hyers-Ulam (HU) stability of Hadamard type fractional integral equations on compact intervals. The stability conditions are developed using a new generalized metric (GM) definition and the fixed point technique by motivating Wang and Lin Ulam's type stability of Hadamard type fractional integral equations. *Filomat* 2014; 28(7): 1323-1331. Moreover, our approach is efficient and ease in use than to the previously studied approaches. Finally, we give two examples to explain our main results.

**Key words:** Hyers-Ulam stability, Hyers-Ulam-Rassias stability, fractional integral equation, fixed point theory, Hadamard type singular kernel

### 1. Introduction

Although the subject fractional calculus was introduced more than 300 years ago, researches are still working in the development of both theory and application [21, 26, 29, 32]. The applications of fractional calculus has been observed in almost every field of sciences, such as mechanics, electricity, biology, economics, physics, biophysics, control theory, signal processing and image processing (see [13, 25, 30, 31, 33]). In recent years, the HU stability of various fractional differential equations has been widely studied (see [5, 7, 14, 19, 20, 24, 38–40]). The study on HU stability was initiated in 1940 and later on, it was extended to Banach spaces [9]. After that, many researchers put their effort to develop the generalized theory to study the HU stability of various type of differential phenomena (see [2–4, 8, 9, 11, 12, 16–18, 22, 23, 27, 28, 36, 38] and the other stability results [15, 34, 35]).

In 2013, Wang et al.[38] gave Ulam's type stability of fractional differential equations involving Hadamard derivative. They obtained some Ulam-Hyers stability conditions by using the method studied in [28].

In 2014, Wang and Lin [37] investigated the Ulam's type stability for fractional integral equations involving Hadamard type singular kernel on a compact interval by using fixed point method. They extended the developed results of [38] by choosing the closed interval based on the method of [27].

In 2016, Abbas et al.[1], developed the Ulam's stability results for partial integral equations using the Schauder's fixed-point results by taking the Hadamard's fractional integral.

\*Correspondence: basci\_y@ibu.edu.tr

2010 *AMS Mathematics Subject Classification*: 26A33,34A34,34D20,45G05,47H10,55M20

Kilbas et al.[21] examined the Cauchy problem for the following nonlinear fractional differential equation with the Hadamard fractional derivative of complex order  $\alpha_1 \in \mathbb{C}$  ( $\Re(\alpha_1) > 0$ ):

$$(D_{a_1+}^{\alpha_1} y)(x) = u(x, y(x)), (a_1 \leq x \leq b_1) \tag{1.1}$$

$$(D_{a_1+}^{\alpha_1-k} y)(a_1) = b_k, (k = 1, \dots, m) \tag{1.2}$$

where  $m = [\Re(\alpha_1)] + 1$  for  $\alpha_1 \notin \mathbb{N}$  and  $m = \alpha_1$  for  $m \in \mathbb{N}$ ,  $[a_1, b_1]$  is a finite interval of  $\mathbb{R}$ , and

$$(D_{a_1+}^{\alpha_1} y)(x) = \left(x \frac{d}{dx}\right)^m \frac{1}{\Gamma(m - \alpha_1)} \int_{a_1}^x \left(\ln \frac{x}{\eta}\right)^{m-\alpha_1+1} y(\eta) \frac{d\eta}{\eta}.$$

They proved the equivalence of (1.1)-(1.2) and a Volterra integral equation in the following form:

$$y(x) = \sum_{j=1}^m \frac{c_j}{\Gamma(\alpha_1 - j + 1)} \left(\ln \frac{x}{a_1}\right)^{\alpha_1-j} + \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left(\ln \frac{x}{\eta}\right)^{\alpha_1-1} u(\eta, y(\eta)) \frac{d\eta}{\eta}, \tag{1.3}$$

and applied this results to establish conditions for a unique solution of the Cauchy problem (1.1)-(1.2). Here  $m - 1 < \alpha_1 \leq m$  ( $m = 1, 2, \dots$ ),  $a_1$  and  $b_1$  are given constants such that  $0 < a_1 \leq x \leq b_1 < \infty$ . Also,  $c_j$  are fixed real numbers for  $j = 1, 2, \dots, m$ ,  $\Gamma(\cdot)$  is the Gamma function and  $u : [a_1, b_1] \times \mathbb{R} \rightarrow \mathbb{R}$ . In this paper, we deal with the Ulam’s type stability for the fractional integral equations (1.3).

We organized this study as follows: In Section 1, we have given introduction. In Section 2, we introduce some definitions and some theorems which will be useful in proofs of our results. In Section 3, by using the motivation of [37], we investigate HUR and HU stability of the Eq.(1.3) on a compact interval with the help of a new GM definition. In Section 4, we give several examples for our results.

### 2. Preliminaries

Below, we give some definitions and some theorems which will be useful in the proofs of our main results.

**Definition 2.1** Let  $M > 0$  be a given constant and  $\psi$  be a nonnegative function. For every function  $y$  satisfying

$$\left| y(x) - \sum_{j=1}^m \frac{c_j}{\Gamma(\alpha_1 - j + 1)} \left(\ln \frac{x}{a_1}\right)^{\alpha_1-j} - \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left(\ln \frac{x}{\eta}\right)^{\alpha_1-1} u(\eta, y(\eta)) \frac{d\eta}{\eta} \right| \times e^{-M(x-a_1)} \leq \psi(x),$$

there is a solution  $y_1$  of the Eq.(1.3) and  $c > 0$  is a constant which is independent of  $y$  and  $y_1$  such that

$$|y(x) - y_1(x)| e^{-M(x-a_1)} \leq c\psi(x), x \in [a_1, b_1],$$

then the Eq.(1.3) is called HUR stable.

In Definition 2.1, if  $\psi(x)$  takes an arbitrary constant function, then the Eq.(1.3) is called HU stable.

Now, the function  $d$  is called a GM on  $Y$  if and only if  $d : Y \times Y \rightarrow [0, \infty]$  satisfies the following conditions for  $Y \neq 0$ :

- (M<sub>1</sub>)  $d(u, v) = 0$  if and only if  $u = v$ ,
- (M<sub>2</sub>)  $d(u, v) = d(v, u)$  for all  $u, v \in Y$ ,
- (M<sub>3</sub>)  $d(u, w) \leq d(u, v) + d(v, w)$  for all  $u, v, w \in Y$ .

**Theorem 2.2** [10] *Let  $(Y, d)$  be a generalized complete metric (GCM) space. Suppose that  $T : Y \rightarrow Y$  is a strictly contractive operator with the Lipschitz constant  $L < 1$ . If there exists a nonnegative integer  $k$  such that  $d(T^{k+1}x, T^kx) < \infty$  for some  $x \in Y$ , then the following are true:*

- (H<sub>1</sub>) *The sequence  $\{T^n x\}$  converges to a fixed point  $x^*$  of  $T$ ,*
- (H<sub>2</sub>)  *$x^*$  is the unique fixed point of  $T$  in*

$$Y^* = \{y \in Y : d(T^k x, y) < \infty\},$$

- (H<sub>3</sub>) *If  $y \in Y^*$ , then*

$$d(y, x^*) \leq \frac{1}{1-L} d(Ty, y).$$

HU stability of various equations is examined using Theorem 2.2 and  $d$  (GM) defined by

$$d(g, h) := \inf \{C \in [0, \infty] : |g(x) - h(x)| \leq C\psi(x), x \in [a_1, b_1]\}, \tag{2.1}$$

see [17, 18, 36] and therein references.

### 3. Main results

Below, we are interested in HUR and HU stability of the Eq.(1.3) on  $[a_1, b_1]$  such that  $[a_1, b_1]$  is a compact interval. Now, we will give the following lemma which will be used in the proofs of our main results. In this lemma, we use a new GM which is different from (2.1).

Let  $I := [a_1, b_1]$  be an interval for  $a_1 > 0$  and the set  $Y$  be such as

$$Y = C(I, \mathbb{R}). \tag{3.1}$$

**Lemma 3.1** [6] *Let  $d : Y \times Y \rightarrow [0, \infty]$  be a function defined by*

$$d(g, h) := \inf \left\{ C \in [0, \infty] : |g(x) - h(x)| e^{-M(x-a_1)} \leq C\psi(x), x \in I \right\}, \tag{3.2}$$

where  $M > 0$  and  $\psi \in C[I, (0, \infty)]$  are given constant and function, respectively. Then  $(Y, d)$  is a GCM space.

**Proof** Firstly, we show that the function  $d$  defined in (3.2) is a GM on  $Y$ . It is clear that the conditions  $M_1$  and  $M_2$  are satisfied. Now, we show that the condition  $M_3$  also satisfies. For some  $g, h, f \in Y$ , we can find an  $x \in I$  such that

$$\begin{aligned} |g(x) - h(x)| e^{-M(x-a_1)} &= |g(x) - f(x) + f(x) - h(x)| e^{-M(x-a_1)} \\ &\leq |g(x) - f(x)| e^{-M(x-a_1)} + |f(x) - h(x)| e^{-M(x-a_1)}. \end{aligned}$$

Then,  $M_3$  holds.

Now, we will show that  $(Y, d)$  is complete metric space. Let  $\{g_n\}$  be a Cauchy sequence on  $(Y, d)$ . Then, we can find an integer  $N_1(\varepsilon) > 0$  for any  $\varepsilon > 0$  such that  $d(g_m, g_n) \leq \varepsilon$  for all  $m, n \geq N_1(\varepsilon)$ . In other words, there exists an integer  $N_1(\varepsilon) > 0$  for any  $\varepsilon > 0$  such that

$$|g_m(x) - g_n(x)| e^{-M(x-a_1)} \leq \varepsilon \psi(x) \tag{3.3}$$

for all  $m, n \geq N_1(\varepsilon)$  and all  $x \in I$ . This means that  $\{g_n(x)\}$  is a Cauchy sequence in  $\mathbb{R}$  for any fixed  $x$ . Because of  $\mathbb{R}$  is complete,  $\{g_n(x)\}$  converges for all  $x \in I$ . Thus, we have a function

$$g(x) := \lim_{n \rightarrow \infty} g_n(x),$$

where  $g : I \rightarrow \mathbb{R}$ . In (3.3), letting  $m \rightarrow \infty$ , we can find an integer  $N_1(\varepsilon) > 0$  for any  $\varepsilon > 0$  such that

$$|g(x) - g_n(x)| e^{-M(x-a_1)} \leq \varepsilon \psi(x) \tag{3.4}$$

for all  $x \in I$  and  $n \geq N_1(\varepsilon)$ . This means that, it can be find an integer  $N_1(\varepsilon) > 0$  for any  $\varepsilon > 0$  such that  $d(g, g_n) \leq \varepsilon$  for all  $n > N_1(\varepsilon)$ . Additionally, because of  $\psi$  is bounded on  $I$ ,  $\{g_n(x)\}$  converges uniformly to  $g$  with the aid of (3.4) and so that  $g \in Y$ . Thus, the proof is complete.  $\square$

Let  $0 < p_1 < 1$ ,  $m - 1 < \alpha_1 \leq m$ ,  $p_1 \leq m$  and the following conditions hold:

[A<sub>1</sub>] : The function  $u : [a_1, b_1] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous and for any  $\eta \in [a_1, b_1]$  and  $y, z \in \mathbb{R}$ ,

$$|u(\eta, y) - u(\eta, z)| \leq L^* \eta^{p_1} |y - z|. \tag{3.5}$$

[A<sub>2</sub>] : For all  $x \in [a_1, b_1]$ , the continuous function  $y : [a_1, b_1] \rightarrow \mathbb{R}$  satisfies the following inequality:

$$\left| y(x) - \sum_{j=1}^m \frac{c_j}{\Gamma(\alpha_1 - j + 1)} \left(\ln \frac{x}{a_1}\right)^{\alpha_1 - j} - \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left(\ln \frac{x}{\eta}\right)^{\alpha_1 - 1} u(\eta, y(\eta)) \frac{d\eta}{\eta} \right| \times e^{-(L^*+1)(x-a_1)} \leq \psi(x), \tag{3.6}$$

and  $\psi : [a_1, b_1] \rightarrow (0, \infty)$  satisfies

$$\left( \int_{a_1}^x (\psi(\eta))^{2/p_1} d\eta \right)^{p_1/2} \leq K^* \psi(x). \tag{3.7}$$

[A<sub>3</sub>] : Let  $0 < K^* L^* M^* < 1$ , where

$$M^* = \frac{1}{\Gamma(\alpha_1)} \left( \frac{1 - p_1}{\alpha_1 - p_1} \right)^{1-p_1} \left( \frac{p_1}{2(L^* + 1)} \right)^{p_1/2} \left( \ln \frac{b_1}{a_1} \right)^{\alpha_1 - p_1}.$$

Here  $K^*$ ,  $L^*$ ,  $M^*$  values were obtained by (3.2) metric. Similar conditions also exist by using (2.1) metric in [39].

**Theorem 3.2** *Suppose that  $[A_1]$ ,  $[A_2]$  and  $[A_3]$  are satisfied. Then, there exists an unique continuous function  $y_1 : [a_1, b_1] \rightarrow \mathbb{R}$  which satisfies (1.3) and*

$$y_1(x) = \sum_{j=1}^m \frac{c_j}{\Gamma(\alpha_1 - j + 1)} \left(\ln \frac{x}{a_1}\right)^{\alpha_1 - j} + \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left(\ln \frac{x}{\eta}\right)^{\alpha_1 - 1} u(\eta, y_1(\eta)) \frac{d\eta}{\eta} \tag{3.8}$$

and

$$|y(x) - y_1(x)| e^{-(L^*+1)(x-a_1)} \leq \frac{\psi(x)}{1 - K^* L^* M^*}, \tag{3.9}$$

for all  $x \in [a_1, b_1]$

**Proof** Let the set  $Y$  be defined as (3.1). Also, let  $d : Y \times Y \rightarrow [0, \infty]$  be a function defined as

$$d(g, h) := \inf \left\{ C \in [0, \infty] : |g(x) - h(x)| e^{-(L^*+1)(x-a_1)} \leq C\psi(x), x \in I \right\}. \tag{3.10}$$

From Lemma 3.1,  $(Y, d)$  is a GCM space. Now, define the operator  $T : Y \rightarrow Y$  by

$$(Ty)(x) = \sum_{j=1}^m \frac{b_j}{\Gamma(\alpha_1 - j + 1)} \left(\ln \frac{x}{a_1}\right)^{\alpha_1 - j} + \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left(\ln \frac{x}{\eta}\right)^{\alpha_1 - 1} u(\eta, y(\eta)) \frac{d\eta}{\eta}, \tag{3.11}$$

for all  $y \in Y$  and  $x \in [a_1, b_1]$ . It is obvious that the operator  $T$  is well defined. Now, we will show that  $T$  is strictly contractive on  $Y$ . From the definition of  $(Y, d)$ , for any  $g, h \in Y$ , we find a  $C_{g,h} \in [0, \infty)$  such that

$$|g(x) - h(x)| e^{-(L^*+1)(x-a_1)} \leq C_{g,h} \psi(x), \tag{3.12}$$

for any  $x \in [a_1, b_1]$ . Then, from the definition  $T$  in (3.11), and (3.5), (3.7), (3.12), Hölder's and Cauchy-Schwarz inequalities, we have

$$\begin{aligned} |(Tg)(x) - (Th)(x)| &= \frac{1}{\Gamma(\alpha_1)} \left| \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1 - 1} \frac{1}{\eta} (u(\eta, g(\eta)) - u(\eta, h(\eta))) d\eta \right| \\ &\leq \frac{L^*}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1 - 1} \eta^{-1} \eta^{p_1} |g(\eta) - h(\eta)| d\eta \\ &\leq \frac{L^* C_{g,h}}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1 - 1} \eta^{p_1 - 1} \psi(\eta) e^{(L^*+1)(\eta-a_1)} d\eta \\ &= \frac{L^* C_{g,h}}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1 - 1} \left(\eta^{\frac{p_1-1}{\alpha_1-1}}\right)^{\alpha_1-1} \psi(\eta) e^{(L^*+1)(\eta-a_1)} d\eta \\ &= \frac{L^* C_{g,h}}{\Gamma(\alpha_1)} \int_{a_1}^x \left((\ln x - \ln \eta) \eta^{\frac{p_1-1}{\alpha_1-1}}\right)^{\alpha_1-1} \psi(\eta) e^{(L^*+1)(\eta-a_1)} d\eta \\ &\leq \frac{L^* C_{g,h}}{\Gamma(\alpha_1)} \left( \int_{a_1}^x \left((\ln x - \ln \eta) \eta^{\frac{p_1-1}{\alpha_1-1}}\right)^{\frac{\alpha_1-1}{1-p_1}} d\eta \right)^{1-p_1} \left( \int_{a_1}^x \left(\psi(\eta) e^{(L^*+1)(\eta-a_1)}\right)^{1/p_1} d\eta \right)^{p_1} \end{aligned}$$

$$\begin{aligned} &\leq \frac{L^*C_{g,h}}{\Gamma(\alpha_1)} \left( \int_{a_1}^x (\ln x - \ln \eta)^{\frac{\alpha_1-1}{1-p_1}} \eta^{-1} d\eta \right)^{1-p_1} \left( \int_{a_1}^x (\psi(\eta))^{2/p_1} d\eta \right)^{p_1/2} \\ &\qquad \qquad \qquad \times \left( \int_{a_1}^x e^{\frac{2(L^*+1)(\eta-a_1)}{p_1}} d\eta \right)^{p_1/2} \\ &\leq \frac{K^*L^*C_{g,h}\psi(x)}{\Gamma(\alpha_1)} \left( \frac{1-p_1}{\alpha_1-p_1} (\ln x - \ln a_1)^{\frac{\alpha_1-p_1}{1-p_1}} \right)^{1-p_1} \left( \frac{p_1}{2(L^*+1)} e^{\frac{2(L^*+1)(x-a_1)}{p_1}} \right)^{p_1/2} \\ &\leq \frac{K^*L^*C_{g,h}\psi(x)}{\Gamma(\alpha_1)} \left( \frac{1-p_1}{\alpha_1-p_1} \right)^{1-p_1} (\ln b_1 - \ln a_1)^{\alpha_1-p_1} \left( \frac{p_1}{2(L^*+1)} \right)^{p_1/2} e^{(L^*+1)(x-a_1)} \end{aligned}$$

for all  $x \in [a_1, b_1]$ . Thus, for all  $x \in [a_1, b_1]$ , we get

$$|(Tg)(x) - (Th)(x)| e^{-(L^*+1)(x-a_1)} \leq K^*L^*M^*C_{g,h}\psi(x).$$

So, we can write

$$d(Tg, Th) \leq K^*L^*M^*d(g, h)$$

for any  $g, h \in Y$ . By the condition  $[A_3]$ ,  $T$  is a strictly contractive on  $Y$ . Let  $g_1 \in Y$ . Since  $g_1$  and  $Tg_1$  are continuous,  $u$  and  $g_1$  are bounded on  $[a_1, b_1]$  and  $\psi(x) > 0$ , then we can find a constant  $C^* > 0$  such that for

$$\begin{aligned} |(Tg_1)(x) - g_1(x)| e^{-(L^*+1)(x-a_1)} &= \left| \sum_{j=1}^m \frac{c_j}{\Gamma(\alpha_1 - j + 1)} \left( \ln \frac{x}{a_1} \right)^{\alpha_1 - j} \right. \\ &\quad \left. + \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left( \ln \frac{x}{\eta} \right)^{\alpha_1 - 1} u(\eta, g_1(\eta)) \frac{d\eta}{\eta} - g_1(x) \right| e^{-(L^*+1)(x-a_1)} \\ &\leq C^*\psi(x). \end{aligned}$$

Therefore, we show that  $d(Tg_1, g_1) < \infty$ . Now, using Theorem 2.2, there exists  $y_1 \in C([a_1, b_1], \mathbb{R})$  such that  $T^n g_1 \rightarrow y_1$  in  $(Y, d)$  as  $n \rightarrow \infty$  and  $Ty_1 = y_1$ . That is, for every  $x \in [a_1, b_1]$ ,  $y_1$  satisfies (3.8). Now, we will show  $\{g \in Y : d(g_1, g) < \infty\} = Y$ . Since  $g$  and  $g_1$  are bounded on  $[a_1, b_1]$  and  $\min_{x \in [a_1, b_1]} \psi(x) > 0$ , for any  $g \in Y$  and any  $x \in [a_1, b_1]$  we can find a constant  $C_g > 0$  such that

$$|g_1(x) - g(x)| e^{-(L^*+1)(x-a_1)} \leq C_g\psi(x).$$

So, we obtain  $d(g_1, g) < \infty$  for all  $g \in Y$ . That is,  $\{g \in Y : d(g_1, g) < \infty\} = Y$ . Hence, the function  $y_1$  is unique continuous with (3.8). Also, since (3.6) we obtain that

$$d(y, Ty) \leq 1. \tag{3.13}$$

So, we obtain

$$d(y, y_1) \leq \frac{1}{1 - K^*L^*M^*} d(Ty, y) \leq \frac{1}{1 - K^*L^*M^*},$$

for all  $x \in [a_1, b_1]$ . That is, the inequality (3.9) holds. □

**Corollary 3.3** *If  $\psi(x)$  is an arbitrary constant function in Theorem 3.2, then we obtain HU stability of the equation (1.3).*

Now, instead of  $[A_1]$  consider the following condition  $[A_1^*]$ .

$[A_1^*]$ : Let  $u : [a_1, b_1] \times \mathbb{R} \rightarrow \mathbb{R}$  be Carathéodory function. Also, there exists  $M_* > 0$  such that

$$|u(\eta, y)| \leq M_* \eta^{p_1} (\ln b_1 - \ln \eta)^{q_1}, p_1 - \alpha_1 \leq q_1 \leq 0.$$

Also, for any  $\eta \in [a_1, b_1]$  and  $y, z \in \mathbb{R}$ , let

$$|u(\eta, y) - u(\eta, z)| \leq L_1^* \eta^{p_1} (\ln b_1 - \ln \eta) |y - z|, p_1 - \alpha_1 \leq q_1 \leq 0.$$

**Theorem 3.4** *Let*

$$M_1^* = \frac{1}{\Gamma(\alpha_1)} \left( \frac{1 - p_1}{\alpha_1 + q_1 - p_1} \right)^{1-p_1} \left( \frac{p_1}{2(L_1^* + 1)} \right)^{p_1/2} (\ln b_1 - \ln a_1)^{\alpha_1 + q_1 - p_1}.$$

*Suppose that  $[A_1^*]$ ,  $[A_2]$  and  $0 < K^* L_1^* M_1^* < 1$  are satisfied. Then for all  $x \in [a_1, b_1]$ , there exists an unique continuous function  $y_1 : [a_1, b_1] \rightarrow \mathbb{R}$  such that  $y_1$  satisfies (1.3) and*

$$|y(x) - y_1(x)| e^{-(L^*+1)(x-a_1)} \leq \frac{\psi(x)}{1 - K^* L_1^* M_1^*}.$$

**Proof** First, we consider the second integral term in (3.8) and we show that it is bounded. That is, using  $[A_1^*]$ , we obtain

$$\begin{aligned} & \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x \left( \ln \frac{x}{\eta} \right)^{\alpha_1-1} u(\eta, y(\eta)) \frac{d\eta}{\eta} \\ & \leq \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1-1} \eta^{-1} \eta^{p_1} (\ln b_1 - \ln \eta)^{q_1} M_* d\eta \\ & \leq \frac{1}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1-1} \eta^{-1} \eta^{p_1} (\ln x - \ln \eta)^{q_1} M_* d\eta \\ & = \frac{M_*}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1+q_1-1} \left( \eta^{\frac{p_1-1}{\alpha_1+q_1-1}} \right)^{\alpha_1+q_1-1} d\eta \\ & = \frac{M_*}{\Gamma(\alpha_1)} \int_{a_1}^x \left[ (\ln x - \ln \eta) \eta^{\frac{p_1-1}{\alpha_1+q_1-1}} \right]^{\alpha_1+q_1-1} d\eta \\ & \leq \frac{M_*}{\Gamma(\alpha_1)} \left( \int_{a_1}^x \left[ (\ln x - \ln \eta) \eta^{\frac{p_1-1}{\alpha_1+q_1-1}} \right]^{\frac{\alpha_1+q_1-1}{1-p_1}} d\eta \right)^{p_1-1} \left( \int_{a_1}^x 1 d\eta \right)^{p_1} \\ & = \frac{M_* (b_1 - a_1)^{p_1}}{\Gamma(\alpha_1)} \left[ \frac{1 - p_1}{\alpha_1 + q_1 - p_1} (\ln x - \ln a_1)^{\frac{\alpha_1+q_1-p_1}{1-p_1}} \right]^{p_1-1}. \end{aligned}$$



Proof is made by using the similar arguments to the proof of Theorem 3.2. Then, we have

$$\begin{aligned}
 |(Tg)(x) - (Th)(x)| &= \frac{1}{\Gamma(\alpha_1)} \left| \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1-1} \frac{1}{\eta} (u(\eta, g(\eta)) - u(\eta, h(\eta))) d\eta \right| \\
 &\leq \frac{L_1^*}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1-1} \eta^{-1} \eta^{p_1} (\ln b_1 - \ln \eta)^{q_1} |g(\eta) - h(\eta)| d\eta \\
 &\leq \frac{L_1^*}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1-1} \eta^{-1} \eta^{p_1} (\ln x - \ln \eta)^{q_1} |g(\eta) - h(\eta)| \\
 &\quad \times e^{-(L_1^*+1)(\eta-a_1)} e^{(L_1^*+1)(\eta-a_1)} d\eta \\
 &\leq \frac{L_1^* C_{g,h}}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1+q_1-1} \eta^{p_1-1} \psi(\eta) e^{(L_1^*+1)(\eta-a_1)} d\eta \\
 &= \frac{L_1^* C_{g,h}}{\Gamma(\alpha_1)} \int_{a_1}^x (\ln x - \ln \eta)^{\alpha_1+q_1-1} \left( \eta^{\frac{p_1-1}{\alpha_1+q_1-1}} \right)^{\alpha_1+q_1-1} \psi(\eta) e^{(L_1^*+1)(\eta-a_1)} d\eta \\
 &= \frac{L_1^* C_{g,h}}{\Gamma(\alpha_1)} \int_{a_1}^x \left( (\ln x - \ln \eta) \eta^{\frac{p_1-1}{\alpha_1+q_1-1}} \right)^{\alpha_1+q_1-1} \psi(\eta) e^{(L_1^*+1)(\eta-a_1)} d\eta \\
 &\leq \frac{L_1^* C_{g,h}}{\Gamma(\alpha_1)} \left( \int_{a_1}^x \left( (\ln x - \ln \eta) \eta^{\frac{p_1-1}{\alpha_1+q_1-1}} \right)^{\frac{\alpha_1+q_1-1}{1-p_1}} d\eta \right)^{1-p_1} \\
 &\quad \times \left( \int_{a_1}^x \left( \psi(\eta) e^{(L_1^*+1)(\eta-a_1)} \right)^{1/p_1} d\eta \right)^{p_1} \\
 &\leq \frac{L_1^* C_{g,h}}{\Gamma(\alpha_1)} \left( \int_{a_1}^x (\ln x - \ln \eta)^{\frac{p_1-1}{\alpha_1+q_1-1}} \eta^{-1} d\eta \right)^{1-p_1} \left( \int_{a_1}^x (\psi(\eta))^{2/p_1} d\eta \right)^{p_1/2} \\
 &\quad \times \left( \int_{a_1}^x e^{\frac{2(L_1^*+1)(\eta-a_1)}{p_1}} d\eta \right)^{p_1/2} \\
 &\leq \frac{K^* L_1^* C_{g,h} \psi(x)}{\Gamma(\alpha_1)} \left( \frac{1-p_1}{\alpha_1+q_1-p_1} (\ln x - \ln a_1)^{\frac{\alpha_1+q_1-p_1}{1-p_1}} \right)^{1-p_1} \\
 &\quad \times \left( \frac{p_1}{2(L_1^*+1)} e^{\frac{2(L_1^*+1)(x-a_1)}{p_1}} \right)^{p_1/2} \\
 &\leq \frac{K^* L_1^* C_{g,h} \psi(x)}{\Gamma(\alpha_1)} \left( \frac{1-p_1}{\alpha_1+q_1-p_1} \right)^{1-p_1} \left( \frac{p_1}{2(L_1^*+1)} \right)^{p_1/2} \\
 &\quad \times e^{(L_1^*+1)(x-a_1)} (\ln b_1 - \ln a_1)^{\alpha_1+q_1-p_1} \\
 &= K^* L_1^* M_1^* C_{g,h} \psi(x),
 \end{aligned}$$

for all  $x \in [a_1, b_1]$ . Thus, we get

$$|(Tg)(x) - (Th)(x)| e^{-(L^*+1)(x-a_1)} \leq K^* L_1^* M_1^* C_{g,h} \psi(x),$$

for all  $x \in [a_1, b_1]$ . Therefore, we can complete the rest of the proof by applying standard process in proof of Theorem 3.2 which is quite similar.  $\square$

**4. Examples**

In this section, we will give the application of our main results with two examples. We will see that examples are based on the results Section 3, although these examples cannot be demonstrated with results of Wang and Lin [37].

**Example 4.1.** Consider the equation (1.3) with  $u(\eta, y(\eta)) = \eta^2 y(\eta)$ . Let  $a_1 \in \mathbb{R}^+$ ,  $b_1 = a_1 e$ ,  $p_1 = \frac{1}{2}$ ,  $\alpha_1 = 1$ ,  $n = 1$ . Then,  $M^* = \frac{1}{2}$  and  $L^* = 3$ . Also, we assume that for all  $x \in [a_1, a_1 e]$ ,  $y : [a_1, a_1 e] \rightarrow \mathbb{R}$  is a continuous function such that

$$\left| y(x) - c_1 - \int_{a_1}^x \eta^2 y(\eta) \frac{d\eta}{\eta} \right| e^{-4(x-a_1)} \leq e^{4x}.$$

Here  $\psi(x) = e^{4x}$ . So, for all  $x \in [a_1, a_1 e]$  we have

$$\left( \int_{a_1}^x e^{16\eta} d\eta \right)^{1/4} = \left( \frac{1}{16} (e^{16x} - e^{16a_1}) \right)^{1/4} \leq \frac{1}{2} e^{4x}.$$

That is, we find  $K^* = \frac{1}{2}$ . Then, we obtain  $0 < K^* L^* M^* = \frac{3}{4} < 1$  and  $\Gamma(1) = 1$ .

Because of Theorem 3.2, we say that there exists an unique continuous function  $y_1 : [a_1, a_1 e] \rightarrow \mathbb{R}$  such that

$$y_1(x) = c_1 + \int_{a_1}^x \eta^2 y_1(\eta) \frac{d\eta}{\eta}$$

and

$$|y(x) - y_1(x)| e^{-4(x-a_1)} \leq 4e^x.$$

for all  $x \in [a_1, a_1 e]$ . Thus by Theorem 3.2, Eq.(1.3) is HUR stable. However, in this example if we use Theorem 2.1 in [37], we get  $K = \frac{1}{2\sqrt{2}}$ ,  $M = 1$ ,  $L = 3$  and  $KML = \frac{3}{2\sqrt{2}} > 1$ . Therefore, since the  $KLM$  is not smaller than one, this example cannot apply to Theorem 2.1 in [37].

**Example 4.2.** Consider the equation (1.3) with  $u(\eta, y) = 3\eta^{\frac{1}{2}} \sin y(\eta) (\ln a_1 e - \ln \eta)^{-\frac{1}{3}}$ . Let  $a_1 \in \mathbb{R}^+$ ,  $b_1 = a_1 e$ ,  $p_1 = \frac{1}{2}$ ,  $\alpha_1 = 1$ ,  $q_1 = -\frac{1}{3}$ ,  $n = 1$ . Then, we find

$$|u(\eta, y)| \leq 3\eta^{\frac{1}{2}} (\ln a_1 e - \ln \eta)^{-\frac{1}{3}}.$$

Also, for all  $\eta \in [a_1, a_1 e]$  and  $y, z \in \mathbb{R}$ , we obtain

$$|u(\eta, y) - u(\eta, z)| \leq 3\eta^{\frac{1}{2}} (\ln a_1 e - \ln \eta) |y - z|.$$

So,  $[A_1^*]$  satisfies. Since  $L_1^* = 3$ , we get  $M_1^* = \frac{\sqrt{3}}{2}$ . Also, we assume that for all  $x \in [a_1, a_1 e]$ ,  $y \in ([a_1, a_1 e], \mathbb{R})$  satisfies

$$\left| y(x) - c_1 - \int_{a_1}^x 3\eta^{\frac{1}{2}} \sin y(\eta) (\ln a_1 e - \ln \eta)^{-\frac{1}{3}} \frac{d\eta}{\eta} \right| e^{-4(x-a_1)} \leq \frac{1}{2\sqrt{2}} e^x.$$

Here  $\psi(x) = \frac{1}{2\sqrt{2}}e^x$ . So, for all  $x \in [a_1, a_1e]$  we have

$$\left(\int_{a_1}^x \frac{1}{64}e^{4\eta}d\eta\right)^{1/4} = \left(\frac{1}{256}(e^{4x} - e^{4a_1})\right)^{1/4} \leq \frac{1}{4}e^{4x}.$$

That is, we find  $K^* = \frac{1}{4}$ . Thus,  $K^*L_1^*M_1^* = \frac{3\sqrt{3}}{8}$ . Since Theorem 3.4, we say that there exists a unique continuous function  $y_1 \in C([a_1, a_1e], \mathbb{R})$  which satisfies

$$y_1(x) = c_1 + \int_{a_1}^x 3\eta^{\frac{1}{2}} \sin y_1(\eta) (\ln a_1e - \ln \eta)^{-\frac{1}{3}} \frac{d\eta}{\eta}$$

and

$$|y(x) - y_1(x)|e^{-4(x-a_1)} \leq \frac{1}{2\sqrt{2}} \frac{8}{8 - 3\sqrt{3}} e^x.$$

for all  $x \in [a_1, a_1e]$ . Thus by Theorem 3.4, Eq.(1.3) is HUR stable. However, in this example if we use Theorem 2.2 in [37], we get  $K = \frac{1}{4}$ ,  $M = \sqrt{3}$ ,  $L = 3$  and  $KML = \frac{3\sqrt{3}}{4} > 1$ . Therefore, since the  $KLM$  is not smaller than one, this example cannot apply to Theorem 2.2 in [37].

## 5. Conclusion

Studies involving fractional calculus has played an important role in several areas of science and engineering. In recent years, HU stability of differential equations which have fractional derivative and integral in the various fields have been studied by many researchers. HU stability is one of the main topics in the theory of fractional equations. In this paper, we examine the HUR and HU stability of Hadamard type fractional integral equations on compact intervals. The stability conditions are developed by using a new GM definition and the fixed point technique by motivating Wang and Lin [37]. It is then shown that the results of this paper are better for some previous results. Finally, we investigate in detail two examples to show the reported results.

## Acknowledgment

The authors would like to thank to the referees for their useful comments and remarks.

## References

- [1] Abbas S, Albarakati W, Benchohra M, Trujillo JJ. Ulam stabilities for partial Hadamard fractional integral equations. *Arabian Journal of Mathematics* 2016; 5: 1-7.
- [2] Akkouchi M. Stability of certain functional equations via a fixed point of Ciric. *Filomat* 2011; 25: 121-127.
- [3] Andras Sz, Kolumbán JJ. On the Ulam-Hyers stability of first order differential systems with non-local initial conditions. *Nonlinear Analysis: Theory, Methods & Applications* 2013; 82: 1-11.
- [4] Andras Sz, Mészáros AR. Ulam-Hyers stability of dynamic equations on time scales via Picard operators. *Applied Mathematics and Computation* 2013; 219: 4853-4864.
- [5] Başcı Y, Öğrekçi S, Mısır A. On Hyers-Ulam stability for fractional differential equations including the new Caputo-Fabrizio fractional derivative. *Mediterranean Journal of Mathematics* 2019; 16: 130-144.

- [6] Başcı Y, Mısır A, Öğrekçi S. On the stability problem of differential equations in the sense of Ulam. *Results in Mathematics* 2020; 75 (6). doi: 10.1007/s00025-019-1132-6
- [7] Biçer E, Tunç C. New theorems for Hyers-Ulam stability of Lienard equation with variables time lags. *International Journal of Mathematics and Computer Science* 2018; 3 (2): 231-242.
- [8] Brillouet-Belluot N, Brzdek J, Cieplinski K. On some recent developments in Ulam's type stability. *Abstract and Applied Analysis* 2012; 2012. doi: 10.1155/2012/716936
- [9] Cimpean DS, Popa D. Hyers-Ulam stability of Euler's equation. *Applied Mathematics Letters* 2011; 24: 1539-1543.
- [10] Diaz JB, Margolis B. A fixed point theorem of the alternative, for contractions on a generalized complete metric space. *Bulletin of the American Mathematical Society* 1968; 74: 305-309.
- [11] Gordji ME, Savadkouhi MB. Stability of a mixed type additive, quadratic and cubic functional equation in random normed spaces. *Filomat* 2011; 25: 43-54.
- [12] Hegyi B, Jung SM. On the stability of Laplace's equation. *Applied Mathematics Letters* 2013; 26: 549-552.
- [13] Hilfer R. *Applications of fractional calculus in physics*. Singapore: World Scientific Publishing, 2000.
- [14] Ibrahim RW. Ulam-Hyers stability for Cauchy fractional differential equation in the unit disk. *Abstract and Applied Analysis* 2012; 2012. doi: 10.1155/2012/613270
- [15] Jeetendra R, Vernold Vivin J. Stability analysis of uncertain stochastic systems with interval time-varying delays and nonlinear uncertainties via augmented Lyapunov functional. *Filomat* 2012; 26: 1179-1188.
- [16] Jung SM. Hyers-Ulam stability of linear differential equations of first order. *Applied Mathematics Letters* 2004; 17: 1135-1140.
- [17] Jung SM. A fixed point approach to the stability of differential equations  $y' = f(x, y)$ . *Bulletin of the Malaysian Mathematical Sciences Society* 2010; 33 (1): 47-56.
- [18] Jung SM, Kim TS, Lee KS. A fixed point approach to the stability of quadratic functional equation. *Bulletin of the Korean Mathematical Society* 2006; 43: 531-541.
- [19] Khan H, Li Y, Chen W, Baleanu D, Khan A. Existence theorems and Hyers-Ulam stability for a coupled system of fractional differential equations with p-Laplacian operator. *Boundary Value Problems* 2017; 2017: 157. doi: 10.1186/s13661-017-0878-6
- [20] Khan H, Tunç C, Chen W, Khan A. Existence theorems and Hyers-Ulam stability for a class of hybrid fractional differential equations with p-Laplacian operator. *Journal of Applied Analysis and Computation* 2018; 8 (4): 1211-1226.
- [21] Kilbas AA, Srivastava HM, Trujillo JJ. *Theory and applications of fractional differential equations*. In: Van Mill J (editor). *North-Holland Mathematics Studies Vol. 204*. Amsterdam, Netherlands: Elsevier Science Publishers BV, 2006.
- [22] Lungu N, Popa D. Hyers-Ulam stability of a first order partial differential equation. *Journal of Mathematical Analysis and Applications* 2012; 385: 86-91.
- [23] Mlesnite O, Petrusel A. Existence and Ulam-Hyers stability results for multivalued coincidence problems. *Filomat* 2013; 26: 965-976.
- [24] Muniyappan P, Rajan S. Hyers-Ulam-Rassias stability of fractional differential equation. *International Journal of Pure and Applied Mathematics* 2015; 102 (4): 631-642.
- [25] Ortigueira MD, Machado JAT. Fractional calculus applications in signals and systems. *Signal Processing* 2006; 86 (10): 2503-2504. doi: 10.1016/j.sigpro.2006.02.001
- [26] Podlubny I. *Fractional differential equations*. Mathematics in Science and Engineering Vol. 198. Cambridge, MA, USA: Academic Press, 1998.
- [27] Rus IA. Ulam stability of ordinary differential equations. *Studia Universitatis Babes-Bolyai Mathematica* 2009; 54: 125-133.

- [28] Rus IA. Ulam stabilities of ordinary differential equations in a Banach space. *Carpathian Journal of Mathematics* 2010; 26: 103-107.
- [29] Samko SG, Kilbas AA, Marichev OI. *Fractional integrals and derivatives*, translated from the 1987 Russian original. Yverdon, Switzerland: Gordon and Breach, 1993.
- [30] Talib I, Tunç C, Noor ZA. New operational matrices of orthogonal Legendre polynomials and their operational. *Journal of Taibah University for Science* 2019; 13 (1): 377-389. doi: 10.1080/16583655.2019.1580662
- [31] Talib I, Belgacem FBM, Khalil H, Tunç C. Nonlinear fractional partial coupled systems approximate solutions through operational matrices approach. *Nonlinear Studies (NS)* 2019; 26 (4): 955-971.
- [32] Tarasov VE. *Fractional dynamics: Application of fractional calculus to dynamics of particles, fields and media*. Heidelberg, Germany: Springer, 2010.
- [33] Tarasov VE. On History of mathematical economics: Application of fractional calculus. *Mathematics* 2019; 7(6): 509. doi: 10.3390/math7060509
- [34] Tunç C. On the stability and boundedness of solutions of nonlinear third order differential equations with delay. *Filomat* 2010; 24 (3): 1-10.
- [35] Tunç C. Stability and boundedness in multi delay vector Lienard equation. *Filomat* 2013; 27: 435-445.
- [36] Wang J, Feckan M, Zhou Y. Ulam's type stability of impulsive ordinary differential equations. *Journal of Mathematical Analysis and Applications* 2012; 395: 258-264.
- [37] Wang J, Lin Z. Ulam's type stability of Hadamard type fractional integral equations. *Filomat* 2014; 28(7): 1323-1331.
- [38] Wang J, Zhou Y, Medved M. Existence and stability of fractional differential equations with Hadamard derivative. *Topological Methods in Nonlinear Analysis* 2013; 41: 113-133.
- [39] Wei W, Li X, Li, X. New stability results for fractional integral equation. *Computers & Mathematics with Applications* 2012; 64 (10): 3468-3476.
- [40] Zheng A, Feng Y, Wang W. The Hyers-Ulam stability of the conformable fractional differential equation. *Mathematica Aeterna* 2015; 5 (3): 485-492.