

1-1-2019

On the n-strong Drazin invertibility in rings

HONGLIN ZOU

DIJANA MOSIC

KEZHENG ZUO

YINLAN CHEN

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ZOU, HONGLIN; MOSIC, DIJANA; ZUO, KEZHENG; and CHEN, YINLAN (2019) "On the n-strong Drazin invertibility in rings," *Turkish Journal of Mathematics*: Vol. 43: No. 6, Article 1. <https://doi.org/10.3906/mat-1905-50>

Available at: <https://dctubitak.researchcommons.org/math/vol43/iss6/1>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

On the n -strong Drazin invertibility in rings

Honglin ZOU^{1*}, Dijana MOSIĆ², Kezheng ZUO¹, Yinlan CHEN¹

¹School of Mathematics and Statistics, Hubei Normal University, Huangshi, China

²Faculty of Sciences and Mathematics, University of Niš, Niš, Serbia

Received: 14.05.2019

Accepted/Published Online: 26.08.2019

Final Version: 22.11.2019

Abstract: Let R be a ring and n be a positive integer. In this paper, further results on the n -strong Drazin inverse are obtained in a ring. We prove that $a \in R$ is n -strongly Drazin invertible if and only if $a - a^{n+1}$ is nilpotent. In terms of this characterization, the extensions of Cline's formula and Jacobson's lemma for this inverse are proved. Moreover, the n -strong Drazin invertibility for the sums of two elements is considered. We prove that $a, b \in R$ are n -strongly Drazin invertible if and only if $a + b$ is n -strongly Drazin invertible, under the condition $ab = 0$. As applications for the additive results, we obtain some equivalent conditions of the n -strong Drazin invertibility of matrices over a ring.

Key words: Strong Drazin inverse, Hirano inverse, n -strong Drazin inverse, Drazin inverse ring

1. Introduction

Let R^D denote the set of all Drazin invertible elements in a ring R . It is well known that if $a, b \in R$, then

$$ab \in R^D \iff ba \in R^D.$$

In this case, $(ba)^D = b((ab)^D)^2a$ [4]. This formula is called Cline's formula for the Drazin inverse. Many researchers considered Cline's formula for various types of generalized inverses, such as (b, c) -inverse [10], Mary inverse [27], Hirano inverse [2], pseudo-Drazin inverse [20], generalized Drazin inverse [13, 14, 16, 23, 24]. In [23], Zeng et al. extended Cline's formula for the (pseudo, generalized) Drazin inverse to more general case. Namely, if $a, b, c, d \in R$ satisfy $acd = dbd$ and $dba = aca$, then

$$ac \in R^D \iff bd \in R^D.$$

In this case, $(bd)^D = b((ac)^D)^2d$ and $(ac)^D = d((bd)^D)^3bac$. Corresponding to Cline's formula, many researchers paid attention to Jacobson's lemma, that is

$$1 - ab \in R^{-1} \iff 1 - ba \in R^{-1}.$$

In this case, $(1 - ba)^{-1} = 1 + b(1 - ab)^{-1}a$. They investigated Jacobson's lemma for different generalized inverses in different settings [1, 2, 5, 6, 17, 18, 25].

The topic for generalized inverses of the sums was studied by many authors. In 1958, Drazin [9] proved that $a + b \in R^D$ with $(a + b)^D = a^D + b^D$ under the condition $a, b \in R^D$ and $ab = ba = 0$. For $a, b \in \mathbb{C}^{n \times n}$,

*Correspondence: honglinzou@163.com

2010 AMS Mathematics Subject Classification: 15A09; 16S50; 16B99

Hartwig et al. [12] obtained a formula for $(a + b)^D$ under the one-sided condition $ab = 0$, which was extended to the additive category by Chen et al. [3]. In addition, the problem of generalized inverses of $a + b$ was also studied under the condition $ab = ba$. For example, Wei and Deng [21] gave the relations of Drazin inverses of $a + b$ and $1 + a^D b$, where $a, b \in \mathbb{C}^{n \times n}$. Later, Zhuang et al. [26] extended the result of [21] to the ring case. The generalized Drazin invertibility and strong Drazin invertibility of the sums under the commutative condition were also investigated [7, 8, 19].

All results mentioned above were the motivation for further consideration of the n -strong Drazin inverse in a ring. This article consists of five sections. In Section 2, we recall the definitions of some generalized inverses and give related notations. In Section 3, characterizations of n -strongly Drazin invertible elements are given in terms of the nilpotency. Then, we investigate Cline’s formula and Jacobson’s lemma for the n -strong Drazin inverse in a ring. In Section 4, we obtain some equivalent conditions for the n -strong Drazin invertibility of the sum $a + b$ under the hypothesis $ab = 0$ (or $a^2 b = aba, ab^2 = bab$). In Section 5, as applications of the previous additive results, we mainly consider the n -strong Drazin invertibility of matrices over a ring. We remark that some results presented in this paper are different from those of Drazin inverses.

2. Preliminaries

Throughout this paper, R denotes a ring with unity 1. R^{nil} and \mathbb{N} stand for the sets of all nilpotent elements in R and positive integers, respectively. Denote by $\binom{n}{k}$ the binomial coefficient $\frac{n!}{k!(n-k)!}$ ($0 \leq k \leq n$).

For the readers’ convenience, we first recall the definitions of some generalized inverses. The Drazin inverse [9] of $a \in R$ is the element $x \in R$ which satisfies

$$xax = x, \quad ax = xa, \quad \text{and} \quad a - a^2 x \in R^{nil}.$$

The element x above is unique if it exists and is denoted by a^D . The power of nilpotency of $a - a^2 a^D$ is called the index of a , and will be denoted by $\text{ind}(a)$. Drazin [9] proved that $a \in R$ is Drazin invertible if and only if a is both right π -regular (i.e. $a^m \in a^{m+1}R$, for some $m \in \mathbb{N}$) and left π -regular ($a^m \in Ra^{m+1}$, for some $m \in \mathbb{N}$), namely a is strongly π -regular.

In 2017, Wang [19] gave the notion of the strong Drazin inverse in a ring. An element $a \in R$ is said to be strongly Drazin invertible [19] if there exists $x \in R$ such that

$$xax = x, \quad ax = xa, \quad \text{and} \quad a - ax \in R^{nil}.$$

In this case, x is unique if it exists and is called the strong Drazin inverse of a . We will denote the strong Drazin inverse of a by a^{sD} . The strongly Drazin invertible elements are exactly the ones which are strongly nil-clean (see [19, Lemma 2.2]). Let $a \in R$, then a^D exists if and only if there exists $x \in R$ such that

$$x \in aR \cap Ra, \quad ax = xa, \quad \text{and} \quad a - ax \in R^{nil}.$$

Suppose that a^D exists. Then, let $x = aa^D$. Obviously, x satisfies $x \in aR \cap Ra, ax = xa,$ and $a - ax \in R^{nil}$.

On the contrary, we have $(a - ax)^m = 0$ for some $m \in \mathbb{N}$. Hence, $a^m(1 - x)^m = a^m(1 + \sum_{i=1}^m (-1)^i \binom{m}{i} x^i) = 0,$

which implies that $a^m = a^m x u = u x a^m,$ for some $u \in R$. Observe that $x = as = ta,$ where $s, t \in R$. Hence, we deduce that $a^m = a^{m+1} s u = u t a^{m+1} \in a^{m+1} R \cap R a^{m+1}$. Hence, a^D exists.

The definition of the Hirano inverse [2] was introduced by Chen and Sheibani in 2017. The Hirano inverse of $a \in R$ is the unique element x (written $x = a^H$) satisfying

$$xax = x, \quad ax = xa, \quad \text{and} \quad a^2 - ax \in R^{nil}.$$

It is interesting that the Hirano inverse is related to tripotent elements (see [2, Theorem 3.3]). In addition, they obtained the relations of the above three kinds of generalized inverses, that is $R^{sD} \subsetneq R^H \subsetneq R^D$, where R^{sD} and R^H mean the sets of all strongly Drazin invertible and Hirano invertible elements in R , respectively.

Recently, motivated by the concepts of the strong Drazin inverse and Hirano inverse, Mosić [15] introduced the notion of the n -strong Drazin inverse in a ring. Let $n \in \mathbb{N}$. An element $x \in R$ is called the n -strong Drazin inverse of $a \in R$ if it satisfies

$$xax = x, \quad ax = xa, \quad \text{and} \quad a^n - ax \in R^{nil}.$$

The previous x is unique if such element exists, and we denote it by a^{nsD} . Clearly, the n -strong Drazin inverse covers the strong Drazin inverse and Hirano inverse, that is, $a^{1sD} = a^{sD}$ and $a^{2sD} = a^H$. The power of nilpotency of $a^n - aa^{nsD}$ is called the n -strong Drazin index of a , denoted by $n\text{-ind}(a)$. The symbol R^{nsD} denote the set of all n -strongly Drazin invertible elements in R . We note that $R^{nil} \subseteq R^{nsD}$. Indeed, $a \in R^{nil}$ if and only if $a \in R^{nsD}$ with $a^{nsD} = 0$. In addition, $R^{-1} \not\subseteq R^{nsD}$. For example, let $R = \mathbb{C}$. Then, $3 \in R^{-1}$, but $3 \notin R^{nsD}$.

Next, we introduce two known lemmas, which are related to the nilpotency.

Lemma 2.1 *Let $a, b \in R$ with $ab = ba$. Then,*

- (1) *If $a \in R^{nil}$ (or $b \in R^{nil}$), then $ab \in R^{nil}$.*
- (2) *If $a, b \in R^{nil}$, then $a + b \in R^{nil}$.*

Lemma 2.2 [22, Lemma 3.5] *Let $a \in R$. If $a^2 - a \in R^{nil}$, then there exists a monic polynomial $\theta(t) \in \mathbb{Z}[t]$ such that $\theta(a) = \theta(a)^2$ and $a - \theta(a)$ is nilpotent.*

3. Cline’s formula and Jacobson’s lemma

In this section, we give an existence criterion for the n -strong Drazin inverse in a ring. Then, by this characterization we prove Cline’s formula and Jacobson’s lemma for the n -strong Drazin inverse. The results presented extend the corresponding ones of the strong Drazin inverse [19] and Hirano inverse [2].

Firstly, we give the relationship between the n -strong Drazin inverse and Drazin inverse. The proof of the following proposition is similar to that of [15, Lemma 2.1].

Proposition 3.1 *Let $n \in \mathbb{N}$. If $a \in R^{nsD}$ with $n\text{-ind}(a) = m$, then $a \in R^D$ and $a^D = a^{nsD}$. Moreover, $\text{ind}(a) \leq nm$.*

Proof Assume that $a \in R^{nsD}$ with $n\text{-ind}(a) = m$. Let $x = a^{nsD}$. Then we have $xax = x$, $ax = xa$, and $(a^n - ax)^m = 0$, which yield

$$(a - a^2x)^{nm} = (a^n - a^{n+1}x)^m = (a^n - ax)^m(1 - ax)^m = 0.$$

Hence, $a - a^2x \in R^{nil}$. Hence, $a \in R^D$ and $a^D = x$. Moreover, $\text{ind}(a) \leq nm$.

Inspired by [2, Theorem 3.1], we obtain a characterization for the n -strong Drazin invertibility in a ring, which plays an important role in the sequel.

Theorem 3.2 *Let $n \in \mathbb{N}$. Then $a \in R^{nsD}$ if and only if $a - a^{n+1} \in R^{nil}$.*

Proof Suppose that $a \in R^{nsD}$ and $x = a^{nsD}$, i.e. $xax = x$, $ax = xa$, and $a^n - ax \in R^{nil}$. Then we deduce that

$$a^n - a^{2n} = (a^n - ax)(1 - ax - a^n) \in R^{nil},$$

which yields that

$$(a - a^{n+1})^n = (a - a^{n+1})a^{n-1}(1 - a^n)^{n-1} = (a^n - a^{2n})(1 - a^n)^{n-1} \in R^{nil}.$$

Hence, $a - a^{n+1} \in R^{nil}$.

On the contrary, since $a - a^{n+1} \in R^{nil}$, we conclude that $(a^n)^2 - a^n = a^{n-1}(a^{n+1} - a) \in R^{nil}$. By Lemma 2.2, there exists a monic polynomial $\theta(t) \in \mathbb{Z}[t]$ such that $\theta(a^n) = \theta(a^n)^2$ and $a^n - \theta(a^n) \in R^{nil}$. Take $e = \theta(a^n)$. Then we have $e = e^2$, $ea = ae$ and $a^n - e \in R^{nil}$. Hence, we obtain $1 + a^n - e \in R^{-1}$. Let $x = (1 + a^n - e)^{-1}a^{n-1}e$. Next, we show that $a^{nsD} = x$ by the definition of the n -strong Drazin inverse. Obviously, $ax = xa$. Note that $a^ne = (1 + a^n - e)e = e(1 + a^n - e)$. Then, we obtain

$$\begin{aligned} xax &= (1 + a^n - e)^{-1}a^ne(1 + a^n - e)^{-1}a^{n-1}e \\ &= (1 + a^n - e)^{-1}(1 + a^n - e)e(1 + a^n - e)^{-1}a^{n-1}e \\ &= (1 + a^n - e)^{-1}a^{n-1}e \\ &= x \end{aligned}$$

and

$$a^n - ax = a^n - (1 + a^n - e)^{-1}a^ne = a^n - e \in R^{nil}.$$

Therefore, $a \in R^{nsD}$ with $a^{nsD} = x$.

Remark 3.3 (1) *Let $A \in \mathbb{C}^{m \times m}$ (rank $A = r > 0$) have the Jordan form*

$$A = P \begin{bmatrix} D & 0 \\ 0 & N \end{bmatrix} P^{-1},$$

where D is invertible and N is nilpotent. Then, by Theorem 3.2 we have

$$\begin{aligned} A \in (\mathbb{C}^{m \times m})^{nsD} &\iff I - D^n \in (\mathbb{C}^{r \times r})^{nil} \\ &\iff \sigma(A) \subseteq \{0, 1, \varepsilon, \varepsilon^2, \dots, \varepsilon^{n-1}\}, \end{aligned}$$

where $\sigma(A)$ denotes the spectrum of A and $\varepsilon = \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n}$.

(2) *We have the following special case,*

$$a \in R^{nsD} \text{ and } n\text{-ind}(a) = 1 \iff a^n = a^{2n}.$$

The necessity is obvious. In fact, from $(a^n - aa^{nsD})^1 = 0$ it follows that $a^n = aa^{nsD}$ and consequently $a^{2n} = aa^{nsD}aa^{nsD} = aa^{nsD} = a^n$. On the contrary, suppose that $a^n = a^{2n}$. Let $x = a^{2n-1}$. Then, we have $xax = a^{2n}a^{2n-1} = a^n a^{2n-1} = a^{2n}a^{n-1} = a^{2n-1} = x$. In addition, it is clear that $ax = xa$ and $a^n - ax = 0$. Hence, we get $a \in R^{nsD}$ and $a^{nsD} = x$. Moreover, $n\text{-ind}(a) = 1$.

Applying Theorem 3.2, we get some properties of the n -strong Drazin inverse in a ring.

Corollary 3.4 Let $n, k \in \mathbb{N}$. If $a \in R^{nsD}$, then $a^k \in R^{nsD}$ and $(a^k)^{nsD} = (a^{nsD})^k$.

Proof Since $a \in R^{nsD}$, by Theorem 3.2 we have $a - a^{n+1} \in R^{nil}$, which yields

$$a^k - (a^k)^{n+1} = a^k - (a^{n+1})^k = (a - a^{n+1}) \sum_{i=0}^{k-1} a^{ni+k-1} \in R^{nil}.$$

Hence, we obtain $a^k \in R^{nsD}$. In view of Proposition 3.1 and [9, Theorem 2], one can see that $(a^k)^{nsD} = (a^k)^D = (a^D)^k = (a^{nsD})^k$.

Corollary 3.5 Let $n \in \mathbb{N}$. If $a \in R^{nsD}$, then $a^{nsD} \in R^{nsD}$ and $(a^{nsD})^{nsD} = a^2 a^{nsD}$.

Proof Let $x = a^{nsD}$. Then we get $xax = x$, $ax = xa$, and $a^n - ax \in R^{nil}$. Hence,

$$x - x^{n+1} = x^{n+1}(a^n - ax) \in R^{nil}.$$

Hence, by Theorem 3.2 we obtain $x \in R^{nsD}$. From [9, Theorem 3], it follows that

$$x^{nsD} = x^D = (a^D)^D = a^2 a^D = a^2 a^{nsD}.$$

Corollary 3.6 Let $n \in \mathbb{N}$. Then,

- (1) If $a \in R^{sD}$, then $a \in R^{nsD}$ and $a^{nsD} = a^{sD} = a^D$.
- (2) If $a \in R^H$, then $a \in R^{2nsD}$ and $a^{2nsD} = a^H = a^D$.
- (3) If $a \in R^{nsD}$, then $a \in R^{2nsD}$ and $a^{2nsD} = a^{nsD} = a^D$.

Proof (1) Since $a \in R^{sD}$, by Theorem 3.2 we have $a - a^2 \in R^{nil}$, which gives

$$a - a^{n+1} = a(1 - a^n) = a(1 - a) \sum_{i=0}^{n-1} a^i = (a - a^2) \sum_{i=0}^{n-1} a^i \in R^{nil}.$$

Hence, $a - a^{n+1} \in R^{nil}$, i.e. $a \in R^{nsD}$. In addition, $a^{nsD} = a^D = a^{sD}$.

(2) can be proved in the same way as the item (1).

(3) follows directly by the equality $a - a^{2n+1} = (a - a^{n+1})(1 + a^n)$.

In terms of Theorem 3.2, we are now in the position to prove the extension of Cline's formula for the n -strong Drazin inverse when $acd = dbd$ and $dba = aca$.

Theorem 3.7 Let $a, b, c, d \in R$ and $n \in \mathbb{N}$. If $acd = dbd$ and $dba = aca$, then

$$ac \in R^{nsD} \iff bd \in R^{nsD}.$$

In this case, $(bd)^{nsD} = b((ac)^{nsD})^2d$ and $(ac)^{nsD} = d((bd)^{nsD})^3bac$.

Proof It will suffice to prove the sufficiency, since the necessity can be proved similarly. From $dba = aca$, it follows that $(ac)^i = (db)^{i-1}ac$ for any $i \in \mathbb{N}$. Now, we show that

$$(ac - (ac)^{n+1})^{m+1} = d(bd - (bd)^{n+1})^{m-1}(b - (bd)^nb)(ac - (ac)^{n+1})$$

by induction on positive integer m .

For $m = 1$, we have

$$\begin{aligned} (ac - (ac)^{n+1})^2 &= (ac - (ac)^{n+1})(ac - (ac)^{n+1}) \\ &= ((ac)^2 - (ac)^{n+2})(1 - (ac)^n) \\ &= (dbac - (db)^{n+1}ac)(1 - (ac)^n) \\ &= (db - (db)^{n+1})(ac - (ac)^{n+1}) \\ &= d(b - (bd)^nb)(ac - (ac)^{n+1}). \end{aligned}$$

Assume that the conclusion holds for positive integer $m = l$. Now, we check it for $m = l + 1$ as follows:

$$\begin{aligned} (ac - (ac)^{n+1})^{l+2} &= (ac - (ac)^{n+1})^{l+1}(ac - (ac)^{n+1}) \\ &= d(bd - (bd)^{n+1})^{l-1}(b - (bd)^nb)(ac - (ac)^{n+1})^2 \\ &= d(bd - (bd)^{n+1})^{l-1}(b - b(db)^nd(b - (bd)^nb)(ac - (ac)^{n+1})) \\ &= d(bd - (bd)^{n+1})^l(b - (bd)^nb)(ac - (ac)^{n+1}). \end{aligned}$$

Note that $bd \in R^{nsD}$, i.e. $bd - (bd)^{n+1} \in R^{nil}$. Hence, $ac - (ac)^{n+1} \in R^{nil}$, i.e. $ac \in R^{nsD}$. By Proposition 3.1 and the formula of [23, Theorem 2.1], we obtain $(ac)^{nsD} = d((bd)^{nsD})^3bac$.

In Theorem 3.7, let $d = a$ and $c = b$, then it is reduced as the following.

Corollary 3.8 Let $a, b \in R$ and $n \in \mathbb{N}$. Then,

$$ab \in R^{nsD} \iff ba \in R^{nsD}.$$

In this case, $(ba)^{nsD} = b((ab)^{nsD})^2a$.

Corollary 3.9 Let $a, b \in R$ and $n, k \in \mathbb{N}$. If $(ab)^k \in R^{nsD}$, then $(ba)^k \in R^{nsD}$.

Proof Since $a((ba)^{k-1}b) = (ab)^k \in R^{nsD}$, by Corollary 3.8 we deduce that $(ba)^k \in R^{nsD}$.

Under the same hypotheses $acd = dbd$ and $dba = aca$, Jacobson's lemma for the n -strong Drazin inverse is investigated as follows.

Theorem 3.10 Let $a, b, c, d \in R$ and $n \in \mathbb{N}$. If $acd = dbd$ and $dba = aca$, then

$$1 - ac \in R^{nsD} \iff 1 - bd \in R^{nsD}.$$

Proof Suppose that $1 - ac \in R^{nsD}$. Then $(1 - ac) - (1 - ac)^{n+1} \in R^{nil}$. Now, by mathematical induction we prove that

$$((1 - bd) - (1 - bd)^{n+1})^{m+1} = -b((1 - ac) - (1 - ac)^{n+1})^m d(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}),$$

for any $m \in \mathbb{N}$.

Assume that $m = 1$. Since $acd = dbd$, we deduce that $(db)^i d = (ac)^i d$ for any $i \in \mathbb{N}$. Then we have

$$\begin{aligned} ((1 - bd) - (1 - bd)^{n+1})^2 &= (bd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^i)(bd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^i) \\ &= (bdbd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} b(db)^i d)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}) \\ &= b(ac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^i)d(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}) \\ &= -b((1 - ac) - (1 - ac)^{n+1})d(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}). \end{aligned}$$

Assume that the conclusion holds for positive integer $m = l$. Now, we verify it for $m = l + 1$. One can see that

$$\begin{aligned} &((1 - bd) - (1 - bd)^{n+1})^{l+2} \\ &= ((1 - bd) - (1 - bd)^{n+1})^{l+1}((1 - bd) - (1 - bd)^{n+1}) \\ &= b((1 - ac) - (1 - ac)^{n+1})^l d(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1})(bd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^i) \\ &= b((1 - ac) - (1 - ac)^{n+1})^l (bdbd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (db)^i d)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}) \\ &= b((1 - ac) - (1 - ac)^{n+1})^l (acd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^i d)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}) \\ &= -b((1 - ac) - (1 - ac)^{n+1})^{l+1} d(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^{i-1}). \end{aligned}$$

Note that $(1 - ac) - (1 - ac)^{n+1} \in R^{nil}$. Then $(1 - bd) - (1 - bd)^{n+1} \in R^{nil}$, i.e. $1 - bd \in R^{nsD}$.

Conversely, assume that $1 - bd \in R^{nsD}$. In order to prove $1 - ac \in R^{nsD}$, we will prove the following equality

$$((1 - ac) - (1 - ac)^{n+1})^{m+2} = d((1 - bd) - (1 - bd)^{n+1})^m bac(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2,$$

for any $m \in \mathbb{N}$.

For the case $m = 1$. Note that $dba = aca$. Then we obtain $a(ca)^i = (db)^i a$, for any $i \in \mathbb{N}$. Hence, we

deduce that

$$\begin{aligned}
 ((1 - ac) - (1 - ac)^{n+1})^3 &= -(ac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^i)(ac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^i)^2 \\
 &= -((ac)^3 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i+2})(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -(a(ca)^2c + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} a(ca)^{i+1}c)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -((db)^2ac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (db)^{i+1}ac)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -(d(bd)bac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} d(bd)^i bac)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -d(bd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^i) bac(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= d((1 - bd) - (1 - bd)^{n+1}) bac(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2.
 \end{aligned}$$

Assume that the conclusion holds for positive integer $m = l$. Then, for the case $m = l + 1$, we get

$$\begin{aligned}
 &((1 - ac) - (1 - ac)^{n+1})^{l+3} \\
 &= -d((1 - bd) - (1 - bd)^{n+1})^l bac(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 (ac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^i) \\
 &= -d((1 - bd) - (1 - bd)^{n+1})^l b(ac)^2(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -d((1 - bd) - (1 - bd)^{n+1})^l (bacac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} ba(ca)^i c)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -d((1 - bd) - (1 - bd)^{n+1})^l (bdbac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} b(db)^i ac)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -d((1 - bd) - (1 - bd)^{n+1})^l (bdbac + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^i bac)(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= -d((1 - bd) - (1 - bd)^{n+1})^l (bd + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (bd)^i) bac(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2 \\
 &= d((1 - bd) - (1 - bd)^{n+1})^{l+1} bac(1 + \sum_{i=1}^{n+1} (-1)^i \binom{n+1}{i} (ac)^{i-1})^2.
 \end{aligned}$$

Observe that $(1 - bd) - (1 - bd)^{n+1} \in R^{nil}$. Hence, $(1 - ac) - (1 - ac)^{n+1} \in R^{nil}$, as required.

Corollary 3.11 *Let $a, b \in R$ and $n \in \mathbb{N}$. Then,*

$$1 - ab \in R^{nsD} \iff 1 - ba \in R^{nsD}.$$

4. The n -strong Drazin invertibility of the sum

Let $p \in R$ be an idempotent ($p^2 = p$). Then we can represent element $a \in R$ as

$$a = pap + pa(1 - p) + (1 - p)ap + (1 - p)a(1 - p)$$

or in the matrix form

$$a = \begin{bmatrix} a_1 & a_3 \\ a_4 & a_2 \end{bmatrix}_p,$$

where $a_1 = pap$, $a_2 = (1 - p)a(1 - p)$, $a_3 = pa(1 - p)$ and $a_4 = (1 - p)ap$. For

$$x = \begin{bmatrix} x_1 & x_3 \\ x_4 & x_2 \end{bmatrix}_p \text{ and } y = \begin{bmatrix} y_1 & y_3 \\ y_4 & y_2 \end{bmatrix}_p,$$

one can use usual matrix rules to obtain matrix forms of the sum $x + y$ and the product xy .

Remark that if $a \in R^{nsD}$, then we have the following matrix representations relative to $p = aa^{nsD}$:

$$a = \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix}_p \text{ and } a^{nsD} = \begin{bmatrix} a_1^{-1} & 0 \\ 0 & 0 \end{bmatrix}_p = \begin{bmatrix} a_1^{nsD} & 0 \\ 0 & 0 \end{bmatrix}_p,$$

where $a_1 \in (pRp)^{-1} \cap (pRp)^{nsD}$ and $a_2 \in ((1 - p)R(1 - p))^{nil}$.

In this section, our purpose is to investigate the n -strong Drazin invertibility of the sum of two elements in a ring. We start with a crucial auxiliary lemma.

Lemma 4.1 *Let $a, b \in R$ be such that $ab = 0$. Then,*

$$a, b \in R^{nil} \iff a + b \in R^{nil}.$$

Proof Since $ab = 0$, we have

$$(a + b)^m = a^m + ba^{m-1} + b^2a^{m-2} + \dots + b^m$$

for any $m \in \mathbb{N}$.

Suppose that $a, b \in R^{nil}$. Choose $k_1 \in \mathbb{N}$ satisfying $a^{k_1} = b^{k_1} = 0$. Then, we have $(a + b)^{2k_1} = 0$, which gives $a + b \in R^{nil}$.

On the contrary, assume that $(a + b)^{k_2} = 0$, for some $k_2 \in \mathbb{N}$. Then,

$$a^{k_2} + ba^{k_2-1} + b^2a^{k_2-2} + \dots + b^{k_2} = 0.$$

Multiplying the preceding equality by a from the left side (resp. by b from the right side), we obtain $a^{k_2+1} = 0$ (resp. $b^{k_2+1} = 0$). Hence, we have $a, b \in R^{nil}$.

Now, we state the relationship between the n -strong Drazin invertibility of the elements a, b and that of the sum $a + b$, under the condition $ab = 0$.

Theorem 4.2 *Let $n \in \mathbb{N}$ and $a, b \in R$ be such that $ab = 0$. Then,*

$$a, b \in R^{nsD} \iff a + b \in R^{nsD}.$$

Proof By the hypothesis $ab = 0$, we have

$$\begin{aligned} x &:= (a + b) - (a + b)^{n+1} \\ &= (a - a^{n+1}) + (b - b^{n+1}) - (ba^n + b^2a^{n-1} + \dots + b^n a) \\ &:= x_1 + x_2 - x_3. \end{aligned}$$

Note that $x_1(x_2 - x_3) = 0$, $x_3x_2 = 0$ and $x_3^2 = 0$. In view of Lemma 4.1, we get

$$\begin{aligned} x \in R^{nil} &\iff x_1 \in R^{nil} \text{ and } x_2 - x_3 \in R^{nil} \\ &\iff x_1 \in R^{nil} \text{ and } x_2 \in R^{nil}. \end{aligned}$$

Then, by Theorem 3.2 we obtain $a, b \in R^{nsD}$ if and only if $a + b \in R^{nsD}$.

Remark 4.3 (1) For the Drazin invertibility, we have

$$a, b \in R^D \iff a + b \in R^D,$$

under the condition $ab = ba = 0$. In fact, the necessity can be seen from [9, Corollary 1]. Now, suppose that $a + b \in R^D$. Then $(a + b)^m = (a + b)^{m+1}R \cap R(a + b)^{m+1}$, for some $m \in \mathbb{N}$. Hence, we have $a^m + b^m = (a^{m+1} + b^{m+1})u = v(a^{m+1} + b^{m+1})$, for some $u, v \in R$. Multiplying the previous equality by a from the left side and right side respectively, we have $a^{m+1} = a^{m+2}u = va^{m+2}$. Hence, $a \in R^D$. Similarly, we can obtain $b \in R^D$.

(2) By [3, Theorem 2.1], one can see that

$$a, b \in R^D \implies a + b \in R^D,$$

under the condition $ab = 0$. Now, we consider its converse. Assume that $a + b \in R^D$. Then, we can obtain that a is right π -regular and b is left π -regular. Is a left π -regular? Is b right π -regular?

Next, we will consider the n -strong Drazin invertibility of the sum $a + b$ under another new condition $a^2b = aba$ and $ab^2 = bab$, which is weaker than $ab = ba$. Indeed, it is obvious that $ab = ba$ imply $a^2b = aba$ and $ab^2 = bab$. However, the converse does not hold in general, which can be seen from the following example:

Example 4.4 Let $R = M_2(\mathbb{Z}_2)$, where \mathbb{Z}_2 denote the residue class ring modulo 2. Take $a = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ and $b = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$. Clearly, $a^2b = aba$ and $ab^2 = bab$. However, $ab \neq ba$.

In order to prove our main result, we need the following lemmas.

Lemma 4.5 Let $a, b \in R^{nil}$. If $a^2b = aba$ and $ab^2 = bab$, then $a + b \in R^{nil}$.

Proof By the hypothesis $a, b \in R^{nil}$, there exists $m \in \mathbb{N}$ such that $a^m = 0$ and $b^m = 0$. Since $a^2b = aba$ and $ab^2 = bab$, we can see that each of the monomials in the expansion of $(a + b)^{3m}$ is either $a^{k_1}b^{k_2}a^{k_3}$ or $b^{l_1}a^{l_2}b^{l_3}$, where $k_1 + k_2 + k_3 = l_1 + l_2 + l_3 = 3m$. Hence, $(a + b)^{3m} = 0$, which means $a + b \in R^{nil}$.

Lemma 4.6 Let $x \in R$ and $p^2 = p \in R$. If x has the representation $x = \begin{bmatrix} a & c \\ 0 & b \end{bmatrix}_p$, then

$$a \in (pRp)^{nil} \text{ and } b \in ((1 - p)R(1 - p))^{nil} \iff x \in R^{nil}.$$

Proof Assume that $a \in (pRp)^{nil}$ and $b \in ((1 - p)R(1 - p))^{nil}$. By a simple computation, we obtain

$$x^k = \begin{bmatrix} a^k & f_k \\ 0 & b^k \end{bmatrix}_p, \text{ for any } k \in \mathbb{N}, \text{ where } f_k = \sum_{i=0}^{k-1} a^i c b^{k-i-1}. \text{ Let } a^{t_1} = 0 \text{ and } b^{t_2} = 0, \text{ where } t_1, t_2 \in \mathbb{N}.$$

Then, we have $x^{t_1+t_2} = 0$, i.e. $x \in R^{nil}$. Conversely, it is clear.

Lemma 4.7 Let $n \in \mathbb{N}$ and $p^2 = p, x, y \in R$. If x and y have the representations

$$x = \begin{bmatrix} a & c \\ 0 & b \end{bmatrix}_p \quad \text{and} \quad y = \begin{bmatrix} b & 0 \\ c & a \end{bmatrix}_{1-p},$$

then

$$a \in (pRp)^{nsD} \text{ and } b \in ((1-p)R(1-p))^{nsD} \iff x \in R^{nsD} \text{ (resp. } y \in R^{nsD}\text{)}.$$

Proof Observe that

$$x - x^{n+1} = \begin{bmatrix} a & c \\ 0 & b \end{bmatrix}_p - \begin{bmatrix} a & c \\ 0 & b \end{bmatrix}_p^{n+1} = \begin{bmatrix} a - a^{n+1} & * \\ 0 & b - b^{n+1} \end{bmatrix}_p.$$

By Lemma 4.6, it follows that

$$a - a^{n+1} \in (pRp)^{nil} \text{ and } b - b^{n+1} \in ((1-p)R(1-p))^{nil} \iff x - x^{n+1} \in R^{nil}.$$

Using Theorem 3.2, we complete the proof.

Lemma 4.8 Let $n \in \mathbb{N}$ and $a, b \in R^{nsD}$ be such that $a^2b = aba$. Then $ab \in R^{nsD}$.

Proof Since $a^2b = aba$, by induction we can obtain $(ab)^m = a^m b^m$ for any $m \in \mathbb{N}$. Applying Theorem 3.2, we only need to prove

$$ab - (ab)^{n+1} = (a - a^{n+1})b + a^{n+1}(b - b^{n+1}) := x + y \in R^{nil}.$$

Note that

$$\begin{aligned} yx &= a^{n+1}(b - b^{n+1})(a - a^{n+1})b \\ &= (a^n(ab) - (ab)^{n+1})(a - a^{n+1})b \\ &= (a - a^{n+1})(a^n(ab) - (ab)^{n+1})b \\ &= a^{n+1}(a - a^{n+1})b(b - b^{n+1}) \\ &= (a - a^{n+1})ba^{n+1}(b - b^{n+1}) \\ &= xy. \end{aligned}$$

In addition, we can check that $x^m = (a - a^{n+1})^m b^m$ and $y^m = (a^{n+1})^m (b - b^{n+1})^m$ for any $m \in \mathbb{N}$. Note that $a - a^{n+1} \in R^{nil}$ and $b - b^{n+1} \in R^{nil}$, which imply $x \in R^{nil}$ and $y \in R^{nil}$. Hence, $x + y \in R^{nil}$ by Lemma 2.1(2).

Remark 4.9 In view of Lemma 4.8 and [26, Lemma 2], one can see that if $n \in \mathbb{N}$ and $a, b \in R^{nsD}$ be such that $ab = ba$, then $ab \in R^{nsD}$ and $(ab)^{nsD} = b^{nsD} a^{nsD}$.

Now, we state our main result in this section as follows.

Theorem 4.10 Let $n \in \mathbb{N}$ and $a, b \in R^{nsD}$ be such that $a^2b = aba$ and $ab^2 = bab$. Then,

$$1 + a^{nsD}b \in R^{nsD} \iff a + b \in R^{nsD}.$$

Proof We consider the matrix representations of a and b relative to the idempotent $p = aa^{nsD}$:

$$a = \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix}_p \quad \text{and} \quad b = \begin{bmatrix} b_1 & b_3 \\ b_4 & b_2 \end{bmatrix}_p,$$

where $a_1 \in (pRp)^{-1} \cap (pRp)^{nsD}$ and $a_2 \in ((1-p)R(1-p))^{nil}$.

The condition $a^2b = aba$ expressed in matrix form yields

$$\begin{bmatrix} a_1^2b_1 & a_1^2b_3 \\ a_2^2b_4 & a_2^2b_2 \end{bmatrix}_p = a^2b = aba = \begin{bmatrix} a_1b_1a_1 & a_1b_3a_2 \\ a_2b_4a_1 & a_2b_2a_2 \end{bmatrix}_p.$$

Thus, we have $a_1^2b_3 = a_1b_3a_2$, i.e. $b_3 = a_1^{-1}b_3a_2$, which implies $b_3 = a_1^{-m}b_3a_2^m$ for any $m \in \mathbb{N}$. Since $a_2 \in ((1-p)R(1-p))^{nil}$, we have $b_3 = 0$. Moreover, we can get $a_1b_1 = b_1a_1$ and $a_2^2b_2 = a_2b_2a_2$. Similarly, by $ab^2 = bab$ we obtain $a_2b_2^2 = b_2a_2b_2$. Therefore, we have

$$b = \begin{bmatrix} b_1 & 0 \\ b_4 & b_2 \end{bmatrix}_p \quad \text{and} \quad a + b = \begin{bmatrix} a_1 + b_1 & 0 \\ b_4 & a_2 + b_2 \end{bmatrix}_p.$$

Now, we prove that $a_2 + b_2 \in ((1-p)R(1-p))^{nsD}$. Since $b \in R^{nsD}$, by Lemma 4.7 we have $b_2 \in ((1-p)R(1-p))^{nsD}$. Let $p' = b_2b_2^{nsD}$. We consider the matrix representations of b_2 and a_2 relative to the idempotent p' :

$$b_2 = \begin{bmatrix} b'_1 & 0 \\ 0 & b'_2 \end{bmatrix}_{p'} \quad \text{and} \quad a_2 = \begin{bmatrix} a'_1 & a'_3 \\ a'_4 & a'_2 \end{bmatrix}_{p'},$$

where $b'_1 \in (p'Rp')^{-1} \cap (p'Rp')^{nsD}$ and $b'_2 \in ((1-p')R(1-p'))^{nil}$. Note that $a_2^2b_2 = a_2b_2a_2$ and $a_2b_2^2 = b_2a_2b_2$. Then, we can obtain that $a'_4 = 0$, $a'_1b'_1 = b'_1a'_1$, $a'_2(b'_2)^2 = b'_2a'_2b'_2$ and $(a'_2)^2b'_2 = a'_2b'_2a'_2$. Hence,

$$a_2 = \begin{bmatrix} a'_1 & a'_3 \\ 0 & a'_2 \end{bmatrix}_{p'} \quad \text{and} \quad a_2 + b_2 = \begin{bmatrix} a'_1 + b'_1 & a'_3 \\ 0 & a'_2 + b'_2 \end{bmatrix}_{p'}.$$

In order to show that $a'_1 + b'_1 = (p' + a'_1(b'_1)^{-1})b'_1 \in (p'Rp')^{nsD}$, by Lemma 4.8 we only need to prove $p' + a'_1(b'_1)^{-1} \in (p'Rp')^{nsD}$. Since $a_2 \in ((1-p)R(1-p))^{nil}$, by Lemma 4.6 we obtain $a'_1 \in (p'Rp')^{nil}$, which yields

$$\begin{aligned} & (p' + a'_1(b'_1)^{-1}) - (p' + a'_1(b'_1)^{-1})^{n+1} \\ &= a'_1(b'_1)^{-1} \left(p' - \sum_{i=1}^{n+1} \binom{n+1}{i} (a'_1)^{i-1} (b'_1)^{1-i} \right) \in (p'Rp')^{nil}. \end{aligned}$$

Hence, $p' + a'_1(b'_1)^{-1} \in (p'Rp')^{nsD}$. Applying Lemma 4.5 to the nilpotent elements a'_2 and b'_2 , we conclude that $a'_2 + b'_2 \in ((1-p')R(1-p'))^{nil}$, which implies $a'_2 + b'_2 \in ((1-p')R(1-p'))^{nsD}$. In view of Lemma 4.7, we obtain $a_2 + b_2 \in ((1-p)R(1-p))^{nsD}$. Then, by Lemma 4.7 again, it follows that $a + b \in R^{nsD}$ is equivalent to $a_1 + b_1 \in (pRp)^{nsD}$.

Note that

$$1 + a^{nsD}b = \begin{bmatrix} p + a_1^{-1}b_1 & 0 \\ 0 & 1 - p \end{bmatrix}_p.$$

Since $1 - p \in ((1 - p)R(1 - p))^{nsD}$, then $1 + a^{nsD}b \in R^{nsD}$ is equivalent to $p + a_1^{-1}b_1 \in (pRp)^{nsD}$. Note that $a_1 \in (pRp)^{nsD}$. Applying Corollary 3.5 we obtain $a_1^{-1} = a_1^{nsD} \in (pRp)^{nsD}$. Hence, $a_1 + b_1 = a_1(p + a_1^{-1}b_1) \in (pRp)^{nsD}$ is identical to $p + a_1^{-1}b_1 \in (pRp)^{nsD}$ by Lemma 4.8. Hence, we conclude that $a + b \in R^{nsD}$ if and only if $1 + a^{nsD}b \in R^{nsD}$.

Remark 4.11 *In the proof of the necessity of Theorem 4.10, the condition $ab^2 = bab$ was not used. However, if we drop it, then the sufficiency is not true in general, which will be shown in the next example:*

Example 4.12 *Let $n = 1 \in \mathbb{N}$ and $R = M_3(\mathbb{C})$. Choose*

$$a = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \text{ and } b = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

Then, we can check that $a^2b = aba$, $a^2 = 0$ and $b^3 = 0$. Hence, $a, b, 1 + a^{sD}b = 1 \in R^{sD}$. Note that the eigenvalues of $(a + b) - (a + b)^2$ are $0, \sqrt{3}i$ and $-\sqrt{3}i$. Hence, $(a + b) - (a + b)^2 \notin R^{nil}$, which yields that $a + b \notin R^{sD}$. In addition, this example also illustrates that the condition $ab^2 = bab$ of Lemma 4.5 cannot be dropped.

The following corollary can be directly derived from Theorem 4.10.

Corollary 4.13 *Let $a, b \in R^{nsD}$ be such that $ab = ba$. Then,*

$$a + b \in R^{nsD} \iff 1 + a^{nsD}b \in R^{nsD}.$$

5. The n -strong Drazin invertibility of the matrix

In this section, as applications for our additive results of Section 4, we obtain some equivalent conditions for the n -strong Drazin invertibility of the matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ over a ring. For the convenience of expressions, we assume that $\sum_{i=j}^k s(i) = 0$ if $k < j$, where $s(i)$ is a function on i , and $a^0 = 1$ for $a \in R$. For any nonnegative integer k , by $[k/2]$ we denote the integer part of $k/2$.

Firstly, we investigate the n -strong Drazin invertibility of some special antitriangular matrices over a ring.

Proposition 5.1 *Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & b \\ c & 0 \end{bmatrix} \in M_2(R)$ be such that $ab = b$, $a = a^2$. Then,*

- (1) $bc \in R^{nil}$ if and only if $M \in M_2(R)^{sD}$.
- (2) If $bc \in R^{nil}$, then $M \in M_2(R)^{nsD}$.

Proof (1) Since $ab = b$ and $a = a^2$, we have

$$M - M^2 = \begin{bmatrix} -bc & 0 \\ c - ca & -cb \end{bmatrix}.$$

Hence, $M - M^2 \in M_2(R)^{nil}$ is identical to $bc \in R^{nil}$. From Theorem 3.2, it follows that $bc \in R^{nil}$ if and only if $M \in M_2(R)^{sD}$.

(2) This follows from item (1) and Corollary 3.6(1) directly.

Remark 5.2 (1) In Proposition 5.1(1), if we change the condition “ $bc \in R^{nil}$ ” to “ $bc \in R^{sD}$ ”, then the conclusion does not hold in general. For example, take $M = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \in M_2(\mathbb{C})$. Obviously, $1 \in \mathbb{C}^{sD}$. However, $M \notin M_2(\mathbb{C})^{sD}$, since $M - M^2 \notin M_2(\mathbb{C})^{nil}$.

(2) The converse of Proposition 5.1(2) is not true for $n \geq 2$ in general, which will be illustrated by the following example:

Example 5.3 Let $R = M_2(\mathbb{Z}_3)$. Choose $a = b = 1 \in R$ and $c = \begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix} \in R$. Then, it is easy to see $bc \notin R^{nil}$. However, $M = \begin{bmatrix} a & b \\ c & 0 \end{bmatrix} \in M_2(R)^{2sD}$, since we can check that $(M - M^3)^2 = 0$.

Theorem 5.4 Let $n \in \mathbb{N}$ and $a, b \in R$ be such that $aba = 0$. Then,

$$M = \begin{bmatrix} a & a \\ b & 0 \end{bmatrix} \in M_2(R)^{nsD} \iff a \in R^{nsD} \iff M' = \begin{bmatrix} a & b \\ a & 0 \end{bmatrix} \in M_2(R)^{nsD}.$$

Proof We will only prove that $a \in R^{nsD}$ is equivalent to $M \in M_2(R)^{nsD}$, since the case for M' is similar.

Suppose that $n = 1$. Then, by the condition $aba = 0$ we have

$$X := M - M^2 = \begin{bmatrix} a - a^2 - ab & a - a^2 \\ b - ba & -ba \end{bmatrix}$$

and

$$X^{m+1} = \begin{bmatrix} (a - a^2)^m(a - a^2 - ab + b) & (a - a^2)^{m+1} \\ (b - ba)(a - a^2)^{m-1}(a - a^2 - ab + b) & (b - ba)(a - a^2)^m \end{bmatrix}$$

for any $m \geq 2$. Hence, $X \in M_2(R)^{nil}$ if and only if $a - a^2 \in R^{nil}$. Applying Theorem 3.2, we claim that $a \in R^{sD}$ is equivalent to $M \in M_2(R)^{sD}$.

The result for $n \geq 2$ follows analogously.

Next, we present an existence criterion for the n -strong Drazin inverse of the anti-triangular $\begin{bmatrix} a & 1 \\ b & 0 \end{bmatrix}$,

which will be used later.

Theorem 5.5 Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & 1 \\ b & 0 \end{bmatrix} \in M_2(R)$ be such that $ab = 0$. Then,

$$a \in R^{nsD} \text{ and } N = \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} \in M_2(R)^{nil} \iff M \in M_2(R)^{nsD},$$

where

$$\begin{aligned} n_{11} &= n_{22} = \frac{1-(-1)^n}{2} (b^{\lfloor \frac{n+1}{2} \rfloor})^2 + (1 - \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor})(b - \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor}), \\ n_{12} &= ((-1)^n - 1)b^{\lfloor \frac{n+1}{2} \rfloor}, \\ n_{21} &= ((-1)^n - 1)b^{\lfloor \frac{n+1}{2} \rfloor + 1}. \end{aligned}$$

Proof Since $ab = 0$, then by induction we have

$$\begin{aligned} X &:= M - M^{n+1} \\ &= \begin{bmatrix} (a - a^{n+1}) - (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) & (1 - a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}}) - (t_3 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor}) \\ b - (t_2 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor}) & -(t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) \end{bmatrix}, \end{aligned}$$

where

$$\begin{aligned} t_1 &= \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} b^i a^{2(\lfloor \frac{n+1}{2} \rfloor - i) + \frac{1-(-1)^{n+1}}{2}}, \\ t_2 &= \sum_{i=1}^{\lfloor \frac{n+1}{2} \rfloor} b^i a^{2(\lfloor \frac{n+1}{2} \rfloor - i) + \frac{3-(-1)^{n+1}}{2}}, \\ t_3 &= \sum_{i=1}^{\lfloor \frac{n-1}{2} \rfloor} b^i a^{2(\lfloor \frac{n}{2} \rfloor - i) + \frac{1-(-1)^n}{2}}. \end{aligned}$$

By a computation, we obtain $X^2 = \begin{bmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \end{bmatrix}$, where

$$\begin{aligned} u_{11} &= (a - a^{n+1})^2 - (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor})(a - a^{n+1}) + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) \\ &\quad + (1 - \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor})(b - (t_2 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor})), \\ u_{12} &= (a - a^{n+1})(1 - a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}}) - (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor})(1 - a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}}) \\ &\quad - (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor} (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} (t_3 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor}), \\ u_{21} &= b(a - a^{n+1}) - b(t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) - (t_2 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor})(a - a^{n+1}) \\ &\quad + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor} (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) - \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} (b - (t_2 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor})), \\ u_{22} &= b(1 - a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}}) - b(t_3 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor}) - (t_2 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor})(1 - a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}}) \\ &\quad + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor} (t_3 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor}) + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}). \end{aligned}$$

Consider the following splitting:

$$X^2 = \begin{bmatrix} (a - a^{n+1})^2 & p_{12} \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} q_{11} & q_{12} \\ q_{21} & q_{22} \end{bmatrix} + \begin{bmatrix} n_{11} & n_{12} \\ n_{21} & n_{22} \end{bmatrix} := P + Q + N,$$

where

$$\begin{aligned} p_{12} &= (a - a^{n+1})(1 - a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}}), \\ q_{11} &= -(t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor})(a - a^{n+1}) + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} t_1 - (1 - \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor}) t_2, \\ q_{12} &= -2t_1 + (t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor}) a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}} + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n}{2} \rfloor} t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} t_3, \\ q_{21} &= b(a - a^{n+1}) - bt_1 - (t_2 + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor})(a - a^{n+1}) + \frac{1+(-1)^n}{2} b^{\lfloor \frac{n+2}{2} \rfloor} t_1 + \frac{1-(-1)^n}{2} b^{\lfloor \frac{n+1}{2} \rfloor} t_2, \end{aligned}$$

$$q_{22} = -bt_3 - t_2 + (t_2 + \frac{1+(-1)^n}{2}b^{\lfloor \frac{n+2}{2} \rfloor} - b)a^{2\lfloor \frac{n}{2} \rfloor + \frac{1-(-1)^n}{2}} + \frac{1+(-1)^n}{2}b^{\lfloor \frac{n+2}{2} \rfloor}t_3 + \frac{1-(-1)^n}{2}b^{\lfloor \frac{n+1}{2} \rfloor}t_1.$$

Note that $P(Q + N) = 0$, $QN = 0$ and $Q^2 = 0$. In view of Lemma 4.1, we have

$$\begin{aligned} X \in M_2(R)^{nil} &\iff X^2 \in M_2(R)^{nil} \\ &\iff P \in M_2(R)^{nil} \text{ and } Q + N \in M_2(R)^{nil} \\ &\iff P \in M_2(R)^{nil} \text{ and } N \in M_2(R)^{nil} \\ &\iff a - a^{n+1} \in R^{nil} \text{ and } N \in M_2(R)^{nil}. \end{aligned}$$

In view of Theorem 3.2, we can conclude that $a \in R^{nsD}$ and $N \in M_2(R)^{nil}$ if and only if $M \in M_2(R)^{nsD}$.

Now, we state a special case of Theorem 5.5.

Corollary 5.6 *Let $n = 2k$ ($k \in \mathbb{N}$) and let $a, b \in R$ be such that $ab = 0$. Then,*

$$a \in R^{nsD} \text{ and } b \in R^{ksD} \iff M = \begin{bmatrix} a & 1 \\ b & 0 \end{bmatrix} \in M_2(R)^{nsD}.$$

Proof Let $n = 2k$ in Theorem 5.5, we have $N = \begin{bmatrix} b(1 - b^k)^2 & 0 \\ 0 & b(1 - b^k)^2 \end{bmatrix}$. Then, one can see that

$$N \in M_2(R)^{nil} \iff b(1 - b^k)^2 \in R^{nil} \iff (b(1 - b^k))^2 \in R^{nil} \iff b - b^{k+1} \in R^{nil}.$$

Hence, we have that $M \in M_2(R)^{nsD}$ is equivalent to $a \in R^{nsD}$ and $b \in R^{ksD}$.

Remark 5.7 *By Corollary 5.6 and Corollary 3.6(3), we can see that*

$$M = \begin{bmatrix} a & 1 \\ b & 0 \end{bmatrix} \in M_2(R)^{nsD} \implies a \in R^{nsD} \text{ and } b \in R^{nsD},$$

under the condition $ab = 0$ and n is one even number. However, the converse does not hold in general, which can be seen in the next example:

Example 5.8 *Let $R = M_2(\mathbb{C})$ and $n = 2 \in \mathbb{N}$. Setting $M = \begin{bmatrix} 0 & 1 \\ b & 0 \end{bmatrix} \in M_2(R)$, where $b = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \in R$. Obviously, $b - b^3 = 0$, which yields $b \in R^{2sD}$. However, we can check that $M - M^3 \notin M_2(R)^{nil}$, so we have $M \notin M_2(R)^{2sD}$.*

Following the same strategy as in the proof of Theorem 5.5, we derive the equivalent condition for the n -strong Drazin invertibility of the transpose of the matrix M as follows:

Theorem 5.9 *Let $n \in \mathbb{N}$ and $M' = \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix} \in M_2(R)$ be such that $ab = 0$. Then,*

$$a \in R^{nsD} \text{ and } N' = \begin{bmatrix} n_{11} & n_{21} \\ n_{12} & n_{22} \end{bmatrix} \in M_2(R)^{nil} \iff M' \in M_2(R)^{nsD},$$

where n_{11} , n_{12} , n_{21} , and n_{22} are defined as in Theorem 5.5.

Combining Theorem 5.5 and Theorem 5.9, together with the equality $(N')^m = (N^m)'$ for any $m \in \mathbb{N}$, we obtain the relationship between the n -strong Drazin invertibility of the matrix M and that of its transpose M' .

Corollary 5.10 *Let $n \in \mathbb{N}$ and let $a, b \in R$ be such that $ab = 0$. Then,*

$$M = \begin{bmatrix} a & 1 \\ b & 0 \end{bmatrix} \in M_2(R)^{nsD} \iff M' = \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix} \in M_2(R)^{nsD}.$$

In the rest of this section, applying the previous results we obtain some characterizations for the n -strong Drazin invertibility of the matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$, under some conditions.

Theorem 5.11 *Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)$ be such that $abc = 0$ and $bd = 0$. Then,*

$$a, d \in R^{nsD} \text{ and } T = \begin{bmatrix} t_{11} & t_{12} \\ t_{21} & t_{22} \end{bmatrix} \in M_2(R)^{nil} \iff M \in M_2(R)^{nsD},$$

where

$$t_{11} = t_{22} = \frac{1-(-1)^n}{2} ((bc)^{\lfloor \frac{n+1}{2} \rfloor})^2 + (1 - \frac{1+(-1)^n}{2} (bc)^{\lfloor \frac{n}{2} \rfloor}) (bc - \frac{1+(-1)^n}{2} (bc)^{\lfloor \frac{n+2}{2} \rfloor}),$$

$$t_{12} = ((-1)^n - 1)(bc)^{\lfloor \frac{n+1}{2} \rfloor + 1},$$

$$t_{21} = ((-1)^n - 1)(bc)^{\lfloor \frac{n+1}{2} \rfloor}.$$

Proof We write $M = P + Q$, where

$$P = \begin{bmatrix} a & b \\ c & 0 \end{bmatrix} \text{ and } Q = \begin{bmatrix} 0 & 0 \\ 0 & d \end{bmatrix}.$$

The condition $bd = 0$ ensures $PQ = 0$. Note that $Q \in M_2(R)^{nsD}$ if and only if $d \in R^{nsD}$. In view of Theorem 4.2, we obtain $M \in M_2(R)^{nsD}$ is equivalent to $P \in M_2(R)^{nsD}$ and $d \in R^{nsD}$.

Since

$$P = \begin{bmatrix} a & b \\ c & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & c \end{bmatrix} \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix},$$

by Corollary 3.8 we have

$$P \in M_2(R)^{nsD} \iff P' := \begin{bmatrix} a & b \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & c \end{bmatrix} = \begin{bmatrix} a & bc \\ 1 & 0 \end{bmatrix} \in M_2(R)^{nsD}.$$

Since $abc = 0$, by Theorem 5.9 we obtain

$$a \in R^{nsD} \text{ and } T = \begin{bmatrix} t_{11} & t_{12} \\ t_{21} & t_{22} \end{bmatrix} \in M_2(R)^{nil} \iff P' \in M_2(R)^{nsD},$$

as required.

Now, we can derive some special cases of Theorem 5.11.

Corollary 5.12 Let $n = 2k$ ($k \in \mathbb{N}$) and $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)$ be such that $abc = 0$ and $bd = 0$.

Then,

$$a, d \in R^{nsD} \text{ and } bc \in R^{ksD} \iff M \in M_2(R)^{nsD}.$$

Let $k = 1$ in Corollary 5.12, we have

Corollary 5.13 Let $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)$ be such that $abc = 0$ and $bd = 0$. Then,

$$a, d \in R^H \text{ and } bc \in R^{sD} \iff M \in M_2(R)^H.$$

Corollary 5.14 Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)$ be such that $abc = 0$, $bd = 0$ and $bc = bca$.

Then,

$$a, d \in R^{nsD} \iff M \in M_2(R)^{nsD}.$$

Proof Using the condition $abc = 0$ and $bc = bca$, we have $T^2 = 0$, where T is defined as in Theorem 5.11.

Remark 5.15 Let $n \in \mathbb{N}$ and $a, b, c, d \in R$ be such that $abc = 0$ and $bd = 0$. Then

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)^{nsD} \implies M' = \begin{bmatrix} a & c \\ b & d \end{bmatrix} \in M_2(R)^{nsD}$$

does not hold in general, even if $d = 0$. For example:

Example 5.16 Let $R = M_2(\mathbb{Z}_2)$ and $n = 1 \in \mathbb{N}$. Choose $M = \begin{bmatrix} a & b \\ c & 0 \end{bmatrix} \in M_2(R)$, where

$$a = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, b = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \text{ and } c = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}.$$

Then, we can check that $(M - M^2)^2 = 0$ and $M' - (M')^2 \notin M_2(R)^{nil}$. Hence, $M \in M_2(R)^{sD}$. However, $M' \notin M_2(R)^{sD}$.

Similar to the proof of Lemma 4.7 and using the representations of [11, Theorem 1], we can obtain the following lemma.

Lemma 5.17 Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} \in M_2(R)$. Then

$$a \in R^{nsD} \text{ and } d \in R^{nsD} \iff M \in M_2(R)^{nsD}.$$

In this case,

$$M^{nsD} = \begin{bmatrix} a^{nsD} & z \\ 0 & d^{nsD} \end{bmatrix},$$

where

$$z = \sum_{i=0}^{\text{ind}(d)-1} (a^{nsD})^{i+2} b d^i (1 - d d^{nsD}) + \sum_{i=0}^{\text{ind}(a)-1} (1 - a a^{nsD}) a^i b (d^{nsD})^{i+2} - a^{nsD} b d^{nsD}.$$

Applying Lemma 5.17, we may now state the following result.

Theorem 5.18 *Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)$ be such that $ca = 0$ and $cb = 0$. Then,*

$$a, d \in R^{nsD} \iff M \in M_2(R)^{nsD}.$$

Proof The matrix M can be split as

$$M = \begin{bmatrix} 0 & 0 \\ c & 0 \end{bmatrix} + \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} := P + Q.$$

The conditions $ca = 0$ and $cb = 0$ imply $PQ = 0$. Note that $P^2 = 0$. By Theorem 4.2 and Lemma 5.17, we conclude that $a, d \in R^{nsD}$ if and only if $M \in M_2(R)^{nsD}$.

The next theorem presents new conditions under which we give a characterization for the n -strong Drazin invertibility of the matrix M over a ring.

Theorem 5.19 *Let $n \in \mathbb{N}$ and $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in M_2(R)$ be such that $bc = cb = 0$ and $ca = dc$. Then,*

$$a, d \in R^{nsD} \iff M \in M_2(R)^{nsD}.$$

Proof Suppose that $a, d \in R^{nsD}$. Now, we consider the following splitting

$$M = \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ c & 0 \end{bmatrix} := P + Q.$$

The conditions $bc = cb = 0$ and $ca = dc$ imply $PQ = QP$. In view of Lemma 5.17, we deduce that $P \in M_2(R)^{nsD}$ and

$$1 + P^{nsD}Q = \begin{bmatrix} 1 + zc & 0 \\ d^{nsD}c & 1 \end{bmatrix},$$

where z is defined as in Lemma 5.17. Since $ca = dc$, by [10, Theorem 2] we obtain

$$ca^{nsD} = ca^D = d^Dc = d^{nsD}c.$$

Hence, for any $m \in \mathbb{N}$, we have

$$b(d^{nsD})^m c = bc(a^{nsD})^m = 0,$$

and

$$bd^m(1 - dd^{nsD})c = bd^m c - bd^{m+1}d^{nsD}c = bca^m - bca^{m+1}a^{nsD} = 0.$$

In addition, $a^{nsD}bd^{nsD}c = a^{nsD}bca^{nsD} = 0$. Hence, $zc = 0$. Hence, $1 + P^{nsD}Q \in M_2(R)^{nsD}$. Applying Corollary 4.13, we deduce that $M \in M_2(R)^{nsD}$.

Conversely, suppose that $M \in M_2(R)^{nsD}$, i.e. $X := M - M^{n+1} \in M_2(R)^{nil}$. By induction, we can obtain

$$X = \begin{bmatrix} a - a^{n+1} & b - \sum_{i=0}^n a^{n-i}bd^i \\ c - (n+1)d^n c & d - d^{n+1} \end{bmatrix}.$$

By a computation, we have

$$X^m = \begin{bmatrix} (a - a^{n+1})^m & * \\ * & (d - d^{n+1})^m \end{bmatrix},$$

for any $m \in \mathbb{N}$. Therefore, we conclude that $a - a^{n+1} \in R^{nil}$ and $d - d^{n+1} \in R^{nil}$, which yield $a, d \in R^{nsD}$, as required.

Acknowledgments

We are very grateful to the referees for their careful reading and valuable suggestions to the improvement of this paper. This research was supported by the National Natural Science Foundation of China (No. 11961076), China Postdoctoral Science Foundation (No. 2018M632385), Research Project of Hubei Provincial Department of Education (No. B2019128), the Ministry of Education, Science and Technological Development, Republic of Serbia (No. 174007).

References

- [1] Castro-González N, Mendes-Araújo C, Patrício P. Generalized inverses of a sum in rings. *Bulletin of the Australian Mathematical Society* 2010; 82: 156-164. doi: 10.1017/S0004972710000080
- [2] Chen HY, Sheibani M. On Hirano inverses in rings. *Turkish Journal of Mathematics* 2019; 43 (4): 2049-2057. doi: 10.3906/mat-1902-105
- [3] Chen JL, Zhuang GF, Wei YM. The Drazin inverse of a sum of morphisms. *Acta Mathematica Scientia* 2009; 29A (3): 538-552 (in Chinese).
- [4] Cline RE. An application of representation for the generalized inverse of a matrix. *MRC Technical Report* 592, 1965.
- [5] Corach G, Duggal B, Harte RE. Extensions of Jacobson's lemma. *Communications in Algebra* 2013; 41: 520-531. doi: 10.1080/00927872.2011.602274
- [6] Cvetković-Ilić DS, Harte R. On a Jacobson's lemma and Drazin invertibility. *Applied Mathematics Letters* 2010; 23: 417-420. doi: 10.1016/j.aml.2009.11.009
- [7] Cvetković-Ilić DS, Liu XY, Wei YM. Some additive results for the generalized Drazin inverse in a Banach algebra. *Electronic Journal of Linear Algebra* 2011; 22: 1049-1058.
- [8] Deng CY, Wei YM. New additive results for the generalized Drazin inverse. *Journal of Mathematical Analysis and Applications* 2010; 370: 313-321. doi: 10.1016/j.jmaa.2010.05.010
- [9] Drazin MP. Pseudo-inverses in associative rings and semigroups. *American Mathematical Monthly* 1958; 65: 506-514.
- [10] Drazin MP. Commuting properties of generalized inverses. *Linear Multilinear Algebra* 2013; 61: 1675-1681. doi: 10.1080/03081087.2012.753593
- [11] Hartwig RE, Shoaf J. Group inverses and Drazin inverses of bidiagonal and triangular Toeplitz matrices. *Journal of the Australian Mathematical Society* 1977; 24 (Series A): 10-34.
- [12] Hartwig RE, Wang GR, Wei YM. Some additive results on Drazin inverse. *Linear Algebra and its Applications* 2001; 322: 207-217. doi: 10.1016/S0024-3795(00)00257-3
- [13] Lian HF, Zeng QP. An extension of Cline's formula for a generalized Drazin inverse. *Turkish Journal of Mathematics* 2016; 40: 161-165. doi: 10.3906/mat-1505-4
- [14] Liao YH, Chen JL, Cui J. Cline's formula for the generalized Drazin inverse. *Bulletin of the Malaysian Mathematical Sciences Society* 2014; 37: 37-42.

- [15] Mosić D. The generalized and pseudo n -strong Drazin inverses in rings. *Linear Multilinear Algebra* 2019. doi: 10.1080/03081087.2019.1599806
- [16] Mosić D. A note on Cline's formula for the generalized Drazin inverse. *Linear Multilinear Algebra* 2015; 63: 1106-1110. doi: 10.1080/03081087.2014.922965
- [17] Mosić D. Extensions of Jacobson's lemma for Drazin inverse. *Aequationes mathematicae* 2017; 91: 419-428. doi: 10.1007/s00010-017-0476-9
- [18] Patrício P, Hartwig RE. The link between regularity and strong- π -regularity. *Journal of the Australian Mathematical Society* 2010; 89: 17-22. doi: 10.1017/S1446788710001448
- [19] Wang Z. A class of Drazin inverses in rings. *Filomat* 2017; 31: 1781-1789.
- [20] Wang Z, Chen JL. Pseudo Drazin inverses in associative rings and Banach algebras. *Linear Algebra and its Applications* 2012; 437: 1332-1345. doi: 10.1016/j.laa.2012.04.039
- [21] Wei YM, Deng CY. A note on additive results for the Drazin inverse. *Linear Multilinear Algebra* 2011; 59: 1319-1329. doi: 10.1080/03081087.2010.496110
- [22] Ying ZL, Koşan T, Zhou YQ. Rings in which every element is a sum of two tripotents. *Canadian Mathematical Bulletin* 2016; 59: 661-672. doi: 10.4153/CMB-2016-009-0
- [23] Zeng QP, Wu ZW, Wen YX. New extensions of Cline's formula for generalized inverses. *Filomat* 2017; 31: 1973-1980.
- [24] Zeng QP, Zhong HJ. New results on common properties of the products AC and BA . *Journal of Mathematical Analysis and Applications* 2015; 427: 830-840. doi: 10.1016/j.jmaa.2015.02.037
- [25] Zhuang GF, Chen JL, Cui J. Jacobson's lemma for the generalized Drazin inverse. *Linear Algebra and its Applications* 2012; 436: 742-746. doi: 10.1016/j.laa.2011.07.044
- [26] Zhuang GF, Chen JL, Cvetković-Ilić DS, Wei YM. Additive property of Drazin invertibility of elements in a ring. *Linear Multilinear Algebra* 2012; 60: 903-910. doi: 10.1080/03081087.2011.629998
- [27] Zou HL, Chen JL, Li TT, Gao YF. Characterizations and representations of the inverse along an element. *Bulletin of the Malaysian Mathematical Sciences Society* 2018; 41: 1835-1857. doi: 10.1007/s40840-016-0430-3