

1-1-2019

The relation between b-weakly compact operator and KB-operator

BAHRİ TURAN

BİROL ALTIN

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

TURAN, BAHRİ and ALTIN, BİROL (2019) "The relation between b-weakly compact operator and KB-operator," *Turkish Journal of Mathematics*: Vol. 43: No. 6, Article 13. <https://doi.org/10.3906/mat-1908-11>
Available at: <https://dctubitak.researchcommons.org/math/vol43/iss6/13>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

The relation between b -weakly compact operator and KB -operator

Bahri TURAN* , Birol ALTIN 

Department of Mathematics, Faculty of Science, Gazi University, Ankara, Turkey

Received: 05.08.2019

Accepted/Published Online: 01.10.2019

Final Version: 22.11.2019

Abstract: Our aim is to solve the problem asked by Bahramnezhad and Azar in "KB-operators on Banach lattices and their relationships with Dunford-Pettis and order weakly compact operators". We show that a continuous operator R from a Banach lattice N into a Banach space M is a b -weakly compact operator if and only if R is a KB -operator.

Key words: b -weakly compact operator, KB -operator

1. Introduction

In [6], Bahramnezhad and Azar defined a new classes of operators, named KB -operator and they have examined some of their properties and asked the following problem;

Problem 1.1 ([6], Problem 2.27) *Give an operator R from a Banach lattice N into a Banach space M which is a KB -operator but is not b -weakly compact.*

We answer the question in the negative. A lot of properties and results on b -weakly compact operators were given in [2 – –5, 7]. Now, we recall the definitions of b -weakly compact operator and KB -operator.

Definition 1.2 *Let R be a continuous operator from a Banach lattice N into a Banach space M .*

(i) *R is called KB -operator if $R(x_n)$ has a norm convergent subsequence in M for every positive increasing sequence (x_n) of the closed unit ball B_N of N .*

(ii) *R is called b -weakly compact if $R(x_n)$ is norm convergent for every positive increasing sequence (x_n) of the closed unit ball B_N of N .*

For the basic theory on vector lattices and for unexplained terminology we refer to [1, 8].

2. Section

We will prove that the classes of KB -operators and the b -weakly compact operators are the same.

Theorem 2.1 *Let R be an operator from a Banach lattice N into a Banach space M . R is a b -weakly compact operator if and only if R is a KB -operator.*

*Correspondence: bturan@gazi.edu.tr

2010 AMS Mathematics Subject Classification: 46B42, 47B60

Proof It is clear that if R is a b -weakly compact then R is a KB -operator. Let (x_n) be a sequence in N such that $0 \leq x_n \uparrow$ and $\|x_n\| \leq 1$. For an arbitrary subsequence (x_k) of (x_n) , let us define

$$\Psi : N'_+ \rightarrow \mathbb{R}_+, f \rightarrow \Psi(f) = \sup f(x_k).$$

For each k and $f \in N'_+$ we have

$$f(x_k) = |f(x_k)| \leq \|f\| \|x_k\| \leq \|f\|.$$

Then, $\sup f(x_k) \in \mathbb{R}_+$. We claim that Ψ is additive. To see this, let $f, g \in N'_+$.

$$\Psi(f + g) = \sup [(f + g)(x_k)] = \sup [f(x_k) + g(x_k)] \leq \sup f(x_k) + \sup g(x_k) = \Psi(f) + \Psi(g).$$

On the other hand, if $x_m, x_t \in (x_k)$, then pick $x_l \in (x_k)$ with $x_m \leq x_l$ and $x_t \leq x_l$, and note that

$$f(x_m) + g(x_t) \leq f(x_l) + g(x_l) \leq \sup [(f + g)(x_l)] = \Psi(f + g).$$

Using a well-known technique (e.g., [1, p.14]), we have $\Psi(f) + \Psi(g) \leq \Psi(f + g)$. Therefore, Ψ is additive and by Theorem 1.7 in [1] Ψ extends uniquely to a positive operator from N' into \mathbb{R} (we call Ψ again). Hence, $\Psi \in N''$. It is easy to see that Ψ is an upper bound of $(x_k)''$ in N'' (where $(x_k)''$ is the image of (x_k) under the well known canonical emdedding of N into to the bidual N''). There exists G in N'' with $(x_k)'' \uparrow G$ as N'' is Dedekind complete. Since $(x_k)''(f) \rightarrow G(f)$ for each $f \in N'_+$, we have $(x_k)'' \rightarrow G$ with respect to $\sigma(N'', N')$. Thus, all subsequences of (x_n) are convergent to the same limit G with respect to $\sigma(N'', N')$. By the hypothesis, there exists a subsequence (x_{n_k}) of (x_n) and $y \in M$ such that $R(x_{n_k}) \rightarrow y$ with respect to norm-topology. This leads to $[R(x_{n_k})]'' \rightarrow y''$ in M'' with respect to norm-topology which implies $[R(x_{n_k})]'' \rightarrow y''$ in M'' with respect to $\sigma(M'', M')$. The continuity of $R'' : (N'', \sigma(N'', N')) \rightarrow (M'', \sigma(M'', M'))$ yields $R'' [(x_k)'] \rightarrow R''(G)$ and $[R(x_{n_k})]'' \rightarrow y''$ with respect to $\sigma(M'', M')$. Since $R'' [(x_{n_k})'] = [R(x_{n_k})]''$, we have $R''(G) = y''$. So, this means that every norm convergent a subsequence of $R(x_n)$ has the same norm limit. Now, we will show that $R(x_n) \rightarrow y$ in M with respect to norm-topology. We assume that $R(x_n)$ does not convergence to y . Thus, there exist $\varepsilon > 0$ and a subsequence (x_m) of (x_n) such that $\|R(x_m) - y\| > \varepsilon$ for all m . By the hypothesis and the above conclusion there exists a subsequence (x_{m_k}) of (x_m) such that $R(x_{m_k}) \rightarrow y$ with respect to norm-topology, which is a contradiction. \square

References

- [1] Aliprantis CD, Burkinshaw O. Positive Operators. Berlin, Germany: Springer, 2006.
- [2] Alpay Ş, Altın B, Tonyalı C. On property (b) of vector lattices. Positivity 2003; 7: 135-139. doi: 10.1023/A:1025840528211
- [3] Altın B. On b -weakly compact operators on Banach lattices. Taiwanese Journal of Mathematics 2007; 11: 143-150. doi: 10.11650/twjmath/1500404641
- [4] Altın B. Some properties of b -weakly compact operators. Gazi Üniversitesi Fen Bilimleri Dergisi 2005; 391-395.
- [5] Aqzzouz B, Moussa M, Hmichane J. Some Characterizations of b -weakly compact operators on Banach lattices. Mathematical Reports 2010; 62: 315-324.
- [6] Bahramnezhad A, Azar KH. KB -operators on Banach lattices and their relationships with Dunford-Pettis and order weakly compact operators. University Politehnica of Bucharest Scientific Bulletin 2018; 80: 91-98.

- [7] Cheng N, Chen ZL. Some properties of b-weakly compact operators on Banach lattice. World Academy of Science, Engineering and Technology International Journal of Mathematical Computational Physical Electrical and Computer Engineering 2010; 4: No 3.
- [8] Meyer-Nieberg P. Banach Lattices. Berlin, Germany: Springer, 1991.