

1-1-2019

A circulant functional equation for the additive function and its stability

VICHIAN LAOHAKOSOL

WATCHARAPON PIMSERT

KANET PONPETCH

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

LAOHAKOSOL, VICHIAN; PIMSERT, WATCHARAPON; and PONPETCH, KANET (2019) "A circulant functional equation for the additive function and its stability," *Turkish Journal of Mathematics*: Vol. 43: No. 6, Article 14. <https://doi.org/10.3906/mat-1906-88>

Available at: <https://journals.tubitak.gov.tr/math/vol43/iss6/14>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

A circulant functional equation for the additive function and its stability

Vichian LAOHAKOSOL¹, Watcharapon PIMSERT¹, Kanet PONPETCH^{2*}

¹Department of Mathematics, Faculty of Science, Kasetsart University, Bangkok, Thailand

² 27 Moo 2, Khlong Yai Sub-district, Tamod District, Phatthalung, Thailand

Received: 26.06.2019

Accepted/Published Online: 04.10.2019

Final Version: 22.11.2019

Abstract: A general solution of a matrix functional equation involving circulant matrices of the additive function is determined, and its stability is established.

Key words: Circulant, additive function, stability

1. Introduction

The classical additive functional equation

$$f(x + y) = f(x) + f(y) \quad (\text{A})$$

was first studied by Cauchy in 1821 for continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$; the solution takes the form $f(x) = ax$ for some $a \in \mathbb{R}$ (see e.g. [3, Chapter 1]). This result of Cauchy has been extended in various directions. In particular, for $f : \mathbb{C} \rightarrow \mathbb{C}$, we have the following result taken from the book [3, Theorem 1.21]: the general solution $f : \mathbb{C} \rightarrow \mathbb{C}$ of (A) is given by $f(z) = A_1(x) + iA_2(x) + A_3(y) + iA_4(y)$, where $z = x + iy$ ($x, y \in \mathbb{R}$), and $A_j : \mathbb{R} \rightarrow \mathbb{R}$ ($j = 1, 2, 3, 4$) are real additive functions, i.e. satisfying (A). In 1903, Pexider solved the functional equation

$$f(x + y) = g(x) + h(y), \quad (\text{PA})$$

referred to as the pexiderized additive functional equation, for unknown functions $f, g, h : \mathbb{C} \rightarrow \mathbb{C}$, and found that the solutions of (PA) are given by $f(x) = a + A(x) + b$, $g(x) = a + A(x)$, $h(x) = A(x) + b$, where $A : \mathbb{C} \rightarrow \mathbb{C}$ is a complex additive function, and a, b are complex constants (see [3, Chapter 1]).

The classical concepts of odd and even functions have been extended to that of type- j functions by Schwaiger in [6]. These are components f_j of a function f defined by

$$f_j(x) = \frac{1}{n} \sum_{k=0}^{n-1} \omega^{-kj} f(\omega^k x) \quad (j = 0, 1, \dots, n-1)$$

and satisfy $f_j(\omega x) = \omega^j f_j(x)$, $f = \sum_{j=0}^{n-1} f_j$, where $\omega := \exp(2\pi i/n)$ is a primitive n^{th} root of unity. Schwaiger adopted this concept and solved the following system of functional equations satisfied by components of the

*Correspondence: kanet.bkp@gmail.com

2010 AMS Mathematics Subject Classification: 39A10, 39A11, 39A99

exponential function

$$f_j(x + \omega^m y) = \sum_{\ell=0}^j \omega^{(j-\ell)m} f_\ell(x) f_{j-\ell}(y) + \sum_{\ell=j+1}^{n-1} \omega^{(n+j-\ell)m} f_\ell(x) f_{n+j-\ell}(y) \quad (j = 0, 1, \dots, n-1) \quad (1.1)$$

where $m \in \{0, 1, \dots, n-1\}$ is fixed. The stability of the system (1.1) was established one year later by Förg-Rob and Schwaiger in [2]. The results in [6] and [2] were simplified and systematized by Muldoon [4] through the use of circulant matrices. In [5], Laohakosol and Ponpetch used Muldoon’s approach [4] to solve a circulant functional equation for the quadratic function, i.e., a system of functional equations satisfied by components of the quadratic function derived via their corresponding circulant matrices, and established its stability.

In this work, through the concept of type- j function, we derive a circulant matrix functional equation for the additive function. Given such a circulant matrix functional equation, their component solutions are determined, and the stability of such system is established.

2. Preliminaries

This section consists of two parts. The first part lays out the notation and terminology that will be kept fixed throughout, while the second part lists (without proofs) basic results needed later.

2.1. Notation

For a fixed integer $n \geq 2$, let $\omega = \exp(2\pi i/n)$ be a primitive n^{th} root of unity. The $n \times n$ (symmetric) Fourier matrix and its conjugate matrix are defined, respectively, by

$$\mathcal{F} := \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega^{-1} & \omega^{-2} & \dots & \omega^{-(n-1)} \\ 1 & \omega^{-2} & \omega^{-4} & \dots & \omega^{-2(n-1)} \\ \vdots & \vdots & \ddots & & \vdots \\ 1 & \omega^{-(n-1)} & \omega^{-2(n-1)} & \dots & \omega^{-(n-1)^2} \end{bmatrix}, \quad \mathcal{F}^* := \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega & \omega^2 & \dots & \omega^{n-1} \\ 1 & \omega^2 & \omega^4 & \dots & \omega^{2(n-1)} \\ \vdots & \vdots & \ddots & & \vdots \\ 1 & \omega^{n-1} & \omega^{2(n-1)} & \dots & \omega^{(n-1)^2} \end{bmatrix}.$$

Note that \mathcal{F} is unitary, i.e., $\mathcal{F} \mathcal{F}^* = I = \mathcal{F}^* \mathcal{F}$, where I denotes the $n \times n$ identity matrix.

The diagonal matrix Ω is defined by $\Omega = \text{diag}(1, \omega, \omega^2, \dots, \omega^{n-1}) := \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & \omega & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \omega^{n-1} \end{bmatrix}$.

Given a sequence $\{a_0, \dots, a_{n-1}\} \subset \mathbb{C}$, its *circulant matrix* is defined by

$$\text{circ}(a_0, a_1, \dots, a_{n-1}) := \begin{bmatrix} a_0 & a_1 & \dots & a_{n-1} \\ a_{n-1} & a_0 & \dots & a_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_1 & a_2 & \dots & a_0 \end{bmatrix};$$

its diagonal matrix is defined by $\text{diag}(a_0, a_1, \dots, a_{n-1}) := \begin{bmatrix} a_0 & 0 & \cdots & 0 \\ 0 & a_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{n-1} \end{bmatrix}$.

The circulant matrix corresponding to the sequence $\{0, 1, 0, \dots, 0\}$ is

$$\pi := \text{circ}(0, 1, 0, \dots, 0) = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 1 & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

The circulant matrix corresponding to a function f , with j -component f_j , is defined by

$$\mathbf{F}(x) := \text{circ}(f_0(x), f_1(x), \dots, f_{n-1}(x)) = \begin{bmatrix} f_0(x) & f_1(x) & \cdots & f_{n-1}(x) \\ f_{n-1}(x) & f_0(x) & \cdots & f_{n-2}(x) \\ \vdots & \vdots & \ddots & \vdots \\ f_1(x) & f_2(x) & \cdots & f_0(x) \end{bmatrix}.$$

2.2. Basic results

The following results are taken from [4, 5].

- 1) If $A = \text{circ}(a_0, a_1, \dots, a_{n-1})$, then $\mathcal{F}A\mathcal{F}^* = \sqrt{n} \text{diag}(\mathcal{F}^* \bar{a})^T$, $\bar{a} := [a_0, \dots, a_{n-1}]^T$, T denoting transpose.
- 2) For a nonnegative integer m , if A is a circulant matrix, then $\mathcal{F}(\Omega^{-m}A\Omega^m)\mathcal{F}^* = \pi^m(\mathcal{F}A\mathcal{F}^*)\pi^{-m}$.
- 3) Any $f : \mathbb{C} \rightarrow \mathbb{C}$ can be written uniquely as $f(x) = f_0(x) + f_1(x) + \dots + f_{n-1}(x)$, where each j -component f_j ($j = 0, 1, \dots, n - 1$) is a type- j function that can be obtained from

$$\begin{bmatrix} f_0(x) \\ f_1(x) \\ \vdots \\ f_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} f(x) \\ f(\omega x) \\ \vdots \\ f(\omega^{n-1}x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F}^* \begin{bmatrix} f(x) \\ f(\omega^{-1}x) \\ \vdots \\ f(\omega^{-(n-1)}x) \end{bmatrix}.$$

- 4) The circulant matrix $\mathbf{F}(x)$ corresponding to $f : \mathbb{C} \rightarrow \mathbb{C}$ satisfies

$$\mathcal{F}\mathbf{F}(x)\mathcal{F}^* = \text{diag}(f(x), f(\omega x), \dots, f(\omega^{n-1}x)) \tag{2.1}$$

$$\mathbf{F}(\omega^m x) = \Omega^{-m}\mathbf{F}(x)\Omega^m \quad (m \in \mathbb{N}). \tag{2.2}$$

- 5) For a nonnegative integer m , if $B = \text{diag}(b_0, b_1, \dots, b_{n-1})$, then $\pi^m B \pi^{-m} = \text{diag}(b_m, b_{m+1}, \dots, b_{m+n-1})$, where suffixes are taken modulo n .
- 6) If $B = \text{diag}(b_0, b_1, \dots, b_{n-1})$ is a diagonal matrix, then $\mathcal{F}^*B\mathcal{F}$ is a circulant matrix, i.e. $\mathcal{F}^*B\mathcal{F} = \text{circ}(d_0, d_1, \dots, d_{n-1})$, where $d_j = (1/n) \sum_{k=0}^{n-1} \omega^{n-kj} b_k$ ($j = 0, 1, \dots, n - 1$).
- 7) For $m \in \{0, 1, \dots, n - 1\}$, $d = \text{gcd}(n, m)$, we have $s + tm \not\equiv u + vm \pmod{n}$, for every $s, u \in \{0, 1, \dots, d - 1\}$ and $t, v \in \{0, 1, \dots, n/d - 1\}$, except when $s = u, t = v$.

3. A circulant functional equation

We begin by deriving a circulant equation for the additive function.

Theorem 3.1 *If $f, g, h : \mathbb{C} \rightarrow \mathbb{C}$ satisfy (PA), then their circulant matrices $\mathbf{F}, \mathbf{G}, \mathbf{H}$ satisfy*

$$\mathbf{F}(x + y) = \mathbf{G}(x) + \mathbf{H}(y) \quad (x, y \in \mathbb{C}).$$

More generally, we have

$$\mathbf{F}(x + \omega^m y) = \mathbf{G}(x) + \Omega^{-m} \mathbf{H}(y) \Omega^m \quad (x, y \in \mathbb{C}; m \in \{0, \dots, n - 1\}). \tag{3.1}$$

Proof From the basic result (2.1) of property 4) in the last section, we have

$$\begin{aligned} \mathbf{G}(x) + \mathbf{H}(y) &= \mathcal{F}^* \text{diag}(g(x), g(\omega x), \dots, g(\omega^{n-1}x)) \mathcal{F} + \mathcal{F}^* \text{diag}(h(y), h(\omega y), \dots, h(\omega^{n-1}y)) \mathcal{F} \\ &= \mathcal{F}^* \text{diag}(g(x) + h(y), g(\omega x) + h(\omega y), \dots, g(\omega^{n-1}x) + h(\omega^{n-1}y)) \mathcal{F} \\ &= \mathcal{F}^* \text{diag}(f(x + y), f(\omega(x + y)), \dots, f(\omega^{n-1}(x + y))) \mathcal{F} = \mathbf{F}(x + y). \end{aligned}$$

Using this last relation together with the basic result (2.2) of property 4), the last assertion is immediate. \square

Our next step is to solve the circulant equation (3.1)

Theorem 3.2 *Let $\mathbf{F}, \mathbf{G}, \mathbf{H}$ be circulant matrices whose first rows are, respectively,*

$$(f_0(x), f_1(x), \dots, f_{n-1}(x)), (g_0(x), g_1(x), \dots, g_{n-1}(x)), (h_0(x), h_1(x), \dots, h_{n-1}(x)),$$

where $f_i, g_i, h_i : \mathbb{C} \rightarrow \mathbb{C}$ are arbitrary functions which need not be components of any functions. Let $m \in \{0, 1, \dots, n - 1\}$, $d := \text{gcd}(n, m)$. If $\mathbf{F}, \mathbf{G}, \mathbf{H}$ satisfy

$$\mathbf{F}(x + \omega^m y) = \mathbf{G}(x) + \Omega^{-m} \mathbf{H}(y) \Omega^m, \tag{3.2}$$

then

- when $d = n$, we have

$$\begin{bmatrix} f_0(x) \\ f_1(x) \\ \vdots \\ f_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} A_0(x) + w_0(0) + v_0(0) \\ A_1(x) + w_1(0) + v_1(0) \\ \vdots \\ A_{n-1}(x) + w_{n-1}(0) + v_{n-1}(0) \end{bmatrix} \tag{3.3}$$

$$\begin{bmatrix} g_0(x) \\ g_1(x) \\ \vdots \\ g_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} A_0(x) + v_0(0) \\ A_1(x) + v_1(0) \\ \vdots \\ A_{n-1}(x) + v_{n-1}(0) \end{bmatrix} \tag{3.4}$$

$$\begin{bmatrix} h_0(x) \\ h_1(x) \\ \vdots \\ h_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} A_0(x) + w_0(0) \\ A_1(x) + w_1(0) \\ \vdots \\ A_{n-1}(x) + w_{n-1}(0) \end{bmatrix}, \tag{3.5}$$

where $A_i : \mathbb{C} \rightarrow \mathbb{C}$ ($i = 0, 1, \dots, n - 1$) are additive functions and

$$v_i(x) := \sum_{\ell=0}^{n-1} \omega^{i\ell} g_\ell(x), \quad w_i(x) := \sum_{\ell=0}^{n-1} \omega^{i\ell} h_\ell(x) \quad (i = 0, \dots, n - 1); \tag{3.6}$$

• when $1 \leq d < n$, we have

$$\begin{bmatrix} f_0(x) \\ f_1(x) \\ \vdots \\ f_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} \alpha_0(x) + C_m(0) + B_0(0) \\ \alpha_m(x) + C_{2m}(0) + B_m(0) \\ \vdots \\ \alpha_{(n/d-1)m}(x) + C_{(n/d)m}(0) + B_{(n/d-1)m}(0) \end{bmatrix} \tag{3.7}$$

$$\begin{bmatrix} g_0(x) \\ g_1(x) \\ \vdots \\ g_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} \alpha_0(x) + B_0(0) \\ \alpha_m(x) + B_m(0) \\ \vdots \\ \alpha_{(n/d-1)m}(x) + B_{(n/d-1)m}(0) \end{bmatrix} \tag{3.8}$$

$$\begin{bmatrix} h_0(x) \\ h_1(x) \\ \vdots \\ h_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} \alpha_0(\omega^m x) + C_m(0) \\ \alpha_m(\omega^m x) + C_{2m}(0) \\ \vdots \\ \alpha_{(n/d-1)m}(\omega^m x) + C_{(n/d)m}(0) \end{bmatrix} \tag{3.9}$$

where, for $j = 0, 1, \dots, n/d - 1$,

$$\alpha_{jm}(x) = \begin{bmatrix} A_{0+jm}(x) \\ A_{1+jm}(x) \\ \vdots \\ A_{d-1+jm}(x) \end{bmatrix}, \quad B_{jm}(0) = \begin{bmatrix} v_{0+jm}(0) \\ v_{1+jm}(0) \\ \vdots \\ v_{d-1+jm}(0) \end{bmatrix}, \quad C_{(j+1)m}(0) = \begin{bmatrix} w_{0+(j+1)m}(0) \\ w_{1+(j+1)m}(0) \\ \vdots \\ w_{d-1+(j+1)m}(0) \end{bmatrix}$$

and $A_{k+jm} : \mathbb{C} \rightarrow \mathbb{C}$ ($k = 0, 1, \dots, d - 1$) are additive functions.

Proof From (3.2), we have

$$\mathcal{F}\mathbf{F}(x + \omega^m y)\mathcal{F}^* = \mathcal{F}\mathbf{G}(x)\mathcal{F}^* + \mathcal{F}\Omega^{-m}\mathbf{H}(y)\Omega^m\mathcal{F}^*.$$

Using the basic results 1) and 2), this relation takes the form

$$\mathbf{U}(x + \omega^m y) = \mathbf{V}(x) + \mathbf{W}_m(y), \tag{3.10}$$

where

$$\text{diag}(u_0(x), u_1(x), \dots, u_{n-1}(x)) = \mathbf{U}(x) = \mathcal{F}\mathbf{F}(x)\mathcal{F}^* = \sqrt{n} \text{diag}(\mathcal{F}^* \bar{f}(x))^T \tag{3.11}$$

$$\text{diag}(v_0(x), v_1(x), \dots, v_{n-1}(x)) = \mathbf{V}(x) = \mathcal{F}\mathbf{G}(x)\mathcal{F}^* = \sqrt{n} \text{diag}(\mathcal{F}^* \bar{g}(x))^T \tag{3.12}$$

$$\text{diag}(w_0(x), w_1(x), \dots, w_{n-1}(x)) = \mathbf{W}(x) = \mathcal{F}\mathbf{H}(x)\mathcal{F}^* = \sqrt{n} \text{diag}(\mathcal{F}^* \bar{h}(x))^T \tag{3.13}$$

$$\mathbf{W}_m(x) = \pi^m \mathbf{W}(x) \pi^{-m}. \tag{3.14}$$

Equation (3.10) and the basic result 5) yield the following system of n equations

$$\begin{aligned} u_0(x + \omega^m y) &= v_0(x) + w_m(y) \\ &\vdots \\ u_{n-1}(x + \omega^m y) &= v_{n-1}(x) + w_{m+n-1}(y). \end{aligned}$$

Using the basic result 7), we subdivide these n equations into d different classes, each with n/d equations, briefly written as

$$u_{k+jm}(x + \omega^m y) = v_{k+jm}(x) + w_{k+(j+1)m}(y) \quad (j = 0, 1, \dots, n/d - 1; k = 0, 1, \dots, d - 1). \tag{3.15}$$

For each $j \in \{0, 1, \dots, n/d - 1\}$ and $k \in \{0, 1, \dots, d - 1\}$, replacing $y = 0$ in (3.15), we have

$$u_{k+jm}(x) = v_{k+jm}(x) + w_{k+(j+1)m}(0), \tag{3.16}$$

while replacing $x = 0$ in (3.15), we have

$$u_{k+jm}(\omega^m y) = v_{k+jm}(0) + w_{k+(j+1)m}(y). \tag{3.17}$$

Substituting (3.16) and (3.17) into (3.15), we get

$$u_{k+jm}(x + \omega^m y) = u_{k+jm}(x) - w_{k+(j+1)m}(0) + u_{k+jm}(\omega^m y) - v_{k+jm}(0).$$

Replacing y by $\omega^{-m}y$ in the last equation and adding $-w_{k+(j+1)m}(0) - v_{k+jm}(0)$ to both sides, we obtain

$$\begin{aligned} &u_{k+jm}(x + y) - w_{k+(j+1)m}(0) - v_{k+jm}(0) \\ &= u_{k+jm}(x) - w_{k+(j+1)m}(0) - v_{k+jm}(0) + u_{k+jm}(y) - w_{k+(j+1)m}(0) - v_{k+jm}(0). \end{aligned} \tag{3.18}$$

Defining

$$A_{k+jm}(x) := u_{k+jm}(x) - w_{k+(j+1)m}(0) - v_{k+jm}(0) \quad (j \in \{0, 1, \dots, n/d - 1\}, k \in \{0, 1, \dots, d - 1\}),$$

the relation (3.18) shows that it is an additive function. Substituting back into (3.16) and (3.17), we have, for $j = 0, 1, \dots, n/d - 1, k = 0, 1, \dots, d - 1,$

$$u_{k+jm}(x) = A_{k+jm}(x) + w_{k+(j+1)m}(0) + v_{k+jm}(0), \tag{3.19}$$

$$v_{k+jm}(x) = A_{k+jm}(x) + v_{k+jm}(0), \tag{3.20}$$

$$w_{k+(j+1)m}(x) = A_{k+jm}(\omega^m x) + w_{k+(j+1)m}(0), \tag{3.21}$$

where A_{k+jm} are additive functions. If $d = n$, then $m = 0$ and from (3.19), (3.20), and (3.21), we get, for $k = 0, 1, \dots, n - 1,$

$$u_k(x) = A_k(x) + w_k(0) + v_k(0), \quad v_k(x) = A_k(x) + v_k(0), \quad w_k(x) = A_k(x) + w_k(0). \tag{3.22}$$

From (3.11), (3.12), (3.13), and (3.22), we get the three solution matrices (3.3), (3.4), (3.5).

If $1 \leq d < n$, then using (3.11)–(3.13), (3.19)–(3.21), we similarly get the three solution matrices (3.7)–(3.9). □

Remark. If the functions f_i, g_i, h_i ($i = 0, 1, \dots, n - 1$) in Theorem 3.2 are identical, then (3.2) becomes

$$\mathbf{F}(x + \omega^m y) = \mathbf{F}(x) + \Omega^{-m} \mathbf{F}(y) \Omega^m.$$

The solutions thus take the form:

- when $d = n$,

$$\begin{bmatrix} f_0(x) \\ f_1(x) \\ \vdots \\ f_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} A_0(x) \\ A_1(x) \\ \vdots \\ A_{n-1}(x) \end{bmatrix} \tag{3.23}$$

where $A_i : \mathbb{C} \rightarrow \mathbb{C}$ ($i = 0, 1, \dots, n - 1$) are additive functions;

- when $1 \leq d < n$,

$$\begin{bmatrix} f_0(x) \\ f_1(x) \\ \vdots \\ f_{n-1}(x) \end{bmatrix} = \frac{1}{\sqrt{n}} \mathcal{F} \begin{bmatrix} \alpha(x) \\ \alpha(\omega^m x) \\ \vdots \\ \alpha(\omega^{(n/d-1)m} x) \end{bmatrix} \tag{3.24}$$

where $\alpha(x) = [A_0(x), A_1(x), \dots, A_{d-1}(x)]^T$ and $A_k : \mathbb{C} \rightarrow \mathbb{C}$ ($k = 0, 1, \dots, d - 1$) are additive functions.

4. Stability

In this section, we establish the stability of the circulant equation (3.1). As in [4], we use the usual 1-norm for a square matrix $A = (a_{i,j})_{0 \leq i,j \leq n-1}$ defined by

$$\|A\| = \max_{0 \leq i \leq n-1} \sum_{j=0}^{n-1} |a_{i,j}|.$$

Throughout this section, let $\phi : \mathbb{C} \times \mathbb{C} \rightarrow [0, \infty)$ be a mapping satisfying the condition that the sum $\sum_{\ell=0}^{\infty} t^{-(\ell+1)} (\phi(t^\ell x, t^\ell y) + \phi(t^\ell x, 0) + \phi(0, t^\ell y))$ converges for all $x, y \in \mathbb{C}$ and for all integers $t \geq 2$.

Theorem 4.1 *Let $\mathbf{F}, \mathbf{G}, \mathbf{H}$ be circulant matrices whose first rows are, respectively,*

$$(f_0(x), f_1(x), \dots, f_{n-1}(x)), (g_0(x), g_1(x), \dots, g_{n-1}(x)), (h_0(x), h_1(x), \dots, h_{n-1}(x)),$$

where $f_i, g_i, h_i : \mathbb{C} \rightarrow \mathbb{C}$ are arbitrary functions which need not be components of any functions, let $m \in \{0, 1, \dots, n - 1\}$. If $\mathbf{F}, \mathbf{G}, \mathbf{H}$ satisfy

$$\|\mathbf{F}(x + \omega^m y) - \mathbf{G}(x) - \Omega^{-m} \mathbf{H}(y) \Omega^m\| \leq \phi(x, y), \tag{4.1}$$

then there exists a circulant matrix $\mathbf{A}(x) := \text{circ}(a_0(x), a_1(x), \dots, a_{n-1}(x))$, $a_j : \mathbb{C} \rightarrow \mathbb{C}$ ($j = 0, 1, \dots, n - 1$) satisfying the matrix additive equation

$$\mathbf{A}(x + y) = \mathbf{A}(x) + \mathbf{A}(y) \tag{4.2}$$

and

$$\|\mathbf{F}(x) - \mathbf{A}(x)\| \leq n \sum_{\ell=1}^{t-1} \Phi_t(x, \ell x) \tag{4.3}$$

$$\|\mathbf{G}(x) - \mathbf{A}(x)\| \leq n^2\phi(x, 0) + n \sum_{\ell=1}^{t-1} \Phi_t(x, \ell x) + \|\mathbf{H}(0)\| \tag{4.4}$$

$$\|\mathbf{H}(x) - \bar{\mathbf{A}}(x)\| \leq n^2\phi(0, x) + n \sum_{\ell=1}^{t-1} \Phi_t(\omega^m x, \ell \omega^m x) + \|\mathbf{G}(0)\|, \tag{4.5}$$

where

$$\Phi_t(x, y) := \sum_{\ell=0}^{\infty} t^{-(\ell+1)} \{n(\phi(t^\ell x, t^\ell \omega^{-m} y) + \phi(t^\ell x, 0) + \phi(0, t^\ell \omega^{-m} y)) + \|\mathbf{V}(0)\| + \|\mathbf{W}_m(0)\|\}$$

with $\mathbf{V}(0) = \text{diag}(v_0(0), v_1(0), \dots, v_{n-1}(0))$, $\mathbf{W}_m(0) = \text{diag}(w_m(0), w_{m+1}(0), \dots, w_{m+n-1}(0))$, v_i, w_i are as defined in (3.6) with suffixes taken modulo n , and $\bar{\mathbf{A}}(x) := \Omega^m \mathbf{A}(\omega^m x) \Omega^{-m}$ is a circulant matrix satisfying (4.2), and whose first row is $(\bar{a}_0(x), \bar{a}_1(x), \dots, \bar{a}_{n-1}(x))$, $\bar{a}_j(x) = \omega^{(n-j)m} a_j(\omega^m x)$ ($j = 0, 1, \dots, n-1$).

Proof Multiplying the expressions in (4.1) by $\|\mathcal{F}\|$ on the left side and by $\|\mathcal{F}^*\|$ on the right side, we obtain

$$\|\mathcal{F}\mathbf{F}(x + \omega^m y)\mathcal{F}^* - \mathcal{F}\mathbf{G}(x)\mathcal{F}^* - \mathcal{F}\Omega^{-m}\mathbf{H}(y)\Omega^m\mathcal{F}^*\| \leq \|\mathcal{F}\|\phi(x, y)\|\mathcal{F}^*\|.$$

By the basic results 1) and 2), this last inequality is of the form

$$\|\mathbf{U}(x + \omega^m y) - \mathbf{V}(x) - \mathbf{W}_m(y)\| \leq n\phi(x, y), \tag{4.6}$$

where U, V, W, W_m are as defined in (3.11)–(3.14). Substituting $y = 0$ into (4.6) we have

$$\|\mathbf{V}(x) + \mathbf{W}_m(0) - \mathbf{U}(x)\| \leq n\phi(x, 0), \tag{4.7}$$

and similarly substituting $x = 0$ into (4.6), we have

$$\|\mathbf{V}(0) + \mathbf{W}_m(y) - \mathbf{U}(\omega^m y)\| \leq n\phi(0, y). \tag{4.8}$$

Using (4.6)–(4.8), we get

$$\begin{aligned} \|\mathbf{U}(x + \omega^m y) - \mathbf{U}(x) - \mathbf{U}(\omega^m y)\| - \|\mathbf{V}(0) + \mathbf{W}_m(0)\| &\leq \|\mathbf{U}(x + \omega^m y) - \mathbf{U}(x) - \mathbf{U}(\omega^m y) + \mathbf{V}(0) + \mathbf{W}_m(0)\| \\ &\leq n(\phi(x, y) + \phi(x, 0) + \phi(0, y)). \end{aligned}$$

Replacing y by $\omega^{-m}y$, we get

$$\|\mathbf{U}(x + y) - \mathbf{U}(x) - \mathbf{U}(y)\| \leq n(\phi(x, \omega^{-m}y) + \phi(x, 0) + \phi(0, \omega^{-m}y)) + \|\mathbf{V}(0)\| + \|\mathbf{W}_m(0)\|.$$

Since $\mathbf{U}(x) = \text{diag}(u_0(x), u_1(x), \dots, u_{n-1}(x)) = (u_{i,j}(x))_{0 \leq i,j \leq n-1}$, where $u_{i,j}(x) = \begin{cases} u_i(x) & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$, by the

definition of norm, we get

$$\max_{0 \leq i \leq n-1} \sum_{j=0}^{n-1} |u_{i,j}(x + y) - u_{i,j}(x) - u_{i,j}(y)| \leq n(\phi(x, \omega^{-m}y) + \phi(x, 0) + \phi(0, \omega^{-m}y)) + \|\mathbf{V}(0)\| + \|\mathbf{W}_m(0)\|.$$

For each $i = 0, 1, \dots, n - 1$, we have

$$|u_i(x + y) - u_i(x) - u_i(y)| \leq n (\phi(x, \omega^{-m}y) + \phi(x, 0) + \phi(0, \omega^{-m}y)) + \|\mathbf{V}(0)\| + \|\mathbf{W}_m(0)\|.$$

By [1, Theorem 13.3, p. 131], there exist unique additive functions $s_i : \mathbb{C} \rightarrow \mathbb{C}$ satisfying (A) such that

$$|u_i(x) - s_i(x)| \leq \sum_{\ell=1}^{t-1} \Phi_t(x, \ell x) \quad (i = 0, 1, \dots, n - 1).$$

Let $\mathbf{S}(x) := \text{diag}(s_0(x), s_1(x), \dots, s_{n-1}(x)) = (s_{i,j}(x))_{0 \leq i,j \leq n-1}$, where $s_{i,j}(x) = \begin{cases} s_i(x) & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$. Using the definition of norm, we have

$$\|\mathbf{U}(x) - \mathbf{S}(x)\| = \max_{0 \leq i \leq n-1} \sum_{j=0}^{n-1} |u_{i,j}(x) - s_{i,j}(x)| \leq \sum_{\ell=1}^{t-1} \Phi_t(x, \ell x). \tag{4.9}$$

Multiplying the expressions in (4.9) by $\|\mathcal{F}^*\|$ on the left side and by $\|\mathcal{F}\|$ on the right side, and using (3.11), we get (4.3), where $\mathbf{A}(x) = \mathcal{F}^* \mathbf{S}(x) \mathcal{F}$. Since $\mathbf{S}(x)$ is a diagonal matrix, the basic result 6) shows that $\mathbf{A}(x)$ is a circulant matrix whose first row is $(a_0(x), a_1(x), \dots, a_{n-1}(x))$, where

$$a_j(x) = \frac{1}{n} \sum_{k=0}^{n-1} \omega^{n-kj} s_k(x) \quad (j = 0, 1, \dots, n - 1).$$

Since each s_k satisfies (A), the function elements a_j satisfy (A), i.e. $\mathbf{A}(x)$ satisfies (4.2).

Multiplying the expressions in (4.7) by $\|\mathcal{F}^*\|$ on the left side and by $\|\mathcal{F}\|$ on the right side, and using (3.11)–(3.13), and the basic result 2), we get

$$\begin{aligned} \|\mathbf{G}(x) - \mathbf{A}(x)\| - \|\mathbf{F}(x) - \mathbf{A}(x)\| - \|\Omega^{-m} \mathbf{H}(0) \Omega^m\| &\leq \|\mathbf{G}(x) + \Omega^{-m} \mathbf{H}(0) \Omega^m - \mathbf{F}(x) + \mathbf{A}(x) - \mathbf{A}(x)\| \\ &\leq n^2 \phi(x, 0). \end{aligned}$$

and using (4.3) the assertion (4.4) follows.

Multiplying the expressions in (4.8) by $\|\mathcal{F}^*\|$ on the left side and by $\|\mathcal{F}\|$ on the right side, and using (3.11)–(3.13), and the basic result 2), we get

$$\begin{aligned} \|\Omega^{-m} \mathbf{H}(y) \Omega^m - \mathbf{A}(\omega^m y)\| - \|\mathbf{F}(\omega^m y) - \mathbf{A}(\omega^m y)\| - \|\mathbf{G}(0)\| \\ \leq \|\mathbf{G}(0) + \Omega^{-m} \mathbf{H}(y) \Omega^m - \mathbf{F}(\omega^m y) + \mathbf{A}(\omega^m y) - \mathbf{A}(\omega^m y)\| \leq n^2 \phi(0, y). \end{aligned}$$

Using (4.3), we get

$$\|\Omega^{-m} \mathbf{H}(y) \Omega^m - \mathbf{A}(\omega^m y)\| \leq n^2 \phi(0, y) + n \sum_{m=1}^{t-1} \Phi_t(\omega^m y, \ell \omega^m y) + \|\mathbf{G}(0)\|. \tag{4.10}$$

Multiplying the expressions in (4.10) by $\|\Omega^m\|$ on the left side and by $\|\Omega^{-m}\|$ on the right side, we obtain (4.5). □

Taking $\phi(x, y) = \varepsilon$ in Theorem 4.1, we get the following

Corollary 4.2 *Let $\varepsilon > 0$ be fixed. If $\mathbf{F}, \mathbf{G}, \mathbf{H}$ satisfy*

$$\|\mathbf{F}(x + \omega^m y) - \mathbf{G}(x) - \Omega^{-m} \mathbf{H}(y) \Omega^m\| \leq \varepsilon, \tag{4.11}$$

then there exists a circulant matrix \mathbf{A} satisfying (4.2) and

$$\|\mathbf{F}(x) - \mathbf{A}(x)\| \leq 3n^2\varepsilon + n\|\mathbf{V}(0)\| + n\|\mathbf{W}_m(0)\| \tag{4.12}$$

$$\|\mathbf{G}(x) - \mathbf{A}(x)\| \leq 4n^2\varepsilon + n\|\mathbf{V}(0)\| + n\|\mathbf{W}_m(0)\| + \|\mathbf{H}(0)\| \tag{4.13}$$

$$\|\mathbf{H}(x) - \bar{\mathbf{A}}(x)\| \leq 4n^2\varepsilon + n\|\mathbf{V}(0)\| + n\|\mathbf{W}_m(0)\| + \|\mathbf{G}(0)\|. \tag{4.14}$$

5. Examples

We end the paper by working out two examples, first for Theorem 3.2, and second for Theorem 4.1.

Example 1. For $n = 3$, the matrix equation (3.2) yields the following three systems of functional equations, corresponding to the three possible values of $m \in \{0, 1, 2\}$,

- $m = 0$; $f_0(x + y) = g_0(x) + h_0(y)$, $f_1(x + y) = g_1(x) + h_1(y)$, $f_2(x + y) = g_2(x) + h_2(y)$
- $m = 1$; $f_0(x + \omega y) = g_0(x) + h_0(y)$, $f_1(x + \omega y) = g_1(x) + \omega h_1(y)$, $f_2(x + \omega y) = g_2(x) + \omega^2 h_2(y)$
- $m = 2$; $f_0(x + \omega^2 y) = g_0(x) + h_0(y)$, $f_1(x + \omega^2 y) = g_1(x) + \omega^2 h_1(y)$, $f_2(x + \omega^2 y) = g_2(x) + \omega h_2(y)$.

(Observe that these systems are equivalent in the sense that each one of them can be transformed to (PA) by variable changing.)

In any case, $d = \gcd(3, m) = 1$ and the solution functions take the form

$$\begin{aligned} \begin{bmatrix} f_0(x) \\ f_1(x) \\ f_2(x) \end{bmatrix} &= \frac{1}{\sqrt{3}} \mathcal{F} \begin{bmatrix} \alpha_0(x) + C_m(0) + B_0(0) \\ \alpha_m(x) + C_{2m}(0) + B_m(0) \\ \alpha_{2m}(x) + C_{3m}(0) + B_{2m}(0) \end{bmatrix} \\ \begin{bmatrix} g_0(x) \\ g_1(x) \\ g_2(x) \end{bmatrix} &= \frac{1}{\sqrt{3}} \mathcal{F} \begin{bmatrix} \alpha_0(x) + B_0(0) \\ \alpha_m(x) + B_m(0) \\ \alpha_{2m}(x) + B_{2m}(0) \end{bmatrix} \\ \begin{bmatrix} h_0(x) \\ h_1(x) \\ h_2(x) \end{bmatrix} &= \frac{1}{\sqrt{3}} \mathcal{F} \begin{bmatrix} \alpha_0(\omega^m x) + C_m(0) \\ \alpha_m(\omega^m x) + C_{2m}(0) \\ \alpha_{2m}(\omega^m x) + C_{3m}(0) \end{bmatrix} \end{aligned}$$

where $\alpha_{jm}(x) = A_{jm}(x)$, $B_{jm}(0) = v_{jm}(0)$, $C_{(j+1)m}(0) = w_{(j+1)m}(0)$ ($j = 0, 1, 2$), and $A_{jm} : \mathbb{C} \rightarrow \mathbb{C}$ are additive functions.

Our second example also treats the case $n = 3$ with constant bound in Theorem 4.1.

Example 2. Let

$$\phi(x, y) = K, \quad f_i(x) = A_i(x) + a_i, \quad g_i(x) = A_i(x) + b_i, \quad h_i(x) = \omega^{-im} A_i(\omega^m x) + c_i \quad (i = 1, 2, 3),$$

where A_i are additive functions and K, a_i, b_i, c_i are constants, and let

$$\begin{aligned} \mathbf{F}(x) &= \text{circ}(f_0(x), f_1(x), f_2(x)), \quad \mathbf{G}(x) = \text{circ}(g_0(x), g_1(x), g_2(x)), \\ \mathbf{H}(x) &= \text{circ}(h_0(x), h_1(x), h_2(x)), \quad \mathbf{A}(x) = \text{circ}(A_0(x), A_1(x), A_2(x)). \end{aligned}$$

Clearly, \mathbf{A} satisfies (4.2). The hypothesis (4.1) here takes the form

$$\|\mathbf{F}(x + \omega^m y) - \mathbf{G}(x) - \Omega^{-m} \mathbf{H}(y) \Omega^m\| \leq K,$$

which is equivalent to

$$\|\text{circ}(f_0(x + \omega^m y) - g_0(x) - h_0(y), f_1(x + \omega^m y) - g_1(x) - \omega^m h_1(y), f_2(x + \omega^m y) - g_2(x) - \omega^{2m} h_2(y))\| \leq K.$$

Using the definitions of norm and f_i, g_i, h_i , this last inequality yields

$$\sum_{i=0}^2 |a_i - b_i - c_i| = \|\text{circ}(a_0 - b_0 - c_0, a_1 - b_1 - c_1, a_2 - b_2 - c_2)\| \leq K. \tag{5.1}$$

From (3.12) and (3.14), we see that $\mathbf{V}(0)$ and $\mathbf{W}_m(0)$ are diagonal matrices, while from their definitions, we see that $\mathbf{H}(0)$ and $\mathbf{G}(0)$ are circulant matrices. By direct computation, we get

$$\Phi_t(x, y) := \sum_{\ell=0}^{\infty} t^{-(\ell+1)} (9K + \|\mathbf{V}(0)\| + \|\mathbf{W}_m(0)\|) = \frac{1}{t-1} (9K + \|\mathbf{V}(0)\| + \|\mathbf{W}_m(0)\|),$$

$$\|\mathbf{V}(0)\| = \max_{0 \leq i \leq 2} |v_i(0)|, \text{ where } v_i(0) = \sum_{\ell=0}^2 \omega^{i\ell} b_i,$$

$$\|\mathbf{W}_m(0)\| = \max_{0 \leq i \leq 2} |w_i(0)|, \text{ where } w_i(0) = \sum_{\ell=0}^2 \omega^{i\ell} c_i,$$

$$\|\mathbf{H}(0)\| = \sum_{\ell=0}^2 |c_i|, \text{ and } \|\mathbf{G}(0)\| = \sum_{\ell=0}^2 |b_i|.$$

Regarding the assertions in (4.3)–(4.5), we get

$$\begin{aligned} \sum_{i=0}^2 |a_i| &= \|\text{circ}(f_0(x) - A_0(x), f_1(x) - A_1(x), f_2(x) - A_2(x))\| = \|\mathbf{F}(x) - \mathbf{A}(x)\| \\ &\leq 27K + 3 \max_{0 \leq i \leq 2} \left| \sum_{\ell=0}^2 \omega^{i\ell} b_i \right| + 3 \max_{0 \leq i \leq 2} \left| \sum_{\ell=0}^2 \omega^{i\ell} c_i \right|, \end{aligned} \tag{5.2}$$

$$\begin{aligned} \sum_{i=0}^2 |b_i| &= \|\text{circ}(g_0(x) - A_0(x), g_1(x) - A_1(x), g_2(x) - A_2(x))\| = \|\mathbf{G}(x) - \mathbf{A}(x)\| \\ &\leq 36K + 3 \max_{0 \leq i \leq 2} \left| \sum_{\ell=0}^2 \omega^{i\ell} b_i \right| + 3 \max_{0 \leq i \leq 2} \left| \sum_{\ell=0}^2 \omega^{i\ell} c_i \right| + \sum_{\ell=0}^2 |c_i|, \end{aligned} \tag{5.3}$$

$$\begin{aligned} \sum_{i=0}^2 |c_i| &= \|\text{circ}(h_0(x) - A_0(\omega^m x), h_1(x) - \omega^{2m} A_1(\omega^m x), h_2(x) - \omega^m A_2(\omega^m x))\| = \|\mathbf{H}(x) - \bar{\mathbf{A}}(x)\| \\ &\leq 36K + 3 \max_{0 \leq i \leq 2} \left| \sum_{\ell=0}^2 \omega^{i\ell} b_i \right| + 3 \max_{0 \leq i \leq 2} \left| \sum_{\ell=0}^2 \omega^{i\ell} c_i \right| + \sum_{\ell=0}^2 |b_i|. \end{aligned} \tag{5.4}$$

The hypothesis (5.1) and the assertions (5.2)–(5.4) are satisfied by choosing for example $a_i = 0, b_i = c_i$ ($i = 0, 1, 2$).

References

- [1] Czerwik S. *Functional Equations and Inequalities in Several Variables*. Singapore: World Scientific, 2002.
- [2] Förg-Rob W, Schwaiger J. On the stability of a system of functional equations characterizing generalized hyperbolic and trigonometric functions. *Aequationes Mathematicae* 1993; 45: 285-296.
- [3] Kannappan Pl. *Functional Equations and Inequalities with Applications*. New York, NY, USA: Springer, 2009.
- [4] Muldoon ME. Generalized hyperbolic functions, circulant matrices and functional equations. *Linear Algebra and its Applications* 2005; 406: 272-284. doi: 10.1016/j.laa.2005.04.011
- [5] Ponpetch K, Laohakosol V, Mavecha S. A system of functional equations satisfied by components of a quadratic function and its stability. *Bulletin of the Australian Mathematical Society* 2019; 100: 304-316. doi: 10.1017/S000497271900025X
- [6] Schwaiger J. On generalized hyperbolic functions and their characterization of functional equations. *Aequationes Mathematicae* 1992; 43: 198-210.