

1-1-2015

Genetic improvement of rice for biotic and abiotic stress tolerance

MAHMOOD UR RAHMAN ANSARI

TAYYABA SHAHEEN

SHAZAI BUKHARI

TAYYAB HUSNAIN

Follow this and additional works at: <https://dctubitak.researchcommons.org/botany>



Part of the [Botany Commons](#)

Recommended Citation

ANSARI, MAHMOOD UR RAHMAN; SHAHEEN, TAYYABA; BUKHARI, SHAZAI; and HUSNAIN, TAYYAB (2015) "Genetic improvement of rice for biotic and abiotic stress tolerance," *Turkish Journal of Botany*. Vol. 39: No. 6, Article 3. <https://doi.org/10.3906/bot-1503-47>
Available at: <https://dctubitak.researchcommons.org/botany/vol39/iss6/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals.

Genetic improvement of rice for biotic and abiotic stress tolerance

Mahmood-ur-Rahman ANSARI^{1*}, Tayyaba SHAHEEN¹, Shazia Anwer BUKHARI², Tayyab HUSNAIN³

¹Plant Research Group, Department of Bioinformatics and Biotechnology, GC University, Faisalabad, Pakistan

²Department of Applied Chemistry and Biochemistry, GC University, Faisalabad, Pakistan

³National Centre of Excellence in Molecular Biology, Lahore, Pakistan

Received: 25.03.2015 • Accepted/Published Online: 06.10.2015 • Printed: 21.12.2015

Abstract: Rice (*Oryza sativa* L.) is among the most important food crops that provide a staple food for nearly half of the world's population. Rice crops are prone to various types of stresses, both biotic and abiotic. Biotic stresses include insect pests, fungus, bacteria, viruses, and herbicide toxicity. Among abiotic stresses, drought, cold, and salinity are also well studied in rice. Various genes have been identified, cloned, and characterized to combat these stresses and protect rice crops. The identified genes are successfully transformed into rice plants to produce transgenic plants. These transgenic rice plants are being evaluated under field conditions in different countries. Genetic engineering has a very positive impact on improvement of rice crops. The development of rice with improved traits of biotic and abiotic stress tolerance is discussed in this review article. The objective of this review is to provide an overview of recent research and development in the field of rice biotechnology.

Key words: Biotic stress, Bt rice, genetically modified rice, drought, fungal resistance, herbicide tolerance

1. Introduction

Rice (*Oryza sativa* L.) is one of the oldest cultivated crops. It has been cultivated in India and China for several thousands of years (Poehlman and Sleper, 1995). The major cultivated species of rice, *Oryza sativa* ($2n = 2x = 24$), originated in southern and southwestern tropical Asia. The other species of cultivated rice, *Oryza glaberrima* ($2n = 2x = 24$), is indigenous to the upper valley of the Niger River and it is cultivated only in western tropical Africa. Twenty-three species and 10 recognized genome types (AA, BB, CC, BBCC, CCDD, EE, FF, GG, HHJJ, and HHKK; Gramene: http://www.gramene.org/species/oryza/rice_taxonomy.html) of *Oryza* are recognized. Close relatives of *O. sativa* are the wild perennial species *O. rufipogon* and the wild annual species *O. nivara*. Both are diploid weedy species with the AA genome.

Rice is an important food crop and it needs continuous improvement due to the continuous increase in population. The major objectives to improve rice crops using biotechnological techniques include: 1) high yield potential, 2) early maturity, 3) resistance to lodging and shattering, 4) resistance to stress environments, 5) disease resistance, 6) insect resistance, 7) grain quality, and 8) enhancement of nutritional components.

2. Economic importance of rice

Rice (*Oryza sativa* L.) is the most important crop of world. About 90% of the world's rice is grown in China, India, Pakistan, Japan, Korea, Southeast Asia, and other adjacent areas (USDA, 2014). Outside of Asia, Brazil and the United States produce the largest amounts of rice (Poehlman and Sleper, 1995). Rice is the staple food for over one-half of the world's people (FAO, 2008). In Pakistan, rice is a highly valued food crop and it is also a major export item. It accounts for 3.2% of the total value added in agriculture and 0.7% of the GDP. Area sown for rice is estimated at 2.891×10^6 ha and the production was 7.005×10^6 t in 2014–2015 (Ministry of Finance, 2015). Rice is cultivated in diversified climatic conditions of Pakistan. Basmati rice (Indica) is grown in the traditional rice-growing belt of Punjab Province. In Swat, in high altitude alpine valleys, temperate Japonica rice is grown. In the south of Khyber Pakhtunkhwa, Sindh and Baluchistan provinces, IRR1-type long-grain heat-tolerant tropical rice is mainly grown.

3. Transformation of rice crops

The world will need about 25% more rice by the year 2030 to meet the estimated demand of an increasing global population (Wani and Sah, 2014). One way to meet this challenge is to grow rice on more area, which is difficult due to increasing urbanization and escalating population in

* Correspondence: mahmoodansari@gcuf.edu.pk

underdeveloped countries. The other option is to improve varieties and increase per hectare yield by breeding efforts through conventional methods as well as modern biotechnology. Biotechnology has a promise to increase yield as well as decrease crop loss due to various biotic and abiotic stresses (Gelvin, 2010; Ozawa and Takaiwa, 2010; Mahmood-ur-Rahman et al., 2014b).

4. Insect-resistant rice

Insect-resistant crops have revolutionized modern agriculture and have become a major tool of integrated pest management programs, leading to reduction in insecticide use while protecting the environment and human health (Brooks and Barfoot, 2013). Like in other crops, insect-resistant rice plants were also developed about two decades ago (Fujimoto et al., 1993). Now genetically modified rice lines expressing a gene from *Bacillus thuringiensis* (Bt rice plants) are under field trials in various countries (Tu JM et al., 2000; High et al., 2004; Wang et al., 2014). There are numerous reports indicating that Bt rice can minimize losses due to lepidopteran pests in Asia (High et al., 2004). Shu et al. (2000) reported that transgenic rice transformed with a synthetic *cry1Ab* gene was found significantly tolerant to eight lepidopteran insects, including striped stem borer (SSB; *Chilo suppressalis*) and yellow stem borer (YSB; *Scirpophaga incertulas*). Furthermore, two lines from Bt rice plants were highly resistant to lepidopteran

pests under field conditions (Kumar et al., 2008; Deka and Barthakur, 2010; Wang et al., 2014). The major milestones in development of insect-resistant rice plants are reviewed in Table 1.

Insect-resistant hybrid rice plants were evaluated in a field in China and were highly tolerant to rice leaf folder (RLF; *Cnaphalocrocis medinalis*) and YSB (Tu JM et al., 2000; Deka and Barthakur, 2010; Chen et al., 2011). Insect-resistant Bt rice has also been produced in Pakistan (Mahmood-ur-Rahman et al., 2007) and in the Mediterranean region (Breitler et al., 2004). Results of recent field trials in both locations showed significant resistance against target insects, i.e. YSB and RLF (Breitler et al., 2004; Bashir et al., 2005; Mahmood-ur-Rahman et al., 2007, 2012, 2013, 2014a, 2014b; Tabashnik et al., 2009) (Table 2). The rice plants were artificially infested (Figures 1A–1C) with the target insects and their attack was measured quantitatively. Some transgenic rice lines/varieties resistant to YSB have also been developed in India (Ramesh et al., 2004). Transgenic rice plants have also been developed in China and their efficacy was tested in the laboratory as well as in the field (Wang et al., 2012; Li et al., 2013; Li F et al., 2014).

Pyramiding of multiple genes against the same pest or a range of pests has proved to be very effective to induce sustainable resistance against insect pests. Research has been carried out to pyramid *cry1Ab* or *cry1Ac* with either

Table 1. Genetic improvement of rice for insect resistance.

S. no.	Gene(s)	Targets	References
1	<i>cry1Ab</i> or <i>cry1Ac</i>	YSB*, SSB**	Shu et al. (2000)
2	<i>cry1Aa</i> or <i>cry1Ab</i>	SSB	Breitler et al. (2004)
3	<i>cry1Ab</i> and <i>cry1Ac</i>	YSB	Ramesh et al. (2004)
4	<i>cry1Ab</i>	SSB	Cotsaftis et al. (2002)
5	<i>cry1Ab</i>	YSB and RLF***	Bashir et al. (2005)
6	<i>cry</i> , <i>Xa21</i> , and <i>RC7</i>	YSB, bacterial blight, sheath blight	Datta et al. (2003)
7	<i>gna</i> and <i>cry1Ac</i>	Homopteran, coleopteran, and lepidopteran insects	Nagadhara et al. (2003)
8	<i>Itr1</i>	Rice weevil	Alfonso-Rubi et al. (2003)
9	<i>cry1Ac</i> and <i>cry2A</i>	YSB and RLF	Mahmood-ur-Rahman et al. (2007)
10	Bt and <i>CpT1</i>	Insect resistance	Rong et al. (2007)
11	Bt, protease inhibitors, enzymes, and plant lectins	Insect resistance	Deka and Barthakur 2010
12	<i>cry2Aa</i>	Insect resistance	Wang et al. (2012)
13	<i>cry1Ab</i>	Insect resistance	Wang et al. (2014)

*YSB = yellow stem borer, **SSB = stripe stem borer, ***RLF = rice leaf folder.

Table 2. Field evaluation of transgenic rice.

S. no.	Gene(s)	Trait	Location	References
1	<i>cry1Ab</i> and <i>cry1Ac</i>	Insect resistance	China	Tu JM et al. (2000)
2	<i>cry1Ab</i>	Insect resistance	China	Shu et al. (2000)
3	<i>cry1Ac</i> and <i>cry2A</i>	Insect resistance	Pakistan	Bashir et al. (2005)
4	<i>cry1Aa</i> and <i>cry1B</i>	Insect resistance	Spain	Breitler et al. (2004)
5	<i>cry2A</i>	Insect resistance	China	Chen et al. (2005)
6	<i>cry1Ac</i> and <i>cry2A</i>	Insect resistance	Pakistan	Mahmood-ur-Rahman et al. (2007, 2012, 2014a)
7	<i>cry1Ac</i> and <i>CpT1</i>	Insect resistance	China	Han et al. (2006)
8	<i>cry1Ab</i>	Insect resistance	China	Wang et al. (2014)
9	<i>Xa21</i>	Bacterial blight resistance	China	Tu J et al. (2000)
10	<i>bar</i>	Herbicide resistance	USA	Oard et al. (2000)
11	<i>bar</i>	Herbicide resistance	Spain	Messeguer et al. (2004)
12	<i>bar</i>	Herbicide resistance	USA	Zhang et al. (2004)




Figure 1. Artificial infestation by yellow stem borer (YSB) in transgenic and nontransgenic rice plants to evaluate the efficiency of Bt genes against target insects. A) Symptoms of YSB attack at the vegetative stage of plant growth; the central tiller is dead due to insect attack, which is called “dead heart”. B) Symptoms of YSB attack at the reproductive stage of plant growth; the panicle becomes white and there is no grain development inside, a condition called “white head”. C) Artificial infestation by YSB.

cry2A or *cry9C* for improved and sustainable resistance in Bt rice. *cry1Ab* and *cry1Ac* have the same binding site in insects like SSB and YSB (Alcantara et al., 2004) and may be very effective in pyramiding of genes. Use of *cry* genes along with the snowdrop (*Galanthus nivalis*) lectin gene, *gna*, provided elevated levels of tolerance to a broad range of pests, including sucking insects (Ramesh et al., 2004), as

the Bt genes have no effect on sucking insect pests (Bernal et al., 2002).

It is important to pyramid *cry* genes with other insecticidal genes to induce sustainable resistance. These non-Bt insect resistant genes may be obtained from plants such as lectins, ribosome-inactivating proteins, or protease inhibitors (Sharma et al., 2003). The lectin gene (*gna*)

has been widely used to develop insect-resistant rice for built-in resistance against various species of homopteran, lepidopteran, and coleopteran insect pests (Nagadhara et al., 2003). Transgenic rice strains producing proteinase inhibitor proteins have been developed and evaluated in the field (Mochizuki et al., 1999).

Recent developments such as an inducible and tissue-specific expression of the transgene, pyramiding of multiple genes for a broader range of protection, and delayed resistance development of insect pests could assist in the commercial application of genetically modified rice (Kumar et al., 2008). Protease inhibitors (Hernandez et al., 2003) and lectins are more effective than *gna* (Lopez et al., 2002). The cloning protease inhibitors and lectins have the potential to allow the development of insect-resistant rice (Kumar et al., 2008), which is more effective against sucking insects.

5. Fungus-resistant rice

Genetic improvement of rice for fungus resistance is the need of the day due to vulnerability of this crop to various pathogens, including fungi, bacteria, and viruses (Gust et al., 2010; Miah et al., 2013; Sattari et al., 2014). There are several genes that were isolated and cloned, and then transformation was carried out in rice plants to confer resistance against various species of fungi. The *Pi-ta* gene was recently cloned and reported to have significant resistance against rice blast (Dai et al., 2010; Delteil et al., 2010). Indigenous resistance to sheath blight was also studied and mapped (Liu et al., 2009). The molecular markers were also found by crossing between transgenic and susceptible/nontransgenic varieties (Liu et al., 2009). Indica rice generations harboring the *PR-3* rice chitinase gene were found tolerant to sheath blight (Datta et al., 2003). The *Rir1b* gene is a member of defense-related genes family. It has been identified and characterized only in cereals (Mauch et al., 1998). Transgenic rice containing *Rir1b* was reported to have an enhanced resistance to rice blast (Li et al., 2009).

Several proteins have also been identified as good candidates to confer resistance against various species of fungi in rice plants, which are involved in tolerance of pathogen attack. Examples include lipid transfer protein (Guiderdoni et al., 2002), selenium-binding protein homolog (Sawada et al., 2004), genes taking part in flavonoid pathways (Gandikota et al., 2001), puroindoline proteins (Krishnamurthy et al., 2001), rice homolog of maize HC-toxin reductase (Uchimiya et al., 2002), defensins (Kanzaki et al., 2002), trichosanthins (Yuan et al., 2002), phytoalexins (Lee et al., 2004), protease inhibitor protein genes (Qu et al., 2003), genes involved in cell death (Matsumura et al., 2003), antifungal protein from *Aspergillus flavus* (Coca et al., 2004), and mycotoxin detoxifying compounds (Higa et al., 2003).

6. Bacteria-resistant rice

Bacterial blight-resistant transgenic rice lines were developed by transforming an endogenous gene, *Xa21* (Song et al., 1995). *Xa21* has been introduced into different rice cultivars by genetic transformation as well as through conventional breeding techniques (Tyagi and Mohanty, 2000). Transgenic plants with *Xa21* were evaluated under field conditions with promising results (Tu J et al., 2000). This gene was found to be the best candidate so far to induce resistance against bacterial blight. It was also reported that the gene could provide multiple stress tolerance along with other genes (Datta et al., 2003). Another resistant gene, *Xa26*, was isolated from rice and encodes a similar protein with similar effects in transgenic plants as the plants having the *Xa21* gene (Sun et al., 2004; Zhang, 2009).

Some other genes have also been transformed to develop bacterial-resistant rice (Zhou et al., 2011). The cecropins are a family of genes having antibacterial activity. They express peptides in the hemolymph of *Cecropia* moths. They have been transformed and expressed in plants with satisfactory results (Huang et al., 1997). Transgenic rice plants producing ferredoxin-like protein, AP1, expressed tolerance to *X. oryzae* (Tang et al., 2001). Rice has also been transformed to induce resistance against *Burkholderia plantarii*. Field studies of rice plants expressing *Xa21* showed a significant increase in yield due to less damage by the pathogen (Tu J et al., 2000). *Xa21* is a good candidate gene to be transformed in rice for sustainable resistance against bacterial diseases and it should be released commercially for the general farming community (Kumar et al., 2008).

7. Virus-resistant rice

Crop damage due to viruses has been a serious problem worldwide for several decades. The risk of damage due to rice viruses, especially rice stripe virus (RSV) and rice dwarf virus (RDV), which caused greater yield losses in the 1960s (Toriyama, 2010), is increasing. The use of insecticides to control the vector insects is one possible way to control rice crops but the high costs of insecticide and the risk to the environment are the major limiting factors. Genetic resistance against rice viruses or their insect vectors is also one of the most effective methods of protecting rice plants from virus infection. Rice plants having resistance against RDV (Shimizu et al., 2009; Sasaya et al., 2013) and RSV (Xiong et al., 2009) were developed and evaluated.

Rice tungro disease is very dangerous viral disease of rice. It is caused by infection by two viruses, rice tungro bacilliform virus (RTBV) and rice tungro spherical virus (RTSV). Green leafhopper (*Nephotettix virescens*) is the vector of RTSV and assists in the transmission of the virus. Rice plants have been transformed by using the coat

protein-mediated resistance strategy. Huet et al. (1999) produced transgenic rice plants containing the RTSV replicase gene. The transgenic plants showed moderate resistance to RTSV.

8. Herbicide-tolerant rice

Herbicide tolerance is an important agronomic trait that has been used to control weeds very efficiently for several decades. Many genes are being used to develop herbicide-tolerant plants, including the *bar* gene and the *EPSPS* gene. The *bar* gene is isolated from *Streptomyces hygroscopicus*, which detoxifies herbicide glufosinate, while the *EPSPS* gene is isolated from *Agrobacterium* strain CP4 and detoxifies glyphosate herbicides (Kumar et al., 2008). Herbicide-resistant GM rice plants encoding the *bar* gene were produced during the early stages of rice transformation research (Tyagi and Mohanty, 2000) and are currently under field trials (Oard et al., 2000).

Considerable efforts have been carried out to identify other sources of genes that could be used for sustainable resistance against herbicides in transgenic rice. One such example is *Bacillus subtilis* protoporphyrinogen oxidase, which is transformed in rice plants and is an efficient source of resistance against oxyfluorfen herbicide (Jung et al., 2004). Other example is human cytochrome P450s; overexpression in rice showed variable responses to herbicides (Kawahigashi et al., 2003).

9. Abiotic stress-tolerant rice

Abiotic stresses including drought, high and low temperatures, salinity, submergence, and oxidative stress contribute significantly to reduce crop yield. More than 50% crop damage has been reported due to these stresses worldwide (Bray et al., 2000; Iqbal et al., 2013; Li Y et al.,

2014). They are often interlinked and cause similar cellular as well as physiological damage. Moreover, they also activate similar cell-signaling pathways (Nakashima et al., 2009; Qin et al., 2011). Several proteins, antioxidants, and compatible solutes are produced in response to stress conditions. Many crop plants have been developed by overexpression of genes responsible for these compounds and evaluated for various abiotic stresses under laboratory and field conditions (Luo et al., 2010).

Tolerance to water shortage and salt stress are the most damaging factors that inhibit yield in rice crops. GM technology is one of the available options to increase abiotic stress tolerance in crop plants (Flowers, 2004). Development of transgenic plants containing various genes that induce tolerance to drought and/or salinity tolerance in different plants (Tyagi and Mohanty, 2000; Iqbal et al., 2013) has been extended to crop species such as rice (Table 3). Furthermore, many new genes have been identified and isolated that are responsible for providing salt and drought stress tolerance in model plants (reviewed by Flowers, 2004; Zhang et al., 2004).

Hoshida et al. (2000) transformed rice with the chloroplastic glutamine synthase (*GS2*) gene and successfully developed transgenic plants. Transgenic plants with high *GS2* levels demonstrated an enhanced photorespiration capability and they were highly tolerant to salt stress and chilling stress (Hoshida et al., 2000). Xiong and Yang (2003) isolated and characterized a stress-responsive *MAPK* gene, *OsMAPK5*, from rice whose expression could be regulated by ABA and various other biotic and abiotic stresses. Plants overexpressing *OsMAPK5* showed elevated tolerance to drought, salt, and cold stress (Xiong and Yang, 2003; Osakabe et al., 2014; Savchenko et al., 2014).

Table 3. Abiotic stress-tolerant genes in rice.

S. no.	Gene(s)	Function	Source	Characteristics of transgenic plant	References
1	<i>GS2</i>	Glutamine synthase	Rice	Tolerant to salt and cold stresses	Hoshida et al. (2000)
2	<i>OsCDPK7</i>	Calcium-dependent protein kinase	Rice	Tolerant to salt and drought stresses	Saijo et al. (2000)
3	<i>OsMAPK5</i>	MAP kinase	Rice	---	Xiong and Yang (2003)
4	<i>Adc, Samdc</i>	Polyamine biosynthesis	<i>Datura</i> , oat	Tolerant to salt and drought stresses	Capell et al. (2004)
5	<i>HVA1</i>	LEA protein	Barley, wheat	Tolerant to salt and dehydration stress	Babu et al. 2004
6	<i>OtsA</i>	Trehalose biosynthesis	<i>E. coli</i>	Tolerant to salt, drought, and cold stress	Jung et al. (2003)
7	<i>p5cs</i>	Proline biosynthesis	Moth bean	Transgene expressed when stress induced	Hur et al. (2004)
8	<i>pdcl, adc</i>	Pyruvate decarboxylase	Rice	Submergence tolerance	Rahman et al. (2001)
9	<i>AGPAT, SGPAT</i>	Fatty acid biosynthesis	<i>Arabidopsis</i> , spinach	Improved photosynthesis at low temperatures	Ariizumi et al. (2002)
10	<i>Cat</i>	Catalase	Wheat	Cold-tolerant	Matsumura et al. (2003)
11	<i>spl7</i>	Heat-stress transcription factor	Rice	Tolerance to heat stress	Yamanouchi et al. (2002)

The dehydration-responsive elements (DREs) regulate the expression of various genes under abiotic stress conditions (Nakashima et al., 2014; Wani and Sah, 2014). Dubouzet et al. (2003) isolated 5 rice *DREB* homologs of *Arabidopsis*, *OsDREB1A*, *OsDREB1B*, *OsDREB1C*, *OsDREB1D*, and *OsDREB2A*, and characterized them in *Arabidopsis*. Researchers generated transgenic rice plants overexpressing various *OsDREB* genes driven by the combination of several promoters. They suggested that *OsDREB1A* proved very useful for the development of transgenic dicot and monocot plants with increased tolerance to drought, salt, and/or cold stresses (Wang et al., 2007; Nawaz et al., 2014).

The *ACS* gene is responsible for submergence resistance in rice plants. The mRNA level was found to be higher under

completely submerged conditions in vascular bundles of young stems and leaf sheaths. Overexpression of the *YK1* gene in rice also resulted in the same results (Uchimiya et al., 2002). A transcription factor (*Spl7*) was identified and cloned in rice plants. Its overexpression in *spl7* mutants inhibited leaf spot development induced by high temperature (Yamanouchi et al., 2002). Heat-tolerant rice plants have also been developed by overexpressing either heat shock proteins (Murakami et al., 2004) or enzymes involved in oxidative stress tolerance (Kouril et al., 2003).

Acknowledgment

The authors gratefully acknowledge the support from the Higher Education Commission of the Government of Pakistan.

References

- Alcantara EP, Aguda RM, Curtiss A, Dean DH, Cohen MB (2004). *Bacillus thuringiensis* δ -endotoxin binding to brush border membrane vesicles of rice stem borers. *Arch Insect Biochem* 55: 169–177.
- Alfonso-Rubi J, Ortego F, Castanera P, Carbonero P, Diaz I (2003). Transgenic expression of trypsin inhibitor *CMe* from barley in indica and japonica rice, confers resistance to the rice weevil *Sitophilus oryzae*. *Transgenic Res* 12: 23–31.
- Ariizumi T, Kishitani S, Inatsugi R, Nishida I, Murata N, Toriyama K (2002). An increase in unsaturation of fatty acids in phosphatidylglycerol from leaves improves the rates of photosynthesis and growth at low temperatures in transgenic rice seedlings. *Plant Cell Physiol* 43: 751–758.
- Babu RC, Zhang JX, Blum A, Ho THD, Wu R, Nguyen HT (2004). *HVA1*, an LEA gene from barley confers dehydration tolerance in transgenic rice (*Oryza sativa* L.) via cell membrane protection. *Plant Sci* 166: 855–862.
- Bashir K, Husnain T, Fatima T, Latif Z, Riaz N, Riazuddin S (2005). Novel indica basmati line (B-370) expressing two unrelated genes of *Bacillus thuringiensis* is highly resistant to two lepidopteran insects in the field. *Crop Prot* 24: 870–879.
- Bernal CC, Aguda RM, Cohen MB (2002). Effect of rice lines transformed with *Bacillus thuringiensis* toxin genes on the brown planthopper and its predator *Cyrtorhinus lividipennis*. *Entomol Exp Appl* 102: 21–28.
- Bray EA, Bailey-Serres J, Weretilnyk E (2000). Responses to abiotic stresses. In: Gruissem W, Buchannan B, Jones R, editors. *Biochemistry and Molecular Biology of Plants*. Rockville, MD, USA: American Society of Plant Physiologists. pp. 1158–1249.
- Breitler JC, Vassal JN, Catala MDM, Meynard D, Marfa V, Mele E, Royer M, Murillo I, Segundo SB, Guiderdoni E et al. (2004). Bt rice harbouring *cry* genes controlled by a constitutive or wound-inducible promoter, protection and transgene expression under Mediterranean field conditions. *Plant Biotechnol J* 2: 417–430.
- Brooks G, Barfoot P (2013). Key environmental impacts of global genetically modified (GM) crop use 1996–2011. *GM Crops Food* 4: 109–119.
- Capell T, Bassie L, Christou P (2004). Modulation of the polyamine biosynthetic pathway in transgenic rice confers tolerance to drought stress. *P Natl Acad Sci USA* 101: 9909–9914.
- Chen H, Tang W, Xu C, Li X, Lin Y, Zhang Q (2005). Transgenic indica rice plants harboring a synthetic *cry2A* gene of *Bacillus thuringiensis* exhibit enhanced resistance against lepidopteran rice pests. *Theor Appl Genet* 111: 1330–1337.
- Chen M, Shelton A, Ye GY (2011). Insect-resistant genetically modified rice in China: from research to commercialization. *Annu Rev Entomol* 56: 81–101.
- Coca M, Bortolotti C, Rufat M, Penas G, Eritja R, Tharreau D, del Pozo AM, Messeguer J, San Segundo B (2004). Transgenic rice plants expressing the antifungal AFP protein from *Aspergillus giganteus* show enhanced resistance to the rice blast fungus *Magnaporthe grisea*. *Plant Mol Biol* 54: 245–259.
- Cotsaftis O, Sallaud C, Breitler JC, Meynard D, Greco R, Pereira A, Guiderdoni E (2002). Transposon-mediated generation of tDNA-free and marker-free rice plants expressing a Bt endotoxin gene. *Mol Breeding* 10: 165–180.
- Dai Y, Jia Y, Correll J, Wang X, Wang Y (2010). Diversification and evolution of the avirulence gene AVR-pita 1 in field isolates of *Magnaporthe oryzae*. *Fungal Genet Biol* 47: 974–980.
- Datta K, Baisakh N, Oliva N, Torrizo L, Abrigo E, Tan J, Rai M, Rehana S, Al-Babili S, Beyer P et al. (2003). Bioengineered 'golden' indica rice cultivars with betacarotene metabolism in the endosperm with hygromycin and mannose selection systems. *Plant Biotechnol J* 1: 81–90.
- Deka S, Barthakur S (2010). Overview on current status of biotechnological interventions on yellow stem borer *Scirpophaga incertulas* (Lepidoptera: Crambidae) resistance in rice. *Biotechnol Adv* 28: 70–81.

- Delteil A, Zhang J, Lessard P, Morel JB (2010). Potential candidate genes for improving rice disease resistance. *Rice* 3: 56–71.
- Dubouzet JG, Sakuma Y, Ito Y, Kasuga M, Dubouzet EG, Miura S, Seki M, Shinozaki K, Yamaguchi-Shinozaki K (2003). *OsDREB* genes in rice, *Oryza sativa* L., encode transcription activators that function in drought, high-salt and cold-responsive gene expression. *Plant J* 33: 751–763.
- Flowers TJ (2004). Improving crop salt tolerance. *J Exp Bot* 55: 307–319.
- Food and Agriculture Organization (FAO) (2008). *FAO and Sustainable Intensification of Rice Production for Food Security*. Rome, Italy: FAO.
- Fujimoto H, Itoh K, Yamamoto M, Kyojuka J, Shimamoto K (1993). Insect resistant rice generated by introduction of a modified δ -endotoxin gene of *Bacillus thuringiensis*. *Biotechnology* 11: 1151–1155.
- Gandikota M, de Kochko A, Chen L, Ithal N, Fauquet C, Reddy AR (2001). Development of transgenic rice plants expressing maize anthocyanin genes and increased blast resistance. *Mol Breeding* 7: 73–83.
- Gelvin SB (2010). Plant proteins involved in *Agrobacterium*-mediated genetic transformation. *Annu Rev Phytopathol* 48: 45–68.
- Guiderdoni E, Cordero MJ, Vignols F, Garcia-Garrido JM, Lescot M, Tharreau D, Meynard D, Ferriere N, Notteghem JL, Delseny M (2002). Inducibility by pathogen attack and developmental regulation of the rice *Ltp1* gene. *Plant Mol Biol* 49: 683–699.
- Gust AA, Brunner F, Nürnberger T (2010). Biotechnological concepts for improving plant innate immunity. *Curr Opin Biotech* 21: 204–210.
- Han L, Wu K, Peng Y, Wang F, Guo Y (2006). Evaluation of transgenic rice expressing *Cry1Ac* and *CpTI* against *Chilo suppressalis* and intrapopulation variation in susceptibility to *Cry1Ac*. *Entomol* 35: 1453–1459.
- Hernandez CA, Pujol M, Alfonso-Rubi J, Armas R, Coll Y, Perez M, Gonzalez A, Ruiz M, Castanera P, Ortego F (2003). Proteolytic gut activities in the rice water weevil, *Lissorhoptrus brevisrostris* Suffrian (*Coleoptera*, Curculionidae). *Arch Insect Biochem* 53: 19–29.
- Higa A, Kimura M, Mimori K, Ochiai-Fukuda T, Tokai T, Takahashi-Ando N, Nishiuchi T, Igawa T, Fujimura M, Hamamoto H et al. (2003). Expression in cereal plants of genes that inactivate *Fusarium* mycotoxins. *Biosci Biotech Bioch* 67: 914–918.
- High SM, Cohen MB, Shu QY, Altosaar I (2004). Achieving successful deployment of *Bt* rice. *Trends Plant Sci* 9: 286–292.
- Hoshida H, Tanaka Y, Hibino T, Hayashi Y, Tanaka A, Takabe T (2000). Enhanced tolerance to salt stress in transgenic rice that overexpresses chloroplast glutamine synthetase. *Plant Mol Biol* 43: 103–111.
- Huang N, Angeles ER, Domingo J, Magpantay G, Singh S, Zhang G, Kumaravadivel N, Bennett J, Khush GS (1997). Pyramiding of bacterial blight resistance genes in rice: marker-assisted selection using RFLP and PCR. *Theor Appl Genet* 95: 313–320.
- Huet H, Mahendra S, Wang J, Sivamani E, Ong CA, Chen L, de Kochko A, Beachy RN, Fauquet C (1999). Near immunity to rice tungro spherical virus achieved in rice by a replicase mediated resistance strategy. *Phytopathology* 89: 1022–1027.
- Hur J, Jung KH, Lee CH, An GH (2004). Stress-inducible *OsP5CS2* gene is essential for salt and cold tolerance in rice. *Plant Sci* 167: 417–426.
- Iqbal M, Khan MA, Naeem M, Aziz U, Afzal J, Latif M (2013). Inducing drought tolerance in upland cotton (*Gossypium hirsutum* L.), accomplishments and future prospects. *World Applied Science Journal* 21: 1062–1069.
- Jung SY, Chung JS, Chon SU, Kuk YI, Lee HJ, Guh JO, Back K (2004). Expression of recombinant protoporphyrinogen oxidase influences growth and morphological characteristics in transgenic rice. *Plant Growth Regul* 42: 283–288.
- Kanzaki H, Nirasawa S, Saitoh H, Ito M, Nishihara M, Terauchi R, Nakamura I (2002). Overexpression of the wasabi defensin gene confers enhanced resistance to blast fungus (*Magnaporthe grisea*) in transgenic rice. *Theor Appl Genet* 105: 809–814.
- Kawahigashi H, Hirose S, Ohkawa H, Ohkawa Y (2003). Transgenic rice plants expressing human *CYP1A1* exude herbicide metabolites from their roots. *Plant Sci* 165: 373–381.
- Kouril R, Lazar D, Lee H, Jo J, Naus J (2003). Moderately elevated temperature eliminates resistance of rice plants with enhanced expression of glutathione reductase to intensive photooxidative stress. *Photosynthetica* 41: 571–578.
- Krishnamurthy K, Balconi C, Sherwood JE, Giroux MJ (2001). Wheat puroindolines enhance fungal disease resistance in transgenic rice. *Mol Plant Microbe In* 14: 1255–1260.
- Kumar S, Chandra A, Pandey KC (2008). *Bacillus thuringiensis* (Bt) transgenic crop: an environment friendly insect-pest management strategy. *J Environ Biol* 29: 641–653.
- Lee HJ, Lee DE, Ha SB, Jang SW, Lee IJ, Ryu SB, Back KW (2004). The characterization of transgenic rice plants expressing a pepper 5-epi aristolochene synthase, the first committed step enzyme for capsidiol synthesis in isoprenoid pathway. *Plant Sci* 166: 881–887.
- Li H, Zhou SY, Zhao WS, Su SC, Peng YL (2009). A novel wall-associated receptor-like kinase gene, OsWAK1, plays important roles in rice blast disease resistance. *Plant Mol Biol* 69: 337–346.
- Li F, Fan G, Wang K, Sun F, Yuan Y, Song G, Li Q, Ma Z, Lu C, Zou C et al. (2014). Genome sequence of the cultivated cotton *Gossypium arboreum*. *Nat Genet* 46: 567–572.
- Li Y, Hu L, Romeis J, Wang Y, Han L, Chen X, Peng Y (2014). Use of an artificial diet system to study the toxicity of gut-active insecticidal compounds on larvae of the green lacewing *Chrysoperla sinica*. *Biol Control* 69: 45–51.
- Li Y, Wang Y, Romeis J, Liu Q, Lin K, Chen X, Peng Y (2013). *Bt* rice expressing *Cry2A2* does not cause direct detrimental effects on larvae of *Chrysoperla sinica*. *Ecotoxicology* 22: 1413–1421.

- Liu G, Jia Y, Correa-Victoria F, Prado GA, Yeater KM, McClung A, Correll JC (2009). Mapping quantitative trait loci responsible for resistance to sheath blight in rice. *Phytopathology* 99: 1078–1084.
- Lopez S, Codina C, Bastida J, Viladomat F, Davidson E, Stewart D (2002). Biodiversity of mannose-specific lectins within *Narcissus* species. *J Agric Food Chem* 50: 2507–2513.
- Luo M, Liu J, Lee RD, Scully BT, Guo B (2010). Monitoring the expression of maize genes in developing kernels under drought stress using oligo-microarray. *J Integr Plant Biol* 52: 1059–1074.
- Mahmood-ur-Rahman, Noor M, Shahid AA, Rao AQ, Husnain T (2014a). Field performance and biosafety studies of segregating generations of transgenic basmati rice (*Oryza sativa* L.) containing multiple genes from *Bacillus thuringiensis*. *Annals of Agriculture Science (Special Issue)* 1: 39–42.
- Mahmood-ur-Rahman, Qasim M, Bokhari SA, Shaheen T (2014b). Bt crops: a sustainable approach towards biotic stress tolerance. In: Ahmad P, Rasool S, editors. *Emerging Technologies and Management of Crop Stress Tolerance. Volume 1 - Biological Techniques*. Amsterdam, the Netherlands: Elsevier, pp. 125–142.
- Mahmood-ur-Rahman, Qasim M, Bukhari SA (2013). Bt crops: from laboratory to field. *NAYS e-Magazine* 5: 3.
- Mahmood-ur-Rahman, Rao AQ, Batool F, Azam S, Shahid AA, Husnain T (2012). Transgene copy number and phenotypic variations in transgenic basmati rice. *J Anim Plant Sci* 22: 1004–1013.
- Mahmood-ur-Rahman, Rashid H, Shahid AA, Bashir K, Husnain T, Riazuddin S (2007). Insect resistance and risk assessment studies of advanced generations of basmati rice expressing two genes of *Bacillus thuringiensis*. *Electron J Biotechnol* 10: 241–251.
- Matsumura H, Nirasawa S, Kiba A, Urasaki N, Saitoh H, Ito M, Kawai-Yamada M, Uchimiya H, Terauchi R (2003). Overexpression of Bax inhibitor suppresses the fungal elicitor-induced cell death in rice (*Oryza sativa* L.) cells. *Plant J* 33: 425–434.
- Mauch F, Reimann C, Freydl E, Schaffrath U, Dudler R (1998). Characterization of the rice pathogen-related protein Rir1a and regulation of the corresponding gene. *Plant Mol Biol* 38: 577–586.
- Messeguer J, Marfa V, Catala MM, Guiderdoni E, Mele E (2004). A field study of pollen-mediated gene flow from Mediterranean GM rice to conventional rice and the red rice weed. *Mol Breeding* 13: 103–112.
- Miah M, Rafii MY, Ismail MR, Puteh AB, Rahim HA, Asfaliza R, Latif MA (2013). Blast resistance in rice: a review of conventional breeding to molecular approaches. *Mol Biol Rep* 40: 2369–2388.
- Ministry of Finance (2015). *Economic Survey of Pakistan*. Islamabad, Pakistan: Ministry of Finance, Economic Advisor's Wing.
- Mochizuki A, Nishizawa Y, Onodera H, Tabei Y, Toki S, Habu Y, Ugaki M, Ohashi Y (1999). Transgenic rice plants expressing a trypsin inhibitor are resistant against rice stem borers, *Chilo suppressalis*. *Entomol Exp Appl* 93: 173–178.
- Murakami T, Matsuba S, Funatsuki H, Kawaguchi K, Saruyama H, Tanida M, Sato Y (2004). Over-expression of a small heat shock protein, *sHSP17.7*, confers both heat tolerance and UV-B resistance to rice plants. *Mol Breeding* 13: 165–175.
- Nagadhara D, Ramesh S, Pasalu IC, Rao YK, Krishnaiah NV, Sarma NP, Bown DP, Gatehouse JA, Reddy VD, Rao KV (2003). Transgenic indica rice resistant to sap-sucking insects. *Plant Biotechnol J* 1: 231–240.
- Nakashima K, Ito Y, Yamaguchi-Shinozaki K (2009). Transcriptional regulatory networks in response to abiotic stresses in *Arabidopsis* and grasses. *Plant Physiol* 149: 88–95.
- Nakashima K, Yamaguchi-Shinozaki K, Shinozaki K (2014). The transcriptional regulatory network in the drought response and its crosstalk in abiotic stress responses including drought, cold, and heat. *Frontiers in Plant Science* 5: 170.
- Nawaz M, Iqbal N, Idrees S, Ullah I (2014). *DREB1A* from *Oryza sativa* var. IR6: homology modelling and molecular docking. *Turk J Bot* 38: 1095–02.
- Oard J, Cohn MA, Linscombe S, Gealy D, Gravois K (2000). Field evaluation of seed production, shattering, and dormancy in hybrid populations of transgenic rice (*Oryza sativa*) and the weed, red rice (*Oryza sativa*). *Plant Sci* 157: 13–22.
- Osakabe Y, Osakabe K, Shinozaki K, Tran LR (2014). Response of plants to water stress. *Frontiers in Plant Science* 5: 86.
- Ozawa K, Takaiwa F (2010). Highly efficient *Agrobacterium*-mediated transformation of suspension-cultured cell clusters of rice (*Oryza sativa* L.). *Plant Sci* 179: 333–337.
- Poehlman JM, Sleper DA (1995). *Breeding Field Crops*. 4th ed. Ames, IA, USA: Iowa State University Press.
- Qin F, Shinozaki K, Yamaguchi-Shinozaki K (2011). Achievements and challenges in understanding plant abiotic stress responses and tolerance. *Plant Cell Physiol* 52: 1569–1582.
- Qu LJ, Chen J, Liu MH, Pan NS, Okamoto H, Lin ZZ, Li CY, Li DH, Wang JL, Zhu GF et al. (2003). Molecular cloning and functional analysis of a novel type of Bowman-Birk inhibitor gene family in rice. *Plant Physiol* 133: 560–570.
- Rahman M, Grover A, Peacock WJ, Dennis ES, Ellis MH (2001). Effects of manipulation of pyruvate decarboxylase and alcohol dehydrogenase levels on the submergence tolerance of rice. *Aust J Plant Physiol* 28: 1231–1241.
- Ramesh S, Nagadhara D, Reddy VD, Rao KV (2004). Production of transgenic indica rice resistant to yellow stem borer and sap-sucking insects, using super-binary vectors of *Agrobacterium tumefaciens*. *Plant Sci* 166: 1077–1085.
- Rong J, Lu BR, Song Z, Su J, Snow AA, Zhang X, Sun S, Chen R, Wang F (2007). Dramatic reduction of crop-to-crop gene flow within a short distance from transgenic rice fields. *New Phytol* 173: 346–353.
- Saijo Y, Hata S, Kyozuka J, Shimamoto K, Izui K (2000). Over-expression of a single Ca²⁺ dependent protein kinase confers both cold and salt/drought tolerance on rice plants. *Plant J* 23: 319–327.

- Sasaya T, Aoki H, Omura T, Yatou O, Saito K (2013). Development of virus-resistant transgenic forage rice cultivars. *Front Microbiol* 4: 409.
- Sattari A, Fakheri B, Hassan FSC, Noroozi M (2014). Blast resistance in rice: a review of breeding and biotechnology. *International Journal of Agriculture and Crop Science* 7: 329–333.
- Savchenko T, Kolla VA, Wang CQ, Nasafi Z, Hicks DR, Phadungchob B, Chehab WE, Brandizzi F, Froehlich J, Dehesh K (2014). Functional convergence of oxylipin and abscisic acid pathways controls stomatal closure in response to drought. *Plant Physiol* 164: 1151–1160.
- Sawada K, Hasegawa M, Tokuda L, Kameyama J, Kodama O, Kohchi T, Yoshida K, Shinmyo A (2004). Enhanced resistance to blast fungus and bacterial blight in transgenic rice constitutively expressing *OsSBP*, a rice homologue of mammalian selenium-binding proteins. *Biosci Biotech Bioch* 68: 873–880.
- Sharma M, Charak KS, Ramanaiah TV (2003). Agricultural biotechnology research in India: status and policies. *Curr Sci* 84: 297–302.
- Shimizu T, Yoshii M, Wei T, Hirochika H, Omura T (2009). Silencing by RNAi of the gene for Pns12, a viroplasm matrix protein of rice dwarf virus, results in strong resistance of transgenic rice plants to the virus. *Plant Biotechnol J* 7: 24–32.
- Shu QU, Ye GY, Cui HR, Cheng XY, Xiang YB, Wu DX, Gao MW, Xia YW, Hu C, Sardana R et al. (2000). Transgenic rice plants with a synthetic *cry1Ab* gene from *Bacillus thuringiensis* were highly resistant to eight lepidopteran rice pest species. *Mol Breeding* 6: 433–439.
- Song WY, Wang GL, Chen LL, Kim HS, Pi LY, Holsten T, Gardner J, Wang B, Zhai WX, Zhu LH et al. (1995). A receptor kinase-like protein encoded by the rice disease resistance gene, *Xa21*. *Science* 270: 1804–1806.
- Sun XL, Cao YL, Yang ZF, Xu CG, Li XH, Wang SP, Zhang QF (2004). *Xa26*, a gene conferring resistance to *Xanthomonas oryzae* pv. *oryzae* in rice, encodes an LRR receptor kinase-like protein. *Plant J* 37: 517–527.
- Tabashnik BE, Van Rensburg JB, Carrière Y (2009). Field-evolved insect resistance to Bt crops: definition, theory, and data. *J Econ Entomol* 102: 2011–2025.
- Tang KX, Sun XF, Hu QN, Wu AZ, Lin CH, Lin HJ, Twyman RM, Christou P, Feng TY (2001). Transgenic rice plants expressing the ferredoxin-like protein (API) from sweet pepper show enhanced resistance to *Xanthomonas oryzae* pv. *Oryzae*. *Plant Sci* 160: 1035–1042.
- Toriyama S, editor (2010). *Rice Stripe: A Serious Threat to Rice Crops of Japan*. Tokyo, Japan: Soufusha Press.
- Tu J, Datta K, Khush GS, Zhang Q, Datta SK (2000). Field performance of *Xa21* transgenic indica rice (*Oryza sativa* L.), IR72. *Theor Appl Genet* 101: 15–20.
- Tu JM, Zhang GA, Datta K, Xu CG, He YQ, Zhang QF, Khush GS, Datta SK (2000). Field performance of transgenic elite commercial hybrid rice expressing *Bacillus thuringiensis* δ -endotoxin. *Nat Biotechnol* 18: 1101–1104.
- Tyagi AK, Mohanty A (2000). Rice transformation for crop improvement and functional genomics. *Plant Sci* 158: 1–18.
- Uchimiya H, Fujii S, Huang JR, Fushimi T, Nishioka M, Kim KM, Yamada MK, Kurusu T, Kuchitsu K, Tagawa M (2002). Transgenic rice plants conferring increased tolerance to rice blast and multiple environmental stresses. *Mol Breeding* 9: 25–31.
- USDA (2014). Foreign Agricultural Service, Production Supply and Distribution (PS&D) and Grain: World Markets and Trade (Grain Circular). Washington, DC, USA: USDA.
- Wang H, Zhang H, Gao F, Li J, Li Z (2007). Comparison of gene expression between upland and lowland rice cultivars under water stress using cDNA microarray. *Theor Appl Genet* 115: 1109–1126.
- Wang Y, Li Y, Romeis J, Chen X, Zhang J, Chen H, Peng Y (2012). Consumption of Bt rice pollen expressing Cry2Aa does not cause adverse effects on adult *Chrysoperla sinica* Tjeder (Neuroptera: Chrysopidae). *Biol Control* 61: 246–251.
- Wang Y, Zhang L, Li H, Han L, Liu Y, Zhu Z, Wang F, Peng Y (2014). Expression of Cry1Ab protein in a marker-free transgenic Bt rice line and its efficacy in controlling a target pest *Chilo suppressalis* (Lepidoptera: Crambidae). *Environ Entomol* 43: 528–536.
- Wani SH, Sah SK (2014). Biotechnology and abiotic stress tolerance in rice. *Journal of Rice Research* 2: e105.
- Xiong LZ, Yang YN (2003). Disease resistance and abiotic stress tolerance in rice are inversely modulated by an abscisic acid inducible mitogen-activated protein kinase. *Plant Cell* 15: 745–759.
- Xiong R, Wu J, Zhou Y, Zhou X (2009). Characterization and subcellular localization of an RNA silencing suppressor encoded by rice stripe *Tenuivirus*. *Virology* 387: 29–40.
- Yamanouchi U, Yano M, Lin H, Ashikari M, Yamada K (2002). A rice spotted leaf gene *Spl7*, encodes a heat stress transcription factor protein P Natl Acad Sci USA 99: 7530–7535.
- Yuan H, Ming X, Wang L, Hu P, An C, Chen Z (2002). Expression of a gene encoding trichosanthin in transgenic rice plants enhances resistance to fungus blast disease. *Plant Cell Rep* 20: 992–998.
- Zhang JZ, Creelman RA, Zhu JK (2004). From laboratory to field. Using information from *Arabidopsis* to engineer salt, cold, and drought tolerance in crops. *Plant Physiol* 135: 615–621.
- Zhang Q (2009). Genetics and improvement of bacterial blight resistance of hybrid rice in China. *Rice Science* 16: 83–92.
- Zhou YL, Uzokwe VN, Zhang CH, Cheng LR, Wang L, Chen K, Gao XQ, Sun Y, Chen JJ, Zhu LH et al. (2011). Improvement of bacterial blight resistance of hybrid rice in China using the *Xa23* gene derived from wild rice (*Oryza rufipogon*). *Crop Prot* 30: 637–644.