

1-1-2021

Immune dysfunction in COVID-19 and judicious use of antirheumatic drugs for the treatment of hyperinflammation

ABDURRAHMAN TUFAN

MARCO MATUCCI-CERINIC

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

TUFAN, ABDURRAHMAN and MATUCCI-CERINIC, MARCO (2021) "Immune dysfunction in COVID-19 and judicious use of antirheumatic drugs for the treatment of hyperinflammation," *Turkish Journal of Medical Sciences*: Vol. 51: No. 7, Article 25. <https://doi.org/10.3906/sag-2110-179>

Available at: <https://journals.tubitak.gov.tr/medical/vol51/iss7/25>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Immune dysfunction in COVID-19 and judicious use of antirheumatic drugs for the treatment of hyperinflammation

Abdurrahman TUFAN^{1*}, Marco MATUCCI CERINIC^{2,3}

¹Division of Rheumatology, Department of Internal Medicine, Gazi University, Ankara, Turkey

²Division of Rheumatology AOUC, Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy

³Unit of Immunology, Rheumatology, Allergy and Rare Diseases (UnIRAR), IRCCS San Raffaele Hospital, Milan, Italy

Received: 17.10.2021 • Accepted/Published Online: 27.11.2021 • Final Version: 17.12.2021

Abstract: In the Wuhan province of China, almost two years ago, in December 2019, the novel Coronavirus 2019 has caused a severe involvement of the lower respiratory tract leading to an acute life-threatening respiratory syndrome, coronavirus disease-19 (COVID-19). Subsequently, coronavirus 2 (SARS-CoV-2) rapidly spread to the entire world causing a pandemic and affected every single person on earth either directly or indirectly with destroying all facets of social life and economy. Since the announcement of COVID-19 as a global pandemic, we have witnessed tremendous scientific work on all aspects of COVID-19 across the globe, which has never been witnessed before. The most remarkable achievement would be the introduction of vaccines, which provide protection from the severe infection and is the only premise for the control of disease. However, despite the tremendous work, the number of treatments either antiviral or immunomodulatory for infected patients are considerably limited, yet disease is causing substantial morbidity and mortality. COVID-19 follows heterogenous disease course among infected individuals, and dysregulated immune system is primarily responsible for the worse outcomes. Immune deficiency, being on corticosteroids for inflammatory diseases, delayed interferon response and advanced age adversely influence prognosis with impairing viral clearance. On the other hand, exuberant immune response with features of cytokine storm is the leading cause of death, which can be alleviated by use of either general immunosuppression with corticosteroids or selective neutralization of potent pro-inflammatory cytokines such as interleukin (IL)-1 and IL-6. Herein, we summarized the potential effective immunomodulatory treatments emphasizing in which patient population it is the most suitable, which dose should be administered, and which is the most appropriate timepoint to administer the drug during the course of the disease.

Key words: SARS-CoV-2, COVID-19, inflammation, cytokine storm, treatment, rheumatology

1. Introduction

In December 2019, almost two years ago, in the Wuhan province of China, the novel Coronavirus 2019 (COVID-19) has caused an outbreak with severe involvement of the lower respiratory tract leading to a new acute life-threatening respiratory syndrome. Subsequently, coronavirus 2 (SARS-CoV-2) rapidly expand to other continents causing a pandemic, which affected every single person on the earth either directly or indirectly with destroying all facets of social life and economy. Since the announcement of COVID-19 as a global pandemic, we have witnessed tremendous scientific work on all aspects of COVID-19 across the globe, which has never been witnessed before. The most remarkable achievement would be the introduction of vaccines, which provide protection from severe infection and is the only premise for the control of the disease. However, despite the tremendous work, the number of treatments either

antiviral or immunomodulatory for infected patients is considerably limited, yet the disease is causing substantial morbidity and mortality.

COVID-19 follows a heterogenous disease course among infected individuals and dysregulated immune system is primarily responsible for the worse outcomes [1]. According to the disease progression, patients may be roughly divided into two groups; asymptomatic or mild cases that usually recover and severe cases (approximately 15%) that develop serious lung inflammation, acute respiratory distress syndrome (ARDS), cardiac and renal injury, multiorgan failure and thromboembolic events, especially in patients with older age, comorbidities, and yet little known genetically susceptible individuals [2-5]. Immune deficiency, being on corticosteroids for inflammatory diseases, delayed interferon response, and advanced age are adverse prognostic factors that impair viral clearance [6]. On the other hand, exuberant

* Correspondence: atufan@gazi.edu.tr

immune response with features of Cytokine Storm (CS) or Multisystem Inflammatory Syndrome (MIS-) is the leading cause of death which can be alleviated by use of either general immunosuppression with corticosteroids or selective neutralization of potent pro-inflammatory cytokines such as interleukin (IL)-1 and IL-6. However, neither approach has achieved high clinical success rates in patients with severe COVID-19 necessitating urgent development of more effective agents.

An efficient immune response against SARS-CoV-2 may be considered fundamental for the resolution of COVID-19. However, studies have shown a significant relationship between the disease severity and the dysregulated immune system, including exhausted T cell response, lymphopenia, neutrophilia, dysregulation (overactivation) of macrophages, impaired type I interferon (IFN-I) response, antibody-dependent enhancement, and especially, cytokine storm (CS) [3, 7]. It has been suggested that, during the response to SARS-CoV-2, the immune dysregulation and the high level of pro-inflammatory cytokines are the main cause of tissue injury. Eventually, the exact immunopathogenesis of COVID-19 remains to be elucidated, but, in brief, overactivated innate and impaired adaptive immune responses characterize severe COVID-19.

The main challenge at current is the identification of patients who would progress into critical disease and whether a specific subset of patients might benefit most from the immunomodulatory treatments. Several well-designed clinical trials demonstrated the efficacy of certain drugs in COVID-19 that are used in daily rheumatology practice, including corticosteroids, tocilizumab, anakinra, Janus Kinase (JAK) inhibitors, colchicine, and intravenous immunoglobulin. Herein, we discussed altered immune responses in COVID-19 patients and current treatment options for severe COVID-19, particularly emphasizing the rational use of drugs, on the basis of either the timing of the therapy, relevant inflammatory characteristics of the patient, or both.

2. The origin and structural features of SARS-CoV2

SARS-CoV-2 has nonsegmented, single-stranded positive-sense RNA (+ssRNA) with 5'-cap structure and 3'-poly-A tail, which is the typical genomic structure of the betacoronaviruses like SARS-CoV and MERS-CoV [8]. The genome of CoV contains six major open reading frames (ORFs) and numerous accessory genes. First ORFs (ORF1a/b), which encompasses the two-third of viral RNA, encode two large proteins of CoVs, polyprotein 1a (pp1a) and pp1ab. These polyproteins are divided into 16 nonstructural proteins (nsps), responsible for viral RNA replication and transcription, by virally encoded chymotrypsin-like protease (3CLpro) or main protease

(Mpro) and papain-like protease (PLpro) [9, 10]. The remaining ORFs on the one-third of the genome encode major structural proteins, including spike (S), envelope (E), membrane (M), and nucleocapsid (N) proteins, all of which are crucial for the viral infectivity. CoVs possess a lipid bilayer envelope with S, M, and E proteins.

Viral N protein acts as an antagonist to the interferon pathway by regulating the signaling and synthesis of type I interferon (IFN), which is one of the most critical responses in the innate immunity to viral infection [11]. The M protein is the most abundant component of the viral envelope that gives the shape of the virus and promotes the membrane curvature and the virus assembly by interacting with the S protein and the ribonucleoprotein [12–14]. The E protein is a small integral membrane protein that facilitates the assembly, the budding, and the envelope formation as well as the M protein [15]. Moreover, the E protein has an ion-channel activity and activates inflammasomes [16].

The S glycoproteins on the surface of CoVs are the receptor binding proteins, which are responsible for the attachment to host cells, viral-host cell membrane fusion, and the internalization of the virus. The S glycoprotein consists of two domains: S1 domain, which includes receptor-binding domain (RBD), interacting with angiotensin converting enzyme 2 (ACE2) on the human host cells as SARS-CoV, and S2 domain, which mediates virus-cell membrane fusion and viral entry [17, 18]. After attachment of SARS-CoV-2 with S protein to ACE2 on the host cells, S protein is cleaved by host cell proteases to reveal the S2 domain for viral-host membrane fusion and viral entry, which is coupled with TNF- α production [8, 19, 20].

3. The immune response and cytokine storm in COVID-19

The effective antiviral responses of the host innate and adaptive immunity, including the production of various proinflammatory cytokines, the activation of T cells, CD4 and CD8+ T cells, are essential for inhibition of viral replication, resolution of virus-induced inflammation, and clearance of infected cells [21, 22]. Nevertheless, the tissue injury caused by the virus could stimulate abnormal production of proinflammatory cytokines, the recruitment of proinflammatory macrophages and granulocytes. This results in the CS with shared features with macrophage activation syndrome (MAS) or secondary hemophagocytic lymphohistiocytosis (sHLH), thus leading to extensive tissue damage [23–25]. Data obtained from SARS-CoV-2 infected patients have shown that severe cases may be characterized by a cytokine storm inexorably progressing to ARDS [26–29]. Several features of COVID-19, such as the cytokine profile, serological markers, and clinical symptoms, resemble viral infection triggered sHLH

[3, 24]. Furthermore, severity of COVID-19 is closely associated with the level of the proinflammatory cytokines and subsets of immune cells [3, 30].

COVID-19 possesses different levels of various cytokines and chemokines through the mild to severe stage of the disease. The retrospective analysis has demonstrated that initial plasma levels of IL-1 β , IL-1RA, IL-7, IL-8, IL-10, IFN- γ , monocyte chemoattractant peptide (MCP)-1, macrophage inflammatory protein (MIP)-1A, MIP-1B, granulocyte-colony stimulating factor (G-CSF), and tumor necrosis factor-alpha (TNF- α) are increased in patients with COVID-19. The further analysis has shown that the plasma concentrations of IL-2, IL-7, IL-17, IL-10, MCP-1, MIP-1A, and TNF- α in intensive care unit (ICU) patients are higher than that of non-ICU patients [26]. Moreover, the plasma levels of IL-2, IL-6, IL-8, IL-10, and TNF- α , observed in severe infection, are prominently greater than those with nonsevere infection [27]. Few retrospective studies have revealed that the lung injury is strongly associated with the levels of IL-1 α , IL-1ra, IL-2, IL-7, IL-10, IL-17, IFN- γ , inducible interferon protein (IP)-10, G-CSF, and MCP-3 and all of these cytokines and chemokines excluding MCP-3 are positively related to SARS-CoV-2 viral load [7, 28]. The plasma level of IL-6, a notorious cytokine contributing to MAS, increases both in mild and severe patient groups of COVID-19 with being remarkably higher among the latter group of patients [3, 27, 29, 31]. Furthermore, based on the assessment of pulmonary infiltration in patients with ARDS, the large area of lung injury ($\geq 50\%$) is closely correlated with the increased level of IL-6 and the subgroup of lymphocytes in the peripheral blood [32].

During the infection, both innate and adaptive immune cells synergistically participate in the antiviral response [33]. The important increment in the number of neutrophils, leukocytes, and the neutrophil-lymphocyte-ratio (NLR) has been observed in severe COVID-19 compared to mild cases. The prominent lymphopenia, indicating an impairment of immune system, develops in most COVID-19 patients, especially in severe ones [3, 27]. Therefore, it seems that neutrophils and leukocytes might reinforce the CS rather than lymphocytes in COVID-19.

The level of lymphocytes and subsets of T cells, which play a significant role in the balancing of the immune response, varies according to the type of the virus due to possible viral pathogenetic mechanisms. Data from recent studies have suggested that SARS-CoV-2 infection can lead to immune dysregulation by affecting the subsets of T cells. In patients with COVID-19, the levels of T helper cells (CD3+, CD4+), cytotoxic suppressor T cells (CD3+, CD8+), and regulatory T cells are below normal levels while T helper cells and T regulatory cells in severe patients are remarkably lower than nonsevere patients. Regulatory

T cells are responsible for the maintenance of the immune homeostasis by suppressing the activation, proliferation, and proinflammatory function of most lymphocytes, including CD4+ T cells, CD8+ T cells, NK cells, and B cells [34, 35]. Furthermore, the percentage of naïve helper T cells amplifies, while the percentage of memory T helper cells and CD28+ cytotoxic suppressor T cells decreases in severe COVID-19 [3, 27]. The equilibrium between the naïve T cells and memory T cells is fundamental for maintaining an efficient immune response [36]. In addition to T cells, the reduction of B cells and NK cells are seen in COVID-19. Another important result is the confirmed strong relationship between inflammatory markers, including erythrocyte sedimentation rate (ESR), C-reactive protein (CRP) and IL-6, and the subset of lymphocytes [30]. However, there is no significant correlation between IL-6 with subsets of lymphocytes [3] and CD4+/CD8+ T cell ratio in SARS-CoV-2 infection is similar to the healthy group, the increase in this ratio and the decline of CD8+ T cells and B cells are considered poor predictors for the assessment of post-treatment clinical follow-up [3, 30]. Taken together, these results indicate that SARS-CoV-2 is responsible for an immune dysregulation with the induction of aberrant cytokine and chemokine response, alteration in levels of the lymphocyte subsets, all of which result in cytokine storm and sharpish tissue damage.

4. Antiinflammatory treatments used for combatting COVID-19

CS is an acute life-threatening dynamic inflammatory condition resulting from chaotic activation of a wide array of cytokines from initiation, immune hyperactivation, and eventually progressive multiorgan dysfunction/failure (MODS/MOF). Prompt diagnosis and effective therapeutic interventions are crucial for halting CS. Treatment is directed at controlling chaotic inflammatory response with specifically or nonspecifically targeting inflammatory cytokines or associated effector signaling pathways for restoring the host immune system (Table). Herein, we reviewed the role of the potential immuno-suppressive and immuno-modulatory treatments currently used in rheumatology practice to control hyper-inflammation encountered in COVID-19.

4.1. Corticosteroids

Initial case-based observations and, thereafter, the report of COVID-19 Global Rheumatology Alliance (GRA), which showed adverse outcomes in those patients receiving corticosteroids, precluded use of these drugs which is deemed to accelerate viral replication [37]. However, with a better understanding of the pathogenesis of severe COVID-19, corticosteroids are employed for their broad-spectrum antiinflammatory actions on production of

Table. Anti-rheumatic drugs with proven efficacy in the treatment of COVID-19.

Drug	Regimen	When to consider	Recommendations/precautions
Corticosteroids	For 10 days (or until discharge, if earlier) with daily doses of · Dexamethasone 6 mg IV or PO or · Prednisone 40 mg or · Methylprednisolone 32 mg Alternatively, 250 to 500 mg pulses for 3 consecutive days	Hospitalized patients with severe COVID-19 who require respiratory support (SpO ₂ ≤94% on room air) or with end organ dysfunction as seen in sepsis	<ul style="list-style-type: none"> · Concomitant use of antivirals is recommended · Beware of secondary viral, bacterial, fungal (i.e. Mucor mycosis) infections · Beware of reactivation of latent viral (i.e. Hepatitis B) and mycobacterial infections · Blood glucose monitoring · Steroid myopathy
Colchicine	0.5 mg PO BID for the first 3 days, and then once per day for 27 consecutive days	PCR positive outpatients with at least one adverse prognostic factor	<ul style="list-style-type: none"> · Evidence is weak for the use of colchicine · Bone marrow suppression, liver toxicity, diarrhea, myopathy, and serious drug interactions may occur with Cyp3A4 metabolites.
Anakinra	Not well determined, varied between 100 mg daily subc to 5 mg/kg twice a day IV. Doses used to treat HLH would be a reasonable approach (doses >4 mg per kg of body weight)	<ul style="list-style-type: none"> · Patients with obvious hyperinflammation (CRP >100 mg/L) who could not tolerate corticosteroids such as uncontrolled diabetes and with liver disease, · Refractory MIS-C and MIS-A 	<ul style="list-style-type: none"> · Evidence is weak for the use of anakinra · Low subcutaneous doses (less than 200 mg/day) might not be effective · Not effective in advanced disease, ie in those who need ventilatory support on admission · More effective if used early before organ dysfunction · Not effective when co-administered with dexamethasone · Risk of neutropenia
Tocilizumab	400 mg iv single infusion, in case of respiratory decline a repeat dose of 400 mg 12–24 h after the first infusion	On top of corticosteroids when hypoxemia progressively worsens	<ul style="list-style-type: none"> · More effective if administered promptly at the time of rapidly progressive severe hypoxemia · Benefit might be unlikely if patients have received ventilatory support for several days or more · Beware of secondary infections and bowel perforation
Baricitinib	4 mg for 14 days or until hospital discharge	<ul style="list-style-type: none"> · Hospitalized patients requiring respiratory support · Should not be used for whom do not require oxygen support and for the COVID-19 prophylaxis 	<ul style="list-style-type: none"> · 2 mg once daily for patients with estimated glomerular filtration rate of less than 60 ml/min · Greater efficacy if combined with corticosteroids · Beware of increased risk of thromboembolic events · Beware of secondary infections, particularly in patients who co-administered corticosteroids · Might increase the risk of a worse outcome if used for prophylaxis
Intravenous immunoglobulin (IVIg)	<ul style="list-style-type: none"> · 1–2 g per kg of body weight for up to 4 days, · MIS-C 2 g/kg single day along with pulse methyl prednisolone 	<ul style="list-style-type: none"> · MIS-C & MIS-A patients · Refractory patients that meet WHO definition of critical COVID-19 	<ul style="list-style-type: none"> · Low quality of evidence for its efficacy · Limited drug supply and extensive cost limits its use · Cardiac function and fluid status should be assessed before infusion. If abnormal, the rate of IVIg infusion may be slowed, or the treatment may be given in divided doses over 2 days, and/or diuretics may be considered to avoid volume overload · Increased risk of hemolytic anemia · A second dose of IVIg is not recommended in patients with refractory MIS-C

CRP: C reactive protein, HLH: Hemophagocytic lymphohistiocytosis, IV: intravenous, MIS-A: Multisystem Inflammatory Syndrome in Adults, MIS-C: Multisystem Inflammatory Syndrome in Children, Subc: subcutaneous, PCR: polymerase chain reaction, PO: per oral, WHO: World Health Organization.

proinflammatory cytokines and activation of immune cells, including T cells, monocytes and macrophages. Several well-designed clinical trials investigated the role of corticosteroids on various stages of the disease with different application regimens.

The largest corticosteroid trial is the open-label RECOVERY trial in which hospitalized COVID-19 patients were randomized to receive either standard of care (SOC, n=4321) or 6 mg/day dexamethasone (n=2104) for up to 10 days [38]. Dexamethasone reduced 28-day mortality rate by 20% (age-adjusted rate ratio 0.8, 95% CI 0.75–0.93) in patients requiring respiratory support as nasal, high-flow oxygen or invasive/noninvasive mechanical ventilation. However, a trend towards harm was noted with the use of dexamethasone in those patients who did not need respiratory support at the time of randomization. Therefore, guidelines¹ do not recommend the use of corticosteroids in patients with nonsevere COVID-19 (SpO₂ >94%, not requiring supplemental oxygen). Studies regarding the effects of corticosteroids on viral clearance are conflicting, but there is a possibility that early administration of corticosteroids in nonseverely ill patients might be deleterious due to an increase of viral shedding or a delay in viral clearance [39, 40]. There is also evidence showing that impaired viral clearance is associated with severe disease and increased age rather than early use of corticosteroids [40].

Efficacy of intravenous dexamethasone was also investigated in COVID-19 patients with moderate to severe ARDS who required ICU admission. The CoDEX trial randomly allocated patients to receive either SOC alone (n=148) or 20 mg daily dexamethasone intravenously for 5 days, 10 mg of dexamethasone daily for 5 days or until ICU discharge plus SOC (n=151)[3, 41]. Dexamethasone provided more ventilator-free days during the first 28 days (6.6; 95% CI, 5.0–8.2 days vs. 4.0; 95% CI, 2.9–5.4 days; difference 2.26; 95% CI, 0.2–4.38 days) compared with SOC. However, all-cause mortality at 28 days was found similar between groups (56.3% in the dexamethasone group vs 61.5% the SOC group; hazard ratio, 0.97; 95% CI, 0.72–1.31). Dexamethasone was well-tolerated without increasing the rate of serious infections or hyperglycemia [3, 41].

Despite there were numerous clinical trials and several meta-analyses, the optimal corticosteroid type, dose, and treatment duration in patients with COVID-19 remains unclear. Dexamethasone is preferred over other corticosteroids for its shown efficacy in HLH but there are several other trials reporting positive results with the use of other agents like methylprednisolone in equivalent total daily doses of dexamethasone 6 mg/day (methylprednisolone 32 mg daily or prednisone 40 mg

daily) [42–44]. Methylprednisolone 250 to 500 mg daily pulses for 3–6 consecutive days has also been shown to improve survival in severe COVID-19 patients [45, 46]. Hydrocortisone would be one of the options although its efficacy seemed to be lower than dexamethasone[46]. The open-label STOIC trial showed that, an inhaled steroid, budesonide 400 µg twice a day until symptom resolution, decreased the need for urgent medical care in mild patients with early COVID-19 [47].

Another possible role of corticosteroids would be the treatment of post-COVID inflammatory lung diseases, as suggested by preliminary data from small observational studies that report symptomatic and radiological improvement in a subset of patients with organizing pneumonia and lung fibrosis [48].

4.2. Chloroquine and hydroxychloroquine

Chloroquine (CQ) and hydroxychloroquine (HCQ) are 4-aminoquinoline derivatives that are approved by the U.S. Food and Drug Administration (FDA) for the treatment of systemic lupus erythematosus, rheumatoid arthritis (RA), and they have been used for these disorders for decades. HCQ does not increase the risk of infection and has lipid-lowering, antithrombotic and antineoplastic properties [49]. CQ and HCQ may prolong QT interval constituting potential risk of fatal arrhythmia in higher doses or if combined with QT prolonging medications as well as those with cardiac diseases [50].

It has been known that CQ and HCQ have nonspecific antimicrobial activity against *Coxiella burnetii*, plasmodium, and many viruses, including hepatitis B, HIV, H1N1, and Zika virus [51]. Therefore, it had rapidly attracted attention at the beginning of COVID-19 pandemic. However, several randomized controlled trials clearly demonstrated that HCQ neither protected nor improved the outcomes of COVID-19 patients [52], leaving some important messages for the future pandemics. First, some trials showed a trend towards harm with use of HCQ [53]. Second, massive demand for HCQ caused drug shortage leading to the flare of disease in patients with lupus. Third, in-vitro studies and animal models do not necessarily mean in vivo efficacy of drugs when used for unapproved indications.

4.3. Intravenous immunoglobulin (IVIg)

IVIg is a blood product containing polyclonal immunoglobulin G isolated and pooled from healthy donors used to treat Immune Thrombocytopenic Purpura (ITP), Kawasaki disease and various inflammatory neurologic and myositis syndromes. It has immunomodulatory functions with unknown mechanisms of action. One of the proposed mechanisms is the interaction of IgG-Fc with Fc gamma receptors located on almost all immune

¹ <https://www.idsociety.org/covid-19-real-time-learning-network/therapeutics-and-interventions/corticosteroids/>

cells, resulting in pleiotropic functional consequences including the expansion of regulatory T cell population, phagocytosis, antibody-dependent cellular cytotoxicity (ADCC), immune cell differentiation and maturation, apoptosis, expression of pro-inflammatory cytokines, chemokines, and antigen-presentation [54, 55].

Studies on efficacy of IVIg are largely retrospective cohort studies, and IVIg was utilized as a rescue medication following other medications like corticosteroids and tocilizumab [56, 57]. Prospective RCTs are scarce and included a small number of patients with conflicting results [56, 58, 59]. Metaanalyses of published studies suggest survival advantage in only critical COVID-19 patients characterized by ARDS, sepsis, septic shock that require intensive care admission [60]. Moreover, its costliness and limited supply restrict its general use. The methodological limitations of all the studies preclude the extraction of any definitive conclusion about when and how to use IVIg in the context of COVID-19. However, patients with transplantation, pregnancy, secondary infections, immune thrombocytopenia, muscular, myocardial, and neurologic manifestations would be suitable candidates for IVIg treatment. It should be remembered that the efficacy of IVIg comes from immunomodulatory actions rather than viral neutralizing actions [61] which can be obtained at higher doses than neutralizing doses such as 1 g per kg body weight daily for up to 4 consecutive days [62].

American College of Rheumatology recommends the use of IVIg as the first-line treatment in patients complicated with Multisystem Inflammatory Syndrome in Children and Adults (MIS-C and MIS-A). In these patients IVIg is given 2 gr of per kg body weight for a single dose along with 1–2 mg/kg methylprednisolone, aspirin, and anti-coagulants [63–66].

4.4. IL-6 antagonists

IL-6 receptors are expressed in almost all immune cells, and IL-6 acts as a master player inducing proliferation and differentiation of immune cells. IL-6 exerts pleiotropic effects on immune cells, which are manifested as promoted differentiation of T-helper type 17 (Th17), CD8⁺ T, and B cells, increased migration of neutrophils, and reduced development of Tregs. Moreover, IL-6 modulates endothelial cells and vascular smooth muscle cells leading to increased vascular permeability and leakage, endothelial activation, and accelerated atherogenesis, which are manifested clinically as cardiovascular events, hypotension, and pulmonary dysfunction in COVID-19. Collectively, IL-6 promotes both immune cell hyperactivation and target organ dysfunction in severe COVID-19 [67].

In healthy individuals, the IL-6 levels in circulation are extremely low and are in the range of 1–5 pg/mL,

marked elevations reported in many inflammatory conditions including cytokine release syndrome [68]. Several therapeutic agents have been developed to inhibit the cytokine itself (sirikumab, siltuximab, clazakizumab, and olokizumab), the signaling via the IL-6 receptor (tocilizumab, sarilumab, levilimab), or its postreceptor downstream signaling pathways (JAK/STAT). Tocilizumab (TCZ) is approved for the treatment of rheumatoid arthritis (RA), juvenile idiopathic arthritis, giant cell arteritis, cytokine release syndrome, idiopathic multicentric Castleman's disease (iMCD)².

COVID-19 patients have high plasma IL-6 levels, especially those with a more severe disease presentation [27]. IL-6 production can be stimulated by SARS-CoV-2 itself or by stimulation of other immune cells [69]. Indeed, it has been shown that, during COVID-19, CD4⁺T lymphocytes are rapidly activated to differentiate into pathogenic Th1 cells, generating GM-CSF and other pro-inflammatory cytokines, which further induce activation of monocytes with high expression of IL-6 [70]. In clinical point of view, there is striking correlation between serum IL-6 levels and SARS-CoV-2 RNAemia, which strongly indicates worse outcome [71]. Hence, blocking IL-6 would potentially reduce the detrimental immune response caused by SARS-CoV-2. There are dozens of clinical trials assessing the efficacy of IL-6 antagonists in patients hospitalized for COVID-19 that variously reported benefit, no effect or harm, probably due to different study designs and enrolled patient characteristics.

A recent metaanalysis evaluated 27 clinical trials performed on almost 11,000 patients with diverse disease characteristics at different dosages of IL-6 antagonists, either alone or combined with corticosteroids [72]. IL-6 antagonist use, compared to usual care or placebo, was associated with lower 28-day all-cause mortality, lower progression to invasive mechanical ventilation (IMV), or death without increasing the risk of infection at the same time period. Concurrent administration of IL-6 antagonists and corticosteroids at randomization provided greater benefit than either drug given individually. The association of IL-6 antagonists with lower mortality was more marked in those patients requiring oxygen flow rate of ≤ 15 L/min and noninvasive ventilation and who did not receive IMV at randomization (70).

TCZ is the most widely studied IL-6 antagonist for the treatment of severe COVID-19, and its efficacy seemed to be more prominent than sarilimumab, while data are limited for other IL-6 antagonists [72]. Similarly, due to limited data, associations could not be determined for the efficacy of low-dose IL-6 antagonists, those who did not require respiratory support, use of concomitant antiviral

² FDA. Actemra (tocilizumab) injection, for intravenous or subcutaneous use: highlights of prescribing information.2010.www.accessdata.fda.gov/drug-satfda_docs/label/2017/125276s114lbl.pdf

treatment, and the risk of secondary infections in advance. Hence, it is prudent to monitor patients for the incidence of secondary/opportunistic infections and the other well-known complications such as bowel perforation. After accumulated data, FDA issued an emergency use authorization for the tocilizumab for the treatment of hospitalized adults and pediatric (≥ 2 years of age) COVID-19 patients who are requiring respiratory support and receiving systemic corticosteroids in June 2021.

4.5. Janus kinase (JAK) inhibitors

JAK inhibitors are potent inhibitors of one or more of the JAK family of enzymes (JAK1, JAK2, JAK3, TYK2), interfering with the JAK-STAT signaling pathway, which mediates the effects of many interleukins (IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, IL-12, IL-15, IL-21, IL-23), IFN-(α , β , γ) and growth factors (GM-CSF, TGF- β , erythropoietin and thrombopoietin) [73]. Like the other respiratory viruses, SARS-CoV-2 triggers inflammation via the JAK/STAT pathway leading to the recruitment of inflammatory cells, including macrophages, monocytes, lymphocytes, natural killer cells, and dendritic cells, as well as activation of pneumocytes and endothelial cells [74].

JAK inhibitors are currently approved for the treatment of RA, psoriatic arthritis, ulcerative colitis, polyarticular juvenile arthritis, and their use in other inflammatory disorders are continuously expanding [75]. Idea in COVID-19 treatment is that many pro-inflammatory cytokines involved in cytokine storm of COVID-19 might be inhibited by a single agent, JAK inhibitors (tofacitinib, baricitinib, ruxolitinib). Besides, these shared properties of JAK inhibitors, baricitinib blocks host Numb-associated kinases (AP-2-associated protein kinase 1, AAK1; cyclin G-associated kinase, GAK; BMP2 inducible kinase, BIKE), which all mediate viral entry [76]. This effect is only restricted to baricitinib among other JAK inhibitors with theoretic potential of blocking viral entry and assembly of virus particles into pneumocytes in therapeutic doses used to treat RA [77]. Given this hypothesis, baricitinib was tested in a large multi-center RCT, the Adaptive COVID-19 Treatment Trial 2 (ACTT-2)[78]. In this trial, a combination of remdesivir with baricitinib (4 mg/day for 14 days or until hospital discharge) showed improved outcomes (days for recovery, clinical status at day 15, and the 28-day mortality) compared to remdesivir used alone in hospitalized COVID-19 patients [78]. These clinical benefits were consistent across different age groups, sexes, and ethnic groups, as well as independent of the duration of symptoms and severity of disease at enrollment, though benefits of baricitinib combination were greater in moderate to severe COVID-19 patients requiring respiratory support as nasal, high flow oxygen or noninvasive ventilation. Although combination did not reveal a significant 28-day mortality difference (5.1% with remdesivir plus baricitinib vs. 7.8% with remdesivir alone), the composite

outcome progression to mechanical ventilation and death was prevented in one third of patients (HR 0.69; 95% CI, 0.50–0.95). Moreover, remdesivir plus baricitinib reduced the mechanical ventilation or extracorporeal membrane oxygenation (ECMO) requirements by median of 11 days among patients who required these interventions after enrollment [78]. With the results of this trial, FDA issued an emergency use authorization for remdesivir plus baricitinib in hospitalized patients with COVID-19 who required oxygen supplementation in November 2020.

COV-BARRIER study investigated the role of baricitinib on a heterogeneous group of patients [79]. Patients were randomly assigned to receive either baricitinib 4 mg 14 days (n=764) or placebo (n=761) on top of standard of care determined by local clinical practice. Only 20% of patients co-administered remdesivir and more than 90% of patients received concomitant dexamethasone. Although baricitinib did not reach the composite primary endpoint (the proportion of patients who progressed to high-flow oxygen, noninvasive ventilation, invasive mechanical ventilation, or death by day 28), the 28-day all-cause mortality was 8% for baricitinib and 13% for placebo (HR 0.57; 95% CI, 0.41–0.78) revealing a 38% relative reduction in mortality. Consistently, the 60-day all-cause mortality was 10% for baricitinib vs 15% for placebo (HR 0.62; 95% CI, 0.47–0.83). According to intention to treat analysis, one additional death was prevented per 20 baricitinib-treated patients, a number like ACTT-2 trial. Benefits were consistent across different age groups, sexes, and ethnic groups, duration of symptoms and baseline corticosteroid use. Observed benefits were greater in patients with severe disease and who did not receive remdesivir at baseline. After the result of this trial, FDA has revised the Emergency Use Authorization, authorizing baricitinib alone for the treatment of COVID-19 in hospitalized adults and pediatric patients (≥ 2 years of age), requiring respiratory support or ECMO in July 2021 [79].

Efficacy of another JAK inhibitor, tofacitinib in COVID-19 pneumonia was investigated an industry-supported trial in 289 patients in Brazil (STOP-COVID) [80]. Tofacitinib was administered 10 mg twice daily for up to 14 days or until hospital discharge to patients requiring respiratory support. Although the STOP-COVID trial was not powered to detect a difference in mortality, it has reached the composite primary endpoint, which was the reduction of death or respiratory failure at 28 days (18.1% of the tofacitinib vs 29.0% of the placebo, risk ratio 0.63; 95% CI, 0.41–0.97). Of note, while 20% of the patients were co-administered corticosteroids in the ACTT-2 trial, almost 90% of patients received them in STOP-COVID trial, suggesting the superiority of combination. Studies for another JAK inhibitor, ruxolitinib, are ongoing and preliminary results are promising [81].

JAK inhibitors increase the risk of secondary infections and thromboembolic events, particularly in patients with underlying cardiovascular disease. However, compared to placebo, baricitinib and tofacitinib had a significantly lower incidence of severe/nonsevere adverse events including infections [78, 80], though patients who received concomitant glucocorticoids after randomization had a higher rate of incident infections [78]. COVID-19 Global Rheumatology Alliance reported a two-fold increased risk of mechanical ventilation and death in RA patients who were under treatment with JAK inhibitors compared to TNF inhibitors [82]. In the same study, investigators also find more than 4-fold increased risk of worse outcomes for rituximab, whereas IL-6 inhibitors and abatacept did not increase adverse outcomes compared to anti-TNF users [78]. Hence, timing of JAK-inhibitor initiation during COVID-19 might have pivotal importance in determining the outcome. During initial phases of SARS-CoV-2 infection, symptomatic patients present with impaired type I/III IFN-mediated antiviral responses. Critically ill COVID-19 patients show genetic polymorphisms in one IFN receptor gene (IFNRA2) and in a gene locus near the TYK2, which is the key for IFN, interleukin (IL)-12 and IL-23 signaling, and Th1/Th17 cell-mediated antiviral immune responses [83]. Pretreatment with JAK inhibitors might blunt IFNs' antiviral responses increasing risk of severe infection; therefore, these group of drugs should not be used for prophylaxis or those not requiring respiratory support [83]. In patients with moderate to severe SARS-CoV-2 pneumonia, combination of baricitinib with corticosteroids provided greater improvement in pulmonary functions than corticosteroids alone; therefore, it would be better to co-administer JAK inhibitors with corticosteroids [80, 84].

4.6. IL-1 inhibitors: anakinra and canakinumab

IL-1 family are pleiotropic cytokines, have roles in inflammation, hematopoiesis, and fibrosis. IL-1 β and TNF- α promote vascular permeability and leakage. Both IL-1 β and IL-18 fuel cytokine storm and MAS and IL-1 cytokines (except IL-18) can be successfully inhibited by IL-1 antagonists [85]. Nod-like receptor family pyrin domain-containing 3 (NLRP3) is a critical inflammasome in the acute protection of the body against a wide variety of noxious stimuli, including RNA viruses [86]. SARS-COV-2 has been shown to induce NLRP3 by its ion channel-forming M protein and ORF8b [87] and activates caspase-1, a molecule responsible for the activation and exuberant release of IL-1 β and IL-18 [26, 27, 88].

Anakinra³ is a recombinant antagonist of human IL-1 and approved for the treatment of RA and certain autoinflammatory disorders with recommended doses of 1–2 mg/kg/day with a maximum daily dose of 8 mg/kg. In

³ FDA. Kineret® (anakinra) for injection, for subcutaneous use: Highlights Of prescribing Information. 2001. [https://www.accessdata.fda.gov/drug-](https://www.accessdata.fda.gov/drug-satfda_docs/label/2012/103950s51361bl.pdf)

terms of sepsis and MAS, the use of extremely high doses of anakinra (2 mg/kg/h for 72 h continuous infusion) was shown to be safe [89]. Because of its proven safety, anakinra is one of the first studied immunomodulatory treatments showing benefit when it's used at higher doses but not with low doses [90]. Currently, there are several anakinra studies registered for COVID-19 and only one RCT; the CORIMUNO-ANA-1 trial has published results [91]. The rest of the available data come from low-quality studies with a significant risk of bias. The CORIMUNO-ANA-1 trial included mild-to-moderate COVID-19 inpatients who required at least 3 L/min of oxygen but not mechanical ventilation and elevated CRP (≥ 25 mg/L). Patients (n=116) were 1:1 randomized to receive either SOC or SOC plus intravenous anakinra (200 mg twice a day on days 1–3, 100 mg twice on day 4, and 100 mg once on day 5, a similar dose used to treat critically ill patients with HLH. The trial was early terminated following an interim analysis that showed any evidence to support anakinra on clinical improvement, ventilator support and death over SOC alone. However, this trial is criticized for use of corticosteroids in more than half of patients in both arms that might shadow efficacy of anakinra. A recent metaanalysis revealed improved survival in patients treated with anakinra (38 deaths in 342) than in those who received SOC (137 deaths in 553; adjusted OR 0.32 95%CI, 0.20–0.51) [92]. Subgroup analyzes showed that anakinra was more effective in reducing mortality in patients with CRP concentrations of >100 mg/L and when given without dexamethasone, but it was not effective when co-administered with dexamethasone. Metaanalysis also reported a nonsignificant increase in the risk of adverse events with anakinra [92]. The SAVE study determined high-risk patients, which potentially progress into severe respiratory failure by measuring serum soluble urokinase plasminogen activator receptor (suPAR) levels as an early indicator for clinical worsening. Patients were treated with 100 mg/day subcutaneous anakinra for 10 days if they had suPAR of >6 ng/mL reporting improved respiratory and survival outcomes suggesting superior efficacy when administered early [93, 94]. The other IL-1 β antagonist canakinumab (450–750 mg single iv infusion) was failed to show efficacy in improving survival without IMV [95].

4.7. Colchicine

Colchicine has multiple effects on the function of the immune system, particularly on neutrophils, including inhibition of tubulin polymerization and microtubule generation, chemotaxis, superoxide anion production, suppression of cellular adhesion molecules, inflammatory chemokines, and cytokines (TNF- α and IL-6) as well as NLRP3 inflammasome activation that mediates

interleukin 1 β production [96]. Colchicine is approved for the treatment of gout and familial Mediterranean fever (FMF) but has attracted attention in recent years for the management of atherosclerotic cardiovascular disease and idiopathic pericarditis [97]. Beyond gout and FMF, given its anti-inflammatory/immuno-modulatory properties and good safety profile, colchicine is widely used to treat a variety of inflammatory conditions. Several studies investigated the efficacy of colchicine in COVID-19, yielding conflicting results.

The first small RCT (n=36 in both arms) by Lopes et al, suggested superiority of colchicine on reduction in length of both supplemental oxygen therapy and hospitalization compared to placebo in patients with moderate to severe COVID-19 but fail to show survival benefit due to the low number of deaths in both groups [98]. The COLCORONA trial included outpatients, diagnosed within 24 h of randomization, over 40 years with at least one high-risk adverse prognostic feature to determine the composite endpoint of death or hospital admission. Patients were 1:1 randomized to receive either colchicine (n=2235) 0.5 mg twice per day for 3 days followed by once per day for 27 more days or placebo (n=2253) [99]. The primary endpoint evolved in 104 (4.7%) in the colchicine group and 131 (5.8%) in the placebo group (OR: 0.79, 95% CI, 0.61-1.03; p=0.08), reaching significance when only PCR confirmed 4159 cases were included (OR 0.75, 95%CI, 0.57–0.99; p=0.042). Death was occurred in 5 (0.2%) and 9 (0.4%) patients (OR 0.56, 95%CI, 0.19–1.66) in colchicine and placebo groups, respectively. In RECOVERY, colchicine arm of the trial was terminated, since the colchicine failed to show a significant difference in the primary endpoint of the 28-day mortality rate of colchicine vs. SOC alone (21% in both colchicine and SOC arms; risk ratio 1.01, 95%CI: 0.93-1.1; P=0.77) [100].

Metaanalyses of observational and RCTs showed an advantage of colchicine use; however, subgroup analysis with randomized controlled trials showed no statistically significant difference in the mortality (OR: 0.80, 95%CI, 0.44–1.46) [101]. Moreover, colchicine neither prevented COVID-19-related hospital admissions nor disease course among rheumatic patients who were receiving it for their underlying diseases [102, 103]. Colchicine has potential toxicities, particularly in those with kidney and liver impairment and elders, myelosuppression, elevation in transaminases, and myopathy, which are commonly observed during the course of COVID-19 [97, 104]. Therefore, routine use of colchicine is not recommended until well designed trials show its clear benefit on specific populations.

4.8. Anti-TNF agents

TNF- α is one of the most potent pro-inflammatory cytokines with a broad spectrum of actions. TNF- α is

produced by macrophages, monocytes, and T cells that promote expression or other inflammatory cytokines via NF κ B pathway. Marked elevations were reported in many inflammatory conditions including cytokine release syndrome. [26]. SARS-CoV viral spike protein can modulate TNF- α -converting enzyme (TACE)-dependent shedding of the ACE2 ectodomain, required for the viral entry which is coupled to TNF- α production [105]. However, studies assessing serum TNF- α levels in COVID-19 patients revealed conflicting results. Some studies found that it elevated, whereas some others did not [67]. High serum TNF concentrations at the time of admission, thus, at very early phase of infection, predicted poor outcomes and found significantly elevated in severe COVID patients [106]. Therefore, it's hypothesized that the use of TNF inhibitors might be effective in blocking viral entry and detrimental effects of exuberant TNF- α , as shown in preclinical studies on severe respiratory syncytial virus and influenza infections [106, 107]. The COVID-19 Global Rheumatology Alliance (GRA) registry showed fewer hospitalizations among people treated with TNF antagonists than non-TNF inhibitor biologic/JAK inhibitor users (adjusted OR 0.40, 95%CI, 0.19–0.81) that indicate the potential favorable role of anti-TNFs [82].

Since TNF fuels other potent pro-inflammatory cytokines, anti-TNF therapy might be more effective when applied early in the disease course to block excessive production of TNF. Intravenous infliximab and subcutaneous adalimumab with loading doses might be preferred over other anti-TNFs. Although there are few observational reports regarding the successful use of anti-TNFs, the limitations of observational data need to be considered when translating these findings to clinical practice. Trials on the use of anti-TNFs in COVID-19 such as the “adalimumab in COVID-19 to prevent respiratory failure in community care, (AVID-CC)” are ongoing, but none of them have published results.

4.9. Anti-IL-17 antagonists

One of the cytokines found abundant in COVID-19 patients is IL-17 and reported to be associated with severe lung inflammation [26]. IL-17 has wide-ranging pro-inflammatory effects on induction of cytokines; IL-1 β , IL-6, TNF- α ; growth factors, G-CSF; chemokines; and matrix metalloproteinases. IL-17 inhibitors did not increase the risk of progressive COVID-19 [108]. For the use of IL-17 antagonists in COVID-19, there are very scarce data, limited to a recent study from Russia, which fail to show clinical benefit of netakimab [109].

4.10. GM-CSF inhibitors

GM-CSF is one of the key molecules involved in cytokine storm which is excessively released in COVID-19 patients [70]. GM-CSF is crucial for driving both innate and adaptive immune responses and blockade of this growth

factor may halt immunopathology caused by the virus. Mavrilimumab is a GM-CSF inhibitor developed for the refractory RA [110]. Studies evaluating the efficacy of mavrilimumab revealed conflicting results [41, 111]. Lenzilumab is another GM-CSF inhibitor, which show promise in the treatment of severe refractory COVID-19 [112].

5. Conclusion

Excessive inflammatory response with features of cytokine storm causes severe disease course and adversely affects the outcome of COVID-19. Successful vaccination

campaigns and antiviral drugs in development programs would be the most effective tools in the fight against the COVID-19. Larger trials are ongoing, and their results are urgently needed to ascertain the most effective treatment options, and some of such novel agents show promise in their early clinical studies. Until the discovery of curative medications, drugs that are used in daily rheumatology practice constitute life-saving treatment options in COVID-19 patients with an extenuating severe detrimental inflammatory response. Herein, we discussed altered immune responses in COVID-19 and presented current treatment options in the treatment of severe COVID-19.

References

1. Kuri-Cervantes L, Pampena MB, Meng W, Rosenfeld AM, Ittner CAG et al. Comprehensive mapping of immune perturbations associated with severe COVID-19. *Science Immunology*. 2020;5(49). doi:10.1126/sciimmunol.abd7114
2. Xu Z, Shi L, Wang Y, Zhang J, Huang L et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet Respiratory Medicine*. 2020. doi:10.1016/s2213-2600(20)30076-x
3. Tomazini BM, Maia IS, Cavalcanti AB, Berwanger O, Rosa RG et al. Effect of Dexamethasone on Days Alive and Ventilator-Free in Patients With Moderate or Severe Acute Respiratory Distress Syndrome and COVID-19: The CoDEX Randomized Clinical Trial. *Journal of the American Medical Association*. 2020;324(13):1307-1316. doi:10.1001/jama.2020.17021
4. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*. 2020 ;382(18):1708-1720. doi:10.1056/NEJMoa2002032
5. Gustine JN, Jones D. Immunopathology of Hyperinflammation in COVID-19. *American Journal of Pathology*. 2021;191(1):4-17. doi:10.1016/j.ajpath.2020.08.009
6. Schultze JL, Aschenbrenner AC. COVID-19 and the human innate immune system. *Cell*. 2021;184(7):1671-1692. doi:10.1016/j.cell.2021.02.029
7. Yang, Y., Shen, C., Li, J., Yuan, J., Wei, J. et al. Plasma IP-10 and MCP-3 levels are highly associated with disease severity and predict the progression of COVID-19. *The Journal of allergy and clinical immunology*. 2020; 146 (1), 119–127.e4. doi.org/10.1016/j.jaci.2020.04.027
8. Ashour HM, Elkhatib WF, Rahman MM, Elshabrawy HA. Insights into the Recent 2019 Novel Coronavirus (SARS-CoV-2) in Light of Past Human Coronavirus Outbreaks. *Pathogens*. 2020;9(3). doi:10.3390/pathogens9030186
9. Ziebuhr J, Snijder EJ, Gorbalenya AE. Virus-encoded proteinases and proteolytic processing in the Nidovirales. *Journal of General Virology*. 2000;81(Pt 4):853-879. doi:10.1099/0022-1317-81-4-853
10. Baez-Santos YM, St John SE, Mesecar AD. The SARS-coronavirus papain-like protease: structure, function and inhibition by designed antiviral compounds. *Antiviral Research*. 2015;115:21-38. doi:10.1016/j.antiviral.2014.12.015
11. Lu X, Pan J, Tao J, Guo D. SARS-CoV nucleocapsid protein antagonizes IFN-beta response by targeting initial step of IFN-beta induction pathway, and its C-terminal region is critical for the antagonism. *Virus Genes*. 2011;42(1):37-45. doi:10.1007/s11262-010-0544-x
12. Ujike M, Taguchi F. Incorporation of spike and membrane glycoproteins into coronavirus virions. *Viruses*. 2015;7(4):1700-1725. doi:10.3390/v7041700
13. Masters PS. The molecular biology of coronaviruses. *Advances in Virus Research*. 2006;66:193-292. doi:10.1016/s0065-3527(06)66005-3
14. Neuman BW, Kiss G, Kunding AH, Bhella D, Baksh MF et al. A structural analysis of M protein in coronavirus assembly and morphology. *Journal of Structural Biology*. 2011;174(1):11-22. doi:10.1016/j.jsb.2010.11.021
15. Schoeman D, Fielding BC. Coronavirus envelope protein: current knowledge. *Virology Journal*. 2019;16(1):69. doi:10.1186/s12985-019-1182-0
16. Nieto-Torres JL, DeDiego ML, Verdía-Báguena C, Jimenez-Guardeño JM, Regla-Nava JA et al. Severe acute respiratory syndrome coronavirus envelope protein ion channel activity promotes virus fitness and pathogenesis. *PLoS Pathogens*. 2014;10(5):e1004077. doi:10.1371/journal.ppat.1004077
17. Xia S, Zhu Y, Liu M, Lan Q, Xu W et al. Fusion mechanism of 2019-nCoV and fusion inhibitors targeting HR1 domain in spike protein. *Cellular & Molecular Immunology*. 2020. doi:10.1038/s41423-020-0374-2
18. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *bioRxiv*. 2020:2020.2001.2022.914952. doi:10.1101/2020.01.22.914952
19. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. 2020. doi:10.1016/j.cell.2020.02.052

20. Ou X, Liu Y, Lei X, Li P, Mi D et al. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nature Communications*. 2020;11(1):1620. doi:10.1038/s41467-020-15562-9
21. Ivashkiv LB, Donlin LT. Regulation of type I interferon responses. *Nature Reviews Immunology*. 2014;14(1):36-49. doi:10.1038/nri3581
22. Li G, Fan Y, Lai Y, Han T, Li Z et al. Coronavirus infections and immune responses. *Journal of Medical Virology*. 2020;92(4):424-432.
23. George MR. Hemophagocytic lymphohistiocytosis: review of etiologies and management. *Journal of Blood Medicine*. 2014;5:69-86. doi:10.2147/jbm.s46255
24. Ramos-Casals M, Brito-Zeron P, Lopez-Guillermo A, Khamashta MA, Bosch X. Adult haemophagocytic syndrome. *Lancet*. 2014;383(9927):1503-1516. doi:10.1016/s0140-6736(13)61048-x
25. McGonagle D, Sharif K, O'Regan A, Bridgewood C. Interleukin-6 use in COVID-19 pneumonia related macrophage activation syndrome. *Autoimmunity Reviews*. 2020:102537. doi:10.1016/j.autrev.2020.102537
26. Huang C, Wang Y, Li X, Ren L, Zhao J et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. doi:10.1016/s0140-6736(20)30183-5
27. Qin C, Zhou L, Hu Z, Zhang S, Yang S et al. Dysregulation of immune response in patients with COVID-19 in Wuhan, China. *Clinical Infectious Diseases*. 2020. doi:10.1093/cid/ciaa248
28. Liu Y, Yang Y, Zhang C, Huang F, Wang F et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Science China Life Sciences*. 2020 Mar;63(3):364-374. doi: 10.1007/s11427-020-1643-8.
29. Özger HS, Aysert Yıldız P, Gaygısız Ü, Uğraş Dikmen A, Demirbaş Gülmez Z et al. The factors predicting pneumonia in COVID-19 patients: preliminary results from a university hospital in Turkey. *Turkish Journal of Medical Sciences*. 2020 ;50(8):1810-1816. doi: 10.3906/sag-2005-385.
30. Wang F, Nie J, Wang H, Zhao Q, Xiong Y et al. Characteristics of peripheral lymphocyte subset alteration in COVID-19 pneumonia. *Journal of Infectious Diseases*. 2020; 221 (11) : 1762-1769. doi:10.1093/infdis/jiaa150
31. Crayne CB, Albeituni S, Nichols KE, Cron RQ. The Immunology of Macrophage Activation Syndrome. *Frontiers in Immunology*. 2019;10:119. doi:10.3389/fimmu.2019.00119
32. Wang W, Liu X, Wu S, Chen S, Li Y et al. Definition and Risks of Cytokine Release Syndrome in 11 Critically Ill COVID-19 Patients With Pneumonia: Analysis of Disease Characteristics. *Journal of infectious diseases*. 2020; 222(9):1444-1451. doi. org/10.1093/infdis/jiaa387
33. Zinkernagel RM. Immunology taught by viruses. *Science*. 1996;271(5246):173-178. doi:10.1126/science.271.5246.173
34. Sakaguchi S, Yamaguchi T, Nomura T, Ono M. Regulatory T cells and immune tolerance. *Cell*. 2008;133(5):775-787. doi:10.1016/j.cell.2008.05.009
35. Sakaguchi S, Miyara M, Costantino CM, Hafler DA. FOXP3+ regulatory T cells in the human immune system. *Nature Reviews Immunology*. 2010;10(7):490-500. doi:10.1038/nri2785
36. Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature*. 1999;401(6754):708-712. doi:10.1038/44385
37. Gianfrancesco M, Hyrich KL, Al-Adely S, Carmona L, Danila MI et al. Characteristics associated with hospitalisation for COVID-19 in people with rheumatic disease: data from the COVID-19 Global Rheumatology Alliance physician-reported registry. *Annals of Rheumatic Diseases*. 2020;79(7):859-866. doi:10.1136/annrheumdis-2020-217871
38. RECOVERY Collaborative Group. Dexamethasone in Hospitalized Patients with Covid-19. *New England Journal of Medicine*. 2021;384(8):693-704. doi:10.1056/NEJMoa2021436
39. Romanou V, Koukaki E, Chantziara V, Stamou P, Kote A et al. Dexamethasone in the Treatment of COVID-19: Primus Inter Pares? *Journal of Personalized Medicine* 2021;11(6). doi:10.3390/jpm11060556
40. Spagnuolo V, Guffanti M, Galli L, Poli A, Querini PR et al. Viral clearance after early corticosteroid treatment in patients with moderate or severe covid-19. *Scientific Reports* 2020;10(1):21291. doi:10.1038/s41598-020-78039-1
41. De Luca G, Cavalli G, Campochiaro C, Della-Torre E, Angelillo P et al. GM-CSF blockade with mavrilimumab in severe COVID-19 pneumonia and systemic hyperinflammation: a single-centre, prospective cohort study. *Lancet Rheumatology*. 2020;2(8) :e465-e473. doi:10.1016/S2665-9913(20)30170-3
42. Fatima SA, Asif M, Khan KA, Siddique N, Khan AZ. Comparison of efficacy of dexamethasone and methylprednisolone in moderate to severe covid 19 disease. *Annals of Medicine and Surgery (Lond)*. 2020;60:413-416. doi:10.1016/j.amsu.2020.11.027
43. Du Plessis EM, Lalla U, Allwood BW, Louw EH, Nortje A et al. Corticosteroids in critical COVID-19: are all corticosteroids equal? *The South African Medical Journal*. 2021;111(6):550-553.
44. WHO Rapid Evidence Appraisal for COVID-19 Therapies (REACT) Working Group. Association Between Administration of Systemic Corticosteroids and Mortality Among Critically Ill Patients With COVID-19: A Meta-analysis. *Journal of the American Medical Association*. 2020;324(13):1330-1341. doi:10.1001/jama.2020.17023
45. Edalatfard M, Akhtari M, Salehi M, Naderi Z, Jamshidi A et al. Intravenous methylprednisolone pulse as a treatment for hospitalised severe COVID-19 patients: results from a randomised controlled clinical trial. *European Respiratory Journal*. 2020;56(6). doi:10.1183/13993003.02808-2020

46. Cui Y, Sun Y, Sun J, Liang H, Ding X et al. Efficacy and Safety of Corticosteroid Use in Coronavirus Disease 2019 (COVID-19): A Systematic Review and Meta-Analysis. *Infectious Diseases and Therapy*. 2021. doi:10.1007/s40121-021-00518-3
47. Ramakrishnan S, Nicolau DV, Jr., Langford B, Mahdi M, Jeffers H et al. Inhaled budesonide in the treatment of early COVID-19 (STOIC): a phase 2, open-label, randomised controlled trial. *Lancet Respiratory Medicine*. 2021;9(7):763-772. doi:10.1016/s2213-2600(21)00160-0
48. Myall KJ, Mukherjee B, Castanheira AM, Lam JL, Benedetti G et al. Persistent Post-COVID-19 Interstitial Lung Disease. An Observational Study of Corticosteroid Treatment. *Annals of the American Thoracic Society*. 2021;18(5):799-806. doi:10.1513/AnnalsATS.202008-1002OC
49. Al-Bari MA. Chloroquine analogues in drug discovery: new directions of uses, mechanisms of actions and toxic manifestations from malaria to multifarious diseases. *Journal of Antimicrobial Chemotherapy*. 2015;70(6):1608-1621. doi:10.1093/jac/dkv018
50. Costedoat-Chalumeau N, Hulot JS, Amoura Z, Leroux G, Lechat P et al. Heart conduction disorders related to antimalarials toxicity: an analysis of electrocardiograms in 85 patients treated with hydroxychloroquine for connective tissue diseases. *Rheumatology (Oxford)*. 2007;46(5):808-810. doi:10.1093/rheumatology/kel402
51. D'Alessandro S, Scaccabarozzi D, Signorini L, Perego F, Ilboudo DP et al. The Use of Antimalarial Drugs against Viral Infection. *Microorganisms*. 2020;8(1). doi:10.3390/microorganisms8010085
52. RECOVERY Collaborative Group. Effect of Hydroxychloroquine in Hospitalized Patients with Covid-19. *New England Journal of Medicine*. 2020;383(21):2030-2040. doi:10.1056/NEJMoa2022926
53. Axfors C, Schmitt AM, Janiaud P, Van't Hooft J, Abd-Elsalam S et al. Mortality outcomes with hydroxychloroquine and chloroquine in COVID-19 from an international collaborative meta-analysis of randomized trials. *Nature Communications*. 2021;12(1):2349. doi:10.1038/s41467-021-22446-z
54. Arumugham VB, Rayi A. *Intravenous Immunoglobulin (IVIg)*. StatPearls. Treasure Island (FL): StatPearls Publishing LLC.; 2020.
55. Rodriguez de la Concepcion ML, Ainsua-Enrich E, Reynaga E, Avila-Nieto C, Santos JR et al. High-dose intravenous immunoglobulins might modulate inflammation in COVID-19 patients. *Life Science Alliance*. 2021;4(9). doi:10.26508/lsa.202001009
56. Liu J, Chen Y, Li R, Wu Z, Xu Q et al. Intravenous immunoglobulin treatment for patients with severe COVID-19: a retrospective multicentre study. *Clinical Microbiology and Infection*. 2021; 27(10): 1488-1493. doi:10.1016/j.cmi.2021.05.012
57. Omma A, Erden A, Armagan B, Guven SC, Karakas O et al. A single center experience of intravenous immunoglobulin treatment in Covid-19. *International Immunopharmacology*. 2021;98:107891. doi:10.1016/j.intimp.2021.107891
58. Tabarsi P, Barati S, Jamaati H, Haseli S, Marjani M et al. Evaluating the effects of Intravenous Immunoglobulin (IVIg) on the management of severe COVID-19 cases: A randomized controlled trial. *International Immunopharmacology*. 2021;90:107205. doi:10.1016/j.intimp.2020.107205
59. Gharebaghi N, Nejadrahim R, Mousavi SJ, Sadat-Ebrahimi SR, Hajizadeh R. The use of intravenous immunoglobulin gamma for the treatment of severe coronavirus disease 2019: a randomized placebo-controlled double-blind clinical trial. *BMC Infectious Diseases*. 2020;20(1):786. doi:10.1186/s12879-020-05507-4
60. Xiang HR, Cheng X, Li Y, Luo WW, Zhang QZ et al. Efficacy of IVIG (intravenous immunoglobulin) for corona virus disease 2019 (COVID-19): A meta-analysis. *International Immunopharmacology*. 2021;96:107732. doi:10.1016/j.intimp.2021.107732
61. Kubota-Koketsu R, Terada Y, Yunoki M, Sasaki T, Nakayama EE et al. Neutralizing and binding activities against SARS-CoV-1/2, MERS-CoV, and human coronaviruses 229E and OC43 by normal human intravenous immunoglobulin derived from healthy donors in Japan. *Transfusion*. 2021;61(2):356-360. doi:10.1111/trf.16161
62. Mascolo S, Carleo MA, Contieri M, Izzo S, Perna A et al. SARS-CoV-2 and inflammatory responses: From mechanisms to the potential therapeutic use of intravenous immunoglobulin. *Journal of Medical Virology*. 2021;93(5):2654-2661. doi:10.1002/jmv.26651
63. Henderson LA, Canna SW, Friedman KG, Gorelik M, Lapidus SK et al. American College of Rheumatology Clinical Guidance for Multisystem Inflammatory Syndrome in Children Associated With SARS-CoV-2 and Hyperinflammation in Pediatric COVID-19: Version 2. *Arthritis Rheumatology*. 2021;73(4):e13-e29. doi:10.1002/art.41616
64. Ouldali N, Toubiana J, Antona D, Javouhey E, Madhi F et al. Association of Intravenous Immunoglobulins Plus Methylprednisolone vs Immunoglobulins Alone With Course of Fever in Multisystem Inflammatory Syndrome in Children. *Journal of the American Medical Association*. 2021;325(9):855-864. doi:10.1001/jama.2021.0694
65. Emeksiz S, Celikel Acar B, Kibar AE, Ozkaya Parlakay A, Perk O et al. Algorithm for the diagnosis and management of the multisystem inflammatory syndrome in children associated with COVID-19. *International Journal of Clinical Practice*. 2021;75(9):e14471. doi:10.1111/ijcp.14471
66. Son MBE, Murray N, Friedman K, Young CC, Newhams MM et al. Multisystem Inflammatory Syndrome in Children - Initial Therapy and Outcomes. *New England Journal of Medicine*. 2021;385(1):23-34. doi:10.1056/NEJMoa2102605
67. Yang L, Xie X, Tu Z, Fu J, Xu D et al. The signal pathways and treatment of cytokine storm in COVID-19. *Signal Transduction and Targeted Therapy*. 2021;6(1):255. doi:10.1038/s41392-021-00679-0
68. Uciechowski P, Dempke WCM. Interleukin-6: A Masterplayer in the Cytokine Network. *Oncology*. 2020;98(3):131-137. doi:10.1159/000505099

69. Liao Y, Wang X, Huang M, Tam JP, Liu DX. Regulation of the p38 mitogen-activated protein kinase and dual-specificity phosphatase 1 feedback loop modulates the induction of interleukin 6 and 8 in cells infected with coronavirus infectious bronchitis virus. *Virology*. 2011;420(2):106-116. doi:10.1016/j.virol.2011.09.003
70. Zhou Y, Fu B, Zheng X, Wang D, Zhao C, Qi Y et al. Pathogenic T-cells and inflammatory monocytes incite inflammatory storms in severe COVID-19 patients, *National Science Review*. 2020; 7: 998–1002, <https://doi.org/10.1093/nsr/nwaa041>.
71. Chen X, Zhao B, Qu Y, Chen Y, Xiong J et al. Detectable Serum Severe Acute Respiratory Syndrome Coronavirus 2 Viral Load (RNAemia) Is Closely Correlated With Drastically Elevated Interleukin 6 Level in Critically Ill Patients With Coronavirus Disease 2019. *Clinical Infectious Disease*. 2020;71(8):1937-1942. doi: 10.1093/cid/ciaa449.
72. WHO Rapid Evidence Appraisal for COVID-19 Therapies (REACT) Working Group. Association Between Administration of IL-6 Antagonists and Mortality Among Patients Hospitalized for COVID-19: A Meta-analysis. *Journal of the American Medical Association*. 2021;326(6):499-518. doi:10.1001/jama.2021.11330
73. A TV, Haikarainen T, Raivola J, Silvennoinen O. Selective JAKinibs: Prospects in Inflammatory and Autoimmune Diseases. *BioDrugs*. 2019;33(1):15-32. doi:10.1007/s40259-019-00333-w
74. Satarker S, Tom AA, Shaji RA, Alosious A, Luvis M et al. JAK-STAT Pathway Inhibition and their Implications in COVID-19 Therapy. *Postgraduate Medical Journal*. 2021;133(5):489-507. doi:10.1080/00325481.2020.1855921
75. Fragoulis GE, McInnes IB, Siebert S. JAK-inhibitors. New players in the field of immune-mediated diseases, beyond rheumatoid arthritis. *Rheumatology (Oxford)*. 2019;58(Suppl 1):i43-i54. doi:10.1093/rheumatology/key276
76. Stebbing J, Sanchez Nieves G, Falcone M, Youhanna S, Richardson P et al. JAK inhibition reduces SARS-CoV-2 liver infectivity and modulates inflammatory responses to reduce morbidity and mortality. *Science Advances*. 2021;7(1). doi:10.1126/sciadv.abe4724
77. Richardson P, Griffin I, Tucker C, Smith D, Oechsle O et al. Baricitinib as potential treatment for 2019-nCoV acute respiratory disease. *Lancet*. 2020;395(10223):e30-e31. doi:10.1016/s0140-6736(20)30304-4
78. Kalil AC, Patterson TF, Mehta AK, Tomashek KM, Wolfe CR et al. Baricitinib plus Remdesivir for Hospitalized Adults with Covid-19. *New England Journal of Medicine*. 2021;384(9):795-807. doi:10.1056/NEJMoa2031994
79. Marconi VC, Ramanan AV, de Bono S, Kartman CE, Krishnan V et al. Efficacy and safety of baricitinib for the treatment of hospitalised adults with COVID-19 (COV-BARRIER): a randomised, double-blind, parallel-group, placebo-controlled phase 3 trial. *Lancet Respiratory Medicine*. 2021. doi:10.1016/S2213-2600(21)00331-3
80. Guimaraes PO, Quirk D, Furtado RH, Maia LN, Saraiva JF et al. Tofacitinib in Patients Hospitalized with Covid-19 Pneumonia. *New England Journal of Medicine*. 2021;385(5):406-415. doi:10.1056/NEJMoa2101643
81. Chen CX, Wang JJ, Li H, Yuan LT, Gale RP et al. JAK-inhibitors for coronavirus disease-2019 (COVID-19): a meta-analysis. *Leukemia*. 2021;35(9):2616-2620. doi:10.1038/s41375-021-01266-6
82. Sparks JA, Wallace ZS, Seet AM, Gianfrancesco MA, Izadi Z et al. Associations of baseline use of biologic or targeted synthetic DMARDs with COVID-19 severity in rheumatoid arthritis: Results from the COVID-19 Global Rheumatology Alliance physician registry. *Annals of the Rheumatic Diseases*. 2021;80(9):1137-1146. doi:10.1136/annrheumdis-2021-220418
83. Solimani F, Meier K, Ghoreschi K. Janus kinase signaling as risk factor and therapeutic target for severe SARS-CoV-2 infection. *European Journal of Immunology*. 2021;51(5):1071-1075. doi:10.1002/eji.202149173
84. Rodriguez-Garcia JL, Sanchez-Nievas G, Arevalo-Serrano J, Garcia-Gomez C, Jimenez-Vizuet JM et al. Baricitinib improves respiratory function in patients treated with corticosteroids for SARS-CoV-2 pneumonia: an observational cohort study. *Rheumatology (Oxford)*. 2021;60(1):399-407. doi:10.1093/rheumatology/keaa587
85. Conti P, Gallenga CE, Tete G, Caraffa A, Ronconi G et al. How to reduce the likelihood of coronavirus-19 (CoV-19 or SARS-CoV-2) infection and lung inflammation mediated by IL-1. *Journal of Biological Regulators & Homeostatic Agents*. 2020;34(2):333-338. doi:10.23812/Editorial-Conti-2
86. Chen IY, Moriyama M, Chang MF, Ichinohe T. Severe Acute Respiratory Syndrome Coronavirus Viroporin 3a Activates the NLRP3 Inflammasome. *Frontiers in Microbiology*. 2019;10:50. doi:10.3389/fmicb.2019.00050
87. Zhao N, Di B, Xu LL. The NLRP3 inflammasome and COVID-19: Activation, pathogenesis and therapeutic strategies. *Cytokine Growth Factor Reviews*. 2021;61:2-15. doi:10.1016/j.cytogfr.2021.06.002
88. Satis H, Ozger HS, Aysert Yildiz P, Hizel K, Gulbahar O et al. Prognostic value of interleukin-18 and its association with other inflammatory markers and disease severity in COVID-19. *Cytokine*. 2021;137:155302. doi:10.1016/j.cyto.2020.155302
89. Opal SM, Fisher CJ, Jr., Dhainaut JF, Vincent JL, Brase R et al. Confirmatory interleukin-1 receptor antagonist trial in severe sepsis: a phase III, randomized, double-blind, placebo-controlled, multicenter trial. The Interleukin-1 Receptor Antagonist Sepsis Investigator Group. *Critical Care Medicine*. 1997;25(7):1115-1124. doi:10.1097/00003246-199707000-00010
90. Cavalli G, De Luca G, Campochiaro C, Della-Torre E, Ripa M et al. Interleukin-1 blockade with high-dose anakinra in patients with COVID-19, acute respiratory distress syndrome, and hyperinflammation: a retrospective cohort study. *Lancet Rheumatology*. 2020;2(6):e325-e331. doi:10.1016/S2665-9913(20)30127-2

91. The CORIMUNO-19 Collaborative group. Effect of anakinra versus usual care in adults in hospital with COVID-19 and mild-to-moderate pneumonia (CORIMUNO-ANA-1): a randomised controlled trial. *Lancet Respiratory Medicine*. 2021;9(3):295-304. doi:10.1016/S2213-2600(20)30556-7
92. Kyriazopoulou E, Huet T, Cavalli G, Gori A, Kyprianou M et al. Effect of anakinra on mortality in patients with COVID-19: a systematic review and patient-level meta-analysis. *Lancet Rheumatology*. 2021;3(10):e690-e697. doi:10.1016/S2665-9913(21)00216-2
93. Kyriazopoulou E, Panagopoulos P, Metallidis S, Dalekos GN, Poulakou G et al. An open label trial of anakinra to prevent respiratory failure in COVID-19. *Elife*. 2021;10. doi:10.7554/eLife.66125
94. Kyriazopoulou E, Poulakou G, Milionis H, Metallidis S, Adamis G et al. Early treatment of COVID-19 with anakinra guided by soluble urokinase plasminogen receptor plasma levels: a double-blind, randomized controlled phase 3 trial. *Nature Medicine*. 2021. doi:10.1038/s41591-021-01499-z
95. Caricchio R, Abbate A, Gordeev I, Meng J, Hsue PY et al. Effect of Canakinumab vs Placebo on Survival Without Invasive Mechanical Ventilation in Patients Hospitalized With Severe COVID-19: A Randomized Clinical Trial. *Journal of the American Medical Association*. 2021;326(3):230-239. doi:10.1001/jama.2021.9508
96. Devereos SG, Siasos G, Giannopoulos G, Vrachatis DA, Angelidis C et al. The GReek study in the Effects of Colchicine in COvid-19 complications prevention (GRECCO-19 study): rationale and study design. *Hellenic Journal of Cardiology*. 2020;61:42-45. doi:10.1016/j.hjc.2020.03.002
97. Tardif JC, Kouz S, Waters DD, Bertrand OF, Diaz R et al. Efficacy and Safety of Low-Dose Colchicine after Myocardial Infarction. *New England Journal of Medicine*. 2019;381(26):2497-2505. doi:10.1056/NEJMoa1912388
98. Lopes MI, Bonjorno LP, Giannini MC, Amaral NB, Menezes PI et al. Beneficial effects of colchicine for moderate to severe COVID-19: a randomised, double-blinded, placebo-controlled clinical trial. *RMD Open*. 2021;7(1). doi:10.1136/rmdopen-2020-001455
99. Tardif JC, Bouabdallaoui N, L'Allier PL, Gaudet D, Shah B et al. Colchicine for community-treated patients with COVID-19 (COLCORONA): a phase 3, randomised, double-blinded, adaptive, placebo-controlled, multicentre trial. *Lancet Respiratory Medicine*. 2021;9(8):924-932. doi:10.1016/S2213-2600(21)00222-8
100. RECOVERY Collaborative Group. Colchicine in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet Respiratory Medicine*. 2021; 9(12):1419-1426. doi:10.1016/S2213-2600(21)00435-5
101. Lien C H, Lee MD, Weng S L, Lin CH, Liu LY et al. Repurposing Colchicine in Treating Patients with COVID-19: A Systematic Review and Meta-Analysis. *Life (Basel, Switzerland)*. 2021; 11(8): 864. <https://doi.org/10.3390/life11080864>
102. Madrid-Garcia A, Perez I, Colomer JI, Leon-Mateos L, Jover JA et al. Influence of colchicine prescription in COVID-19-related hospital admissions: a survival analysis. *Therapeutic Advances in Musculoskeletal Disease*. 2021;13:1759720X211002684. doi:10.1177/1759720X211002684
103. Kharouf F, Ishay Y, Kenig A, Bitan M, Ben-Chetrit E. Incidence and course of COVID-19 hospitalizations among patients with familial Mediterranean fever. *Rheumatology (Oxford)*. 2021. doi:10.1093/rheumatology/keab577
104. Satis H, Armagan B, Bodakci E, Atas N, Sari A et al. Colchicine intolerance in FMF patients and primary obstacles for optimal dosing. *Turkish Journal of Medical Sciences*. 2020;50(5):1337-1343. doi:10.3906/sag-2001-261
105. Haga S, Yamamoto N, Nakai-Murakami C, Osawa Y, Tokunaga K et al. Modulation of TNF-alpha-converting enzyme by the spike protein of SARS-CoV and ACE2 induces TNF-alpha production and facilitates viral entry. *Proceedings of the National Academy of Sciences*. 2008;105(22):7809-7814. doi:10.1073/pnas.0711241105
106. Robinson PC, Liew DFL, Liew JW, Monaco C, Richards D et al. The Potential for Repurposing Anti-TNF as a Therapy for the Treatment of COVID-19. *Med (N Y)*. 2020; 1(1):90-102. doi:10.1016/j.medj.2020.11.005.
107. Feldmann M, Maini RN, Woody JN, Holgate ST, Winter G et al. Trials of anti-tumour necrosis factor therapy for COVID-19 are urgently needed. *The Lancet*. 2020. doi:10.1016/S0140-6736(20)30858-8
108. Kridin K, Schonmann Y, Solomon A, Damiani G, Tzur Bitan D et al. Risk of COVID-19 Infection, Hospitalization, and Mortality in Patients with Psoriasis Treated by Interleukin-17 Inhibitors. *Journal of Dermatological Treatment*. 2021;1-28. doi:10.1080/09546634.2021.1905766
109. Avdeev SN, Trushenko NV, Tsareva NA, Yaroshetskiy AI, Merzhoeva ZM et al. Anti-IL-17 monoclonal antibodies in hospitalized patients with severe COVID-19: A pilot study. *Cytokine*. 2021;146:155627. doi:10.1016/j.cyto.2021.155627
110. Crotti C, Biggioggero M, Becciolini A, Agape E, Favalli EG. Mavrilimumab: a unique insight and update on the current status in the treatment of rheumatoid arthritis. *Expert Opinion on Investigational Drugs*. 2019;28(7):573-581. doi:10.1080/13543784.2019.1631795
111. Cremer PC, Abbate A, Hudock K, McWilliams C, Mehta J et al. Mavrilimumab in patients with severe COVID-19 pneumonia and systemic hyperinflammation (MASH-COVID): an investigator initiated, multicentre, double-blind, randomised, placebo-controlled trial. *Lancet Rheumatology*. 2021;3(6):e410-e418. doi:10.1016/S2665-9913(21)00070-9
112. Temesgen Z, Burger CD, Baker J, Polk C, Libertin C et al. Lenzilumab Efficacy and Safety in Newly Hospitalized Covid-19 Subjects: Results from the Live-Air Phase 3 Randomized Double-Blind Placebo-Controlled Trial. *medRxiv*. 2021. doi:10.1101/2021.05.01.21256470