

1-1-2015

Investigation of root-knot nematode (*Meloidogyne* spp.) resistance in almond rootstocks with DNA markers

SULTAN BAY TÜRKOĞLU

NERGİZ ÇOBAN

ALİ ÖZKAN

HALİT SEYFETTİN ATLI

CANAN CAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

Recommended Citation

TÜRKOĞLU, SULTAN BAY; ÇOBAN, NERGİZ; ÖZKAN, ALİ; ATLI, HALİT SEYFETTİN; and CAN, CANAN (2015) "Investigation of root-knot nematode (*Meloidogyne* spp.) resistance in almond rootstocks with DNA markers," *Turkish Journal of Agriculture and Forestry*. Vol. 39: No. 4, Article 5. <https://doi.org/10.3906/tar-1408-97>

Available at: <https://journals.tubitak.gov.tr/agriculture/vol39/iss4/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Investigation of root-knot nematode (*Meloidogyne* spp.) resistance in almond rootstocks with DNA markers

Sultan BAY TÜRKOĞLU¹, Nergiz ÇOBAN², Ali ÖZKAN³, Halit Seyfettin ATLI⁴, Canan CAN^{1*}

¹Department of Biology, Faculty of Arts and Sciences, Gaziantep University, Gaziantep, Turkey

²Pistachio Research Station, Ministry of Food, Agriculture, and Livestock, Gaziantep, Turkey

³Department of Gastronomy and Culinary Arts, Faculty of Fine Arts, Gaziantep University, Gaziantep, Turkey

⁴Department of Horticulture, Faculty of Agriculture, Siirt University, Siirt, Turkey

Received: 26.08.2014 • Accepted: 04.04.2015 • Published Online: 08.07.2015 • Printed: 30.07.2015

Abstract: Seven almond varieties, Ferragnes, Nemaguard, AB3 (*Amygdalus orientalis* Mill.), GN22, GF677, Myrobalan (*Prunus cerasifera*), and *Pissardi nigra*, and F₁ crossbreed almonds, Ferragnes × *Pissardi nigra*, Ferragnes × Myrobalan, Ferragnes × GN22, Nemaguard × GF677, Myrobalan × AB3 (*Prunus dulcis* Mill.), Myrobalan × Ferragnes, *Pissardi nigra* × AB3, and GF677 × Myrobalan, were used to determine resistance against root-knot nematode (RKN) (*Meloidogyne* spp.) using DNA markers. Parental lines and 316 F₁ plants were tested with SSR (plgms8, plgms9, and plgms19) and STS (STS-OPS14a, STS-834b) markers. It was found that plgms19 exhibited 76%, 77.01%, 73.13%, and 86.6% inheritance of resistance in the Ferragnes × *Pissardi nigra*, Ferragnes × Myrobalan, Ferragnes × GN22, and Nemaguard × GF677 F₁ populations, respectively. No resistance was detected in the Nemaguard × GF677 population with the plgms8 marker. Since the plgms19 and STS-834b primers were found to be effective at high ratios in determining resistance to RKN in the Nemaguard × GF677 F₁ population, the use of these markers could be included in breeding studies. It was also detected that the STS-OPS14a primer is appropriate to use for determining the sensitivity to RKN in almond rootstocks.

Key words: Almond, breeding, DNA marker, nematode

1. Introduction

Almond (*Amygdalus communis* L. syn. *Prunus dulcis* (Miller) D.A. Webb) is placed within the subfamily Prunoideae of Rosaceae. It is consumed as fresh and dry nuts and is also used in the food industry (Kester et al., 1991). In Turkey, fresh and dry consumption of almond is very common and almond nuts are widely used in the food industry (Akalin, 1952; Bayrak and Yilmaz, 2009). Cultivation of almond in Turkey is increasing and almonds are grown in all the regions except for the Black Sea and high plateaus. A total of 69,838 t of fruit production was reported in 2011 and the yield has been stated to increase steadily since the 1990s (<http://www.turkstat.gov.tr/>). Almond was grown as a hedge plant before the 1990s and from that time orchards have been planted because of the government's increasing support for almond producers.

There are serious diseases and pests in almonds that cause considerable yield losses. Among them, root-knot nematodes (RKNs) *Meloidogyne arenaria*, *M. incognita*, and *M. javanica* have been reported from the Mediterranean countries that have large almond plantations (Sasser, 1977; Lamberti, 1979).

M. arenaria was reported from the northern part of the Mediterranean countries while *M. incognita* and *M. javanica* are distributed throughout the southern Mediterranean (Trudgill and Blok, 2001). In Turkey, *M. incognita*, *M. javanica*, and *M. arenaria* were determined as the most widespread and economically important RKN species, while *M. hapla* and *M. chitwoodi* were reported as rare species (Söğüt and Elekçioğlu, 2000; Devran and Söğüt, 2009, 2011; Özarıslan and Elekçioğlu, 2010). Söğüt and Elekçioğlu (2010) used classical and DNA-based techniques to diagnose 79 RKN population collected from different regions and hosts. They identified *M. javanica*, *M. arenaria*, *M. incognita*, and *M. chitwoodi* at rates of 24.05%, 22.78%, 17.72%, and 10.13%, respectively. Devran and Söğüt (2009) collected 95 RKN samples from vegetable growing areas of the West Mediterranean region to determine *Meloidogyne* spp. with species-specific primers and the proportions were 64.2%, 28.4%, and 7.3% for *M. incognita*, *M. javanica*, and *M. arenaria*, respectively. *M. incognita* race 2 and *M. javanica* race 1 were determined to be widespread in protected vegetables grown in the West Mediterranean region (Devran and Söğüt, 2011).

* Correspondence: can@gantep.edu.tr

Meloidogyne spp. has been reported at a rate of 52% from the world's agricultural areas, including vegetables, fruit trees, and ornamental plants grown in Mediterranean climates (Trudgill and Blok, 2001). Besides the yield loss that *Meloidogyne* spp. cause in stone fruits orchards, their control is cumbersome since the banning of methyl bromide (MeBr) by the Kyoto protocol due to its high toxicity (Layne, 1987, Nyczepir, 1991). In Turkey MeBr usage was prohibited in 2007 except for critical use, and therefore other fumigant applications such as dazomet (98% a.i. Basamid) in combination with solarization have been used for controlling soil-borne fungal pathogens and nematodes (Yücel et al., 2007).

An effective and environmentally friendly approach in stone fruit trees for controlling nematodes is to develop resistant rootstocks (Boerma and Hussey, 1992; Vrain, 1999, Lopez-Perez et al., 2006). Resistance specificities to *Meloidogyne* spp. in plum, almond, and peach have been well documented (Esmenjaud et al., 1997), and the *Ma* gene from the Myrobalan plum (*Prunus cerasifera*) was cloned (Claverie et al., 2011). The *Ma* gene exhibits a complete-spectrum resistance to *M. arenaria*, *M. incognita*, and *M. javanica* with several R alleles having a dominant nature (Esmenjaud et al., 1996; Lecouls et al., 1997).

Simple sequence repeats (SSRs), namely plgms8, plgms9, and plgms19, were reported to exhibit tight linkage with the *Ma* gene; among them, plgms19 cosegregated with the *Ma* loci in Myrobalan plum (Claveire et al., 2004a, 2004b). It was found that plgms8 amplified a 316-bp band in Myrobalan and was mapped to the *Ma* loci (Claveire et al., 2004b). Sequence-tagged site (STS) markers were detected to be linked to *Mia/mia* and *Mja/mja* and among them STS-OPS14a and STS-834b exhibited close linkages (Yamamoto and Hayashi, 2002).

Marker-assisted selection has great potential in plant breeding studies and selection of agriculturally important characters with DNA markers has opened new possibilities (Collard et al., 2005). The DNA markers exhibiting tight linkage to desired genes have been used as molecular tools in many crop plants (Ribaut and Hoisington, 1998), called 'molecular breeding' in plant and animal breeding (Rafalski and Tingey, 1993). Exploitation of germplasm resources and characterization of genetic variation have been studied more effectively by use of molecular markers (Tanksley et al., 1989; Tanksley and McCouch, 1997; Gur and Zamir, 2004). Techniques in genomics also have useful applications in polymorphism detection and homozygosity testing besides protection of plant biodiversity (Heckenberger et al., 2006).

In Turkey, rootstock breeding against nematodes in almonds through controlled hybridization has not been conducted and such studies are very important since almond and peach plantation productions are increasing

every year. DNA-based techniques may shorten the time necessary for RKN-resistant almond rootstock breeding. In this study 7 different parents and 316 F₁ plants were tested with 5 DNA-based markers to define the effectiveness of DNA markers in RKN resistance determination. Parental lines and F₁ individuals were screened with SSR (plgms8, plgms9, and plgms19) and STS (STS-OPS14a and STS-834b) markers to explore inheritance of resistance.

2. Materials and methods

2.1. Plant materials

Ferragnes, Nemaguard, AB3 (*Amygdalus orientalis* Mill.), GN22, GF677, Myrobalan plum (*Prunus cerasifera*), and *Pissardi nigra* were used as parental lines. *P. cerasifera*, *P. nigra*, Nemaguard, and GN22 lines used in this study were selected as resistant parental lines to RKN and Ferragnes, AB3, and GF677 as susceptible to RKN (Esmenjaud et al., 1993; Athl, 2010).

The F₁ plants screened in this study were Ferragnes × *Pissardi nigra*, Ferragnes × Myrobalan, Ferragnes × GN22, Nemaguard × GF677, Myrobalan × Ferragnes, *Pissardi nigra* × AB3, Myrobalan × AB3, and GF677 × Myrobalan. Leaves of 5-year-old parental lines and 2-year-old F₁ plants were collected from the Pistachio Research Station (Gaziantep, Turkey) experimental plots in April and May, frozen in liquid nitrogen, and kept at -80 °C until use.

2.2. Genomic DNA isolation

Young leaves were used for genomic DNA isolation and the modified CTAB protocol of Doyle and Doyle (1987) was used. The amount of DNA was measured by spectrophotometric absorbance at wavelengths of 260–280 nm and the concentration was adjusted to 40–60 ng/μL.

2.3. Markers and PCR analysis

Parents and F₁ plants were tested with SSR and STS markers that were reported to have tight linkages with *Ma* genes (Lecouls et al., 1997, 2004; Claverie et al., 2004a, 2004b). Sequences of SSR (plgms8, plgms9, plgms19) and STS (STS-OPS14a, STS-834b) markers, product sizes, and binding temperatures are given in Table 1. These markers were reported to be linked to RKN resistance by Lecouls et al. (2004), Claverie et al. (2004b), and Yamamoto and Hayashi (2002). Polymerase chain reaction (PCR) analyses of parents with SSR and STS markers were conducted on RKN-resistant and RKN-susceptible parental lines and band profiles were evaluated. According to band profiles of parental lines, F₁ individuals were tested with markers. PCR amplifications with STS and SSR markers were performed in a 20-μL final volume containing 40–60 ng of genomic DNA, 0.7 U Taq polymerase (Fermentas), 0.2 μM primer, 2 mM dNTPs (Fermentas), 1.5 mM MgCl₂, 1X reaction buffer, and ddH₂O (Claveire et al., 2004a). PCR conditions consisted of denaturation at 95 °C for 1 min, followed by 40 cycles of 95 °C for 70 s, 56 °C for 15 s, and

Table 1. The sequence, product sizes and binding temperatures of DNA markers (Yomamoto and Hayashi, 2002; Claverie et al., 2004b; Lecouls et al., 2004).

Markers	Primer sequences (5'-3')	Product size	Temperature
plgms8	F:AAACAGCCAGATCCGGAGTA	316	56 °C
	R:TATAAGTCCGCCATCGCTTG		
plgms9	F:GGTGGGAGAATTCGACTATCA	156	56 °C
	R:CAACCCAATACCACGTACCC		
plgms19	F:CGGGGTTCAAACCTCAACAAG	327	56 °C
	R:AGACGTGCTGCTTTGTTCAC		
STS-OPS14a	F:TTACGTGAAACAGGGAAATGAA	402	55 °C
	R:GCCTTAGTACAGCTCCGCC		
STS-834b	F:GCAGTCAAAAATTTCAAACC	227	55 °C
	R:TCCGATTCGAGCCACTACA		

72 °C for 75 s, with final elongation at 72 °C for 4 min. PCR products were run on 2% agarose gel and visualized using a gel documentation system under UV light. Evaluations of paternal lines and F₁ individuals were conducted by the presence (+) or absence (-) of expected band profiles and coded as R for resistant or S for susceptible, respectively. Chi-square values were determined by using SPSS 16.

3. Results

Eight different crosses (Ferragnes × *Pissardi nigra*, Ferragnes × Myrobalan, Ferragnes × GN22, Nemaguard × GF677, Myrobalan × Ferragnes, *Pissardi nigra* × AB3, Myrobalan × AB3, and GF677 × Myrobalan) were screened with DNA markers to define resistance to RKN. Genotypes and observed numbers are presented in Tables 2–6. Distribution of markers for resistance and susceptibility were calculated with chi-square and P values (Tables 2–5). Myrobalan × AB3, Myrobalan × Ferragnes, *Pissardi nigra* × AB3, and GF677 × Myrobalan hybrids were not included in chi-square analyses due to incomplete numbers of F₁ individuals (Table 6).

plgms19 segregated with resistance in the Ferragnes × *Pissardi nigra* F₁ populations. A total of 125 F₁ individuals were screened in this cross and the rate of resistance was 76% (Table 2). Similarly, segregation with resistance was higher with plgms19 in Ferragnes × Myrobalan, Ferragnes × GN22, and Nemaguard × GF677 F₁ individuals (Tables 3–5). The highest rate of resistance (86.6%) was obtained with Nemaguard × GF677 F₁ individuals (Table 5).

plgms8 and plgms9 segregated with susceptibility in the Ferragnes × *Pissardi nigra* and Ferragnes × Myrobalan F₁ populations (Tables 2 and 3). In Ferragnes × Myrobalan, plgms8 detected 87 (100%) F₁ individuals as susceptible

and the rate of susceptibility was 99.2% in Ferragnes × *Pissardi nigra*. In Ferragnes × GN22, plgms8 exhibited 100% susceptibility, whereas plgms9 detected 68.6% resistance (Table 4).

P. cerasifera, *P. nigra*, and Nemaguard and GN22 have RKN resistance but AB3, GF677, and Ferragnes were reported to be susceptible to RKN (Esmenjaud et al., 1993; Athi, 2010). PCR profiles of parental lines and hybrid individuals are presented in Figures 1 and 2. Hybrid individuals exhibited 327-bp and 156-bp fragments with the plgms19 and plgms9 primers, respectively.

Inheritance of resistance with STS-834b was 15.2%, 9.1%, and 93.3% in Ferragnes × *Pissardi nigra*, Ferragnes × Myrobalan, and Nemaguard × GF677 F₁ individuals, respectively (Tables 2, 4, and 5). STS-834b detected 100% susceptibility in Ferragnes × GN22 F₁ individuals (Table 4). STS-OPS14a found 11.5%, 16%, 31.4%, and 66.6% resistant F₁ individuals in the Ferragnes × Myrobalan, Ferragnes × *Pissardi nigra*, Ferragnes × GN22, and Nemaguard × GF677 populations, respectively (Tables 2–5).

4. Discussion

This study tested Ferragnes, Nemaguard, AB3 (*Amygdalus orientalis* Mill.), GN22, GF677, Myrobalan, and *Pissardi nigra* along with F₁ individuals with DNA markers to define the inheritance of resistance to RKNs. Three SSR (plgms8, plgms19, and plgms9) and two STS (STS-OPS14a and STS-834b) markers that were reported to have tight linkage to the RKN resistance specificities were used to screen 7 parental lines and 316 F₁ individuals (Yomamoto and Hayashi, 2002; Claverie et al., 2004a, 2004b).

Table 2. Distribution of RKN resistance and susceptibility in Ferragnes × *Pissardi nigra* F₁ individuals with chi-square and P values.

Markers	Genotype*	Number observed (%)	Percentage expected
plgms19	R	95 (76%)	50
	S	30 (24%)	
	Chi-square (P value)	33.800 (P < 0.0001)	
plgms8	R	1 (0.8%)	50
	S	124 (99.2%)	
	Chi-square (P value)	121.032 (P < 0.0001)	
plgms9	R	23 (18.4%)	50
	S	102 (81.6%)	
	Chi-square (P value)	49.928 (P < 0.0001)	
STS-OPS14a	R	20 (16%)	50
	S	105 (84%)	
	Chi-square (P value)	57.800 (P < 0.0001)	
STS-834b	R	19 (15.2%)	50
	S	106 (84.8%)	
	Chi-square (P value)	60.552 (P < 0.0001)	

*R: Resistant, S: Susceptible.

Table 3. Distribution of RKN resistance and susceptibility in Ferragnes × *Myrobalan* F₁ individuals with chi-square and P values.

Markers	Genotype*	Number observed (%)	Percentage expected
plgms19	R	67 (77.01%)	50
	S	20 (22.99%)	
	Chi-square (P value)	25.391 (P < 0.0001)	
plgms8	R	0 (0.0%)	50
	S	87 (100.0%)	
	Chi-square (P value)	87.000 (P < 0.0001)	
plgms9	R	16 (18.4%)	50
	S	71 (81.6%)	
	Chi-square (P value)	34.77 (P < 0.0001)	
STS-OPS14a	R	10 (11.5%)	50
	S	77 (88.5%)	
	Chi-square (P value)	51.59 (P < 0.0001)	
STS-834b	R	8 (9.1%)	50
	S	79 (90.9%)	
	Chi-square (P value)	57.94 (P < 0.0001)	

*R: Resistant, S: Susceptible.

Table 4. Distribution of RKN resistance and susceptibility in Ferragnes × GN22 F₁ individuals with chi-square and P values.

Markers	Genotype*	Number observed (%)	Percentage expected
plgms19	R	49 (73.13%)	50
	S	18 (26.87%)	
	Chi-square (P value)	14.343 (P < 0.0001)	
plgms8	R	0 (0.0%)	50
	S	67 (100.0%)	
	Chi-square (P value)	67 (P < 0.0001)	
plgms9	R	46 (68.6%)	50
	S	21 (31.4%)	
	Chi-square (P value)	9.328 (P = 0.002)	
STS-OPS14a	R	21 (31.4%)	50
	S	46 (68.6%)	
	Chi-square (P value)	9.328 (P = 0.002)	
STS-834b	R	0 (0%)	50
	S	67 (100.0%)	
	Chi-square (P value)	67 (P < 0.0001)	

*R: Resistant, S: Susceptible.

Table 5. Distribution of RKN resistance and susceptibility in Nemaguard × GF677 F₁ individuals with chi-square and P values.

Markers	Genotype*	Number observed (%)	Percentage expected
plgms19	R	13 (86.6%)	50
	S	2 (13.4%)	
	Chi-square (P value)	8.067 (P = 0.004)	
plgms8	R	0 (0.0%)	50
	S	15 (100.0%)	
	Chi-square (P value)	15 (P < 0.0001)	
plgms9	R	4 (26.6%)	50
	S	11 (73.4%)	
	Chi-square (P value)	3.267 (P = 0.071)	
STS-OPS14a	R	10 (66.6%)	50
	S	5 (33.4%)	
	Chi-square (P value)	1.667 (P = 0.197)	
STS-834b	R	14 (93.3%)	50
	S	1 (6.7%)	
	Chi-square (P value)	11.267 (P = 0.001)	

*R: Resistant, S: Susceptible.

Table 6. Distribution of RKN resistance in Myrobalan × AB3, Myrobalan × Ferragnes, *Pissardi nigra* × AB3, and GF677 × Myrobalan F₁ individuals with plgms19.

	Genotype*	Number observed**	Percentage expected
Myrobalan × AB3	R	2	50
	S	2	
Chi-square (P value)		n/a	
Myrobalan × Ferragnes	R	0	50
	S	4	
Chi-square (P value)		n/a	
<i>Pissardi nigra</i> × AB3	R	3	50
	S	4	
Chi-square (P value)		n/a	
GF677 × Myrobalan	R	6	50
	S	1	
Chi-square (P value)		n/a	

*R: Resistant, S: Susceptible. ** n/a: Not tested.




Figure 1. PCR profiles of parents and hybrid individuals with plgms19. M: 50-bp DNA molecular marker. R: Resistant, S: Susceptible.




Figure 2. PCR profiles of parents and hybrid individuals with plgms9. M: 50-bp DNA molecular marker. R: Resistant, S: Susceptible.

The plgms19 marker that is linked to the *Ma* loci (Claverie et al., 2004b) resulted in statistically significant positive difference for RKN resistance in the *Ferragnes* × *Pissardi nigra* and *Ferragnes* × *Myrobalan* F₁ populations (Tables 2 and 3). The resistance rates were 73.13% and 86% in the *Ferragnes* × GN22 and *Nemaguard* × GF677 F₁ populations, respectively (Tables 4 and 5). Claverie et al. (2004b) used SSR markers linked to RKN resistance and determined the inheritance rate of markers with the *Ma* loci. The results obtained with the STS markers in our study exhibited rates of RKN resistance inheritance similar to that of SSR markers found by Claverie et al. (2004b).

plgms8 showed segregation with susceptible genotype in *Ferragnes* × *Pissardi nigra*, *Ferragnes* × *Myrobalan*, and *Nemaguard* × GF677 F₁ populations (Tables 2, 3, and 5). Similarly, plgms9 segregation with susceptible genotype was observed in *Ferragnes* × *Pissardi nigra*, *Ferragnes* × *Myrobalan*, and *Ferragnes* × GN22 F₁ individuals (Tables 2–4). The plgms8 and plgms9 markers were mapped at both sides of the *Ma* region and away from the plgms19

marker defined with recombination (Claverie et al., 2004b). The segregation of plgms8 and plgms9 along with susceptibility in our F₁ population may be due to the distance of these markers from the *Ma* region compared to that of plgms19.

Yamamoto and Hayashi (2002) reported STS marker analyses to explore RKN resistance for marker-assisted analyses. Their study identified four STS markers that exhibited tight linkage in the *Joseitou* variety and STS-834b was reported to lie at loci that determined resistance to *M. javanica*. In our study, we have screened 125, 87, and 67 F₁ individuals with STS-834b in crosses of *Ferragnes* × *Pissardi nigra*, *Ferragnes* × *Myrobalan*, and *Ferragnes* × GN22, respectively (Tables 2–5). In all the crosses the marker segregated with RKN resistance as reported by Yamamoto and Hayashi (2002). STS-834b could be included in marker-assisted selection of almond rootstock breeding studies.

Chi-square and P value calculations were conducted for a total of 316 F₁ individuals derived from *Ferragnes* × *Myrobalan*, *Ferragnes* × *Pissardi nigra*, *Ferragnes* × GN22,

and Nemaguard × GF677 crosses (Tables 2–5). However, chi-square values were not calculated for Myrobalan × AB3, Myrobalan × Ferragnes, *Pissardi nigra* × AB3, or GF677 × Myrobalan F_1 individuals due to small numbers of F_1 individuals (Table 6). Hybridization studies should be conducted in future studies to define RKN resistance in these crosses.

We concluded that plgms19 could be used to determine RKN resistance in Ferragnes × *Pissardi nigra* and Nemaguard × GF677 hybrid individuals due to high rates of resistance genotypes obtained in this study. Further studies in F_2 and F_3 populations are necessary to determine Mendelian segregation of plgms19 in resistant individuals. In addition, in vitro RKN testing should also be conducted

in these populations. Since plgms19 and STS-834b primers were found to be effective in determining resistance to RKN in the Nemaguard × GF677 F_1 population, usage of these markers could be beneficial in further studies. It was also detected that the STS-OPS14a primer is appropriate to use for determining sensitivity to RKNs in almond rootstock breeding.

Acknowledgments

The authors thank the Directorate of the Pistachio Research Station (Gaziantep, Turkey) for maintenance of plant materials. This study was supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) with project number 111O133.

References

- Akalın Ş (1952). Büyük Bitkiler Kılavuzu. Cilt 1. Ankara, Turkey: Güzel Sanatlar Matbaası (in Turkish).
- Atlı HS (2010). GF677 ve GN22 Badem Anaçlarının Çelikle Çoğaltılma Durumlarının Karşılaştırılması. Gaziantep, Turkey: Antepfıstığı Araştırma Enstitüsü Müdürlüğü (in Turkish).
- Bayrak S, Yılmaz Ö (2009). Ceviz-badem yetiştiriciliği. Ankara, Turkey: Rekmay Reklam ve Tanıtım Ltd. (in Turkish).
- Boerma HR, Hussey RS (1992). Breeding plants for resistance to nematodes. *J Nematol* 24: 242–252.
- Clavierie M, Bosselut N, Lecouls AC, Voisin R, Poizat C, Dirlwanger E, Kleinhentz M, Lafargue B, Laigret F, Esmenjaud D (2004a). Location of independent root-knot nematode resistance genes in plum and peach. *Theor Appl Genet* 108: 765–773.
- Clavierie M, Dirlwanger E, Bosselut N, Ghelder VC, Voisin R, Kleinhentz M, Lafargue B, Abad P, Rosso MN, Chalhoub B et al. (2011). The *Ma* gene for complete-spectrum resistance to *Meloidogyne* species in *Prunus* is a TNL with a huge repeated C-terminal post-LRR region. *Plant Physiol* 156: 779–792.
- Clavierie M, Dirlwanger E, Cosson P, Bosselut N, Lecouls AC, Voisin R, Kleinhentz M, Lafargue B, Caboche C, Chalhoub B et al. (2004b). High-resolution mapping and chromosome landing at the root-knot nematode resistance locus *Ma* from Myrobalan plum using a large-insert BAC DNA library. *Theor Appl Genet* 109: 1318–1327.
- Collard BCY, Jahufer MZZ, Brouwer JB, Pang ECR (2005). An introduction to markers, quantitative trait loci (QTL) mapping, and marker-assisted selection for crop improvement: the basic concepts. *Euphytica* 142: 169–196.
- Devran Z, Söğüt MA (2009). Distribution and identification of root-knot nematodes from Turkey. *J Nematol* 41: 128–133.
- Devran Z, Söğüt MA (2011). Characterizing races of *Meloidogyne incognita*, *M. javanica* and *M. arenaria* in the West Mediterranean region of Turkey. *Crop Prot* 30: 451–455.
- Doyle JJ, Doyle JL (1987). A rapid isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* 19: 11–15.
- Esmenjaud D, Minot JC, Voisin R (1996). Effect of durable inoculum pressure and high temperature on root gall, nematode numbers and survival of Myrobalan plum genotypes (*Prunus cerasifera* Ehr.) highly resistant to *Meloidogyne* spp. *Theor Appl Genet* 92: 873–879.
- Esmenjaud D, Minot JC, Voisin R, Pinochet J, Simard MH, Salesses G (1997). Differential response to root-knot nematodes in *Prunus* species and correlative genetic implications. *J Nematol* 29: 372–380.
- Esmenjaud D, Minot JC, Voisin R, Salesses G, Poupet R, Onesto JP (1993). Assessment of a method using plantlets grown from in vitro for studying resistance of *Prunus cerasifera* Ehr. (Myrobalan plum) to *Meloidogyne* spp. *Nematropica* 23: 41–48.
- Gur A, Zamir D (2004). Unused natural variation can lift yield barriers in plant breeding. *PLoS Biol* 2: e245.
- Heckenberger M, Muminovic J, Ruppevan der Voort J, Peleman J, Bohn M, Melchinger AE (2006). Identification of essentially derived varieties from biparental crosses of homogenous lines: III. AFLP data from maize inbreds and comparison with SSR data. *Mol Breeding* 17: 111–125.
- Kester D, Gradziel T, Graselley C (1991). Almonds (*Prunus*). In: Moore J, Ballington J, editors. Genetic Resources of Temperate Fruit and Nut Crops. Wageningen, the Netherlands: ISHS, pp. 701–758.
- Lamberti F (1979). Economic importance of *Meloidogyne* spp. in subtropical and Mediterranean climates. In: Lamberti F, Taylor CE, editors. Root-Knot Nematodes (*Meloidogyne* spp.): Systematics, Biology and Control. New York, NY, USA: Academic Press, pp. 342–357.
- Layne REC (1987). Peach rootstocks. In: Rom RC, Carlson RF, editors. Rootstocks for Fruit Crops. New York, NY, USA: Wiley, pp. 185–216.

- Lecouls AC, Bergougnoux V, Rubio-Cabetas MJ, Bosselut N, Voisin R, Poessel JL, Faurobert M, Bonnet A, Salesses G, Dirlwanger E et al. (2004). Marker-assisted selection for the wide-spectrum resistance to root-knot nematodes conferred by the *Ma* gene from Myrobalan plum (*Prunus cerasifera*) in interspecific *Prunus* material. *Mol Breeding* 13: 113–124.
- Lecouls AC, Salesses G, Minot JC, Voisin R, Bonnet A, Esmenjaud D (1997). Spectrum of the *Ma* genes for resistance to *Meloidogyne* spp. in Myrobalan plum. *Theor Appl Genet* 95: 1325–1334.
- Lopez-Perez JA, Strange ML, Kaloshian I, Ploeg AT (2006). Differential response of MI gene resistant tomato rootstocks to root-knot nematodes (*Meloidogyne incognita*). *Crop Prot* 25: 382–388.
- Nyczepir AP (1991). Nematode management strategies in stone fruits in the United States. *J Nematol* 23: 334–341.
- Özarslandan A, Elekçioğlu İH (2010). Identification of root-knot nematode species (*Meloidogyne* spp.) (Nemata: Meloidogynidae) collected from different parts of Turkey by molecular and morphological methods. *Turk J Entomol* 34: 323–335.
- Rafalski J, Tingey S (1993). Genetic diagnostics in plant breeding: RAPDs, microsatellites and machines. *Trends Genet* 9: 275–280.
- Ribaut JM, Hoisington D (1998). Marker-assisted selection: new tools and strategies. *Trends Plant Sci* 3: 236–239.
- Sasser JN (1977). Worldwide dissemination and importance of the root-knot nematodes *Meloidogyne* spp. *J Nematol* 22: 585–589.
- Sögüt MA, Elekçioğlu İH (2000). *Meloidogyne incognita* Chitwood Nemata: Heteroderidae) ırk-2'nin Farklı Domates Çeşitlerinde Bazı Biyolojik Özellikleri Üzerine Araştırmalar. *Türkiye Entomoloji Dergisi* 24: 113–124 (in Turkish).
- Tanksley SD, McCouch SR (1997). Seed banks and molecular maps: unlocking genetic potential from the wild. *Science* 277: 1063–1066.
- Tanksley SD, Young ND, Paterson AH, Bonierbale MW (1989). RFLP mapping in plant breeding: New tools for an old science. *Nat Biotechnol* 7: 257–263.
- Trudgill DL, Blok VC (2001). Apomictic polyphagous root knot nematodes: exceptionally successful and damaging biotrophic root pathogens. *Annu Rev Phytopathol* 39: 53–77.
- Vrain TC (1999). Engineering natural and synthetic resistance for nematode management. *J Nematol* 31: 424–436.
- Yamamoto T, Hayashi T (2002). New root-knot nematode resistance genes and their STS markers in peach. *Sci Hortic-Amsterdam* 96: 81–90.
- Yücel S, Özarslandan A, Çolak A, Ay T, Can C (2007). Effect of solarization and fumigant applications on soil borne pathogens and root-knot nematodes in greenhouse-grown tomato in Turkey. *Phytoparasitica* 35: 450–456.