

1-1-2019

Fixed point properties for a degenerate Lorentz-Marcinkiewicz space

VEYSEL NEZİR

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

NEZİR, VEYSEL (2019) "Fixed point properties for a degenerate Lorentz-Marcinkiewicz space," *Turkish Journal of Mathematics*: Vol. 43: No. 4, Article 9. <https://doi.org/10.3906/mat-1802-79>
Available at: <https://dctubitak.researchcommons.org/math/vol43/iss4/9>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Fixed point properties for a degenerate Lorentz–Marcinkiewicz space

Veysel NEZİR*

Department of Mathematics, Faculty of Science and Letters, Kafkas University, Kars, Turkey

Received: 19.02.2018

Accepted/Published Online: 27.05.2019

Final Version: 31.07.2019

Abstract: We construct an equivalent renorming of ℓ^1 , which turns out to produce a degenerate ℓ^1 -analog Lorentz–Marcinkiewicz space $\ell_{\delta,1}$, where the weight sequence $\delta = (\delta_n)_{n \in \mathbb{N}} = (2, 1, 1, 1, \dots)$ is a decreasing positive sequence in $\ell^\infty \setminus c_0$, rather than in $c_0 \setminus \ell^1$ (the usual Lorentz situation). Then we obtain its isometrically isomorphic predual $\ell_{\delta,\infty}^0$ and dual $\ell_{\delta,\infty}$, corresponding degenerate c_0 -analog and ℓ^∞ -analog Lorentz–Marcinkiewicz spaces, respectively. We prove that both spaces $\ell_{\delta,1}$ and $\ell_{\delta,\infty}^0$ enjoy the weak fixed point property (w-fpp) for nonexpansive mappings yet they fail to have the fixed point property (fpp) for nonexpansive mappings since they contain an asymptotically isometric copy of ℓ^1 and c_0 , respectively. In fact, we prove for both spaces that there exist nonempty, closed, bounded, and convex subsets with invariant fixed point-free affine, nonexpansive mappings on them and so they fail to have fpp for affine nonexpansive mappings. Also, we show that any nonreflexive subspace of $\ell_{\delta,\infty}^0$ contains an isomorphic copy of c_0 and so fails fpp for strongly asymptotically nonexpansive maps. Finally, we get a Goebel and Kuczumow analogy by proving that there exists an infinite dimensional subspace of $\ell_{\delta,1}$ with fpp for affine nonexpansive mappings.

Key words: Nonexpansive mapping, nonreflexive Banach space, fixed point property, weak fixed point property, closed, bounded, convex set, asymptotically isometric copy of c_0 , asymptotically isometric copy of ℓ^1 , Lorentz–Marcinkiewicz spaces

1. Introduction and preliminaries

A Banach space is said to have the fixed point property (fpp) for nonexpansive mappings [fpp(ne)] if every nonexpansive invariant mapping defined on any nonempty closed, bounded, and convex subset has a fixed point. Wondering how nonexpansive mappings behave if smaller sets are picked, some other fixed point properties such as w-fpp and w*-fpp have become fixed point theorists' center of interest. A Banach space (its dual) is said to have w-fpp (w*-fpp) if every nonexpansive invariant mapping defined on any nonempty weakly (weak*, respectively) compact and convex subset has a fixed point. There have been many strategies to check whether or not a Banach space X (dual space X^*) possesses w-fpp (w*-fpp, respectively) and fpp, and different types of Banach spaces have been the subject of many papers in the last 50 or so years.

In 1965, Browder [4] gave the first example of the largest class of Banach spaces with fpp(ne) until then by proving that Hilbert spaces have fpp(ne). In the same year, Browder [5] and Göhde [14] each independently generalized this to all uniformly convex Banach spaces; for example, $X = L^p$, $1 < p < \infty$, with its usual norm $\|\cdot\|_p$.

*Correspondence: veyselnezir@yahoo.com

2010 *AMS Mathematics Subject Classification:* Primary 46B45, 47H09; Secondary 46B42, 46B10

Again in 1965, Kirk [16] generalized Browder's theorem to all reflexive Banach spaces with normal structure.

There is a strong relationship between the concept of reflexive Banach spaces and Banach spaces having $\text{fpp}(\text{ne})$. As has been mentioned above, it was showed that under the conditions of some geometric properties for Banach spaces such as uniform convexity or normal structure, reflexivity implies $\text{fpp}(\text{ne})$. Less contingent relations have become a matter of curiosity and the role of the norm of the Banach space in the complete equivalence of the two concepts has recently become the center of interest. It has been observed that most nonreflexive classical Banach spaces such as ℓ^1 , Banach space of absolutely summable scalar sequences with its usual absolute summing norm, and c_0 , the Banach space of real valued sequences converging to 0 with its usual absolute supremum norm, fail $\text{fpp}(\text{ne})$ and it is still and has been an open question for over 50 years whether or not all nonreflexive Banach spaces can be renormed to have the fixed point property.

In 2008, Lin [19] gave the first example of a nonreflexive space that can be renormed to have $\text{fpp}(\text{ne})$. He verified this fact by renorming ℓ^1 .

Banach spaces containing nice copies (asymptotically isometric (ai) copies) of ℓ^1 or c_0 cannot have $\text{fpp}(\text{ne})$. Furthermore, if a Banach space is a Banach lattice or has an unconditional basis then it is nonreflexive if and only if it contains either an isomorphic copy of ℓ^1 or c_0 [12, 15, 21]. Because these two nonreflexive Banach spaces ℓ^1 and c_0 share many common properties and they are the fundamental examples of nonreflexive Banach spaces, in order to investigate whether or not nonreflexive Banach spaces can be renormed to have the fixed point property, investigating the question of the second example c_0 has become one of the most important subjects for researchers. In my recent joint paper with Mustafa [25], we gave a positive answer to this question under affinity condition. Unconditionally resolving this problem is still a research topic, and the reason for the difficulty of solving it is that c_0 does not offer adequate tools for researchers; on the contrary, the ℓ^1 space provides possibilities facilitating the research, such as having the Schur property or having weak Opial property, although it has many common properties with c_0 . Thus, for the aim of solving the aforementioned important question, the c_0 -analogue of Lin's result, or to construct bridges that can yield a solution method for that, due to the possibility of offering more tools, researchers have showed interest in working on ℓ^1 -like and c_0 -like spaces. In the literature, c_0 -analogue and ℓ^1 -analogue spaces have been considered, and fixed-point theory oriented properties have been questioned, even by constructing new equivalent norms on ℓ^1 again. By Gamboa de Buen and Nuñez-Medina [11], it has been showed that Lin's result is not the only example for nonreflexive Banach spaces. In their study, first of all, they constructed an equivalent norm on c_0 and obtained isometrically isomorph dual ℓ^1 , and then they proved that there exists another equivalent norm on ℓ^1 with $\text{fpp}(\text{ne})$ by using Lin's properties [20] generalizing [19].

A Banach space, which is very similar to ℓ^1 and offers exactly the same features, but without a norm obtained only by a linear expansion of the usual one, is rarely included in the literature (even the renorming of Gamboa de Buen and Nuñez-Medina is different, too, since it has $\text{fpp}(\text{ne})$). There are some similarities such as Orlicz spaces or Lebesgue space ($L^1[0, 1], \|\cdot\|_1$) but different properties are observed. For example, by Alspach's result [1], $L^1[0, 1]$ does not have w-fpp, contrary to ℓ^1 .

Here, we can note that Maurey [23] showed that $(c_0, \|\cdot\|_\infty)$ and reflexive subspaces of $L^1[0, 1]$ do have w-fpp using ultrafilter techniques. Conversely to Maurey's result, Downling and Lennard [8] showed that every nonreflexive subspace of $L^1[0, 1]$ fails the fixed point property. Before their result, in 1996, by Carothers et al. [6], the analogous result for nonreflexive subspaces of the Lorentz function space $L_{w,1}(0, \infty)$ was established,

but even before that, in 1991, Carothers et al. [7] showed that the Lorentz space $L_{w,1}(\mu)$ enjoys the weak* fixed point property if (X, Σ, μ) is a σ -finite measure space; that is, if C is a weak-star compact convex subset of $L_{w,1}(\mu)$, then every nonexpansive mapping on C has a fixed point.

One of Banach spaces offering the closest features to ℓ^1 's is the ℓ^1 -analogue Lorentz–Marcinkiewicz space $\ell_{w,1}$, where the weight sequence $w = (w_n)_{n \in \mathbb{N}}$ is a decreasing positive sequence in $c_0 \setminus \ell^1$ that I studied in my PhD thesis [24], written under the supervision of Chris Lennard, and its fixed point theory oriented properties were studied, but due to troubles inherited by complexity of its norm depending on its weight sequence's approaching to 0, it has not been proven whether there are completely similar features even in this space.

In this paper, as a result of valuable discussions with Lennard*, by constructing an equivalent renorming of ℓ^1 , we produce a degenerate ℓ^1 -analog Lorentz–Marcinkiewicz space $\ell_{\delta,1}$, where the weight sequence $\delta = (\delta_n)_{n \in \mathbb{N}} = (2, 1, 1, 1, \dots)$ is a decreasing positive sequence in $\ell^\infty \setminus c_0$, rather than in $c_0 \setminus \ell^1$ (the usual Lorentz situation). Then we obtain its isometrically isomorphic predual $\ell_{\delta,\infty}^0$ and dual $\ell_{\delta,\infty}$, corresponding degenerate c_0 -analog and ℓ^∞ -analog Lorentz–Marcinkiewicz spaces, respectively. We prove that both Banach spaces $\ell_{\delta,1}$ and $\ell_{\delta,\infty}^0$ enjoy w-fpp(ne) yet they fail to have fpp(ne) since, respectively, they contain an ai copy of ℓ^1 and c_0 . In fact, we prove that for both spaces, there exist nonempty, closed, bounded, and convex subsets with invariant fixed point-free affine, nonexpansive mappings on them and so they fail to have fpp for affine nonexpansive mappings. Finally, we prove that there exists an infinite dimensional subspace of $\ell_{\delta,1}$ with fpp(ne) for affine mappings so we obtain an analogue result of Goebel and Kuczumow's on ℓ^1 [13] under affinity condition.

Throughout the study, we denote the set of all positive integers and the set of all real numbers by \mathbb{N} and \mathbb{R} , respectively. Throughout this paper our scalar field is \mathbb{R} .

Let $(X, \|\cdot\|)$ be a Banach space and $E \subseteq X$. We will denote the convex hull of E by $\text{co}(E)$.

As usual, we define the Banach space $(c_0, \|\cdot\|_\infty)$ by the vector space of all scalar sequences converging to 0 such that $\|x\|_\infty := \sup_{n \in \mathbb{N}} |x_n|$, for all $x = (x_n)_{n \in \mathbb{N}} \in c_0$; and $(\ell^1, \|\cdot\|_1)$ is the vector space of all absolutely summable scalar sequences such that $\|x\|_1 := \sum_{n=1}^{\infty} |x_n|$ for all $x = (x_n)_{n \in \mathbb{N}} \in \ell^1$.

Definition 1.1 Assume that $(X, \|\cdot\|)$ is a Banach space and K is a nonempty closed, bounded, convex subset.

1. If $U : K \rightarrow K$ is a mapping such that for all $\lambda \in [0, 1]$ and for all $x, y \in K$, $U((1 - \lambda)x + \lambda y) = (1 - \lambda)U(x) + \lambda U(y)$, then U is said to be an affine mapping.
2. If $U : K \rightarrow K$ is a mapping such that $\|U(x) - U(y)\| \leq \|x - y\|$, for all $x, y \in K$ then U is said to be a nonexpansive mapping.

Also, if for every nonexpansive mapping $U : K \rightarrow K$, there exists $z \in K$ with $U(z) = z$, then K is said to have the fixed point property for nonexpansive mappings [fpp(ne)].

3. If $U : K \rightarrow K$ is a mapping such that $\exists \{\beta_{n,m} : n, m \in \mathbb{N}, n \geq m \geq 0\} \subseteq [1, \infty)$ such that $[\forall x, y \in K$ and $\forall n \geq m$, $\|U^n x - U^n y\| \leq \beta_{n,m} \|U^m x - U^m y\|]$, $[\beta_{n,m} \rightarrow 1$ as $n \geq m \rightarrow \infty]$, and $[\beta_{n,m} \rightarrow 1$ as $n \rightarrow \infty, \forall m]$, then U is said to be a strongly asymptotically nonexpansive [17].

*Lennard CJ. Personal communication, 2017.

We should note that in our PhD thesis [24], written under the supervision of Chris Lennard, we studied the usual Lorentz–Marcinkiewicz spaces and their fixed point properties; hence, we can give their definitions below to understand how different the degenerate ones are.

Let $w \in (c_0 \setminus \ell^1)^+$, $w_1 = 1$ and $(w_n)_{n \in \mathbb{N}}$ be decreasing; that is, consider a scalar sequence given by $w = (w_n)_{n \in \mathbb{N}}$, $w_n > 0, \forall n \in \mathbb{N}$ such that $1 = w_1 \geq w_2 \geq w_3 \geq \dots \geq w_n \geq w_{n+1} \geq \dots, \forall n \in \mathbb{N}$ with $w_n \rightarrow 0$ as $n \rightarrow \infty$ and $\sum_{n=1}^{\infty} w_n = \infty$. This sequence is called a weight sequence. For example, $w_n = \frac{1}{n}, \forall n \in \mathbb{N}$.

Definition 1.2

$$l_{w,\infty} := \left\{ x = (x_n)_{n \in \mathbb{N}} \in c_0 \mid \|x\|_{w,\infty} := \sup_{n \in \mathbb{N}} \frac{\sum_{j=1}^n x_j^*}{\sum_{j=1}^n w_j} < \infty \right\} .$$

Here, x^* represents the decreasing rearrangement of sequence x , which is the sequence of $|x| = (|x_j|)_{j \in \mathbb{N}}$, arranged in nonincreasing order, followed by infinitely many zeros when $|x|$ has only finitely many nonzero terms.

This space is nonseparable and an analogue of l_∞ space.

Definition 1.3

$$l_{w,\infty}^0 := \left\{ x = (x_n)_{n \in \mathbb{N}} \in c_0 \mid \limsup_{n \rightarrow \infty} \frac{\sum_{j=1}^n x_j^*}{\sum_{j=1}^n w_j} = 0 \right\} .$$

This is a separable subspace of $l_{w,\infty}$ and an analogue of c_0 space.

Definition 1.4

$$l_{w,1} := \left\{ x = (x_n)_{n \in \mathbb{N}} \in c_0 \mid \|x\|_{w,1} := \sum_{j=1}^{\infty} w_j x_j^* < \infty \right\} .$$

This is a separable subspace of $l_{w,\infty}$ and an analogue of l_1 space with the following facts: $(l_{w,\infty}^0)^* \cong l_{w,1}$ and $(l_{w,1})^* \cong l_{w,\infty}$, where the star denotes the dual of a space while \cong denotes isometrically isomorphic.

More information about Lorentz spaces can be seen in [21].

We also need the following theorems and definitions to obtain our results.

Theorem 1.5 [18] Let X be a Banach space. If X has an unconditional basis (e_n) with unconditional constant $\lambda < \frac{\sqrt{33}-3}{2}$, then X has the w-fpp.

Theorem 1.6 [17] If X is a Banach space containing an isomorphic copy of c_0 , then X fails the fixed point property for affine strongly asymptotically nonexpansive mappings.

Definition 1.7 [9] We say that a Banach space $(X, \|\cdot\|)$ contains an ai copy of ℓ^1 if there exist a sequence $(x_n)_n$ in X and a null sequence $(\varepsilon_n)_n$ in $(0, 1)$ so that

$$\sum_{n=1}^{\infty} (1 - \varepsilon_n) |a_n| \leq \left\| \sum_{n=1}^{\infty} a_n x_n \right\| \leq \sum_{n=1}^{\infty} |a_n| ,$$

for all $(a_n)_n \in \ell^1$.

Definition 1.8 [9] We say that a Banach space $(X, \|\cdot\|)$ contains an ai copy of c_0 if there exist a sequence $(x_n)_n$ in X and a null sequence $(\varepsilon_n)_n$ in $(0, 1)$ so that

$$\sup_n (1 - \varepsilon_n) |a_n| \leq \left\| \sum_{n=1}^{\infty} a_n x_n \right\| \leq \sup_n |a_n|,$$

for all $(a_n)_n \in c_0$.

Theorem 1.9 [9] If a Banach space $(X, \|\cdot\|)$ contains an ai copy of ℓ^1 or an ai copy of c_0 , then X fails fpp(ne).

2. Producing some degenerate Lorentz–Marcinkiewicz spaces $\ell_{\delta,1}$, $\ell_{\delta,\infty}^0$, and $\ell_{\delta,\infty}$, where the weight sequence is in $\ell^\infty \setminus c_0$

For all $x = (x_n)_{n \in \mathbb{N}} \in \ell^1$, we define

$$\|x\| := \|x\|_1 + \|x\|_\infty = \sum_{n=1}^{\infty} |x_n| + \sup_{n \in \mathbb{N}} |x_n|.$$

Clearly $\|\cdot\|$ is an equivalent norm on ℓ^1 with $\|x\|_1 \leq \|x\| \leq 2\|x\|_1, \forall x \in \ell^1$.

We shall call $\|\cdot\|$ the $1 \boxplus \infty$ -norm on ℓ^1 .

Note that $\forall x \in \ell^1, \|x\| = 2x_1^* + x_2^* + x_3^* + x_4^* + \dots$, where z^* is the decreasing rearrangement of $|z| = (|z_n|)_{n \in \mathbb{N}}, \forall z \in c_0$.

Let $\delta_1 := 2, \delta_2 := 1, \delta_3 := 1, \dots, \delta_n := 1, \forall n \geq 4$.

We see that $(\ell^1, \|\cdot\|)$ is a (degenerate) Lorentz space $\ell_{\delta,1}$, where the weight sequence $\delta = (\delta_n)_{n \in \mathbb{N}}$ is a decreasing positive sequence in $\ell^\infty \setminus c_0$, rather than in $c_0 \setminus \ell^1$ (the usual Lorentz situation).

This suggests that $\ell_{\delta,\infty}^0 = (c_0, \|\cdot\|)$ is an isometric predual of $(\ell^1, \|\cdot\|)$ where for all $z \in c_0$,

$$\|z\| := \sup_{n \in \mathbb{N}} \frac{\sum_{j=1}^n z_j^*}{\sum_{j=1}^n \delta_j} = \sup_{n \in \mathbb{N}} \frac{1}{n+1} \sum_{j=1}^n z_j^*.$$

There is a way to rewrite $\|z\|$ without using decreasing rearrangements of $|z|$. This may help with calculations involving this norm.

Fix $z \in c_0$, arbitrary.

$$\|z\| = \sup_{n \in \mathbb{N}} \frac{1}{n+1} \sum_{j=1}^n z_j^*.$$

Note that $\forall n \in \mathbb{N}, \sum_{j=1}^n z_j^* = \sup_{\substack{K \subseteq \mathbb{N} \\ \#(K)=n}} \sum_{i \in K} |z_i|$, where $\#(K)$ is the number of elements in K for all finite subsets

$K \subseteq \mathbb{N}$.

Thus,

$$\begin{aligned} \|z\| &= \sup_{n \in \mathbb{N}} \frac{1}{n+1} \sup_{\substack{K \subseteq \mathbb{N} \\ \#(K)=n}} \sum_{i \in K} |z_i| \\ &= \sup_{n \in \mathbb{N}} \sup_{\substack{K \subseteq \mathbb{N} \\ \#(K)=n}} \frac{1}{\#(K)+1} \sum_{i \in K} |z_i|. \end{aligned}$$

Hence, for all $z \in c_0$,

$$\|z\| = \sup_{\substack{\emptyset \neq K \subseteq \mathbb{N} \\ \#(K) < \infty}} \frac{1}{\#(K)+1} \sum_{i \in K} |z_i|.$$

Also, noting that formula 2.1 can be extended to $\ell^\infty: \forall w = (w_i)_{i \in \mathbb{N}} \in \ell^\infty$, we define

$$\|w\| := \sup_{\substack{\emptyset \neq K \subseteq \mathbb{N} \\ \#(K) < \infty}} \frac{1}{\#(K)+1} \sum_{i \in K} |w_i|.$$

Lemma 2.1 *Dual space of $(\ell^1, \|\cdot\|)$ is isometrically isomorphic to $(\ell^\infty, \|\cdot\|)$; i.e. $(\ell^1, \|\cdot\|)^* \cong (\ell^\infty, \|\cdot\|)$.*

Proof First of all, considering canonical basis $(e_n)_{n \in \mathbb{N}}$, define the sequence $(Q_n)_{n \in \mathbb{N}}$ in ℓ^1 by $Q_1 := \frac{1}{2}e_1$, $Q_2 := \frac{1}{3}(e_2 + e_3)$, $Q_3 := \frac{1}{4}(e_4 + e_5 + e_6)$, $Q_4 := \frac{1}{5}(e_7 + e_8 + e_9 + e_{10}), \dots$ and note that $\|Q_n\| = 1, \forall n \in \mathbb{N}$, and that for any $x := (\xi_n)_{n \in \mathbb{N}} \in \ell^1$, there is unique representation of x with scalars $\alpha_1 = 2\xi_1, \alpha_2 = 3\xi_2, \alpha_3 = 3\xi_3, \alpha_4 = 4\xi_4, \alpha_5 = 4\xi_5, \alpha_6 = 4\xi_6, \alpha_7 = 5\xi_7, \alpha_8 = 5\xi_8, \alpha_9 = 5\xi_9, \alpha_{10} = 5\xi_{10}, \dots$ such that $x = \sum_{n=1}^{\infty} \alpha_n Q_n$.

Now, consider any $f \in \ell^{1*}$. Then, since f is linear and bounded, $f(x) = \sum_{n=1}^{\infty} \alpha_n f(Q_n) = \sum_{n=1}^{\infty} \alpha_n \gamma_n, \gamma_n := f(Q_n), \forall n \in \mathbb{N}$.

Thus, $(\gamma_n)_{n \in \mathbb{N}} \in \ell^\infty$.

Also, $\sup_{\substack{\emptyset \neq K \subseteq \mathbb{N} \\ \#(K) < \infty}} \frac{1}{\#(K)+1} \sum_{i \in K} |\gamma_i| \leq \sup_{k \in \mathbb{N}} |\gamma_k| \leq \|f\|_* = \sup_{\substack{x \in \ell^1 \\ \|x\|=1}} |f(x)|$. \heartsuit where $\|\cdot\|_*$ denotes the operator

norm.

Furthermore, if $(\gamma_n)_{n \in \mathbb{N}} \in \ell^\infty$ is given arbitrarily, we can construct linear functionals in ℓ^{1*} with the elements of ℓ^∞ as follows:

$$g(x) = \sum_{k=1}^{\infty} \alpha_k \gamma_k.$$

Showing linearity is no problem, and for the boundedness of the linear functional g , we will use the inequality 5.2(5) in [22, page 52]. Then we have

$$\begin{aligned}
 |g(x)| &\leq \sum_{k=1}^{\infty} |\alpha_k \gamma_k| \\
 &\leq \sup_{\substack{\emptyset \neq K \subseteq \mathbb{N} \\ \#(K) < \infty}} \frac{1}{\#(K) + 1} \sum_{i \in K} |\gamma_i| \sum_{k=1}^{\infty} \alpha_k^* \\
 &\leq \sup_{\substack{\emptyset \neq K \subseteq \mathbb{N} \\ \#(K) < \infty}} \frac{1}{\#(K) + 1} \sum_{i \in K} |\gamma_i| \|x\|.
 \end{aligned}$$

Thus, indeed the functional g is bounded, and by the linearity, $g \in \ell^{1*}$. Also, the above inequality says that for any $f \in \ell^{1*}$ and for any $x \in \ell^1$,

$$\|f\|_* = \sup_{\|x\|=1} |f(x)| \leq \sup_{\substack{\emptyset \neq K \subseteq \mathbb{N} \\ \#(K) < \infty}} \frac{1}{\#(K) + 1} \sum_{i \in K} |\gamma_i| = \|(\gamma_k)_{k \in \mathbb{N}}\|. \quad \heartsuit\heartsuit$$

Hence, \heartsuit together with $\heartsuit\heartsuit$ tells us that the norm is preserved; i.e., $\|f\|_* = \|(\gamma_k)_{k \in \mathbb{N}}\|$. Therefore, the isomorphism between the two given normed spaces ℓ^{1*} and ℓ^∞ is a fact. Taking an element out of ℓ^{1*} is thus in certain sense the same as speaking about an element out of ℓ^∞ . \square

Remark 2.2 $B_{(\ell^\infty, \|\cdot\|_\infty)} = \{x \in \ell^\infty : \|x\|_\infty \leq 1\}$ has the fixed point property for nonexpansive mappings since it is a hyperconvex metric space and hyperconvex metric spaces have fpp(ne) by Soardi [29].

3. Fixed point properties for $\ell_{\delta, \infty}^0$

Here we take the degenerate c_0 -analog Lorentz–Marcinkiewicz space $\ell_{\delta, \infty}^0$ and begin our section by showing that $\ell_{\delta, \infty}^0$ has w-fpp. Then, in the following two subsections, first we show that $\ell_{\delta, \infty}^0$ contains an ai copy of c_0 and so it fails to have fpp(ne), but in the next subsection, in fact, we see that it fails to have fpp for affine nonexpansive mappings. Then, in the final subsection, we show that any nonreflexive vector subspace of $\ell_{\delta, \infty}^0$ contains an isomorphic copy of c_0 and so fails fpp for strongly asymptotically nonexpansive maps.

3.1. $\ell_{\delta, \infty}^0$ has w-fpp

Theorem 3.1 Let $X := \ell_{\delta, \infty}^0$ and let $\|\cdot\| = \|\cdot\|_{\delta, \infty}$ as we discussed in the above notes. Then Banach space $(X, \|\cdot\|)$ has the weak fixed point property.

Proof To prove this theorem, we will use Theorem 1.5. First, we need to recall the definition of the unconditional constant of an unconditional basis $(u_n)_{n \in \mathbb{N}}$ and for this we can assume the sequence $(u_n)_{n \in \mathbb{N}}$ is normalized; i.e. $\|u_n\| = 1, \forall n \in \mathbb{N}$. Then $(u_n)_{n \in \mathbb{N}}$ is said to be an unconditional basis if for every convergent series $\sum_{j=1}^{\infty} s_j u_j$ (where $(s_j)_{j \in \mathbb{N}}$ is a sequence of scalars) and for every sequence of signs $(\varepsilon_n)_{n \in \mathbb{N}} (\varepsilon_n = \pm 1)$ there exists a constant $\lambda \geq 1$ such that

$$\left\| \sum_{j=1}^{\infty} s_j \varepsilon_j u_j \right\| \leq \lambda \left\| \sum_{j=1}^{\infty} s_j u_j \right\|.$$

Then, the unconditionality constant of the unconditional basis is the smallest λ satisfying the above condition; that is,

$$\lambda := \sup_{\substack{\varepsilon = (\varepsilon_n)_{n \in \mathbb{N}} \\ \varepsilon_n = \pm 1}} \frac{\|\sum_{j=1}^{\infty} s_j \varepsilon_j u_j\|}{\|\sum_{j=1}^{\infty} s_j u_j\|}.$$

Now, for our space, we have a normalized basis $(u_n)_{n \in \mathbb{N}} = ((n + 1)\varepsilon_n)_{n \in \mathbb{N}}$. Define $S_{\mathbb{N}} := \{\pi : \mathbb{N} \rightarrow \mathbb{N} \mid \pi \text{ 1-1 and onto map}\}$ set of all permutations. Then $\forall \varepsilon_j = \pm 1, \forall \pi \in S_{\mathbb{N}}$, and for any sequence of scalars $(s_j)_{j \in \mathbb{N}}$,

$$\left\| \sum_{n=1}^{\infty} \varepsilon_n s_{\pi(n)} u_n \right\|_{\delta, \infty} = \left\| \sum_{n=1}^{\infty} s_n u_n \right\|_{\delta, \infty} = \left\| \sum_{n=1}^{\infty} s_n^* u_n \right\|_{\delta, \infty}.$$

Hence, $\lambda = 1 < \frac{\sqrt{33}-3}{2}$. □

3.2. $\ell_{\delta, \infty}^0$ contains an ai copy of c_0

Theorem 3.2 $\forall \delta \in c_0 \setminus \ell^1, \exists Z \subseteq \ell_{\delta, \infty}^0$, such that Z is an ai copy of c_0 and so $\ell_{\delta, \infty}^0$ fails the fixed point property for affine, $\|\cdot\|_{\delta, \infty}$ -nonexpansive mappings.

Proof Let $\delta \in c_0 \setminus \ell^1$ be given. Fix $\varepsilon_j \downarrow_j 0, \varepsilon_j \in (0, 1)$. Then choose a sequence $(r_n)_{n \in \mathbb{N}}$ in \mathbb{N} so that $1 \leq r_1 < r_2 < r_3 < \dots$ and each term is large enough such that

$$\frac{\sum_{j=1}^{r_n} \delta_j}{\sum_{j=1}^{r_{n+1}} \delta_j} \leq \varepsilon_{n+1}, \forall n \in \mathbb{N} \text{ (since } \sum_{j=1}^{\infty} \delta_j = \infty \text{) ,}$$

i.e. increasing to ∞ fast enough so that

$$\frac{\sum_{j=1}^{r_n} \delta_j}{\sum_{j=1}^{r_{n+1}} \delta_j} \leq \varepsilon_{n+1}, \forall n \in \mathbb{N}.$$

Define $\eta_1 = g_1 = (\delta_1, \delta_2, \delta_3, \dots, \delta_{r_1}, 0, 0, 0, \dots)$.

Then notice $\|\eta_1\|_{\delta, \infty} = \|g_1\|_{\delta, \infty} = 1$, and next define

$$g_2 = (0, 0, 0, \dots, 0, \delta_{r_1+1}, \delta_{r_1+2}, \dots, \delta_{r_2-1}, \delta_{r_2}, 0, 0, 0, \dots)$$

↑

r_1 st term,

and then define $\eta_2 := g_1 + g_2$ and notice $\|\eta_2\|_{\delta, \infty} = \|g_1 + g_2\|_{\delta, \infty} = 1$, and next define

$$g_3 = (0, 0, 0, \dots, 0, \delta_{r_2+1}, \delta_{r_2+2}, \dots, \delta_{r_3-1}, \delta_{r_3}, 0, 0, 0, \dots)$$

↑

r_2 nd term,

and then define $\eta_3 := g_1 + g_2 + g_3$ and notice $\|\eta_3\|_{\delta,\infty} = \|g_1 + g_2 + g_3\|_{\delta,\infty} = 1$. Continuing in this way, we obtain a sequence $(g_n)_{n \in \mathbb{N}}$ and so $(\eta_n)_{n \in \mathbb{N}}$ such that $\|\eta_n\|_{\delta,\infty} = \|g_1 + g_2 + g_3 + \dots + g_n\|_{\delta,\infty} = 1, \forall n \in \mathbb{N}$. Then define the set

$$Z := \left\{ x = t_1 g_1 + t_2 g_2 + t_3 g_3 + \dots = \sum_{n=1}^{\infty} t_n g_n \mid t = (t_n)_{n \in \mathbb{N}} \in c_0 \right\}.$$

Then let $t \in c_0$ and $x \in Z$ be arbitrarily given. Then,

$$\begin{aligned} \|x\|_{\delta,\infty} &= \|t_1 g_1 + t_2 g_2 + t_3 g_3 + \dots\|_{\delta,\infty} = \left\| \sum_{n=1}^{\infty} t_n g_n \right\|_{\delta,\infty} \\ &= \| |t_1| g_1 + |t_2| g_2 + |t_3| g_3 + \dots \|_{\delta,\infty} = \left\| \sum_{n=1}^{\infty} |t_n| g_n \right\|_{\delta,\infty} \\ &\leq \| |t| \|_{\infty} \|g_1 + g_2 + g_3 + \dots\|_{\delta,\infty} = \left\| \sum_{n=1}^{\infty} |t| \|_{\infty} g_n \right\|_{\delta,\infty} \\ &= \|t\|_{\infty} \|g_1 + g_2 + g_3 + \dots\|_{\delta,\infty} = \|t\|_{\infty} \left\| \sum_{n=1}^{\infty} g_n \right\|_{\delta,\infty} \\ &= \|t\|_{\infty} \|\delta\|_{\delta,\infty} \\ &= \|t\|_{\infty}. \end{aligned}$$

Also,

$$\|x\|_{\delta,\infty} \geq \frac{|t_1| \sum_{j=1}^{r_1} \delta_j}{\sum_{j=1}^{r_1} \delta_j} = |t_1| \geq |t_1|(1 - \varepsilon_1).$$

Next,

$$\begin{aligned} \|x\|_{\delta,\infty} &\geq \|t_2 g_2\|_{\delta,\infty} \\ &\geq \frac{|t_2| \sum_{j=r_1+1}^{r_2} \delta_j}{\sum_{j=1}^{r_2-r_1} \delta_j} = |t_2| \frac{\sum_{j=1}^{r_2} \delta_j - \sum_{j=1}^{r_1} \delta_j}{\sum_{j=1}^{r_2-r_1} \delta_j} \\ &\geq |t_2| \frac{\sum_{j=1}^{r_2} \delta_j - \sum_{j=1}^{r_1} \delta_j}{\sum_{j=1}^{r_2} \delta_j} \\ &\geq |t_2|(1 - \varepsilon_2). \end{aligned}$$

Similarly,

$$\begin{aligned} \|x\|_{\delta,\infty} &\geq \|t_3 g_3\|_{\delta,\infty} \\ &\geq \frac{|t_3| \sum_{j=r_2+1}^{r_3} \delta_j}{\sum_{j=1}^{r_3-r_2} \delta_j} = |t_3| \frac{\sum_{j=1}^{r_3} \delta_j - \sum_{j=1}^{r_2} \delta_j}{\sum_{j=1}^{r_3-r_2} \delta_j} \\ &\geq |t_3| \frac{\sum_{j=1}^{r_3} \delta_j - \sum_{j=1}^{r_2} \delta_j}{\sum_{j=1}^{r_3} \delta_j} \\ &\geq |t_3|(1 - \varepsilon_3). \end{aligned}$$

Then, inductively, we obtain $\|x\|_{\delta,\infty} \geq |t_n|(1 - \varepsilon_n), \forall n \in \mathbb{N}$. Hence,

$$\begin{aligned} \sup_{\nu \in \mathbb{N}} (1 - \varepsilon_\nu) |t_\nu| &\leq \|x\|_{\delta,\infty} \leq \sup_{\nu \in \mathbb{N}} |t_\nu| \leq \sup_{\nu \in \mathbb{N}} (1 + \varepsilon_\nu) |t_\nu| \\ \sup_{\nu \in \mathbb{N}} (1 - \varepsilon_\nu) |t_\nu| &\leq \left\| \sum_{\nu=1}^{\infty} t_\nu \delta_\nu \right\|_{\delta,\infty} \leq \sup_{\nu \in \mathbb{N}} (1 + \varepsilon_\nu) |t_\nu|. \end{aligned}$$

□

3.3. $\ell_{\delta,\infty}^0$ fails fpp for affine $\|\cdot\|_{\delta,\infty}$ -nonexpansive mappings

Theorem 3.3 *Define*

$$E := \{u = (s_1 \delta_1, s_2 \delta_2, s_3 \delta_3, \dots) \mid s \in c_0, 1 = s_1 \geq s_2 \geq \dots \geq 0\}.$$

Then $E \subseteq l_{\delta,\infty}^0$ is a convex, closed, bounded set, and $\exists T : E \rightarrow E$ s.t. T is fixed point free, $\|\cdot\|_{\delta,\infty}$ -nonexpansive, affine mapping.

Proof It is clear that E is convex. We need to show $E \subseteq l_{\delta,\infty}^0$. Let $u \in E$ be given. Then $u^* = u$ and

$$\|u\|_{\delta,\infty} = \sup_{n \in \mathbb{N}} \frac{\sum_{j=1}^n u_j^*}{\sum_{j=1}^n \delta_j}.$$

$$\Psi_n(u) = \frac{\sum_{j=1}^n s_j \delta_j}{\sum_{j=1}^n \delta_j} \leq \frac{\sum_{j=1}^n 1 \delta_j}{\sum_{j=1}^n \delta_j} = 1.$$

Hence, $\|u\|_{\delta,\infty} \leq 1, \forall u \in E$. Thus, $E \subseteq l_{\delta,\infty}$ and, in fact, $E \subseteq \mathcal{B}_{l_{\delta,\infty}}$. Now, to show $E \subseteq l_{\delta,\infty}^0$, we need to show for $u \in E, \Psi_n(u) \xrightarrow{n} 0$, and then we would prove $u \in l_{\delta,\infty}^0$.

Now, first of all,

$$0 \leq \Psi_n(u) = \sum_{j=1}^n s_j \gamma_j^{(n)}$$

where

$$\gamma_j^{(n)} = \frac{\delta_j}{\sum_{k=1}^n \delta_k} > 0 \text{ and } \sum_{j=1}^n \gamma_j^{(n)} = 1.$$

Then fix $\varepsilon > 0$ and choose $N = N_\varepsilon$ s.t. $\forall j \geq N_\varepsilon, s_j < \frac{\varepsilon}{2}$.

Let $n > N_\varepsilon$, n arbitrary.

$$\begin{aligned} \Psi_n(u) &= \sum_{j=1}^N s_j \gamma_j^{(n)} + \sum_{j=N+1}^n s_j \gamma_j^{(n)} \\ &\leq (1) \sum_{j=1}^N \gamma_j^{(n)} + \sum_{j=N+1}^n s_j \gamma_j^{(n)} \\ &< \sum_{j=1}^N \gamma_j^{(n)} + \frac{\varepsilon}{2} \sum_{j=N+1}^n \gamma_j^{(n)} \text{ but since } \sum_{j=N+1}^n \gamma_j^{(n)} < 1 \\ &< \sum_{j=1}^N \gamma_j^{(n)} + \frac{\varepsilon}{2} \\ &= \sum_{j=1}^N \left(\frac{\delta_j}{\sum_{k=1}^n \delta_k} \right) + \frac{\varepsilon}{2} \\ &= \frac{\sum_{j=1}^{N_\varepsilon} \delta_j}{\sum_{k=1}^n \delta_k} + \frac{\varepsilon}{2} =: \frac{K(\varepsilon)}{\sum_{k=1}^n \delta_k} + \frac{\varepsilon}{2}. \end{aligned}$$

Choose $M_\varepsilon > N_\varepsilon$ s.t. $\forall n \geq M_\varepsilon$,

$$\frac{K(\varepsilon)}{\sum_{k=1}^n \delta_k} < \frac{\varepsilon}{2}.$$

Then $\forall n \geq M_\varepsilon (> N_\varepsilon)$,

$$0 \leq \Psi_n(u) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \text{ and so}$$

$$\lim_{n \rightarrow \infty} \Psi_n(u) = 0.$$

Hence, $E \subseteq l_{\delta, \infty}^0$. □

Claim 3.4 E is closed.

Proof Let $(u_n)_{n \in \mathbb{N}}$ be any sequence in E norm convergent to a $u_0 \in l_{\delta, \infty}$, i.e. $u_n \xrightarrow[n]{\|\cdot\|} u_0$ in norm then $u_n \xrightarrow[n]{} u_0$ coordinatewise.

Say $u_0 := (u_{0,1}, u_{0,2}, u_{0,3}, \dots, u_{0,j}, \dots)$ and $u_n := (s_1^n \delta_1, s_2^n \delta_2, s_3^n \delta_3, \dots, s_j^n \delta_j, \dots)$. Then $u_{0,1} = \delta_1$ and $s_j^n \delta_j \xrightarrow[n]{} u_{0,j}$ pointwise. Then $s_j^n \xrightarrow[n]{} \frac{u_{0,j}}{\delta_j} =: s_j$ $s_1 = \frac{u_{0,1}}{\delta_1} = 1$, and each $s_j^n \geq 0 \Rightarrow s_j \geq 0$, by pointwise convergence. Now, fix $j \in \mathbb{N}$, and we can show $s_j \geq s_{j+1}$ by taking the limit as n goes to ∞ using the inequality $s_j^n \geq s_{j+1}^n$, since $s_j^n \xrightarrow[n]{} s_j$. Hence, $s_j \geq s_{j+1}, \forall j \in \mathbb{N}$. Now, we need to show $s_j \xrightarrow[j]{} 0$. Since $u_0 = (u_{0,j})_{j \in \mathbb{N}} \in l_{\delta, \infty}^0 \subseteq c_0$, $u_{0,j} \xrightarrow[j]{} 0$. Now we know that $u_0 = (s_1 \delta_1, s_2 \delta_2, \dots, s_j \delta_j, \dots)$, $1 = s_1 \geq s_2 \geq \dots \geq s_j \geq \dots \geq 0$, and so $u_0^* = u_0$. Recall that $u_0 \in l_{\delta, \infty}^0$ by hypothesis.

$$\Leftrightarrow \lim_{n \rightarrow \infty} \frac{\sum_{j=1}^n (u_0^*)_j}{\sum_{j=1}^n \delta_j} = 0.$$

Set for each $n \in \mathbb{N}$, $\Lambda_n := \frac{\sum_{j=1}^n (u_0^*)_j}{\sum_{j=1}^n \delta_j}$, and then $\Lambda_n = \frac{\sum_{j=1}^n s_j \delta_j}{\sum_{j=1}^n \delta_j} \xrightarrow{n} 0$.

Suppose we get a contradiction that $s_j \not\xrightarrow{j} 0$.

However, we know $s_j \downarrow_j$ and $s_j \geq 0, \forall j \in \mathbb{N}$. Thus, $\exists L > 0$ s.t. $s_j \xrightarrow{j} L$. Then,

$$\Lambda_n = \frac{\sum_{j=1}^n s_j \delta_j}{\sum_{j=1}^n \delta_j} \geq \frac{\sum_{j=1}^n L \delta_j}{\sum_{j=1}^n \delta_k} = L > 0, \forall n \in \mathbb{N}.$$

This is a clear contradiction since we obtain that $\forall n \in \mathbb{N}, \Lambda_n \geq L > 0$, whereas $\Lambda_n \xrightarrow{n} 0$. In conclusion, $E \subseteq l_{\delta, \infty}^0$ is a closed bounded convex subset. □

Now we will prove the following claim.

Claim 3.5 $\exists T : E \rightarrow E \quad \|\cdot\|_{\delta, \infty}$ -nonexpansive and fixed point free.

Proof Indeed, consider T as the right shift.

That is,

$$\begin{aligned} T : u = (s_j \delta_j)_{j \in \mathbb{N}} &\longrightarrow Tu = (1\delta_1, s_1\delta_2, s_2\delta_3, s_3\delta_4, \dots) \\ &= (1\delta_1, 1\delta_2, s_2\delta_3, s_3\delta_4, \dots). \end{aligned}$$

T is clearly fixed point free. Indeed, assume $\exists u \in E$ s.t. $Tu = u$ but then that means $1 = s_1 = s_2 = s_3 = \dots = s_k \Leftrightarrow u = (\delta_1, \delta_2, \delta_3, \dots) \in l_{\delta, \infty} - l_{\delta, \infty}^0 \Rightarrow u \notin E$, which would be a contradiction. Now, let us see that T is nonexpansive. Hence, let $u, z \in E$ and say $u = (s_1\delta_1, s_2\delta_2, s_3\delta_3, \dots)$, $z = (s_1\delta_1, s_2\delta_2, s_3\delta_3, \dots)$ for some scalars s_n and v_n .

Then $Tu = (1\delta_1, s_1\delta_2, s_2\delta_3, s_3\delta_4, \dots) = (1\delta_1, 1\delta_2, s_2\delta_3, s_3\delta_4, \dots)$ and

$Tz = (1\delta_1, v_1\delta_2, v_2\delta_3, v_3\delta_4, \dots) = (1\delta_1, 1\delta_2, v_2\delta_3, v_3\delta_4, \dots)$.

Note that if $\beta = (\beta_j)_{j \in \mathbb{N}} \in l_{\delta, \infty}$ is s.t. $0 \leq \alpha_j \leq \beta_j, \forall j \in \mathbb{N}$,

then $\alpha = (\alpha_j)_{j \in \mathbb{N}} \in l_{\delta, \infty}$ and $[\|\alpha\|_{\delta, \infty} \leq \|\beta\|_{\delta, \infty}] \quad [\diamond \diamond]$

Also,

$$\forall u \in c_0, \forall n \in \mathbb{N}, \sum_{j=1}^n u_j^* = \max_{\substack{F \subseteq \mathbb{N} \\ \#(F) = n}} \sum_{k \in F} |u_k| \quad [\diamond \diamond]$$

and so it follows that $[\diamond \diamond] \Rightarrow [\diamond \diamond]$. Hence,

$$\begin{aligned} \|u - z\|_{\delta, \infty} &= \|(0, (s_2 - v_2)\delta_2, (s_3 - v_3)\delta_3, (s_4 - v_4)\delta_4, \dots)\|_{\delta, \infty} \\ &= \|(|s_2 - v_2| \delta_2, |s_3 - v_3| \delta_3, |s_4 - v_4| \delta_4, \dots)\|_{\delta, \infty} \\ &= \sup_{n \in \mathbb{N}} \left[\frac{\sum_{k=1}^n |s_{k+1} - v_{k+1}| \delta_{k+1}}{\sum_{k=1}^n \delta_j} \right]. \end{aligned}$$

Furthermore,

$$\begin{aligned} \|Tu - Tz\|_{\delta, \infty} &= \|(0, 0, (s_2 - v_2)\delta_3, (s_3 - v_3)\delta_4, (s_4 - v_4)\delta_5, \dots)\|_{\delta, \infty} \\ &= \|(0, 0, |s_2 - v_2|\delta_3, |s_3 - v_3|\delta_4, |s_4 - v_4|\delta_5, \dots)\|_{\delta, \infty} \\ &= \|(|s_2 - v_2|\delta_3, |s_3 - v_3|\delta_4, |s_4 - v_4|\delta_5, \dots)\|_{\delta, \infty} \\ &= \sup_{n \in \mathbb{N}} \left[\frac{\sum_{k=1}^n |s_{k+1} - v_{k+1}| \delta_{k+2}}{\sum_{k=1}^n \delta_k} \right]. \end{aligned}$$

Thus, $\|Tu - Tz\|_{\delta, \infty} \leq \|u - z\|_{\delta, \infty}$. □

The proof of Theorem 3.3 is complete.

3.4. Any closed nonreflexive vector subspace Z of $\ell^0_{\delta, \infty}$ contains an isomorphic copy of c_0

Theorem 3.6 *Let Z be any closed, nonreflexive vector subspace of $\ell^0_{\delta, \infty}$. Then Z contains an isomorphic copy of c_0 and so $(Z, \|\cdot\|_{\delta, \infty})$ fails the fixed point property for strongly asymptotically nonexpansive maps.*

Proof We know that if Z is nonreflexive then, by [21, Theorem 1.c.12], \exists an isomorphic copy of ℓ^1 or c_0 inside Z . □

Claim 3.7 *Z does not contain an isomorphic copy of ℓ^1 and so it contains an isomorphic copy of c_0 .*

Proof By contradiction, assume not, i.e. $\ell^1 \lesssim \ell^0_{\delta, \infty}$. Then, by well-known facts in the literature [3, 24, Proposition 5.1.1, 5.1.2, 5.1.3], there exists a subspace Z of the dual space of $(\ell^0_{\delta, \infty})^*$, i.e. $Z \leq (\ell^0_{\delta, \infty})^*$ such that $(\ell^1)^*$ is isometrically identical to $(\ell^0_{\delta, \infty})^*/Z$, i.e. $(\ell^1)^* \cong (\ell^0_{\delta, \infty})^*/Z$. Moreover, $(\ell^1)^* \cong l^\infty$ and $(\ell^0_{\delta, \infty})^*/Z \cong (l_{\delta, 1}/Z)$. However, we know that $l_{\delta, 1}$ is separable and so is the quotient space $l_{\delta, 1}/Z$. However, l^∞ is nonseparable. Hence, we get a contradiction; that is, $(\ell^1)^* \cong (\ell^0_{\delta, \infty})^*/Z$ cannot be true and so Z cannot contain an isomorphic copy of ℓ^1 . Therefore, Z has to contain an isomorphic copy of c_0 . □

The proof of Theorem 3.6 is complete.

4. Fixed Point Properties for $\ell_{\delta, 1}$

As we discussed in Section 2, recall that ℓ^1 with equivalent norm $\|\cdot\|$ is a (degenerate) Lorentz space $\ell_{\delta, 1}$ with the weight sequence $\delta = (\delta_n)_{n \in \mathbb{N}} = (2, 1, 1, 1, \dots) \in c_0 \setminus \ell^1$ such that for all $x = (x_n)_{n \in \mathbb{N}} \in \ell^1$, $\|x\| = \sum_{n=1}^\infty \delta_n x_n^*$, where $x = (x_n^*)_{n \in \mathbb{N}}$ is decreasing rearrangement of x .

In the following two subsections, first we show that $\ell_{\delta, 1}$ contains an ai of ℓ^1 and so it fails to have fpp(ne), but in the next subsection, in fact, we see that it fails to have fpp for affine nonexpansive mappings. Then, in the final subsection, we show that $\ell_{\delta, 1}$ has w-fpp(ne).

4.0.1. $\ell_{\delta, 1}$ contains an ai copy of ℓ^1

Theorem 4.1 *$\ell_{\delta, 1}$ contains an ai copy of ℓ^1 and so it fails the fixed point property for $\|\cdot\|$ -nonexpansive mappings.*

Proof Consider the sequence $(Q_n)_{n \in \mathbb{N}}$ constructed as in Lemma 2.1 using canonical basis $(e_n)_{n \in \mathbb{N}}$ given by $Q_1 := \frac{1}{2}e_1$, $Q_2 := \frac{1}{3}(e_2 + e_3)$, $Q_3 := \frac{1}{4}(e_4 + e_5 + e_6)$, $Q_4 := \frac{1}{5}(e_7 + e_8 + e_9 + e_{10}), \dots$ and note that $\|Q_n\| = 1, \forall n \in \mathbb{N}$.

Then for all $t = (t_n)_{n \in \mathbb{N}} \in \ell^1$,

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} t_n Q_n \right\| &= \left\| t_1 \frac{1}{2}e_1 + t_2 \frac{1}{3}(e_2 + e_3) + t_3 \frac{1}{4}(e_4 + e_5 + e_6) \right. \\ &\quad \left. + t_4 \frac{1}{5}(e_7 + e_8 + e_9 + e_{10}) + \dots \right\| \\ &= \left[\frac{1}{2}|t_1| + \frac{2}{3}|t_2| + \frac{3}{4}|t_3| + \frac{4}{5}|t_4| + \dots \right] \\ &\quad + \left[\frac{1}{2}|t_1| \sqrt{\frac{1}{3}}|t_2| \sqrt{\frac{1}{4}}|t_3| \sqrt{\frac{1}{5}}|t_4| + \dots \right] \\ &\leq |t_1| + |t_2| + |t_3| + |t_4| + \dots \end{aligned}$$

and

$$\left\| \sum_{n=1}^{\infty} t_n Q_n \right\| \geq \sum_{n=1}^{\infty} (1 - \varepsilon_n) |t_n|$$

where $\varepsilon_n := \frac{1}{n+1}, \forall n \in \mathbb{N}$. □

4.0.2. $\ell_{\delta,1}$ fails fpp for affine $\|\cdot\|$ -nonexpansive mappings

Theorem 4.2 $\ell_{\delta,1}$ fails fpp for affine $\|\cdot\|$ -nonexpansive mappings

Proof Consider the set

$$C := \left\{ x = (x_n)_{n \in \mathbb{N}} \in \ell^1 : \text{each } x_n \geq 0 \ \& \ \sum_{n=1}^{\infty} x_n = 1 \right\}.$$

Clearly, $C \subseteq \ell^1$ is a closed, bounded, and convex subset. Now consider the right shift mapping $U : C \rightarrow C$ defined by $Ux := (0, x_1, x_2, x_3, \dots), \forall x \in C$. Then U is a fixed point free, affine $\|\cdot\|$ -nonexpansive mapping. □

4.0.3. $\ell_{\delta,1}$ has w-fpp(ne)

Lemma 4.3 $\ell_{\delta,1}$ has the weak fixed point property.

Proof Recall that we named the renorming of ℓ^1 with the equivalent norm $\|\cdot\|$ by a degenerate Lorentz–Marcinkiewicz space $\ell_{\delta,1}$. Let $\emptyset \neq K \subseteq \ell^1$ be weakly compact. Let $T : K \rightarrow K$ be a $\|\cdot\|$ -nonexpansive map. Since ℓ^1 has the Schur property, i.e. weak convergence is equivalent to norm convergence, K is norm compact. Also, T is norm-to-norm continuous. Hence, by Schauder’s theorem [28], T has a fixed point. □

5. An infinite-dimensional subspace of $\ell_{\delta,1} = (\ell^1, \|\cdot\|)$ with fpp(ne)

In this section, we will show that there exists a large class of nonweakly* compact, closed, bounded, and convex subsets of $\ell_{\delta,1} = (\ell^1, \|\cdot\|)$ with fpp(ne) under affinity condition using the ideas of Goebel and Kuczumow [13] where they show that there exists a large class of nonweakly* compact, closed, bounded, and convex subsets of $(\ell^1, \|\cdot\|_1)$ with fpp(ne).

In 1979, Goebel and Kuczumow showed the above fact, generalizing the result for the following class of sets.

Example 5.1 Fix $b \in (0, 1)$. Define the sequence $(f_n)_{n \in \mathbb{N}}$ in c_0 by setting $f_1 := be_1$, $f_2 := be_2$, and $f_n := e_n$, for all integers $n \geq 3$. Next, define the closed, bounded, convex subset $E = E_b$ of ℓ^1 by

$$E := \left\{ \sum_{n=1}^{\infty} t_n f_n : \text{each } t_n \geq 0 \text{ and } \sum_{n=1}^{\infty} t_n = 1 \right\} .$$

However, we will be working on the following class.

Example 5.2 Fix $b \in (0, \frac{2}{3}]$. Define the sequence $(f_n)_{n \in \mathbb{N}}$ in c_0 by setting $f_1 := be_1$, $f_2 := be_2$, and $f_n := e_n$, for all integers $n \geq 3$. Next, define the closed, bounded, convex subset $E = E_b$ of ℓ^1 by

$$E := \left\{ \sum_{n=1}^{\infty} t_n f_n : \text{each } t_n \geq 0 \text{ and } \sum_{n=1}^{\infty} t_n = 1 \right\} .$$

In order to obtain our results, we will be using an easy fact below such that its proof is straightforward.

Lemma 5.3 Let $(X, \|\cdot\|)$ be a Banach space.

1. If X has the Banach–Saks property and $x \in X$ is the weak limit of a bounded sequence $(x_n)_n$, then there exists a subsequence $(x_{n_k})_k$ whose Cesaro norm limit is x such that if s is defined by

$$s(y) = \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x_{n_k} - y \right\| , \forall y \in X , \text{ then we have } s(x) = 0 \text{ and } s(y) = \|y - x\| , \forall y \in X .$$

2. If X has the weak Banach–Saks property and $x \in X$ is the weak limit of the sequence $(x_n)_n$, then there exists a subsequence $(x_{n_k})_k$ whose Cesaro norm limit is x such that if s is defined by

$$s(y) = \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x_{n_k} - y \right\| , \forall y \in X , \text{ then we have } s(x) = 0 \text{ and } s(y) = \|y - x\| , \forall y \in X .$$

Hence, due to the weak Banach–Saks property of our space, which can be deduced by the works [2, 26, 27], the above applies.

Theorem 5.4 The set E defined as in Example 5.2 has the fixed point property for $\|\cdot\|$ -nonexpansive mappings where the norm $\|\cdot\|$ on ℓ^1 is given as follows: $\|x\| = \|x\|_1 + \|x\|_\infty, \forall x \in \ell^1$.

Proof We will be using the proof steps of Goebel and Kuczumow given in detail as in Everest’s PhD thesis [10], written under the supervision of Lennard. Let $T : E \rightarrow E$ be a nonexpansive mapping. Then there exists a sequence $(x^{(n)})_{n \in \mathbb{N}} \in E$ such that $\|Tx^{(n)} - x^{(n)}\| \xrightarrow{n} 0$ and so $\|Tx^{(n)} - x^{(n)}\|_1 \xrightarrow{n} 0$. Without loss of generality, passing to a subsequence if necessary, there exists $z \in \ell^1$ such that $x^{(n)}$ converges to z in weak* topology. Then, by Lemma 5.3, we can define a function $s : \ell^1 \rightarrow [0, \infty)$ by

$$s(y) = \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x^{(k)} - y \right\| , \forall y \in \ell^1$$

and so

$$s(y) = \|y - z\|, \forall y \in \ell^1.$$

Next, define

$$W := \bar{E}^{w^*} = \left\{ \sum_{n=1}^{\infty} t_n f_n : \text{each } t_n \geq 0 \text{ and } \sum_{n=1}^{\infty} t_n \leq 1 \right\}.$$

Case 1: $z \in E$.

Then we have $s(Tz) = \|Tz - z\|$ and

$$\begin{aligned} s(Tz) &= \limsup_m \left\| Tz - \frac{1}{m} \sum_{k=1}^m x^{(k)} \right\| \\ &\leq \limsup_m \left\| Tz - T\left(\frac{1}{m} \sum_{k=1}^m x^{(k)}\right) \right\| + \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x^{(k)} - T\left(\frac{1}{m} \sum_{k=1}^m x^{(k)}\right) \right\|. \end{aligned}$$

Then, since T is affine,

$$\begin{aligned} s(Tz) &\leq \limsup_m \left\| Tz - T\left(\frac{1}{m} \sum_{k=1}^m x^{(k)}\right) \right\| + \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x^{(k)} - \frac{1}{m} \sum_{k=1}^m Tx^{(k)} \right\| \\ &\leq \limsup_m \left\| z - \frac{1}{m} \sum_{k=1}^m x^{(k)} \right\| \\ &= s(z). \end{aligned}$$

Therefore, $\|z - Tz\| \leq 0$ and so $Tz = z$.

Case 2: $z \in W \setminus E$.

Then z is of the form $\sum_{n=1}^{\infty} \gamma_n f_n$ such that $\sum_{n=1}^{\infty} \gamma_n < 1$ and $\gamma_n \geq 0, \forall n \in \mathbb{N}$.

Define $\delta := 1 - \sum_{n=1}^{\infty} \gamma_n$ and next define

$$h_\lambda := (\gamma_1 + \lambda\delta)f_1 + (\gamma_2 + (1 - \lambda)\delta)f_2 + \sum_{n=3}^{\infty} \gamma_n f_n.$$

We want h_λ to be in E , so we restrict values of λ to be in $[-\frac{\gamma_1}{\delta}, \frac{\gamma_2}{\delta} + 1]$, and then

$$\begin{aligned} \|h_\lambda - z\| &= \|\lambda\delta f_1 + (1 - \lambda)\delta f_2\| \\ &= \|(\lambda\delta b, (1 - \lambda)\delta b, 0, 0, \dots)\| \\ &= b\delta \max\{|\lambda|, |1 - \lambda|\} + b\delta|\lambda| + b\delta|1 - \lambda| \\ &= \max \begin{cases} 2b\delta - 3b\delta\lambda & \text{if } \lambda \in [-\frac{\gamma_1}{\delta}, 0), \\ 2b\delta - b\delta\lambda & \text{if } \lambda \in [0, \frac{1}{2}), \\ b\delta(1 + \lambda) & \text{if } \lambda \in (\frac{1}{2}, 1], \\ 3b\delta\lambda - b\delta & \text{if } \lambda \in (1, \frac{\gamma_2}{\delta} + 1]. \end{cases} \end{aligned}$$

Define

$$\Gamma := \min_{\lambda \in [-\frac{\gamma_1}{\delta}, \frac{\gamma_2}{\delta} + 1]} \|h_\lambda - z\|.$$

Therefore, $\|h_\lambda - z\|$ is minimized when $\lambda \in [0, 1]$ with unique minimizer such that its minimum value would be $\Gamma = \frac{3b\delta}{2}$.

Now fix $y \in E$ of the form $\sum_{n=1}^{\infty} t_n f_n$ such that $\sum_{n=1}^{\infty} t_n = 1$ with $t_n \geq 0, \forall n \in \mathbb{N}$.

Then,

$$\begin{aligned} \|y - z\| &= \left\| \sum_{k=1}^{\infty} t_k f_k - \sum_{k=1}^{\infty} \gamma_k f_k \right\| \\ &= \left\| (t_1 - \gamma_1)be_1 + (t_2 - \gamma_2)be_2 + (t_3 - \gamma_3)e_3 + (t_4 - \gamma_4)e_4 + \dots \right\| \\ &= \max \left\{ \begin{array}{l} 2b|t_1 - \gamma_1| + b|t_2 - \gamma_2| + |t_3 - \gamma_3| \\ + |t_4 - \gamma_4| + |t_5 - \gamma_5| + \dots, \\ 2b|t_2 - \gamma_2| + b|t_1 - \gamma_1| + |t_3 - \gamma_3| \\ + |t_4 - \gamma_4| + |t_5 - \gamma_5| + \dots, \\ 2|t_3 - \gamma_3| + b|t_1 - \gamma_1| + b|t_2 - \gamma_2| \\ + |t_4 - \gamma_4| + |t_5 - \gamma_5| + \dots, \\ 2|t_4 - \gamma_4| + b|t_1 - \gamma_1| + b|t_2 - \gamma_2| \\ + |t_3 - \gamma_3| + |t_5 - \gamma_5| + \dots, \\ 2|t_5 - \gamma_5| + b|t_1 - \gamma_1| + b|t_2 - \gamma_2| + |t_3 - \gamma_3| \\ + |t_4 - \gamma_4| + |t_6 - \gamma_6| + \dots, \\ \dots \end{array} \right\}. \end{aligned}$$

Subcase 2.1: $|t_1 - \gamma_1| \geq |t_2 - \gamma_2|$ and $b|t_1 - \gamma_1| \geq |t_k - \gamma_k|, \forall k \geq 3$.

Then,

$$\begin{aligned} \|y - z\| &= b|t_1 - \gamma_1| + b \sum_{k=1}^{\infty} |t_k - \gamma_k| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\ &\geq b\delta + b|t_1 - \gamma_1| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\ &\geq b\delta + b|t_1 - \gamma_1| + (1 - b)|\delta - (t_1 - \gamma_1) - (t_2 - \gamma_2)| \\ &\geq b\delta + b|t_1 - \gamma_1| + (1 - b)\delta - 2(1 - b)|t_1 - \gamma_1| \\ &= \delta - (2 - 3b)|t_1 - \gamma_1|. \end{aligned}$$

Subcase 2.1.1: Assume $\frac{\delta}{2} \geq |t_1 - \gamma_1|$.

Then clearly the last inequality from above says that $\|y - z\| \geq \frac{3b\delta}{2}$.

Subcase 2.1.2: Assume $\frac{\delta}{2} < |t_1 - \gamma_1|$.

Then $\|y - z\| \geq b\delta + b|t_1 - \gamma_1| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \geq \frac{3b\delta}{2}$.

Subcase 2.2: $|t_2 - \gamma_2| \geq |t_1 - \gamma_1|$ and $b|t_2 - \gamma_2| \geq |t_k - \gamma_k|, \forall k \geq 3$.

Then,

$$\begin{aligned} \|y - z\| &= b|t_2 - \gamma_2| + b \sum_{k=1}^{\infty} |t_k - \gamma_k| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\ &\geq b\delta + b|t_2 - \gamma_2| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\ &\geq b\delta + b|t_2 - \gamma_2| + (1 - b)|\delta - (t_1 - \gamma_1) - (t_2 - \gamma_2)|. \end{aligned}$$

Therefore,

$$\begin{aligned} \|y - z\| &\geq b\delta + b|t_2 - \gamma_2| + (1 - b)\delta - 2(1 - b)|t_2 - \gamma_2| \\ &\geq \delta - (2 - 3b)|t_2 - \gamma_2|. \end{aligned}$$

Subcase 2.2.1: Assume $\frac{\delta}{2} \geq |t_2 - \gamma_2|$.

Then clearly the last inequality from above says that $\|y - z\| \geq \frac{3b\delta}{2}$.

Subcase 2.2.2: Assume $\frac{\delta}{2} < |t_2 - \gamma_2|$.

Then $\|y - z\| \geq b\delta + b|t_2 - \gamma_2| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \geq \frac{3b\delta}{2}$.

Subcase 2.3: $|t_3 - \gamma_3| \geq b|t_1 - \gamma_1|$, $|t_3 - \gamma_3| \geq b|t_2 - \gamma_2|$, and $|t_3 - \gamma_3| \geq |t_k - \gamma_k|$, $\forall k \geq 4$.

Then,

$$\begin{aligned} \|y - z\| &= |t_3 - \gamma_3| + b \sum_{k=1}^{\infty} |t_k - \gamma_k| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\ &\geq b\delta + |t_3 - \gamma_3| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\ &\geq b\delta + |t_3 - \gamma_3| + (1 - b)|\delta - (t_1 - \gamma_1) - (t_2 - \gamma_2)| \\ &\geq b\delta + |t_3 - \gamma_3| + (1 - b)\delta - \frac{2(1 - b)}{b}|t_3 - \gamma_3| \\ &= \delta - \frac{2 - 3b}{b}|t_3 - \gamma_3|. \end{aligned}$$

Subcase 2.3.1: Assume $\frac{b\delta}{2} \geq |t_3 - \gamma_3|$.

Then clearly the last inequality from above says that $\|y - z\| \geq \frac{3b\delta}{2}$.

Subcase 2.3.2: Assume $\frac{b\delta}{2} < |t_3 - \gamma_3|$.

Then $\|y - z\| \geq b\delta + |t_3 - \gamma_3| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \geq \frac{3b\delta}{2}$.

Subcase 2.4: $|t_4 - \gamma_4| \geq b|t_1 - \gamma_1|$, $|t_4 - \gamma_4| \geq b|t_2 - \gamma_2|$, and $|t_4 - \gamma_4| \geq |t_k - \gamma_k|$, $\forall k \geq 5$ and for $k = 3$.

Then,

$$\begin{aligned}
 \|y - z\| &= |t_4 - \gamma_4| + b \sum_{k=1}^{\infty} |t_k - \gamma_k| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\
 &\geq b\delta + |t_4 - \gamma_4| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \\
 &\geq b\delta + |t_4 - \gamma_4| + (1 - b)|\delta - (t_1 - \gamma_1) - (t_2 - \gamma_2)| \\
 &\geq b\delta + |t_4 - \gamma_4| + (1 - b)\delta - \frac{2(1 - b)}{b}|t_4 - \gamma_4| \\
 &= \delta - \frac{2 - 3b}{b}|t_4 - \gamma_4|.
 \end{aligned}$$

Subcase 2.4.1: Assume $\frac{b\delta}{2} \geq |t_4 - \gamma_4|$.

Then clearly the last inequality from above says that $\|y - z\| \geq \frac{3b\delta}{2}$.

Subcase 2.4.2: Assume $\frac{b\delta}{2} < |t_4 - \gamma_4|$.

Then $\|y - z\| \geq b\delta + |t_4 - \gamma_4| + (1 - b) \sum_{k=3}^{\infty} |t_k - \gamma_k| \geq \frac{3b\delta}{2}$.

Thus, we continue in this way and see that $\|y - z\| \geq \frac{3b\delta}{2}$ from all cases.

Therefore, when λ is chosen to be in $[0, 1]$, for any $y \in E$ and for $z \in W \setminus E$, $\|y - z\| \geq \Gamma$ such that there exists unique $\lambda_0 \in [0, 1]$ with $\|h_{\lambda_0} - z\| = \Gamma$.

Now define a subset in our set by $\Lambda := \{y : \|y - z\| \leq \Gamma\}$. Note that $\Lambda \subseteq E$ is a nonempty compact convex subset such that for any $h \in \Lambda$,

$$\begin{aligned}
 s(Th) &= \limsup_m \left\| Th - \frac{1}{m} \sum_{k=1}^m x^{(k)} \right\| \\
 &\leq \limsup_m \left\| Th - T \left(\frac{1}{m} \sum_{k=1}^m x^{(k)} \right) \right\| + \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x^{(k)} - T \left(\frac{1}{m} \sum_{k=1}^m x^{(k)} \right) \right\|.
 \end{aligned}$$

However, since T is affine,

$$\begin{aligned}
 s(Th) &\leq \limsup_m \left\| Th - T \left(\frac{1}{m} \sum_{k=1}^m x^{(k)} \right) \right\| + \limsup_m \left\| \frac{1}{m} \sum_{k=1}^m x^{(k)} - \frac{1}{m} \sum_{k=1}^m Tx^{(k)} \right\| \\
 &\leq \limsup_m \left\| h - \frac{1}{m} \sum_{k=1}^m x^{(k)} \right\| \\
 &= s(h).
 \end{aligned}$$

Also, $s(Th) = \|z - Th\|$ and $s(h) = \|z - h\|$. Hence,

$$\begin{aligned}
 \|z - Th\| \leq \|z - h\| &\implies \|z - Th\| = \|z - h\| \\
 &\implies Th \in \Lambda.
 \end{aligned}$$

Therefore, $T(\Lambda) \subseteq \Lambda$, and since T is continuous, Schauder's fixed point theorem [28] tells us that T has a fixed point such that $h = h_{\lambda_0}$ is the unique minimizer of $\|y - z\| : y \in E$ and $Th = h$.

Therefore, E has fpp(ne) as desired. \square

Remark 5.5 *Generalizing the sets as Goebel and Kuczumow [13] or Everest [10] did, one can obtain larger classes with fixed point property for nonexpansive mappings, which could be considered as a future project for other researchers in the field and for us as well.*

Acknowledgments

The author is grateful to Chris Lennard for his valuable comments and helpful discussions on the subject. The author also thanks the anonymous reviewers for their helpful and constructive comments that greatly contributed to improving the final version of the paper.

References

- [1] Alspach DE. A fixed point free nonexpansive map. Proceedings of the American Mathematical Society 1981; 82 (3): 423-424.
- [2] Astashkin SV, Sukochev FA. Banach-Saks property in Marcinkiewicz spaces. Journal of Mathematical Analysis and Applications 2007; 336 (2): 1231-1258.
- [3] Beauzamy B. Introduction to Banach Spaces and Their Geometry. Vol. 11. Amsterdam, the Netherlands: Elsevier Science Pub. Co., 1982.
- [4] Browder FE. Fixed-point theorems for noncompact mappings in Hilbert space. Proceedings of the National Academy of Sciences of the United States of America 1965; 53 (6): 1272-1276.
- [5] Browder FE. Fixed-point theorems for noncompact mappings in Hilbert space. Proceedings of the National Academy of Sciences of the United States of America 1965; 54 (4): 1041-1044.
- [6] Carothers NL, Dilworth SJ, Lennard CJ. On a localization of the UKK property and the fixed point property in $L_{w,1}$. Lecture Notes in Pure and Applied Mathematics 1996; 175: 111-124.
- [7] Carothers NL, Dilworth SJ, Lennard CJ, Trautman DA. A fixed point property for the Lorentz space $L_{p,1}(\mu)$. Indiana University Mathematics Journal 1991; 40 (1): 345-352.
- [8] Dowling PN, Lennard CJ. Every nonreflexive subspace of $L_1[0, 1]$ fails the fixed point property. Proceedings of the American Mathematical Society 1997; 125 (2): 443-446.
- [9] Dowling PN, Lennard CJ, Turett B. Renormings of l^1 and c_0 and Fixed Point Properties. Handbook of Metric Fixed Point Theory. Berlin, Germany: Springer, 2001.
- [10] Everest T. Fixed points of nonexpansive maps on closed, bounded, convex sets in l^1 . PhD, University of Pittsburgh, Pittsburgh, PA, USA, 2013.
- [11] Gamboa de Buen B, Nuñez-Medina F. A generalization of a renorming theorem by Lin and a new nonreflexive space with the fixed point property which is nonisomorphic to l^1 . Journal of Mathematical Analysis and Applications 2013; 405 (1): 57-70.
- [12] Goebel K, Kirk WA. Topics in Metric Fixed Point Theory. Cambridge, UK: Cambridge University Press, 1990.
- [13] Goebel K, Kuczumow T. Irregular convex sets with fixed-point property for nonexpansive mappings. Colloquium Mathematicum 1979; 40 (2): 259-264.
- [14] Göhde D. Zum Prinzip der kontraktiven Abbildung. Mathematische Nachrichten 1965; 30 (3-4): 251-258 (in German).

- [15] Khamsi MA, Kirk WA. An Introduction to Metric Spaces and Fixed Point Theory. New York, NY, USA: John Wiley & Sons, 2011.
- [16] Kirk WA. A fixed point theorem for mappings which do not increase distances. American Mathematical Monthly 1965; 72 (9): 1004-1006.
- [17] Lennard CJ, Nezir V. Semi-strongly asymptotically non-expansive mappings and their applications on fixed point theory. Hacettepe Journal of Mathematics and Statistics 2017; 46 (4): 613-620.
- [18] Lin PK. Unconditional bases and fixed points of nonexpansive mappings. Pacific Journal of Mathematics 1985; 116 (1): 69-76.
- [19] Lin PK. There is an equivalent norm on ℓ_1 that has the fixed point property. Nonlinear Analysis: Theory, Methods & Applications 2008; 68 (8): 2303-2308.
- [20] Lin PK. Renorming of ℓ_1 and the fixed point property. Journal of Mathematical Analysis and Applications 2010; 362 (2): 534-541.
- [21] Lindenstrauss J, Tzafriri L. Classical Banach Spaces I. Berlin, Germany: Springer-Verlag, 1977.
- [22] Lorentz GG. Some new functional spaces. Annals of Mathematics 1950; 51: 37-55.
- [23] Maurey B. Points fixes des contractions de certains faiblement compacts de L^1 . Seminaire d'Analyse Fonctionnelle, 1980-1981, Centre de Mathématiques, École Polytech., Palaiseau, Exp. No. VIII, 19 pp., 1981 (in French).
- [24] Nezir V. Fixed point properties for c_0 -like spaces. PhD, University of Pittsburgh, Pittsburgh, PA, USA, 2012.
- [25] Nezir V, Mustafa N. c_0 can be renormed to have the fixed point property for affine nonexpansive mappings. Filomat 2018; 32 (16): 5645-5663.
- [26] Rakov SA. Banach-Saks property of a Banach space. Mathematical notes of the Academy of Sciences of the USSR 1979; 26 (6): 909-916.
- [27] Rakov SA. Banach-Saks exponent of certain Banach spaces of sequences. Mathematical Notes of the Academy of Sciences of the USSR 1982; 32 (5): 791-797.
- [28] Schauder J. Der Fixpunktsatz in Funktionalräumen. Studia Mathematica 1930; 2 (1): 171-180 (in German).
- [29] Soardi PM. Existence of fixed points of nonexpansive mappings in certain Banach lattices. Proceedings of the American Mathematical Society 1979; 73: 25-29.