

1-1-2019

Some applications of differential subordination for certain starlike functions

RAHIM KARGAR

LUCYNA TROJNAR SPELINA

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)



Recommended Citation

KARGAR, RAHIM and SPELINA, LUCYNA TROJNAR (2019) "Some applications of differential subordination for certain starlike functions," *Turkish Journal of Mathematics*: Vol. 43: No. 4, Article 13. <https://doi.org/10.3906/mat-1903-28>

Available at: <https://journals.tubitak.gov.tr/math/vol43/iss4/13>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Some applications of differential subordination for certain starlike functions

Rahim KARGAR¹ , Lucyna TROJNAR-SPELINA^{2*} 

Young Researchers and Elite Club, Ardabil Branch, Islamic Azad University, Ardabil, Iran

²Faculty of Mathematics and Applied Physics, Rzeszów University of Technology, Rzeszów, Poland

Received: 09.03.2019

Accepted/Published Online: 06.06.2019

Final Version: 31.07.2019

Abstract: Let $\mathcal{S}^*(q_c)$ denote the class of functions f analytic in the open unit disc Δ , normalized by the condition $f(0) = 0 = f'(0) - 1$ and satisfying the following inequality

$$\left| \left(\frac{zf'(z)}{f(z)} \right)^2 - 1 \right| < c \quad (z \in \Delta, 0 < c \leq 1).$$

By use of the subordination principle for the univalent functions we have

$$f \in \mathcal{S}^*(q_c) \Leftrightarrow \frac{zf'(z)}{f(z)} \prec \sqrt{1+cz} \quad (z \in \Delta, 0 < c \leq 1).$$

In the present paper, for an analytic function p in Δ with $p(0) = 1$ we give some conditions which imply $p(z) \prec \sqrt{1+cz}$. These conditions are then used to obtain some corollaries for certain subclasses of analytic functions.

Key words: Analytic, univalent, subordination, Janowski starlike functions, Bernoulli lemniscate

1. Introduction

Let Δ be the open unit disc in the complex plane \mathbb{C} , i.e. $\Delta = \{z \in \mathbb{C} : |z| < 1\}$ and $\mathcal{H}(\Delta)$ be the class of functions that are analytic in Δ . Also, let $\mathcal{A} \subset \mathcal{H}(\Delta)$ be the class of functions that have the following Taylor–Maclaurin series expansion

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in \Delta).$$

Thus, if $f \in \mathcal{A}$, then it satisfies the following normalization condition

$$f(0) = 0 = f'(0) - 1.$$

The set of all univalent (one-to-one) functions f in Δ is denoted by \mathcal{U} . Let f and g belong to class $\mathcal{H}(\Delta)$. Then we say that a function f is subordinate to g , written by

$$f(z) \prec g(z) \quad \text{or} \quad f \prec g,$$

*Correspondence: lspelina@prz.edu.pl

2000 AMS Mathematics Subject Classification: 30C45

if there exists a Schwarz function w with the following properties

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad (z \in \Delta),$$

such that $f(z) = g(w(z))$ for all $z \in \Delta$. In particular, if $g \in \mathcal{U}$, then we have

$$f(z) \prec g(z) \Leftrightarrow (f(0) = g(0) \quad \text{and} \quad f(\Delta) \subset g(\Delta)).$$

Furthermore, we say that the function $f \in \mathcal{U}$ is starlike if and only if

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0 \quad (z \in \Delta).$$

The familiar class of starlike functions in Δ is denoted by \mathcal{S}^* . Also the function $f \in \mathcal{U}$ is called convex if and only if

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > 0 \quad (z \in \Delta).$$

We denote by \mathcal{K} the class of convex functions in Δ . A function $f \in \mathcal{A}$ is said to be close-to-convex, if there exists a function $g \in \mathcal{K}$ such that

$$\operatorname{Re} \left\{ \frac{f'(z)}{g'(z)} \right\} > 0 \quad (z \in \Delta).$$

The class of close-to-convex functions is denoted by \mathcal{C} . Note that $\mathcal{C} \subset \mathcal{U}$.

Let $c \in (0, 1]$. We say that the function $f \in \mathcal{A}$ belongs to the class $\mathcal{S}^*(q_c)$, if it satisfies the following condition

$$\left| \left(\frac{zf'(z)}{f(z)} \right)^2 - 1 \right| < c \quad (z \in \Delta).$$

The class $\mathcal{S}^*(q_c)$ was introduced by Sokół, see [19]. Also, the class $\mathcal{S}^*(q_1) \equiv \mathcal{SL}^*$ was considered in [20]. In Geometric Function Theory there are many interesting subclasses of starlike functions which have been defined by subordination, see for example [3–8, 14, 16–18]. In the sequel we give a necessary and sufficient condition for the class $\mathcal{S}^*(q_c)$ by using the subordination.

Define

$$q_c(z) := \sqrt{1 + cz} \quad (z \in \Delta, c \in (0, 1]) \tag{1.1}$$

and Ω_c by

$$\Omega_c := \{\zeta \in \mathbb{C} : \operatorname{Re}\{\zeta\} > 0, |\zeta^2 - 1| < c\}.$$

Then we have $q_c(\Delta) = \Omega_c$, see [19]. Indeed, the function $q_c(z)$ maps Δ onto a set bounded by Bernoulli lemniscate. It is easy to see that $f \in \mathcal{S}^*(q_c)$ if and only if it satisfies the following differential subordination

$$\frac{zf'(z)}{f(z)} \prec q_c(z) \quad (z \in \Delta, c \in (0, 1]),$$

where q_c is defined by (1.1) and the branch of the square root is chosen to be $q_c(0) = 1$. Noting to the above we have $\mathcal{S}^*(q_c) \subset \mathcal{S}^*$. Another class that we are interested to study is the class $\mathcal{U}(c)$ which is defined as follows:

$$\mathcal{U}(c) := \left\{ f \in \mathcal{A} : \left| \left(\frac{zf'(z)}{f(z)} \right)^2 - 1 \right| \leq c, 0 < c \leq 1, z \in \Delta \right\}.$$

For each $c \in (0, 1]$ we have $\mathcal{U}(c) \subset \mathcal{U}$, see [12]. Let A and B be two fixed constants such that $-1 \leq B < A \leq 1$. We denote by $\mathcal{S}^*[A, B]$ the class of Janowski starlike functions $f \in \mathcal{A}$ and satisfying the condition

$$\frac{zf'(z)}{f(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta).$$

This class was introduced by Janowski [2]. We remark that $\mathcal{S}^*[1, -1]$ becomes the class of starlike functions.

Next, we recall a lemma, called Jack's lemma.

Lemma 1.1 (see [1], see also [15, Lemma 1.3, p. 28]) *Let w be a nonconstant function meromorphic in Δ with $w(0) = 0$. If*

$$|w(z_0)| = \max\{|w(z)| : |z| \leq |z_0|\} \quad (z \in \Delta),$$

then there exists a real number k ($k \geq 1$) such that $z_0 w'(z_0) = k w(z_0)$.

In this paper, for an analytic function $p(z)$ in the unit disk Δ we find some conditions that imply $p(z) \prec \sqrt{1 + cz}$. Also, some interesting corollaries are obtained.

2. Main Results

We start with the following.

Theorem 2.1 *Let p be an analytic function in Δ with $p(0) = 1$, $|A| \leq 1$, $|B| < 1$, $0 < c \leq 1$. Also let γ satisfy the following inequality*

$$\gamma \geq \frac{2(|A| + |B|)}{c(1 - |B|)}(1 + c). \quad (2.1)$$

Then the following subordination

$$1 + \gamma \frac{zp'(z)}{p(z)} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta) \quad (2.2)$$

implies that

$$p(z) \prec \sqrt{1 + cz} \quad (z \in \Delta).$$

Proof Let γ satisfy the condition (2.1) and consider

$$F(z) := 1 + \gamma \frac{zp'(z)}{p(z)} \quad (2.3)$$

for all $z \in \Delta$. Define the function w by the relation

$$p(z) = \sqrt{1 + cw(z)} = 1 + p_1 z + p_2 z^2 + \cdots, \quad (2.4)$$

or $w(z) = (p^2(z) - 1)/c = w_1 z + \cdots$. By the hypothesis, since p is analytic and $p(0) = 1$, thus w is meromorphic in Δ and $w(0) = 0$. We shall show that $|w(z)| < 1$ in Δ . With a simple calculation (2.4) gives

$$\gamma \frac{zp'(z)}{p(z)} = \frac{c\gamma zw'(z)}{2(1 + cw(z))}.$$

Using the last equality in (2.3) we get

$$F(z) = 1 + \frac{c\gamma zw'(z)}{2(1+cw(z))}$$

and thus by computation we obtain

$$\frac{F(z) - 1}{A - BF(z)} = \frac{c\gamma zw'(z)}{2A(1+cw(z)) - B[2(1+cw(z)) + c\gamma zw'(z)]}.$$

Now assume that there exists a point $z_0 \in \Delta$ such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1.$$

Therefore, by Lemma 1.1, there exists a number $k \geq 1$ such that $z_0 w'(z_0) = kw(z_0)$. Without loss of generality we may assume that $w(z_0) = e^{i\delta}$ where $\delta \in [-\pi, \pi]$. For this z_0 , we have

$$\begin{aligned} \left| \frac{F(z_0) - 1}{A - BF(z_0)} \right| &= \left| \frac{ck\gamma e^{i\delta}}{2A(1+ce^{i\delta}) - B[2(1+ce^{i\delta}) + c\gamma ke^{i\delta}]} \right| \\ &\geq \frac{ck\gamma}{2|A||1+ce^{i\delta}| + |B|[2 + (2c+c\gamma k)e^{i\delta}]} \\ &= \frac{ck\gamma}{2|A|p_1(\delta) + |B|p_2(\delta)} \\ &=: H(\cos \delta) \end{aligned}$$

where the expressions $p_1(\delta)$ and $p_2(\delta)$ have a form

$$p_1(\delta) = \sqrt{1 + 2c \cos \delta + c^2},$$

$$p_2(\delta) = \sqrt{4 + c(2 + \gamma k)[4 \cos \delta + c(2 + \gamma k)]}$$

and

$$H(t) = \frac{ck\gamma}{2|A|\sqrt{1+2ct+c^2} + |B|\sqrt{4+4(2c+c\gamma k)t+(2c+c\gamma k)^2}}.$$

By a simple computation it can be easily seen that $H'(t) < 0$. Thus, H is a decreasing function when $-1 \leq t = \cos \delta \leq 1$ and consequently

$$H(t) \geq H(1) = \frac{ck\gamma}{2|A|(1+c) + |B|(2+2c+c\gamma k)}. \quad (2.5)$$

Now consider the function

$$L(k) = \frac{ck\gamma}{2|A|(1+c) + |B|(2+2c+c\gamma k)} \quad (k \geq 1). \quad (2.6)$$

It is easy to see that $L'(k) > 0$. In conclusion,

$$L(k) \geq L(1) = \frac{c\gamma}{2|A|(1+c) + |B|(2+2c+c\gamma)}. \quad (2.7)$$

Finally from the definition of H and from (2.5)–(2.7), it follows that

$$\left| \frac{F(z_0) - 1}{A - BF(z_0)} \right| \geq \frac{c\gamma}{2|A|(1+c) + |B|(2(1+c) + c\gamma)} =: T(A, B, c, \gamma).$$

The inequality (2.1) implies that $T(A, B, c, \gamma) > 1$. However, this is a contradiction with the assumption (2.2). This is the end of the proof. \square

If we put $p(z) = zf'(z)/f(z)$ in Theorem 2.1, then we obtain the following result:

Corollary 2.1 *Let $|A| \leq 1$, $|B| < 1$, $0 < c \leq 1$ and let*

$$\gamma \geq \frac{2(|A| + |B|)}{c(1 - |B|)}(1 + c).$$

If f satisfies the subordination

$$1 + \gamma \left(1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta),$$

then $f \in \mathcal{S}^(q_c)$.*

If we let $c = 1$ in Corollary 2.1, then we have:

Corollary 2.2 *Let $|A| \leq 1$, $|B| < 1$ and let*

$$\gamma \geq \frac{4(|A| + |B|)}{1 - |B|}.$$

If f satisfies the following subordination

$$1 + \gamma \left(1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta),$$

then $f \in \mathcal{SL}^$.*

Taking $A = 1$ and $B = 0$ in Corollary 2.2, we obtain:

Corollary 2.3 *Let $\gamma \geq 4$. If f satisfies the following inequality*

$$\operatorname{Re} \left\{ 1 + \gamma \left(1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right\} > 0$$

for all $z \in \Delta$, then $f \in \mathcal{SL}^$.*

If we put $p(z) = z\sqrt{f'(z)}/f(z)$ in Theorem 2.1, then we have:

Corollary 2.4 *Let $|A| \leq 1$, $|B| < 1$, $0 < c \leq 1$ and let*

$$\gamma \geq \frac{2(|A| + |B|)}{c(1 - |B|)}(1 + c).$$

If the function f satisfies the following condition

$$1 + \gamma \left(1 + \frac{1}{2} \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta),$$

then

$$\left| \left(\frac{z}{f(z)} \right)^2 f'(z) - 1 \right| \leq c \quad (z \in \Delta).$$

This means that $f \in \mathcal{U}(c)$, hence it is univalent in Δ .

If we put $p(z) = \sqrt{f'(z)}$ and $c = 1$ in Theorem 2.1, then we have the following result:

Corollary 2.5 Assume that $|A| \leq 1$, $|B| < 1$ and that

$$\gamma \geq \frac{4(|A| + |B|)}{1 - |B|}.$$

If

$$1 + \gamma \left(\frac{1}{2} \frac{zf''(z)}{f'(z)} \right) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta),$$

then f is univalent in Δ by [13].

If we put $p(z) = f(z)/z$ in Theorem 2.1, then we obtain the following result.

Corollary 2.6 Let $|A| \leq 1$, $|B| < 1$, $0 < c \leq 1$ and let

$$\gamma \geq \frac{2(|A| + |B|)}{c(1 - |B|)}(1 + c).$$

If the function f satisfies the following condition

$$1 + \gamma \left(\frac{zf'(z)}{f(z)} - 1 \right) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \Delta),$$

then

$$\left| \left(\frac{f(z)}{z} \right)^2 - 1 \right| < c$$

for all $z \in \Delta$.

Taking $A = 1$ and $B = 0$ in Corollary 2.6, we obtain:

Corollary 2.7 Let $\gamma \geq 2(1 + 1/c)$ with $c \in (0, 1]$. If the following inequality holds

$$\operatorname{Re} \left\{ 1 + \gamma \left(\frac{zf'(z)}{f(z)} - 1 \right) \right\} > 0 \quad (z \in \Delta),$$

then

$$\left| \left(\frac{f(z)}{z} \right)^2 - 1 \right| < c \quad (z \in \Delta).$$

For the proofs of next theorems we need a couple of lemmas.

Lemma 2.1 ([11]) *Let q be univalent in the unit disk Δ and θ and ϕ be analytic in a domain \mathbb{U} containing $q(\Delta)$ with $\phi(w) \neq 0$ when $w \in q(\Delta)$. Set $Q(z) = zq'(z)\phi(q(z))$, $h(z) = \theta(q(z)) + Q(z)$. Suppose that Q is starlike (univalent) in Δ , and*

$$\operatorname{Re} \left\{ \frac{zh'(z)}{Q(z)} \right\} = \operatorname{Re} \left\{ \frac{\theta'(q(z))}{\phi(q(z))} + \frac{zQ'(z)}{Q(z)} \right\} > 0 \quad (z \in \Delta).$$

If p is analytic in Δ , with $p(0) = q(0)$, $p(\Delta) \subset \mathbb{U}$ and

$$\theta(p(z)) + zp'(z)\phi(p(z)) \prec \theta(q(z)) + zq'(z)\phi(q(z)), \tag{2.8}$$

then $p(z) \prec q(z)$, and q is the best dominant of (2.8).

Lemma 2.2 (see [9], see also [10, p. 24]) *Assume that \mathcal{Q} is the set of analytic functions that are injective on $\overline{\Delta} \setminus E(f)$, where $E(f) : \{\omega : \omega \in \partial\Delta \text{ and } \lim_{z \rightarrow \omega} f(z) = \infty\}$, and are such that $f'(\omega) \neq 0$ for $(\omega \in \partial\Delta \setminus E(f))$. Let $\psi \in \mathcal{Q}$ with $\psi(0) = a$ and let $\varphi(z) = a + a_m z^m + \dots$ be analytic in Δ with $\varphi(z) \not\equiv a$ and $m \in \mathbb{N}$. If $\varphi \not\prec \psi$ in Δ , then there exist points $z_0 = r_0 e^{i\theta} \in \Delta$ and $\omega_0 \in \partial\Delta \setminus E(\psi)$, for which $\varphi(|z| < r_0) \subset \psi(\Delta)$, $\varphi(z_0) = \psi(\omega_0)$ and $z_0 \varphi'(z_0) = k \omega_0 \psi'(\omega_0)$, for some $k \geq m$.*

Next we prove the following.

Theorem 2.2 *Let $p \in \mathcal{H}(\Delta)$ with $p(0) = 1$ and $c \in (0, 1]$. If the function p satisfies the subordination*

$$\frac{1}{3}p^3(z) + zp'(z) \prec \frac{1}{3}(\sqrt{1+cz})^3 + \frac{cz}{2\sqrt{1+cz}} \quad (z \in \Delta), \tag{2.9}$$

then

$$p(z) \prec \sqrt{1+cz} \quad (z \in \Delta),$$

and the function $\sqrt{1+cz}$ is the best dominant of (2.9).

Proof Consider

$$q_c(z) = \sqrt{1+cz}, \quad \theta(\omega) = \frac{1}{3}\omega^3, \quad \phi(\omega) = 1.$$

We know that q_c is analytic and univalent in Δ . Also $q_c(0) = p(0) = 1$. Moreover, both functions $\theta(\omega)$ and $\phi(\omega)$ are analytic in the ω -plane with $\phi(\omega) \neq 0$. The function

$$Q(z) = zq'_c(z)\phi(q(z)) = \frac{cz}{2\sqrt{1+cz}} = zq'_c(z),$$

is a starlike function, because q_c is convex. If we put

$$h(z) = \theta(q_c(z)) + Q(z) = \frac{1}{3}q_c^3(z) + zq'_c(z), \tag{2.10}$$

then we have

$$\operatorname{Re} \left\{ \frac{zh'(z)}{Q(z)} \right\} = \operatorname{Re} \left\{ 1 + cz + \left(1 + \frac{zq_c''(z)}{q_c'(z)} \right) \right\} > 1 - c \geq 0$$

for all $z \in \Delta$. Therefore, the function h given by (2.10) is close-to-convex and univalent in Δ . Thus, by the Lemma 2.1 and (2.9), we find that $p(z) \prec q_c(z)$ and $q_c(z)$ is the best dominant of (2.9) so the desired conclusion follows. \square

If we put $p(z) = zf'(z)/f(z)$, then we have the following result:

Corollary 2.8 *Let $c \in (0, 1]$. If a function f satisfies the subordination*

$$\frac{1}{3} \left(\frac{zf'(z)}{f(z)} \right)^3 + \left(1 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \left(\frac{zf'(z)}{f(z)} \right) \prec \frac{1}{3} (\sqrt{1+cz})^3 + \frac{cz}{2\sqrt{1+cz}},$$

then $f \in \mathcal{S}^*(q_c)$ where $z \in \Delta$.

Finally we prove the following:

Theorem 2.3 *Let $k \geq 1$ and $0 < c \leq 1$. If $p \in \mathcal{H}(\Delta)$ with $p(0) = 1$ and it satisfies the condition*

$$\operatorname{Re} \{p(z)(p(z) + zp'(z))\} > 1 + c(1 + k/2) \quad (z \in \Delta), \quad (2.11)$$

then

$$p(z) \prec \sqrt{1+cz} \quad (z \in \Delta).$$

Proof Suppose that $p(z) \not\prec q_c(z) = \sqrt{1+cz}$. Then there exist points $z_0, |z_0| < 1$ and $\omega_0, |\omega_0| = 1, \omega_0 \neq 1$ satisfying the following conditions

$$p(z_0) = q_c(\omega_0), \quad p(|z| < |z_0|) \subset q_c(\Delta) \quad \text{and} \quad |\omega_0| = 1.$$

From Lemma 2.2, we find that there exists a number $k \geq 1$ such that

$$\{p(z_0)(p(z_0) + zp'(z_0))\} = \{q_c(\omega_0)(q_c(\omega_0) + k\omega_0 q_c'(\omega_0))\} = 1 + c(1 + k/2)\omega_0. \quad (2.12)$$

By setting $\omega_0 = e^{i\delta}$, $\delta \in [-\pi, \pi]$ in (2.12), it can be easily seen that

$$\operatorname{Re}\{1 + c(1 + k/2)\omega_0\} = 1 + c(1 + k/2) \cos \delta \leq 1 + c(1 + k/2).$$

However, it contradicts our assumption (2.11) and consequently $p(z) \prec q_c(z)$ in Δ . \square

If we let $p(z) = zf'(z)/f(z)$, then we have the following result:

Corollary 2.9 *Let $0 < c \leq 1$ and let $k \geq 1$. If f satisfies the following inequality*

$$\operatorname{Re} \left\{ \left(\frac{zf'(z)}{f(z)} \right)^2 \left(2 + \frac{zf''(z)}{f'(z)} - \frac{zf'(z)}{f(z)} \right) \right\} > 1 + c(1 + k/2) \quad (z \in \Delta)$$

then $f \in \mathcal{S}^*(q_c)$.

Acknowledgments

The authors would like to thank Professor S. Ponnusamy for providing information about the reference [13].

References

- [1] Jack IS. Functions starlike and convex of order α . *Journal of the London Mathematical Society* 1971; 3: 469-474.
- [2] Janowski W. Some extremal problems for certain families of analytic functions I. *Annales Polonici Mathematici* 1973; 28: 297-326.
- [3] Kargar R, Ebadian A, Sokół J. On Booth lemniscate and starlike functions. *Analysis and Mathematical Physics* 2019; 9: 143. <https://doi.org/10.1007/s13324-017-0187-3>
- [4] Kargar R, Ebadian A, Sokół J. Radius problems for some subclasses of analytic functions. *Complex Analysis and Operator Theory* 2017; 11: 1639-1649.
- [5] Kumar S, Ravichandran V. A subclass of starlike functions associated with a rational function. *Southeast Asian Bulletin of Mathematics* 2016; 40: 199-212.
- [6] Kuroki K, Owa S. Notes on new class for certain analytic functions. *RIMS Kokyuroku Kyoto University* 2011; 1772: 21-25.
- [7] Mendiratta R, Nagpal S, Ravichandran V. A subclass of starlike functions associated with left-half of the lemniscate of Bernoulli. *International Journal of Mathematics* 2014; 25: 17 pp.
- [8] Mendiratta R, Nagpal S, Ravichandran V. On a subclass of strongly starlike functions associated with exponential function. *Bulletin of the Malaysian Mathematical Sciences Society* 2015; 38: 365-386.
- [9] Miller SS, Mocanu PT. *Differential subordinations and univalent functions*. Michigan Mathematical Journal 1981; 28: 151-171.
- [10] Miller SS, Mocanu PT. *Differential Subordination, Theory and Application*. New York, NY, USA: Marcel Dekker, Inc., 2000.
- [11] Miller SS, Mocanu PT. On some classes of first order differential subordinations. *Michigan Mathematical Journal* 1985; 32: 185-195.
- [12] Obradović M, Ponnusamy S. Univalence and starlikeness of certain transforms defined by convolution of analytic functions. *Journal of Mathematical Analysis and Applications* 2007; 336: 758-767.
- [13] Obradović M, Ponnusamy S. Univalence of the average of two analytic functions. In: 60 years of analytic functions in Lublin—in memory of our professors and friends Jan G. Krzyż, Zdzisław Lewandowski and Wojciech Szpiał; 2012. pp. 169-183.
- [14] Raina RK, Sokół J. Some properties related to a certain class of starlike functions. *Comptes Rendus Mathématique* 2015; 353: 973-978.
- [15] Ruscheweyh S. *Convolutions in Geometric Function Theory*. Montreal, Quebec: Les Presses de l'Université de Montréal, 1982.
- [16] Sharma K, Jain NK, Ravichandran V. Starlike functions associated with a cardioid. *Afrika Matematika* 2016; 27: 923-939.
- [17] Sokół J. A certain class of starlike functions. *Computers & Mathematics with Applications* 2011; 62: 611-619.
- [18] Sokół J, Stankiewicz J. Radius of convexity of some subclasses of strongly starlike functions. *Folia Scienten University of Tech Resoviensis* 1996; 19: 101-105.
- [19] Sokół J. On some subclass of strongly starlike functions. *Demonstratio Mathematica* 1998; 21: 81-86.
- [20] Sokół J. Radius problems in the class \mathcal{SL}^* . *Applied Mathematics and Computation* 2009; 214: 569-573.