

1-1-2014

Influence of natural zeolite on nitrogen dynamics in soil

STANISLAV TORMA

JOZEF VILCEK

PETER ADAMISIN

EMILIA HUTTMANOVA

ONDREJ HRONEC

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

Recommended Citation

TORMA, STANISLAV; VILCEK, JOZEF; ADAMISIN, PETER; HUTTMANOVA, EMILIA; and HRONEC, ONDREJ (2014) "Influence of natural zeolite on nitrogen dynamics in soil," *Turkish Journal of Agriculture and Forestry*. Vol. 38: No. 5, Article 17. <https://doi.org/10.3906/tar-1311-13>
Available at: <https://journals.tubitak.gov.tr/agriculture/vol38/iss5/17>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Influence of natural zeolite on nitrogen dynamics in soil

Stanislav TORMA^{1,*}, Jozef VILCEK^{1,2}, Peter ADAMISIN³, Emilia HUTTMANOVA³, Ondrej HRONEC³

¹Soil Science and Conservation Research Institute Bratislava, Regional Workplace Presov, Presov, Slovakia

²Faculty of Humanities and Natural Sciences, University of Presov in Presov, Presov, Slovakia

³Faculty of Management, University of Presov in Presov, Presov, Slovakia

Received: 04.11.2013 • Accepted: 09.02.2014 • Published Online: 15.08.2014 • Printed: 12.09.2014

Abstract: Sorption of the cations NH_4^+ , Al^{3+} , Fe^{2+} , Mn^{2+} , and Mg^{2+} in powdered zeolite and its mixture with soil was monitored. More than 90% of the observed cations were fixed in the zeolite in the first few minutes of the laboratory experiment, which is a very fast speed of sorption, and the ammonia cations reacted fastest. Consequently, the dynamics of the nitrate and ammoniac nitrogen in soil after application of zeolite in different amounts was monitored. A decreased ammonium content was apparent in the variants with zeolite 1 month after zeolite application in comparison with variants without zeolite owing to the specific fixation of NH_4^+ cations in the zeolite lattice (92.5 mg in the control variant and 77.2–81.0 mg per kg of soil in the tested variants). Three months later, there was an increase in ammonium content in the variants with zeolite of 24% to 59% in comparison with the control variant owing to gradual ammonium release from the zeolite lattice. Six months after the zeolite application, statistically significant differences were found not only between the control variant and the variants with zeolite, but also between individual variants with various zeolite doses ($H = 14.201$; $P = 0.003$ according to the Kruskal–Wallis test). The nitrification process in the soil was less intense under the influence of applied zeolite. The nitrate nitrogen contents in the soil decreased by 66% to 78% in comparison with the control variant in the autumn period; therefore, the amount of nitrate leaching from the soil horizons to the groundwater was less. In summary, zeolite can be considered to be a slow-releasing nitrogen fertiliser.

Key words: Ammonium, nitrate, soil nitrogen dynamics, sorptive speed, zeolite

1. Introduction

Zeolites are crystalline hydrated aluminosilicates. Their basic building unit is a crystalline lattice of silicon tetrahedrons and aluminium octahedrons. Zeolites, as individual rocks, were discovered by the Swedish scientist Alex Frederik Cronstedt in 1756 (Cronstedt, 1756). Regardless of their impressive physicochemical properties, zeolites did not find practical application for a long time because investigations concentrated exclusively on zeolites of volcanic origin. The number of zeolites in the total volume of rocks rarely exceeded a few per cent. The discovery of zeolites of sedimentary origin allowed their use to be investigated seriously. Because of their extremely high sorptive capacity, which depends on the dominant mineral, zeolites can be utilised in different fields of national economy, including agriculture.

The utilisation of zeolites in agriculture as a fungicide, herbicide, or carrier of plant nutritional elements is feasible simply because of the high sorptive capacity of this rock. When nutrients are introduced into the soil in this way, their consumption is reduced, so there is no need

for redundant delivery of raw materials and consequently fewer nutrients (mostly nitrogen, which causes eutrophication of water sources) are leached into ground and surface waters. Zeolites from Italy (Ciambelli et al., 1985) and New Zealand (Nguyen and Tanner, 1998) have been used for ammonium removal from wastewater. After researching Turkish zeolites, Işıldar (1999) confirmed that nitrification decreased after zeolite application to soil, albeit not in all observed soil types. Zeolites from eastern Slovakia were used to protect the groundwater at waste disposal sites (Fronczyk and Garbulewski, 2013). Another use for natural zeolites is to decrease the heavy metal content of plants (Petkova et al., 2000) or reduce the soil contamination risk of heavy metals (Michalik and Missik, 1995).

2. Materials and methods

2.1. Natural zeolite

The zeolite used in this work was obtained from the Nizny Hrabovec deposit (48°51'36"N, 21°45'54"E) in Slovakia. Its chemical composition was as follows: SiO_2 , 70.9%; Al_2O_3 ,

* Correspondence: s.torma@vupop.sk

1.1%; Fe₂O₃, 1.4%; CaO, 2.9%; K₂O, 3.3%; MgO, 0.80%; Na₂O, 0.54%; H₂O⁺, 4.6%; H₂O⁻, 4.3%; and ignition loss, 2.78%. The rest was created by traces of other elements (titanium, manganese, phosphorus, sulphur, and others). The content of heavy metals in the zeolite was no higher than was permissible for waste application to agricultural soils in Slovakia (Table 1).

Zeolite with the dominant mineral clinoptilolite was from the Nizny Hrabovec deposit and its granulometric composition was as follows: fraction 0.06–0.02 mm, 11.10%; fraction 0.02–0.01 mm, 16.36%; and fraction <0.01 mm, 72.54%.

This powdered zeolite, containing particles smaller than 45 µm, contains on average 57.2% clinoptilolite (Singliar, 1992). In this work, the chemical composition of the powdered zeolite used was determined after sample analysis of a mixture of inorganic acids (Na, K) or after melting with Na₂CO₃ (Al, Ca, Fe, and Mg), and eventually zeolite heavy metal contents were gravimetrically (SiO₂, unbounded and bounded water) determined in lixivium of 2 M HNO₃.

2.2. Sorption of selected cations in zeolite powder

The sorption speed of NH₄⁺, Ca²⁺, Mg²⁺, Fe²⁺, and Mn²⁺ cations from a water solution in the powdered zeolite was evaluated first. The following procedure was chosen: to 1 L of solution (prepared from NH₄Cl, FeSO₄, MnSO₄, CaCl₂, and MgCl₂ salts) that contained 50 mg NH₄⁺, 30 mg Fe²⁺, 10 mg Mn²⁺, 200 mg Ca²⁺, and 100 mg Mg²⁺, 100 g of powdered zeolite was added and it was mixed for 1, 2, 4, 10, 20, 40, 80, 160, 320, 540, and 960 min. Ca, Mg, Mn, and Fe concentrations were determined by atom absorptive spectrophotometry, and ammonium cations were determined photometrically using the Nessler and Jendrasik method (Hanes, 1999). To determine cation sorptive capacity, a methodology based on direct displacement of the saturated salt NH₄Cl was used, whereby an adsorbed cation NH₄⁺ was directly displaced by KNO₃ solution. The cation NH₄⁺ was identified in the KNO₃ extract after rinsing with an acidified solution of NaCl by way of distillation and titration or photometry (Hanes, 1999).

2.3. Influence of natural zeolite on the mineral nitrogen dynamics in soil

In consequence of the results of the laboratory experiments, the effect of natural zeolite on the dynamics of mineral nitrogen in soil with a depth of 0.0–0.3 m was verified in field experiments. Selected chemical soil properties were determined using the obligatory methods for soil analysis (Fiala, 1999).

There were 4 variants of the experiment: K, control variant without any fertilisation; Z₁, 600 kg of zeolite per hectare; Z₂, 900 kg of zeolite per hectare; and Z₃, 1200 kg of zeolite per hectare.

All variants were carried out 4 times. The experimental field was 50 m² (10 × 5 m). The experimental crop was medium-ripening white cabbage (*Brassica oleracea* L., Slava variety) grown with the usual agricultural techniques. Seedlings were planted on 26 April and harvested on 25 August 2011. Average soil samples were taken and analysed for pH, nitrogen (ammonia and nitrate), and the contents of available phosphorus, potassium, magnesium, and organic carbon before the experiment was set up and after its completion on each site. In addition, once a month, soil samples from a depth of 0.0–0.3 m were taken from each site in order to monitor the dynamics of ammonia and nitrate nitrogen in the soil.

The statistical processing assumed that the zeolite dose influenced both ammonium and nitrate content in the soil. The Kruskal–Wallis test was chosen for verification of this hypothesis. In the case of rejection of the null hypothesis, the nonparametric Mann–Whitney test was used.

3. Results

3.1. Cation sorption speed in zeolite powder

The exchangeable zeolite cations Ca²⁺, Mg²⁺, K⁺, and Na⁺ are already present in natural deposits. These cations can exchange zeolite for NH₄⁺ and other metal cations, depending on the pH value of the solution and their concentrations.

The time-dependent development of NH₄⁺ sorption from solution in zeolite is shown in Figure 1, where the curve reflects the time dependence of the sorption of the initial NH₄⁺ concentration.

Table 1. Content of heavy metals in zeolite (in mg kg⁻¹ of solids in lixivium 2 M HNO₃).

	Pb	Cd	Cr	Cu	Zn	Ni
Zeolite	16.8	0.15	3.0	15.0	111	1.9
Maximum allowed content*	750	10.0	1000	1000	2500	300

*: According to the Act of the National Council of the Slovak Republic No. 188/2003 about waste application into agricultural soils.




Figure 1. Sorption rate of ammonia cations (NH_4^+) on powdered zeolite (0.1 kg L^{-1} , fraction under $45 \mu\text{m}$).

The development of the sorption indicates a very rapid response and, in the first minute, a sorption of more than 90% of the maximum potential ammonia concentration in zeolite on an infinite timescale can be observed. During the next period from 2 min up to several hours, the concentration of NH_4^+ does not change significantly and is almost constant. Figure 2 presents the rate of sorption of the other observed cations. Similar to the sorption rate of the ammonia cations, the iron cation sorption (Fe^{2+}) is very fast. A significant decrease in Fe^{2+} concentration could also be caused by Fe^{2+} oxidation to Fe^{3+} , and it is coagulated at filtration. Manganese (Mn^{2+}) sorption is also fast in the first minute, but as manganese has less sorption selectivity as regards zeolite, no significant reduction in concentration occurs, only approximately 40%. The calcium (Ca^{2+}) concentration in solution with time dependence did not have a tendency to increase. The concentration of Mg^{2+} was initially slightly decreased and then increased slightly later.

On the basis of experiments in which the time dependence of NH_4^+ , Fe^{2+} , Mn^{2+} , Ca^{2+} , and Mg^{2+} sorption in the zeolite powder was measured, it can be concluded that the sorption of the cations is extraordinarily fast. This is illustrated by the fact that in the first minute sorption in zeolite reached more than 90% of the maximum possible concentration of equilibrium, which would stabilise in




Figure 2. Sorption rate of selected cations Mg^{2+} , Fe^{2+} , Mn^{2+} , and Ca^{2+} on powdered zeolite (0.1 kg L^{-1} , fraction under $45 \mu\text{m}$).

infinite time. It can be assumed, however, that this process is dependent on the sizes of the zeolite particles, so when they increase, the stabilisation of sorption equilibrium, and especially that of NH_4^+ , will be slower.

3.2. The effect of zeolite application to soil on mineral nitrogen dynamics

The stabilisation of ionic exchange balance in several minutes or hours can be considered as fast because the zeolite will be in the soil for several years after its application. This knowledge can be applied to liquid nitrogen-based plant nutrition on the basis that zeolite applied to the soil would prevent the escape of redundant ammonium ions, as these would be bound to zeolite in moist soil very quickly and would be gradually released into the soil solution. The observed cation solution was verified by a field experiment based on clay eutric Cambisols, the basic chemical properties of which are shown in Table 2 (nutrient analyses according to the Mehlich method, organic carbon according to the Tjurin method).

In Figure 3, the dynamics of NH_4^+ in the soil is presented. It can be seen that after 1 month of zeolite application to the soil, the eutric Cambisol topsoil contained 14% to 20% more ammonia nitrogen (92.5 mg kg^{-1} of soil) in the control variant compared with the experimental variants ($77.2\text{--}81.0 \text{ mg kg}^{-1}$ of soil). There was a change in this subsequent period; the content of ammonia nitrogen in the topsoil in the variants with applied zeolites increased and after 3 months reached 24% to 59% higher values (depending on the amount of zeolite) than the control variant. At the end of the field experiment (5 months after the application of zeolite), the experimental variants reached $68.5\text{--}86.9 \text{ mg}$, and the control variant reached only 55.2 mg per kilogram of soil.

From the calculated values of the Mann–Whitney test, we can conclude that statistically significant differences in mean NH_4^+ concentrations in the soil were confirmed not only between the control variant and the observed ones, but between all observed variants, as well (Table 3).

Statistical analysis of the results according to the Kruskal–Wallis test confirmed the significant differences between the mean values of all observed variants ($H = 14.201$ with $P = 0.003$). The Mann–Whitney test results are given in Table 4.

After the zeolite application, the nitrate nitrogen content also changed during the observed period (Figure 4). All variants kept the spring maximum in April ($24.3\text{--}27.5 \text{ mg kg}^{-1}$ of soil) and the summer minimum in June and July ($1.4\text{--}4.9 \text{ mg kg}^{-1}$ of soil). At the time of crop harvest (in August) there was an apparent increase in nitrate nitrogen (12 mg kg^{-1} of soil) in the control variant, which corresponded to the so-called autumn maximum. This was not confirmed in the variants with zeolite because the nitrification process of ammonia nitrogen, owing to the

Table 2. Selected chemical soil parameters and clay content in the soil (eutric Cambisols).

Depth (m)	pH	N-NO ₃ ⁻ mg kg ⁻¹	N-NH ₄ ⁺ mg kg ⁻¹	N _{an}	N _{total}	P	K	Mg	C _{org} %	Clay %
0.0–0.3	7.2	34.3	26.8	61.1	2043	118	411	166	1.62	65.4
0.3–0.6	7.3	24.6	27.8	52.4	1717	89	294	173	1.31	58.6

presence of zeolite in the soil, becomes slower. Trapping and keeping nitrogen in the soil after a zeolite application was explained by Barbarick and Pirela (1984): NH₄⁺ cations penetrate the zeolite channels, the small size of which do not allow nitrifying bacteria to penetrate. From this perspective, the application of zeolite holds back the chemical change of ammonium cations into nitrate anions, thereby decreasing the amount of nitrate leaching into the groundwater. In this way, it is possible to significantly reduce nitrogen loss, which reaches, according to several authors, 25%–35% of the total amount of applied nitrogen in mineral fertilizers.

In our case, the content of nitrate nitrogen in the soil in the experimental variants was 66% to 78% lower in comparison with the control variant. Statistical analysis of the results according to the Kruskal–Wallis test confirmed the significant differences between the mean values of all observed variants ($H = 13.708$ with $P = 0.003$). The Mann–Whitney test results are given in Table 5.

4. Discussion

We found very fast cation sorption in zeolite but it should be noted that the particle size of the zeolite was 45 µm. The speed of sorption would be very different for larger particles. This is confirmed by Horvathova and Kachnak (1987), who measured the time dependence of heavy metal sorption in zeolite fraction of 0.2–3.2 mm. Stabilization of the exchange equilibrium was reached in only a few hours.

Our results showed a reduction of ammonia nitrogen content in the soil a short time after zeolite application. This reduction of the contents in the experimental variants can probably be attributed to fixation of ammonium ions by specific areas in the zeolite crystal lattice, because they

bind to the zeolite with the greatest intensity of all cations (Cicisvili and Andronikasvili, 1988). Similar findings were reached in our previous works (Torma and Chimic, 1992).

During the 5 months of the experiment, we found higher contents of ammonia nitrogen in the variants with zeolite, which means ammonia nitrogen is gradually released from zeolite and more is available for cultivated plants. Chelischeva and Chelischev (1984) and Tukvadze (1984) found a similar effect after zeolite application, although their experiments contained significantly higher zeolite amounts. Application of 2 t of zeolite in heavy soils increased the hydrolysable nitrogen content twice, and when they used 4 and 6 t of zeolite, it increased by 4 and 5 times, respectively. Sopkova et al. (1993), Ming and Aleln (2001), Rehakova et al. (2003), and Uher and Balogh (2004) suggested that fertiliser with the addition of zeolite became a 'slow-releasing fertiliser', although Ruzek and Kovanda (1996) debated the use of the term in relation to zeolite.

The fact that zeolite can relatively strongly fix the ammonia nitrogen leads to the reduction of nitrogen leaching. This is confirmed not only by our results, but also by researchers in many countries who have had positive results with zeolite application decreasing nitrogen loss due to leaching (e.g., Işildar, 1999; Sepaskhah and Yousefi, 2007; Gholamhoseini et al., 2009; Bernardi et al., 2010; Ippolito et al., 2011; Aghaalikhani et al., 2011). Zeolite is also used to remove nitrates from water (Mazeikiene et al., 2010).

Similar results were obtained by Bajrakov and Shevchenko (1984). In their experiments there was an observed decrease of the N-NO₃⁻ in the soil only to the level of 41%–48% after the zeolite application compared with the nitrate nitrogen content in the control variant.

Whereas at the beginning of the experiment statistically significant differences were confirmed only between the control variant and observed variants, in the last month of the experiment they were also confirmed between all observed variants. This was confirmed by a Kruskal–Wallis test with a relatively high level of significance ($H = 13.780$ and $P = 0.003$).

Regarding the obtained results, it is possible to conclude that sorption of cations NH₄⁺, Fe²⁺, Mn²⁺, Ca²⁺, and Mg²⁺ in zeolite with a grain size of less than 45 µm takes place very quickly. More than 90% of the maximum





Figure 3. Dynamics of ammonia nitrogen in the soil horizon 0.0–0.3 m after the application of different amounts of zeolite.

Table 3. The main statistical parameters from the last measurement at the end of the field experiment (ammonia nitrogen).

	Control variant	600 kg zeolite ha ⁻¹	900 kg zeolite ha ⁻¹	1200 kg zeolite ha ⁻¹
Minimum	54.50	67.30	69.40	86.10
Maximum	56.10	68.90	70.50	87.40
Median	55.20	68.90	70.25	87.05
Arithmetic mean	55.25	68.50	70.10	86.90
Standard deviation	0.733	0.800	0.483	0.572

Table 4. Mann–Whitney test results of the observed variants (regarding ammonium nitrogen).

	600 kg zeolite ha ⁻¹	900 kg zeolite ha ⁻¹	1200 kg zeolite ha ⁻¹
Control variant	0.000 + (0.029)	0.000 + (0.029)	0.000 + (0.029)
600 kg zeolite ha ⁻¹		0.000 + (0.029)	0.000 + (0.029)
900 kg zeolite ha ⁻¹			0.000 + (0.029)

**Figure 4.** Dynamics of nitrate nitrogen in the soil horizon 0.0–0.3 m after the application of different amounts of zeolite.

possible concentration of those cations is sorbed in the first few minutes. As zeolite has an effect on soil for several years after its application, this can be considered to be a significant fact in the prevention of nitrogen loss from soil.

The dynamics of ammonia nitrogen in the soil during the 5 months of observation justified zeolite's description as a slow-releasing nitrogen fertiliser. A short time after zeolite application, the nitrogen content in the soil decreased as a result of NH_4^+ fixation by zeolite, but after 3 or 4 months the opposite was true, and the increase was

probably caused by the gradual release of nitrogen from the zeolite crystal lattice. Therefore, the zeolite works in the soil as a sort of slow-releasing nitrogen fertiliser with NH_4^+ ions and indirectly limits the intensity of nitrification in the soil. Cultivated plants are thus provided a sufficient amount of nitrogen throughout the vegetation period. On the other hand, the presence of zeolite in the soil inhibits the process of nitrification to a certain extent, so that the nitrate leaching into deeper soil horizons is not so intensive.

Zeolite application to soil favourably affects the environment by preventing the leakage of mineral nutrients (especially nitrogen) into groundwater and increasing the efficiency of nutrients occurring in soil, and it does not negatively affect the quality of the cultivated products. It is fair to say that natural zeolite plays a positive role in plant nutrition.

Acknowledgements

The paper was supported by the Slovak Research and Development Agency under contract no. APVV-0131-11 and the Scientific Grant Agency of the Ministry of Education of the Slovak Republic under contract no. VEGA 1/0008/13.

Table 5. Mann–Whitney test results of the observed variants (regarding nitrate nitrogen).

	600 kg zeolite ha ⁻¹	900 kg zeolite ha ⁻¹	1200 kg zeolite ha ⁻¹
Control variant	0.000 + (0.029)	0.000 + (0.029)	0.000 + (0.029)
600 kg zeolite ha ⁻¹		1.500 - (0.570)	0.000 + (0.029)
900 kg zeolite ha ⁻¹			0.000 + (0.029)

References

- Aghaalikhani M, Gholamhoseini M, Dolatabadian A, Khodaei-Joghan A, Asilan KS (2012). Zeolite influences on nitrate leaching, nitrogen-use efficiency, yield and yield components of canola in sandy soil. *Arch Agr Soil Sci* 58: 1149–1169.
- Barbarick KA, Pirela HJ (1984). Agronomic and horticultural uses of zeolite. In: Pond WG, Mumpton FA, editors. *Zeo-agriculture: Use of Natural Zeolites in Agriculture and Aquaculture*. Boulder, CO, USA: Westview Press, pp. 93–104.
- Bernardi ACC, Oliveira PPA, Monte MBM, Polidoro JC, Souza-Barros F (2010). Brazilian sedimentary zeolite use in agriculture. In: Gilkes R, Prakongkep N, editors. *Proceedings of the 19th World Congress of Soil Science: Soil Solution for a Changing World*, 1–6 August 2010. Brisbane, Australia: Curran Associates, pp. 37–40.
- Chelischeva RV, Chelischev NF (1984). On the influence of clinoptilolite on distribution character of basic acting components of fertilisers in the system soil-plant. In: Cicisvili GV, editor. *Proceedings of the Conference and Symposium on the Utilization of Natural Zeolites in Animal Husbandry and Plant-growing*, 1–5 November 1981. Tbilisi, USSR: Mecniereba, pp. 234–235 (article in Russian with an abstract in English).
- Ciambelli P, Corbo P, Liberti L, Lopez A, Porcelli C (1985). Ammonia removal from municipal water by phillipsite. In: Drzaj B, Hocevar S, Pejovnik S, editors. *Proceedings of the International Symposium on Zeolites. Synthesis, Structure, Technology and Application*, 3–8 September 1984. Ljubljana, Yugoslavia: Elsevier, pp. 539–546.
- Cicisvili GV, Andronikasvili TG (1988). Ispolzovanije prirodnyh ceolitov v rastenijevodstve. In: Samsonija OA, editor. *Primenenije kliniptilolitsoderzhashich tufov v rastenijevodstve*. Tbilisi, USSR: Mecniereba, pp. 5–33 (in Russian).
- Cronstedt AF (1756). Observation and Description of an Unknown Kind of Rock to Be Named Zeolites. Stockholm, Sweden: Kongl Vetenskaps Academiens Handlingar.
- Fiala K (1999). *Metody Analyz Pod*. Bratislava, Slovakia: VUPOP (in Slovak).
- Fronczyk J, Garbulewski K (2013). Evaluation of zeolite-sand mixtures as reactive materials protecting groundwater at waste disposal sites. *J Environ Sci* 25: 1764–1772.
- Gholamhoseini M, Agha AM, Malakouti MJ (2009). Effect of zeolite on reducing nitrogen leaching in canola (*Brassica napus* L.) forage production in a sandy soil. *Iran J Soil Res* 23: 49–60.
- Hanes J (1999). *Analyza Sorpcnych Vlastnosti Pod*. Bratislava, Slovakia: VUPOP (in Slovak).
- Horvathova E, Kachnak S (1987). Heavy metals removal from water with clinoptilolite. *Vod Hosp* 1: 10–13 (article in Slovak with an abstract in English).
- Ippolito JA, Tarkalson DD, Lehrsch GA (2011). Zeolite soil application method affects inorganic nitrogen, moisture and corn growth. *Soil Sci* 176: 136–142.
- Işıldar AA (1999). Effect of addition of zeolite to the soil on nitrification. *Turk J Agric For* 23: 363–368 (article in Turkish with an abstract in English).
- Kovanda F, Ruzek P (1996). The release speed of ammonia ions from zeolites in the soils. *Rostl Vyr* 42: 149–154 (article in Czech with an abstract in English).
- Mazeikiene A, Valentukeviciene M, Rimeika M, Matuzevicius AB, Daukyns R (2010). Removal of nitrates and ammonium ions from water using natural sorbent zeolite (clinoptilolite). *J Environ Eng Landscape Manage* 16: 38–44.
- Michalik J, Missik J (1995). Sorption properties of zeolites from Nizny Hrabovec. *Acta Fytotechnica* 50: 131–141 (article in Slovak with an abstract in English).
- Ming DW, Allen ER (2001). Use of natural zeolites in agronomy, horticulture, and environmental soil remediation. *Rev Mineral Geochem* 45: 619–654.
- Nguyen ML, Tanner CC (1998). Ammonium removal from wastewater using natural New Zealand zeolites. *New Zeal J Agric Res* 41: 427–446.
- Petkova D, Popandova S, Popova T, Chakalov K (2000). Testing of zeolite-glaucanite-phosphorite mixtures for decrease of lead uptake by plants. *Bulg J Agric Sci* 6: 321–326.
- Rehakova M, Cuvanova S, Gavacova Z, Rimar J (2003). Application of natural zeolite of the clinoptilolite type in agrochemistry and agriculture. *Chem Listy* 97: 260–264.
- Sepaskhah AR, Yousefi F (2007). Effect of zeolite application on nitrate and ammonium retention of a loamy soil under saturated conditions. *Soil Res* 45: 368–373.
- Shevchenko LA, Bajrakov VV (1984). The influence of clinoptilolite on agro-chemical properties and fertility of light turf-podsol soils of Polesie region (the Ukrainian SSR). In: Cicisvili GV, editor. *Proceedings of the Conference and Symposium on the Utilization of Natural Zeolites in Animal Husbandry and Plant-growing*. 1–5 November 1981. Tbilisi, USSR: Mecniereba, pp. 106–112 (article in Russian with an abstract in English).
- Singliar M (1992). Use of natural zeolites in agriculture. *Agrochemia (Bratislava)* 32: 91–93 (article in Slovak with an abstract in English).
- Sopkova A, Bubanec J, Cernaj P, Singliar M, Mondik P, Budinska V, Fabian P (1993). The fertilizers with higher efficiency using natural zeolites. *Chem Prum* 43: 145–149 (article in Slovak with an abstract in English).
- Torma S, Chemic J (1992). Soil properties treatment by dressing of the natural zeolite. In: Dzatko M, editor. *Scientific Papers SFRI* 17. Bratislava, Czechoslovakia: VUPU, pp. 147–164 (article in Slovak with an abstract in English).
- Tukvadze ED (1984). Utilization of clinoptilolite on heavy clays. In: Cicisvili GV, editor. *Proceedings of the Conference and Symposium on the Utilization of Natural Zeolites in Animal Husbandry and Plant-growing*. 1–5 November 1981. Tbilisi, USSR: Mecniereba, pp. 223–225 (article in Russian with an abstract in English).
- Uher A, Balogh Z (2004). Use of zeolites for recultivation of sandy soils in horticulture. *Acta Horticulturae et Regiotecturae* 7: 46–52.