

1-1-2014

## Dry matter yield, feeding value, and antioxidant activity in Mediterranean chicory (*Cichorium intybus* L.) germplasm

GIOVANNA PILUZZA

LEONARDO SULAS

SIMONETTA BULLITTA

Follow this and additional works at: <https://journals.tubitak.gov.tr/agriculture>



Part of the [Agriculture Commons](#), and the [Forest Sciences Commons](#)

---

### Recommended Citation

PILUZZA, GIOVANNA; SULAS, LEONARDO; and BULLITTA, SIMONETTA (2014) "Dry matter yield, feeding value, and antioxidant activity in Mediterranean chicory (*Cichorium intybus* L.) germplasm," *Turkish Journal of Agriculture and Forestry*. Vol. 38: No. 4, Article 10. <https://doi.org/10.3906/tar-1309-70>  
Available at: <https://journals.tubitak.gov.tr/agriculture/vol38/iss4/10>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Agriculture and Forestry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## Dry matter yield, feeding value, and antioxidant activity in Mediterranean chicory (*Cichorium intybus* L.) germplasm

Giovanna PILUZZA, Leonardo SULAS\*, Simonetta BULLITTA

Institute for Animal Production System in Mediterranean Environments, National Research Council (CNR ISPAAM), Sassari, Italy

Received: 20.09.2013 • Accepted: 21.12.2013 • Published Online: 27.05.2014 • Printed: 26.06.2014

**Abstract:** Knowledge about the forage yield potential and bioactive compound contents of Mediterranean natural populations of chicory is still scarce, even though it could be very helpful in view of the exploitation of the feeding value of local chicory germplasm for ruminants. Dry matter yield, chemical composition, antioxidant capacity, total polyphenols, flavonoids, and condensed tannins content were determined in 7 natural populations of Mediterranean wild chicory and, as a comparison, in the chicory commercial variety Spadona. Original seed was collected in different pastureland areas of Sardinia (Italy) and plants were grown in North Sardinia. Statistically significant differences were found in the leaf dry matter yield, ranging from 1.3 to 3.8 t ha<sup>-1</sup>, and were not found in protein and fiber contents. The local populations were comparable to the commercial variety; however, the antioxidant activities, detected by means of 2 in vitro assays, significantly differed between populations. Among bioactive compounds, total flavonoids ranged from 27 to 37 g catechin equivalent kg<sup>-1</sup> dry weight and statistically significant differences were also found in total polyphenol content and nontannic polyphenols; condensed tannins were not detected. Considering the importance of some bioactive compounds for the enhancement of animal nutrition and welfare, the exploitation of variability in bioactive compound content and antioxidant capacity of the chicory natural populations can be important for a more complete valorization of its Mediterranean germplasm.

**Key words:** Chicory, dry matter yield, forage quality, natural populations, Trolox equivalent antioxidant capacity

### 1. Introduction

Chicory (*Cichorium intybus* L.) is a typical Mediterranean plant, belonging to the family Asteraceae, indigenous to Europe, West Asia, Egypt, North America, and Italy. There are many varieties of chicory having different commercial uses, with some misunderstanding about the botanical classification of such varieties. Vegetable commercial varieties (e.g., Bruxelles chicory, Treviso red chicory, endive) all belonging to the species *intybus* have remarkable economic importance. Spontaneous chicory plants are also still used for human consumption, cooked or uncooked, in some Mediterranean regions. In Sardinia, gastronomic uses and medicinal properties of chicory have been well known for a long time (Manca Dell'Arca, 1780); 2 other authors have referred to chicory ethnobotany (Moris, 1840; Atzei, 2003). Lancioni et al. (2007) reported on the alimentary and therapeutic uses of chicory in the folk traditions of central Sardinia.

Moreover, chicory is a spontaneous component of pasturelands, crops, and uncultivated land. Seeds of chicory plants, growing as weeds in forage legume crops, after seed harvesting and cleaning have been occasionally

used as a component of forage seed mixtures grazed in Mediterranean basin areas and should be considered for their forage potential (Molle et al., 2008). Chicory is palatable for ruminants, low in fiber, and high in nonstructural carbohydrates (Athanasiadou et al., 2007). In a recent paper, Di Grigoli et al. (2012) remarked on the potential of chicory to provide lasting good-quality herbage and the positive effect of this species on dairy performance in grazing ewes. In fact, in such an experiment, the crude protein (CP) content of the chicory was higher than that of an oat-berseem (*Avena sativa* L. and *Trifolium alexandrinum* L.) mixture and similar to that of sulla (*Hedysarum coronarium* L.). Moreover, the chicory forage had less dry matter (DM) and fewer structural carbohydrates, as it can remain green longer than other forage species. Because of such characteristics, chicory has been highly appreciated by farmers (Barry, 1998; Sitzia et al., 2006).

In spite of its natural distribution in many countries of Europe, forage chicory is a relatively new crop in many countries. In fact, much of the breeding for improved forage characteristics has been done in New Zealand,

\* Correspondence: l.sulas@cspm.ss.cnr.it

where the first variety, Puna, was released in 1985 (Moloney and Milne, 1993; Hume et al., 1995). Other forage chicory varieties, such as Lacerta, Forage Feast, Oasis, and Puna II, have been developed in France, the United States, Australia, and China (Rumball et al., 2003; Wang and Cui, 2011). This confirms the increased interest in forage chicory at a worldwide level. Currently, forage chicory is also being considered as a species with potential for reducing methane emissions from ruminants (Prusty et al., 2012, 2013; Sun et al., 2012). In addition, it has been tested in Turkey as a possible biomonitor of heavy metals in sites with different degrees of pollution (Aksoy, 2008).

According to Clark et al. (1990), a proportion of 70% leaves and 30% stems should be desirable in forage chicory because of the higher digestibility of leaves compared to stems. Therefore, frequent grazing has been proposed to produce a high proportion of leaves, up to 78% of the total aboveground biomass (Li et al., 1997).

Ozturk et al. (2006) studied the evolution of the nutritive value of wild chicory forage, harvested at different maturity stages in Turkey, and demonstrated that chicory should be harvested at the vegetative stage to obtain higher quality forage. Sulas et al. (2007) studied wild chicory plants from Sardinia and found that local populations of forage chicory can represent a valuable opportunity for semiarid environments under rainfed conditions. Moreover, wild chicory was shown to be a moderately salt-tolerant species (Sergio et al., 2012), which allows the use of brackish waters for its irrigation.

In the last 10 years, forage chicory has been studied for its secondary compounds, such as polyphenols, which can reduce intestinal infections in grazing animals (Marley et al., 2003). Polyphenols are among the most significant compounds related to the antioxidant properties of plant materials. Flavonoids have also been shown to act as scavengers of various oxidizing species [i.e. superoxide

anions ( $O_2^-$ ), hydroxyl radicals, and peroxy radicals], and they may also act as quenchers of single oxygen (Harborne and Williams, 2000).

Mulinacci et al. (2001) investigated the content of phenolic compounds, mainly cinnamic acids and flavonoids in *C. intybus* L. var. *silvestre*. Di Venere et al. (2004, 2009) also studied phenolic composition and antioxidant activity in wild chicory; they found the presence of caffeic acid derivatives and flavonoids (quercetin and kaempferol glycosides) and a very high antioxidant activity compared to other wild edible species. However, information about the dry matter yield potential and bioactive compounds contents of natural populations of chicory is still scarce, even if these details could be very helpful in establishing relationships such as that between forage polyphenol contents and their feeding values for ruminants. Within a general activity aimed at the in loco exploitation of Sardinian forage germplasm, the dry matter yield, chemical composition, bioactive compounds content (total polyphenols, flavonoids, condensed tannins), and antioxidant capacity were examined in the leaves of Mediterranean natural populations of chicory.

## 2. Materials and methods

### 2.1. Origin, management, and samplings of plant materials

*Cichorium intybus* seeds utilized in this investigation were previously collected from local natural populations in pastureland areas from 3 up to 962 m a.s.l. (Table 1) of Sardinia, Italy, where the species is usually grazed by ruminants and also used for human consumption (Atzei, 2003); the *C. intybus* Italian commercial variety Spadona was also included as a control.

The experiment was carried out in 2005 and 2006 in North Sardinia (41°N, 8°E, 81 m a.s.l.). The soil at the site is a flat sandy clay loam overlaid on limestone (Xerochrepts),

**Table 1.** List of collection and surveying sites of forage chicory germplasm in Sardinia (Italy).

Populations	FAO Soil Classification*	Altitude (m a.s.l.)	Latitude	Longitude
CIT01	Petric Calcisols	3	40°35'	8°18'
CIT02	Lithic Leptosols	275	40°28'	9°07'
CIT03	Eutric Cambisols	962	40°21'	8°55'
CIT04	Hypocalcic Calcisols	40	40°40'	8°21'
CIT05	Vertic Cambisol	158	39°41'	8°54'
CIT06	Rock outcrop	374	40°55'	8°38'
CIT07	Calcaric Cambisols	85	40°36'	8°29'

\*According to FAO et al. (2006) soil classification manual.

with low organic C ( $12 \text{ g kg}^{-1}$ ) and N ( $0.8 \text{ g kg}^{-1}$ ), pH 7.5, low  $\text{P}_2\text{O}_5$  content, and adequate  $\text{K}_2\text{O}$  content. The climate is typically Mediterranean, with a long-term average annual rainfall of 554 mm and a mean annual air temperature of  $16.2^\circ\text{C}$ .

In November 2005, plots ( $2.5 \text{ m}^2$  each) of 7 Sardinian natural populations of chicory and the Spadona commercial variety were manually transplanted, with 25 cm between rows and 20 cm apart within rows, under a randomized block design with 3 replicates. Fertilization of  $40 \text{ kg ha}^{-1}$  of N and  $100 \text{ kg ha}^{-1}$  of  $\text{P}_2\text{O}_5$  was applied at sowing. In spring 2006, when plants were at the vegetative stage, 3 plants per plot were cut at 5 cm above ground for production and quality determinations. Leaf length was measured in 4 leaves per plant.

## 2.2. Forage quality analysis

Forage samples were dried in an oven at  $65^\circ\text{C}$  for 48 h, then ground through a 1-mm screen to be analyzed for quality traits. Total N was determined using the Kjeldahl method and crude protein was calculated by multiplying the N content by 6.25. Neutral and acid detergent fiber (NDF and ADF) and acid detergent lignin (ADL) were determined using the procedure of Goering and Van Soest (1970). Total digestible nutrients (TDN), digestible dry matter (DDM), dry matter intake (DMI), relative feed value (RFV), and net energy for lactation ( $\text{NE}_l$ ) were estimated according to the following equations adapted from Horrocks and Vallentine (1999):

$$\text{TDN} = (-1.291 \times \text{ADF}) + 101.35,$$

$$\text{DMI} = 120 / \% \text{NDF dry matter basis},$$

$$\text{DDM} = 88.9 - (0.779 \times \% \text{ADF, dry matter basis}),$$

$$\text{RFV} = \% \text{DDM} \times \% \text{DMI} \times 0.775,$$

$$\text{NE}_l = (1.044 - (0.0119 \times \% \text{ADF})) \times 2.205.$$

## 2.3. Bioactive compounds and antioxidant capacity

Plant samples were kept on ice during harvesting, freeze-dried, and ground to a fine powder for the chemical analysis. Ground leaf samples (50 mg) were treated with 2.5 mL of acetone/water (7:3) mixture and shaken for 60 min. The samples were then centrifuged for 10 min at 3900 rpm and the supernatant was stored at  $4^\circ\text{C}$  until use for the following determinations. Total phenolic content (TotP) of extracts was determined using the Folin-Ciocalteu reagent according to Singleton and Rossi (1965), with some modifications by Piluzza and Bullitta (2010). Results were expressed as g gallic acid equivalent  $\text{kg}^{-1}$  dry weight of plant material (g GAE  $\text{kg}^{-1}$  DW). Nontannic phenolics (NTP) were determined after precipitation of tannin components with polyvinylpyrrolidone according to the FAO/IAEA (2000). Results were expressed as g GAE  $\text{kg}^{-1}$  DW. The butanol assay of Porter et al. (1986) was adapted (Piluzza and Bullitta, 2010) for quantification of extractable condensed tannins content from our samples. Total flavonoids (TotF) were quantified by colorimetric


assay with the  $\text{AlCl}_3$  method (Kim et al., 2003). Catechin was used as a standard and the flavonoid content was expressed as g catechin equivalent  $\text{kg}^{-1}$  dry weight of plant material (g CE  $\text{kg}^{-1}$  DW). Antioxidant capacity was determined by means of ABTS [(2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt)] and DPPH (1,1-diphenyl-2-picrylhydrazyl) assays (Surveswaran et al., 2007) with some modifications (Piluzza and Bullitta, 2011). Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid), a water-soluble analog of vitamin E, was used as the reference standard. The results were expressed in terms of Trolox equivalent antioxidant capacity (TEAC), as mmol Trolox equivalent  $100 \text{ g}^{-1}$  dry weight of plant material (mmol TEAC  $100 \text{ g}^{-1}$  DW).

## 2.4. Statistical analysis

Statistical significance was performed by 2-way analysis of variance (ANOVA). Differences between means were assessed with the least significant difference (LSD) test for means separation. The significance level was fixed at  $P \leq 0.05$  for all the statistical analyses. Coefficients of determination ( $R^2$ ) were calculated using Microsoft Excel 2000.

## 3. Results

Considering that a high proportion of leaves is desirable in chicory (Clark et al., 1990) and that phytomass partitioning demonstrated that stems' contribution to total plant dry matter was very limited compared to leaves at spring cutting harvest, the data referred to are for leaves only. Leaf dry matter content was on average 11.5% and it ranged from 10.1% (Spadona) to 13.7% (CIT01). Leaf length (Figure 1), leaf DM yield, and CP yield (Table 2) significantly varied between populations. The commercial variety Spadona exhibited the longest leaves, with an average of 56 cm, and statistically significant variations



**Figure 1.** Average leaf length (cm) of the natural populations of forage chicory and the commercial variety Spadona. Values marked by different letters are significantly different.

**Table 2.** Dry matter (DM), crude protein (CP), neutral detergent fiber (NDF), acid detergent fiber (ADF), and acid detergent lignin (ADL) contents in leaves of Mediterranean natural populations of forage chicory.

Populations	DM (kg ha <sup>-1</sup> )	CP (kg ha <sup>-1</sup> )	CP (g kg <sup>-1</sup> DM)	NDF (g kg <sup>-1</sup> DM)	ADF (g kg <sup>-1</sup> DM)	ADL (g kg <sup>-1</sup> DM)
CIT01	1963.5	363.4	173.9	372.7	303.4	188.1
CIT02	1729.3	334.5	200.7	396.1	332.9	175.2
CIT03	1372.9	255.8	185.2	406.5	336.8	205.8
CIT04	2889.3	515.8	177.0	379.8	309.0	157.9
CIT05	2176.4	400.2	182.6	364.9	290.6	166.2
CIT06	2113.3	407.5	193.0	395.2	343.2	188.2
CIT07	3808.4	689.2	182.5	406.2	336.5	186.3
Spadona*	3945.3	646.3	162.1	366.0	300.9	178.5
LSD <sub>0.05</sub>	809.1	196.7	ns	ns	ns	ns

\*: Commercial variety.

for the length of leaves were found among the wild populations. CIT07 leaves were about twice longer than those of CIT03. Leaf DM ranged from about 1.3 (CIT03) to 4 t ha<sup>-1</sup> of DM (Spadona); leaf DM yield of Spadona and CIT07 significantly differed from the remaining ones. Among Sardinian chicory populations, CIT07 exhibited significantly higher leaf lengths and leaf DM yields.

Chemical composition of chicory leaves did not significantly differ between populations. CP concentration ranged from 162 (Spadona) to 200 (CIT02) g kg<sup>-1</sup> DM, whereas, considering the CP production per hectare, leaf

CP yield varied from 255.8 (CT03) to 689.3 (CT07) kg ha<sup>-1</sup>, showing wide differences between populations due to the combined effect of both DM and CP content, the latter considered one of the most important criteria for forage quality evaluation. Other important characteristics for forage quality are NDF and ADF; NDF content ranged from 366 (Spadona) to 406.5 (CIT03) g kg<sup>-1</sup>; ADF content ranged from 290.6 (CIT05) to 336.8 (CIT03) g kg<sup>-1</sup> DM, whereas ADL content ranged from 157.9 (CIT04) to 205.8 (CIT03) g kg<sup>-1</sup> DM. TDN, DDM, DMI, RFV, and NE<sub>l</sub> (Table 3) did not significantly differ between populations.


**Table 3.** Total digestible nutrients (TDN), digestible dry matter (DDM), digestible dry matter intake (DMI), relative feed value (RFV), and net energy for lactation (NE<sub>l</sub>) in leaves of Mediterranean natural populations of forage chicory.

Populations	TDN (g kg <sup>-1</sup> DM)	DDM (g kg <sup>-1</sup> DM)	DMI (g kg <sup>-1</sup> of body weight)	RFV (%)	NE <sub>l</sub> (Mcal kg <sup>-1</sup> )
CIT01	621.8	652.6	32.4	165.1	1.505
CIT02	583.7	629.7	30.3	148.5	1.428
CIT03	578.7	626.6	29.5	143.7	1.418
CIT04	614.5	648.2	31.6	159.0	1.491
CIT05	638.2	662.6	33.1	170.6	1.539
CIT06	570.4	621.6	30.3	146.3	1.401
CIT07	578.9	626.8	29.7	145.1	1.418
Spadona*	624.9	654.5	32.8	166.4	1.512
LSD <sub>0.05</sub>	ns	ns	ns	ns	ns

\*: Commercial variety.

TDN ranged from 570.4 (CIT06) to 638.2 (CIT05) g kg<sup>-1</sup> DM and variations were related to the ADF concentration. DDM ranged from 621.6 (CIT06) to 662.6 (CIT05). The DMI range was 29.5–33.1 and it was negatively correlated with NDF. RFV exceeded a value of 151, which is indicative of prime forage (Horrocks and Vallentine, 1999), in CIT01, CIT04, CIT05, and Spadona; a similar trend was recorded for NE<sub>l</sub>.

Statistically significant differences between populations were found for antioxidant capacity, TotP, NTP (Table 4), and TotF content (Figure 2). ABTS assay exhibited a variation of antioxidant capacities among populations, from 13.63 (CIT04) to 17.51 (CIT01) mmol TEAC 100 g<sup>-1</sup> DW. However, all Sardinian populations, except CIT04, did not significantly differ from Spadona for antioxidant capacity determined by means of the ABTS method. Statistically significant differences were found for the total antioxidant capacity determined through the DPPH assay, ranging from 15.63 (CIT04) to 23.86 (CIT05) mmol TEAC 100 g<sup>-1</sup> DW and, except for CIT04 and CIT05, natural populations did not significantly differ from Spadona. TotP content (Table 4) showed statistically significant variation ranging from 33.76 (CIT07) to 43.89 (CIT05) g GAE kg<sup>-1</sup> DW; except for CIT04 and CIT07, TotP content of natural populations did not significantly differ from that of Spadona. The NTP content (Table 4) varied significantly from 20.27 to 25.85 g GAE kg<sup>-1</sup> DW in CIT07 and CIT01, respectively. TotF (Figure 2) showed a statistically significant variation and ranged from 27.13 (CIT04) to 36.94 (CIT05) g CE kg<sup>-1</sup> DW. Significant correlations were




**Figure 2.** Total flavonoid content of the natural populations of forage chicory and the commercial variety Spadona. Values marked by different letters are significantly different.

found between the CP (CP kg ha<sup>-1</sup>) and DM yield (DM kg ha<sup>-1</sup>) (Figure 3) and leaf length and CP (Figure 4a) and DM yield (Figure 4b). Significant correlations were also found between the TotF and total antioxidant capacity assayed by means of ABTS (Figure 5a) and DPPH (Figure 5b) methods. ABTS and total phenolics and nontannic phenolics showed a significant correlation of R<sup>2</sup> = 0.5651 and R<sup>2</sup> = 0.7511, respectively, whereas statistical significant correlations were not found between antioxidant capacity and DM yield.

**Table 4.** Trolox equivalent antioxidant capacity (TEAC) determined by means of ABTS and DPPH methods, total phenolic (TotP), and nontannic phenolic (NTP) content in leaves of the Mediterranean natural populations of forage chicory.

Populations	TEAC (mmol 100 g <sup>-1</sup> DW)		TotP (g GAE kg <sup>-1</sup> DW)**	NTP (g GAE kg <sup>-1</sup> DW)**
	ABTS	DPPH		
CIT01	17.51	20.66	41.92	25.85
CIT02	16.03	20.14	39.86	24.65
CIT03	15.75	19.58	38.60	23.49
CIT04	13.63	15.63	34.87	20.60
CIT05	17.49	23.86	43.89	24.01
CIT06	16.23	21.84	40.63	25.17
CIT07	14.10	18.12	33.76	20.27
Spadona*	15.93	20.18	40.60	22.06
LSD <sub>0.05</sub>	2.1	3.3	5.4	2.78

\*: Commercial variety. \*\*: Grams of gallic acid equivalent per kilogram of dry weight.





**Figure 3.** Linear regression between chicory crude protein (CP kg ha<sup>-1</sup>) and dry matter (DM kg ha<sup>-1</sup>).



**4. Discussion**

Notwithstanding the wide variations recorded, our results highlight the remarkable potential of some of the Mediterranean wild populations of chicory under study as a forage pasture species, because of their production of relevant amounts of leaves. In fact, Spadona and CIT07

produced a similar amount of dry matter that was 2 times higher than that of CIT03 and, compared to the remaining populations, showed higher plant vigor in winter, reaching an early complete ground cover rate of the plots (data not shown). With regard to protein and fiber content, Sardinian populations did not significantly differ from the commercial variety Spadona. According to Molle et al. (2008), the chicory commercial cultivar Spadona (positively evaluated for forage purposes), when ingested by sheep (who are able to select the more proteinaceous and less fibrous part of plants), can exhibit an improved nutritional quality compared to other forages available. The statistically nonsignificant variations recorded between Sardinian populations in terms of protein and fiber content suggest that substantial differences can arise from the different production potential of populations in terms of forage yield per hectare. In Sardinia, Nieddu et al. (2000) and Sulas (2004) suggested the use of chicory as a fodder crop; Sitzia et al. (2006) used cultivar Spadona and sulla as forage crops in a dairy sheep system, and they concluded that both chicory and sulla, cultivated under Mediterranean conditions, can serve as high-quality pasture for sheep. According to Sulas et al. (2009), the same



**Figure 4.** Linear regression between chicory leaf length and CP (a) and DM yield (b).



**Figure 5.** Linear regression between chicory total flavonoid content and antioxidant capacity determined by means of ABTS assay (a) and DPPH assay (b).



Spadona cultivar was able to produce up to 6 t ha<sup>-1</sup> of DM per year when harvested at a later stage (early flowering) but with a relevant contribution of stems to the total DM yield. The recorded values for Spadona leaf chemical composition, except for ADL, are quite similar to those reported by Sitzia et al. (2006) and Di Grigoli et al. (2012) for the same chicory cultivar grown in different grazing experiments in Sardinia and Sicily (Italy), respectively. The overall mean values obtained in our quality evaluations were consistent with the values reported for forage chicory quality in Turkey by Ozturk et al. (2006).

Comparisons with other published data about polyphenols and the antioxidant capacity of the species are quite difficult, owing to variations in the methods, procedures, and standards used for analyses. Ivanova et al. (2005) studied the polyphenols and antioxidant capacity of Bulgarian medicinal plants, including chicory; the plant extracts were prepared as an infusion with boiling water and the standard used was quercetin. Kratchanova et al. (2010) found in Bulgarian chicory a polyphenol content of 1821 mg 100 g<sup>-1</sup>, lower than our results. They prepared the plant extract with 80% acetone in 0.2% formic acid and the standard was gallic acid. Sulas et al. (2007) found a content of polyphenols of 30 g kg<sup>-1</sup> DM in Sardinian chicory populations and the Puna variety. It has been recognized that flavonoids show antioxidant activity and their effects on human nutrition and health are considerable. According to Pieroni et al. (2002), nondomesticated food vegetables play a central role as traditional functional foods; specifically, their antioxidant properties are regarded as quite noteworthy. Except for 2, all the other local populations showed total flavonoids content comparable or higher than Spadona and are therefore useful for both animal and human nutrition and health.

## References

- Aksoy A (2008). Chicory (*Cichorium intybus* L.): a possible biomonitor of metal pollution. *Pak J Bot* 40: 791–797.
- Athanasidou S, Gray D, Younie D, Tzamaloukas O, Jackson F, Kyriazakis I (2007). The use of chicory for parasite control in organic ewes and their lambs. *Parasitology* 134: 299–307.
- Atzei AD (2003). *Le piante nella tradizione della Sardegna*. Sassari, Italy: Delfino ed. (in Italian).
- Barry TN (1998). The feeding value of chicory (*Cichorium intybus*) for ruminant livestock. *J Agric Sci* 131: 251–257.
- Bullitta S, Piluzza G (2005). Evolution of condensed tannins content in Mediterranean natural populations of leguminous species and forage chicory. In: Bullitta S, editor. *Bioactive Compounds in Pasture Species for Phytotherapy and Animal Welfare*. Sassari, Italy: The ANFIT-MiPAF Project, pp. 19–29.
- No CT was detected in the natural populations under study, although some authors referred to its presence in forage chicory (Terrill et al., 1992; Molan et al., 2002; Häring et al., 2007, 2008). According to Piluzza et al. (2013), there have been conflicting reports on the presence of CT in forage chicory; in fact, the presence of CT was not detected in other studies on chicory (Waghorn et al., 2002; Bullitta and Piluzza, 2005; Tava, 2005; Sulas et al., 2007).
- Statistically significant variations were not recorded between Sardinian populations in terms of protein and fiber content. However, some Sardinian populations showed statistically significant differences in forage yield, bioactive compound contents, and antioxidant capacity, which can be important for animal feeding and also for human consumption. The results obtained show that some of the examined natural populations of chicory possess productive and qualitative traits comparable to Spadona, a commercial cultivar positively evaluated for forage purposes. The variability in bioactive compound contents and antioxidant capacity could be exploited for a more complete valorization of Mediterranean wild chicories and encourages investigations on Mediterranean germplasms of the species as sources of functional feed and food and natural antioxidants. Additional research needs to be accomplished in further investigation aimed at elucidating the chemical composition of phenolics.

## Acknowledgments

Part of this study was funded within the project CISIA (Integrated Knowledge for Sustainability and Innovation of the Agro-Food Made in Italy, 2011–2013), of the CNR - Dipartimento Scienze Bio-Agroalimentari.

Clark DA, Anderson CB, Berquist T (1990). Growth rates of 'Grasslands Puna' chicory (*Cichorium intybus* L.) at various cutting intervals and heights and rates of nitrogen. *N Z J Agric Res* 33: 213–217.

Di Grigoli A, Todaro M, Di Miceli G, Genna V, Tornambè G, Alicata ML, Giambalvo D, Bonanno A (2012). Effects of continuous and rotational grazing of different forage species on ewe milk production. *Small Rumin Res* 106/5: 529–536.

Di Venere D, Linsalata V, Sergio L, Cardinali A, Pieralice M, Bianco VV (2004). Composizione fenolica ed attività antiossidante di alcune specie spontanee eduli della Macchia Mediterranea. *Italus Hortus* 11: 128–131 (in Italian).

Di Venere D, Sergio L, Linsalata V, Pieralice M, Cardinali A, Cascarano N, Bianco VV (2009). Proprietà antiossidanti di specie erbacee spontanee eduli. *It J Agron/Riv Agron* 4 (Suppl. 4): 635–640 (in Italian).



- FAO/IAEA (2000). Quantification of Tannins in Tree Foliage. Vienna, Austria: Working Document, IAEA.
- FAO, ISSS, ISRIC (2006). World Reference Base for Soil Resources 2006. A Framework for International Classification, Correlation and Communication. World Soil Resource Report N. 103. Rome, Italy: FAO.
- Goering HK, Van Soest PJ (1970). Forage Fiber Analysis: Apparatus, Reagents, Procedures, and Some Applications. Agricultural Handbook 379.b. Washington DC, USA: U.S. Government Printing Office.
- Harborne JB, Williams CA (2000). Advances in flavonoid research since 1992. *Phytochemistry* 55: 481–504.
- Häring DA, Scharenberg A, Heckendorn F, Dohme F, Lüscher A, Maurer V, Suter D, Hertzberg H (2008). Tanniferous forage plants: agronomic performance, palatability and efficacy against parasitic nematodes in sheep. *Renew Agric Food Syst* 23: 19–29.
- Häring DA, Suter, D, Amrhein N, Lüscher A (2007). Biomass allocations is an important determinant of the tannin concentration in growing plants. *Ann Bot* 99: 111–120.
- Horrocks RD, Vallentine JF (1999). Harvested Forages. London, UK: Academic Press.
- Hume DE, Lyons TB, Hay, RJM (1995). Evaluation of 'Grasslands Puna' chicory (*Cichorium intybus* L.) in various grass mixtures under sheep grazing. *NZ J Agric Res* 1995: 38, 317–328.
- Ivanova I, Gerova D, Chervenkov T, Yankova T (2005). Polyphenols and antioxidant capacity of Bulgarian medicinal plants. *J Ethnopharmacol* 96: 145–150.
- Kim DO, Chun OK, Kim YJ, Moon HY, Lee CY (2003). Quantification of polyphenolics and their antioxidant capacity in fresh plums. *J Agric Food Chem* 51: 6509–6515.
- Kratchanova M, Denev P, Ciz M, Lojek A, Mihailov A (2010). Evaluation of antioxidant activity of medicinal plants containing polyphenol compounds. Comparison of two extraction systems. *Acta Biochim Pol* 57: 229–234.
- Lancioni MC, Ballero M, Mura L, Maxia A (2007). Usi popolari e terapeutici nella tradizione popolare del Goceano (Sardegna Centrale). *Atti Soc Tosc Sci Nat Mem Serie B* 114: 45–56 (in Italian).
- Li GD, Kemp D, Hodgson J (1997). Regrowth, morphology and persistence of Grasslands Puna chicory (*Cichorium intybus* L.) in response to grazing frequency and intensity. *Grass Forage Sci* 52: 33–41.
- Manca Dell'Arca A (1870). Agricoltura di Sardegna. Nuoro, Italy: 2000 Illisso ed. (in Italian).
- Marley CL, Cook R, Barrett J, Keatingue R, Lampkin NH, McBride SD (2003). The effects of dietary forage on the development and survival of helminth parasites in ovine faeces. *Vet Parasitol* 118: 93–107.
- Molan AL, Duncan A, Barry TN, McNabb WC (2002). Effects of condensed tannins and sesquiterpene lactones extracted from chicory on the viability of deer lungworm larvae. *Proc NZ Soc Anim Prod* 60, 26–29.
- Moris JH (1840). Flora Sardoia. Naples, Italy: Taurini.
- Molle G, Decandia M, Cabiddu A, Landau SY, Cannas A (2008). An update on the nutrition of dairy sheep grazing Mediterranean pastures. *Small Rumin Res* 77: 93–112.
- Moloney SC, Milne GD (1993). Establishment and management of Grasslands Puna chicory used as a specialist, high quality forage herb. *Proc N Z Grassld Assoc* 55: 113–118.
- Mulinacci N, Innocenti M, Gallori S, la Marca G, Vincieri FF (2001). Optimisation of the chromatographic determination of polyphenols in the aerial parts of *Cichorium intybus* L. *Chromatographia* 54: 455–461.
- Nieddu S, Saba P, Stangoni AP, Sulas L (2000). La cicoria da foraggio, un'interessante alternativa. *L'informatore agrario* 12: 69–71 (in Italian).
- Ozturk D, Bal MA, Erol A, Sahin M, Ozkan CO, Karakas E, Ata M, Karabay P (2006). Determination of nutritive value of wild chicory (*Cichorium intybus*) forage harvested at different maturity stage using in vitro and in situ measurements. *Pak J Biol Sci* 9: 253–259.
- Pieroni A, Nebel S, Quave C, Münz H, Heinrich M (2002). Ethnopharmacology of liakra: traditional weedy vegetables of the Arbëreshë of the Vulture area in southern Italy. *J Ethnopharmacol* 81: 165–185.
- Piluzza G, Bullitta S (2010). The dynamics of phenolics concentration in some pasture species and implications for animal husbandry. *J Sci Food Agr* 90: 1452–1459.
- Piluzza G, Bullitta S (2011). Correlation between phenolic content and antioxidant properties in twenty-four plant species of traditional ethnoveterinary use in the Mediterranean area. *Pharmaceut Biol* 49: 240–247.
- Piluzza G, Sulas L, Bullitta S (2013). Tannins in forage plants and their role in animal husbandry and environmental sustainability: a review. *Grass Forage Sci* 69: 32–48.
- Porter LJ, Hristich LN, Chan BG (1986). The conversion of procyanidins and prodelphinidins to cyanidin and delphinidin. *Phytochemistry* 25: 223–230.
- Prusty S, Jha N, Kundu SS, Bala PA, Kumar A, Datt C (2012). Nutritional evaluation of chicory, berseem and oats fodders in growing buffalo calves. *Indian J Anim Nutr* 29: 157–160.
- Prusty S, Mohimi M, Kundu SS, Kumar A, Datt C (2013). Methane emissions from river buffaloes fed on green fodders in relation to the nutrient intake and digestibility. *Trop Anim Health Prod* 46: 65–70.
- Rumball WRA, Skipp RG, Keogh R, Claydon B (2003). 'Puna II' forage chicory (*Cichorium intybus* L.). *N Z J Agric Res* 46: 53–55.
- Sergio L, De Paola A, Cantore V, Pieralice M, Cascarano NA, Bianco VV, Di Venere D (2012). Effect of salt stress on growth parameters, enzymatic antioxidant system, and lipid peroxidation in wild chicory (*Cichorium intybus* L.). *Acta Physiol Plant* 34: 2349–2358.

- Singleton VL, Rossi JA (1965). Colorimetry of total phenolics with phosphor-molybdic-phosphotungstic acid reagent. *Am J Enol Vitic* 16: 144–158.
- Sitzia M, Ligios S, Fois N (2006). Sulla and chicory production and quality under sheep grazing management. In: Loveras J, Gonzales Rodriguez A, Vázquez-Janez O, Pineiro J, Santamaria O, Olea L, Poblaciones MJ, editors. Sustainable Grassland Productivity. Proceedings of the 21st General Meeting of the European Grassland Federation. 3–6 April 2006; Badajoz, Spain, pp. 448–450.
- Sulas L (2004). Forage chicory: a valuable crop for Mediterranean environments. *Cahiers Options Méditerranéennes* 62: 137–140.
- Sulas L, Canu S, Muresu R, Piluzza G (2007). A new annual forage chicory (*Chicorium intybus* L.) from Sardinia. In: Bullitta S, editor. Plant Genetic Resources of Geographical and “Other” Islands. Sassari, Italy: Sassari Publisher, pp. 115–119.
- Sulas L, Seddaiu G, Muresu R, Roggero PP (2009). Nitrogen fixation of sulla under Mediterranean conditions. *Agron J* 101: 1470–1478.
- Sun XZ, Hoskin SO, Zhang GG, Molano G, Muetzel CS, Pinares-Patiño CS, Clark H, Pacheco D (2012). Sheep fed forage chicory (*Cichorium intybus*) or perennial ryegrass (*Lolium perenne*) have similar methane emissions. *Anim Feed Sci Technol* 172: 217–225.
- Surveswaran S, Cai Y, Corke H, Sun M (2007). Systematic evaluation of natural phenolic antioxidant from 133 Indian medicinal plants. *Food Chem* 102: 938–953.
- Tava A (2005). Biologically active compounds of forage species: their characterization in plants in relation to quality of productions. In: Bullitta S, editor. Bioactive Compounds in Pasture Species for Phytotherapy and Animal Welfare. Sassari, Italy: The ANFIT-MiPAF Project, pp. 3–17.
- Terrill TH, Rowan AM, Douglas GB, Barry TN (1992). Determination of extractable and bound condensed tannin concentration in forage plants, protein concentrate meals and cereal grains. *J Sci Food Agric* 58: 321–329.
- Waghorn GC, Tavendale MH, Woodfield DR (2002). Methane-geneis from forages fed to sheep. *Proc N Z Grassld Assoc* 64: 167–171.
- Wang Q, Cui J (2011). Perspective and utilization technologies of chicory (*Cichorium intybus* L.): a review. *Afr J Biotechnol* 10: 1966–1977.