

1-1-2019

Coefficient bounds and distortion theorems for the certain analytic functions

OSMAN ALTINTAŞ

NİZAMİ MUSTAFA

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ALTINTAŞ, OSMAN and MUSTAFA, NİZAMİ (2019) "Coefficient bounds and distortion theorems for the certain analytic functions," *Turkish Journal of Mathematics*: Vol. 43: No. 2, Article 30. <https://doi.org/10.3906/mat-1812-85>

Available at: <https://journals.tubitak.gov.tr/math/vol43/iss2/30>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Coefficient bounds and distortion theorems for the certain analytic functions

Osman ALTINTAŞ¹, Nizami MUSTAFA^{2,*} ¹Department of Mathematics Education, Başkent University, Ankara, Turkey²Department of Mathematics, Faculty of Science and Letters, Kafkas University, Kars, Turkey

Received: 24.12.2018

Accepted/Published Online: 19.02.2019

Final Version: 27.03.2019

Abstract: In this paper, we introduce and investigate an analytic function class $P_q(\lambda, A, B)$ that we call the class of q -starlike and q -convex functions with respect to the parameter λ . We give coefficient bounds estimates, distortion bound and growth theorems for the functions belonging to this class.

Key words: Starlike function, convex function, q -starlike function, q -convex function, q -derivative

1. Introduction and preliminaries

Let $H(U)$ denote the class of analytic functions on the open unit disk $U = \{z \in \mathbb{C} : |z| < 1\}$ in the complex plane. Also, let T be the class of analytic functions $f \in H(U)$, which are normalized by $f(0) = 0 = f'(0) - 1$ with series expansion

$$f(z) = z - a_2z^2 - a_3z^3 - \dots - a_nz^n - \dots = z - \sum_{n=2}^{\infty} a_nz^n, a_n \geq 0. \quad (1.1)$$

Also, we denote by S the subclass of T consisting of functions which are also univalent in U .

We say that a function $f(z)$ is subordinate to $g(z)$ and written as $f \prec g$, if there exists a (Schwartz) function $\omega(z)$ with $\omega(0) = 0$ and $|\omega(z)| < 1$ such that $f(z) = g(\omega(z))$. In particular, when $g(z)$ is univalent then the above subordination is equivalent to $f(0) = g(0)$ and $f(U) \subset g(U)$ (see [14]).

Some of the important subclasses of S are $S^*(\alpha)$ and $C(\alpha)$, respectively, starlike and convex functions of order $\alpha \geq 0$ (for details, see [11, 14] and [27]).

$$S^*(\alpha) = \left\{ f \in S : \Re \left(\frac{zf'(z)}{f(z)} \right) > \alpha, z \in U \right\} \quad (1.2)$$

and

$$C(\alpha) = \left\{ f \in S : \Re \left(\frac{(zf'(z))'}{f'(z)} \right) > \alpha, z \in U \right\}. \quad (1.3)$$

$S^* = S^*(0)$ and $C = C(0)$ are, respectively, well-known starlike and convex functions in U . As it is known, geometrically, a function $f \in A$ is called starlike in U if $f(U)$ is starlike domain in the complex plane and convex in U if $f(U)$ is convex domain in the complex plane.

*Correspondence: nizamimustafa@gmail.com

2010 AMS Mathematics Subject Classification: 30C45, 30C50, 30C80

Jackson [16] introduced the q -derivative operator D_q of a function $f \in H(U)$ as follows:

$$D_q f(z) = \begin{cases} \frac{f(z)-f(qz)}{(1-q)z}, & \text{if } z \neq 0 \\ f'(0), & \text{if } z = 0 \end{cases}, \tag{1.4}$$

for $q \in (0, 1)$. It is clear that $\lim_{q \rightarrow 1^-} D_q f(z) = f'(z)$ for $f \in H(U)$.

The following is the formula for the q -derivative of the function z^n as follows:

$$D_q z^n = [n]_q z^{n-1}, n \in \mathbb{N}, \tag{1.5}$$

where

$$[n]_q = \sum_{k=1}^n q^{k-1}, [0]_q = 0 \tag{1.6}$$

is the q -analogue of the natural numbers (which is called the basic number n). It follows from (1.6) that $[n]_q = (1 - q^n)/(1 - q)$ and $\lim_{q \rightarrow 1^-} [n]_q = n$ for $q \in (0, 1)$.

Also,

$$D_q [f(z)g(z)] = f(qz)D_q g(z) + g(z)D_q f(z), \tag{1.7}$$

$$D_q [af(z) + bg(z)] = aD_q f(z) + bD_q g(z), a, b \in \mathbb{C}, \tag{1.8}$$

$$D_q \left[\frac{f(z)}{g(z)} \right] = \frac{g(z)D_q f(z) - f(z)D_q g(z)}{g(qz)g(z)}, g(qz)g(z) \neq 0 \tag{1.9}$$

and on a product of functions, the general Leibniz rule for action of powers of the q -derivative operator is

$$D_q^{(n)}(fg)(z) = \sum_{k=0}^n \binom{n}{k}_q D_q^{(k)} f(q^{n-k}z) D_q^{(n-k)} g(q^kz), \tag{1.10}$$

where $\binom{n}{k}_q$ is q -binomial coefficients given by

$$\binom{n}{k}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!},$$

for $k = 0, 1, 2, \dots, n$ such that

$$[n]_q! = \prod_{k=1}^n [k]_q, [0]_q! = 1$$

is q -factorial function. Note that $\lim_{q \rightarrow 1^-} [n]_q! = n!$ is factorial function, and $\lim_{q \rightarrow 1^-} \binom{n}{k}_q = \binom{n}{k}$ are binomial coefficients. We refer to [12, 15, 18] about some properties of difference operator D_q .

It follows from (1.10) that $D_q^2 f(z) = D_q(D_q f(z))$, and $D_q^n f(z) = D_q(D_q^{n-1} f(z))$ for $n = 3, 4, \dots$

It follows from (1.4) and (1.1) that

$$D_q f(z) = 1 - \sum_{n=2}^{\infty} [n]_q a_n z^{n-1} \tag{1.11}$$

and

$$D_q [zD_q f(z)] = D_q f(z) + zD_q^2 f(z).$$

For $q \in (0, 1)$, we define by $S_q^*(\alpha)$ and $C_q(\alpha)$ the subclass of T , which we will call, respectively, q -starlike and q -convex functions of order $\alpha \geq 0$, as follows

$$S_q^*(\alpha) = \left\{ f \in S : \Re \frac{zD_q f(z)}{f(z)} > \alpha, z \in U \right\},$$

$$C_q(\alpha) = \left\{ f \in S : \Re \left(\frac{D_q(zD_q f(z))}{D_q f(z)} \right) > \alpha, z \in U \right\}.$$

It is clear that $\lim_{q \rightarrow 1^-} S_q^*(\alpha) = S^*(\alpha)$ and $\lim_{q \rightarrow 1^-} C_q(\alpha) = C(\alpha)$.

Studies on q -difference equations were initiated by Jackson [17], Carmichael [10], Mason [19], Adams [1], and Trjizinsky [28]. A recent study in this subject was done by Bangerezako [9]. Some properties related with function theory involving q -theory were first introduced by Ismail et al. [15]. Moreover, there are more studies in this matter such as [2, 21, 23, 24]. As the study [2] suggests, there are a lot that can be done for this research topic. For example, q -analogy of starlikeness and convexity of analytic functions in the unit disk and in arbitrary simply connected domains would be interesting for researchers in this field. Recently, in [8], q -convexity for basic hypergeometric functions was studied. In a recent paper [24], Bieberbach conjecture problem for q -close-to-convex functions was estimated optimally. In fact, sharpness of their result is still open to debate as the authors also stated.

Recently, several subclasses of analytic functions have been investigated by many researchers using the q -derivative operator. For example, to define several subclasses of analytic functions and to study the extension of the theory of univalent functions, the theory of q -derivative, q -derivative operator can be used as tools.

In [3], by using applications of the q -derivative, it was shown that Szasz Mirakyan operators are convex when convex functions are taken such that their result generalizes well known results for $q = 1$. Also, in [3], the authors showed that q -derivatives of these operators approach to q -derivatives of approximated functions. Recently, by Uçar et al. [30], some properties of q -close-to-convex functions were studied. In [22], Ramachandran et al. studied certain bound for q -starlike and q -convex functions with respect to symmetric points. In [22], the coefficient estimates for the subclasses of q -starlike and q -convex functions with respect to symmetric points involving q -difference operator were established and the authors provided certain applications based on these results for subclasses of univalent functions defined by convolution.

Very recently, q -starlike functions were investigated and basic characterization, growth theorem, and distortion theorem for this class were given by Polatoğlu [20]. Also, q -close-to-convex function with respect to Janowski starlike functions were studied and basic characterization, growth theorem, and distortion theorem for this class were given by Uçar [29].

Inspired by the studies mentioned above, we introduce the function class $P_q(\lambda, A, B)$ as follows.

Definition 1.1 A function $f \in T$ given by (1.1) is said to be in the class $P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ if the following condition is satisfied

$$\frac{zD_q f(z) + \lambda z^2 D_q^2 f(z)}{\lambda z D_q f(z) + (1 - \lambda) f(z)} \prec \frac{1 + Az}{1 + Bz}, z \in U,$$

where $-1 \leq B < A \leq 1$.

In the special case for $A = -B = 1$ and $A = 1 - 2\alpha$, $B = -1$ from the Definition 1.1, we can give the following definitions, respectively.

Definition 1.2 A function $f \in T$ given by (1.1) is said to be in the class $P_q(\lambda, 1, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ if the following condition is satisfied

$$\frac{zD_q f(z) + \lambda z^2 D_q^2 f(z)}{\lambda z D_q f(z) + (1 - \lambda) f(z)} \prec \frac{1 + z}{1 - z}, z \in U.$$

Definition 1.3 A function $f \in T$ given by (1.1) is said to be in the class $P_q(\lambda, 1 - 2\alpha, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ if the following condition is satisfied:

$$\frac{zD_q f(z) + \lambda z^2 D_q^2 f(z)}{\lambda z D_q f(z) + (1 - \lambda) f(z)} \prec \frac{1 + (1 - 2\alpha)z}{1 - z}, z \in U.$$

Note that classes $\lim_{q \rightarrow 1^-} P_q(\lambda, A, B) = P(\lambda, A, B)$, and $P(\lambda, 1 - 2\alpha, -1)$ with $A = 1 - 2\alpha$, $B = -1$ were studied by Altıntaş [4, 5] and Altıntaş et al. [6, 7].

The main purpose of this paper is to give coefficient bounds estimates and distortion theorems for the functions belonging to the class $P_q(\lambda, A, B)$, $q \in (0, 1)$.

2. Coefficient bound estimates for the class $P_q(\lambda, A, B)$

In this section, we will investigate some inclusion results of the subclass $P_q(\lambda, A, B)$, $q \in (0, 1)$. Here, we provide coefficient estimates and some conditions for the functions belonging to this class.

A necessary condition for the functions belonging to the class $P_q(\lambda, A, B)$ is given by the following theorem.

Theorem 2.1 If $f \in T$ belongs to the class $P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ then the following condition is satisfied

$$\sum_{n=2}^{\infty} \left([n]_q - \beta \right) \left[1 + \left([n]_q - 1 \right) \lambda \right] a_n \leq 1 - \beta, \tag{2.1}$$

where $\beta = \frac{1-A}{1-B}$, $-1 \leq B < A \leq 1$. The result obtained here is sharp.

Proof Let $f \in T$. Throughout this study, we assume that $F(z)$ is defined by

$$F(z) = \lambda z D_q f(z) + (1 - \lambda) f(z), z \in U. \tag{2.2}$$

It is clear that $F(z)$ is an analytic function in U with conditions

$$F(0) = 0 = D_q F(0) - 1.$$

A simple computation using (1.1), (1.11), and (2.2) shows that $F(z)$ has the following power series expansion

$$F(z) = z - \sum_{n=2}^{\infty} A_n z^n, z \in U, \tag{2.3}$$

where

$$A_n = \left[1 + \left([n]_q - 1\right) \lambda\right] a_n, \quad n = 2, 3, \dots \tag{2.4}$$

Assume that $f \in P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$, $-1 \leq B < A \leq 1$. Then,

$$\frac{zD_q F(z)}{F(z)} \prec \frac{1 + Az}{1 + Bz}, \quad z \in U. \tag{2.5}$$

Using the definition of the subordination condition, we have

$$h(z) = \frac{zD_q F(z)}{F(z)} = \frac{1 + A\omega(z)}{1 + B\omega(z)}$$

with $\omega(0) = 0$ and $|\omega(z)| < 1$, $z \in U$.

It follows that

$$|\omega(z)| = \left| \frac{h(z) - 1}{A - Bh(z)} \right| < 1.$$

Then, by taking $h(z) = u + iv$ from the last inequality, we find that

$$1 - A^2 + (1 - B^2)u^2 < 1 - A^2 + (1 - B^2)(u^2 + v^2) < 2u(1 - AB);$$

that is,

$$(1 - B^2)u^2 - 2u(1 - AB)u + (1 - A^2) < 0.$$

The last inequality implies that

$$\frac{1 - A}{1 - B} < u = \Re(h(z)) < \frac{1 + A}{1 + B}$$

(see also inequality (17) in [26]).

Thus, we write

$$\Re\left(\frac{zD_q F(z)}{F(z)}\right) > \beta, \tag{2.6}$$

where $\beta = \frac{1-A}{1-B}$, $-1 \leq B < A \leq 1$.

From (2.3) and (2.6) using (1.11) for the function $F(z)$, we obtain

$$\Re\left(\frac{z - \sum_{n=2}^{\infty} [n]_q A_n z^n}{z - \sum_{n=2}^{\infty} A_n z^n}\right) > \beta, \quad z \in U.$$

The expression

$$\frac{z - \sum_{n=2}^{\infty} [n]_q A_n z^n}{z - \sum_{n=2}^{\infty} A_n z^n}$$

is real if the chosen z is real. Thus, from the last inequality letting $z \rightarrow 1$ through real values, we obtain

$$1 - \sum_{n=2}^{\infty} [n]_q A_n \geq \beta \left(1 - \sum_{n=2}^{\infty} A_n z^n \right),$$

which is equivalent to

$$\sum_{n=2}^{\infty} ([n]_q - \beta) A_n \leq 1 - \beta.$$

Taking values $A_n, n = 2, 3, \dots$ from (2.4), we complete the proof of inequality (2.1).

To see that result obtained in the theorem is sharp, we note that equality is attained in the inequality when $f(z)$ is chosen so that

$$f(z) = f_n(z) = z - \frac{1 - \beta}{([n]_q - \beta) [1 + ([n]_q - 1) \lambda]} z^n, z \in U, n = 2, 3, \dots \tag{2.7}$$

Hence, the result obtained in the theorem is sharp for the functions $f_n(z)$ defined by (2.7). Therefore, the proof of Theorem 2.1 is completed. □

From the Theorem 2.1, using different values of the parameters, we can easily obtain the following results.

Corollary 2.2 *If $f \in T$ belongs to the class $P_q(\lambda, 1, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ then the following condition is satisfied:*

$$\sum_{n=2}^{\infty} [1 + ([n]_q - 1) \lambda] [n]_q a_n \leq 1.$$

The result obtained here is sharp.

Corollary 2.3 *If $f \in T$ belongs to the class $P_q(\lambda, 1 - 2\alpha, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$, $\alpha \in [0, 1)$, then the following condition is satisfied:*

$$\sum_{n=2}^{\infty} ([n]_q - \alpha) [1 + ([n]_q - 1) \lambda] a_n \leq 1 - \alpha.$$

The result obtained here is sharp.

Corollary 2.4 *If $f \in T$ belongs to the class $S_q^*(\alpha)$, $q \in (0, 1)$, $\alpha \in [0, 1)$ then the following condition is satisfied:*

$$\sum_{n=2}^{\infty} ([n]_q - \alpha) a_n \leq 1 - \alpha.$$

The result obtained here is sharp.

Corollary 2.5 *If $f \in T$ belongs to the class $C_q(\alpha)$, $q \in (0, 1)$, $\alpha \in [0, 1)$ then the following condition is satisfied:*

$$\sum_{n=2}^{\infty} \left([n]_q - \alpha \right) [n]_q a_n \leq 1 - \alpha.$$

The result obtained here is sharp.

Note that related results obtained in Corollaries 2.4 and 2.5 for the classes $S^*(\alpha)$ and $C(\alpha)$, respectively, were obtained by Silverman in [25] as follows.

Theorem 2.6 *(see [25], Theorem 2) A function $f \in T$ is in the class $S^*(\alpha)$, $\alpha \in [0, 1)$ if and only if*

$$\sum_{n=2}^{\infty} (n - \alpha) a_n \leq 1 - \alpha.$$

Corollary 2.7 *(see [25], Corollary 2) A function $f \in T$ is in the class $S^*(\alpha)$, $\alpha \in [0, 1)$ if and only if*

$$\sum_{n=2}^{\infty} n(n - \alpha) a_n \leq 1 - \alpha.$$

On the coefficient bounds of the functions belonging to the class $P_q(\lambda, A, B)$, we give the following theorem.

Theorem 2.8 *If $f \in T$ belongs to the class $P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ then the following condition is satisfied:*

$$\begin{aligned} \sum_{n=2}^{\infty} a_n &\leq \frac{1 - \beta}{\left([n]_q - \beta \right) \left[1 + \left([n]_q - 1 \right) \lambda \right]}, \\ \sum_{n=2}^{\infty} [n]_q a_n &\leq \frac{(1 - \beta) [n]_q}{\left([n]_q - \beta \right) \left[1 + \left([n]_q - 1 \right) \lambda \right]}, n = 2, 3, \dots, \end{aligned}$$

where $\beta = \frac{1-A}{1-B}$, $-1 \leq B < A \leq 1$. The results obtained here are sharp.

Proof Let $f \in P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$. Then, using Theorem 2.1, we can write

$$\left([n]_q - \beta \right) \left[1 + \left([n]_q - 1 \right) \lambda \right] \sum_{n=2}^{\infty} a_n \leq \sum_{n=2}^{\infty} \left([n]_q - \beta \right) \left[1 + \left([n]_q - 1 \right) \lambda \right] a_n \leq 1 - \beta,$$

which is equivalent to the first assertion of the theorem.

Similarly, we write

$$\left[1 + \left([n]_q - 1 \right) \lambda \right] \sum_{n=2}^{\infty} \left([n]_q - \beta \right) a_n \leq \sum_{n=2}^{\infty} \left([n]_q - \beta \right) \left[1 + \left([n]_q - 1 \right) \lambda \right] a_n \leq 1 - \beta;$$

that is,

$$\left[1 + \left([n]_q - 1\right) \lambda\right] \sum_{n=2}^{\infty} [n]_q a_n \leq 1 - \beta + \left[1 + \left([n]_q - 1\right) \lambda\right] \beta \sum_{n=2}^{\infty} a_n,$$

In the right side of the last inequality using the first assertion of the theorem, we arrive at the following inequality

$$\left[1 + \left([n]_q - 1\right) \lambda\right] \sum_{n=2}^{\infty} [n]_q a_n \leq \frac{(1 - \beta) [n]_q}{[n]_q - \beta}.$$

From here, we can easily see that the second assertion of the theorem is true.

It is clear that the results obtained in theorem are sharp for the functions defined by (2.7).

Thus, the proof of Theorem 2.8 is completed. □

Remark 2.9 Numerous consequences of the results obtained in Theorem 2.8 can be given for different values of parameters A, B, λ , and β .

The following theorem is direct result of Theorem 2.1.

Theorem 2.10 If $f \in T$ belongs to the class $P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$ then the following inequality for the coefficients is satisfied:

$$a_n \leq \frac{1 - \beta}{\left([n]_q - \beta\right) \left[1 + \left([n]_q - 1\right) \lambda\right]}, \quad n = 2, 3, \dots,$$

where $\beta = \frac{1-A}{1-B}$, $-1 \leq B < A \leq 1$.

From Theorem 2.10, we arrive at the following results.

Corollary 2.11 If $f \in T$ belongs to the class $P_q(\lambda, 1, -1)$, $q \in (0, 1)$,

$\lambda \in [0, 1]$ then the following inequality for the coefficients is satisfied:

$$a_n \leq \frac{1}{\left[1 + \left([n]_q - 1\right) \lambda\right] [n]_q}, \quad n = 2, 3, \dots$$

Corollary 2.12 If $f \in T$ belongs to the class $P_q(\lambda, 1 - 2\alpha, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$, $\alpha \in [0, 1)$ then the following inequality for the coefficients is satisfied:

$$a_n \leq \frac{1 - \alpha}{\left([n]_q - \alpha\right) \left[1 + \left([n]_q - 1\right) \lambda\right]}, \quad n = 2, 3, \dots$$

Corollary 2.13 If $f \in T$ belongs to the class $S_q^*(\alpha)$, $q \in (0, 1)$, $\alpha \in [0, 1)$ then the following inequality for the coefficients is satisfied:

$$a_n \leq \frac{1 - \alpha}{[n]_q - \alpha}, \quad n = 2, 3, \dots$$

Corollary 2.14 *If $f \in T$ belongs to the class $C_q(\alpha)$, $q \in (0, 1)$, $\alpha \in [0, 1)$ then the following inequality for the coefficients is satisfied:*

$$a_n \leq \frac{1 - \alpha}{([n]_q - \alpha) [n]_q}, \quad n = 2, 3, \dots$$

Remark 2.15 *Numerous consequences of the results obtained in the above corollaries can be given for different values of parameters λ and α .*

3. Distortion bound and growth theorems for the class $P_q(\lambda, A, B)$

In this section, we give distortion and growth theorems for the functions belonging to the class $P_q(\lambda, A, B)$.

Our coefficient bounds enable us to prove the following theorem.

Theorem 3.1 *Let $f \in T$ belong to the class $P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$. Then,*

$$r - \frac{1 - \beta}{([2]_q - \alpha) [1 + ([2]_q - 1) \lambda]} r^2 \leq |f(z)| \leq r + \frac{1 - \beta}{([2]_q - \alpha) [1 + ([2]_q - 1) \lambda]} r^2,$$

$$|z| = r, \quad r \leq 1,$$

where $\beta = \frac{1-A}{1-B}$, $-1 \leq B < A \leq 1$, with equality for the function

$$f(z) = z - \frac{1 - \beta}{([2]_q - \beta) [1 + ([2]_q - 1) \lambda]} z^2, \quad |z| = r. \tag{3.1}$$

Proof Assume that $f \in P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$. Then, using Theorem 2.1, we can write

$$([2]_q - \beta) [1 + ([2]_q - 1) \lambda] \sum_{n=2}^{\infty} a_n \leq \sum_{n=2}^{\infty} ([n]_q - \beta) [1 + ([n]_q - 1) \lambda] a_n < 1 - \beta;$$

that is,

$$\sum_{n=2}^{\infty} a_n \leq \frac{1 - \beta}{([2]_q - \beta) [1 + ([2]_q - 1) \lambda]},$$

Using this, we can easily show that

$$|f(z)| \leq r + \sum_{n=2}^{\infty} a_n r^n \leq r + r^2 \sum_{n=2}^{\infty} a_n \leq r + \frac{1 - \beta}{([2]_q - \beta) [1 + ([2]_q - 1) \lambda]} r^2. \tag{3.2}$$

On the other hand,

$$|f(z)| \geq r - \sum_{n=2}^{\infty} a_n r^n \geq r - r^2 \sum_{n=2}^{\infty} a_n \geq r - \frac{1 - \beta}{([2]_q - \beta) [1 + ([2]_q - 1) \lambda]} r^2. \tag{3.3}$$

Thus, the combination of (3.2) and (3.3) completes the proof of the inequalities.

It is clear that equalities are attained in the inequalities when the function $f(z)$ is chosen as in (3.1).

Thus, the proof of Theorem 3.1 is completed. □

From the Theorem 3.1, we arrive at the following results.

Corollary 3.2 *Let $f \in T$ belong to the class $P_q(\lambda, 1, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$. Then,*

$$r - \frac{1}{\left[1 + \left([2]_q - 1\right) \lambda\right] [2]_q} r^2 \leq |f(z)| \leq r + \frac{1}{\left[1 + \left([2]_q - 1\right) \lambda\right] [2]_q} r^2, |z| = r, r \leq 1,$$

such that equality is achieved for the function

$$f(z) = z - \frac{1}{\left[1 + \left([2]_q - 1\right) \lambda\right] [2]_q} z^2, |z| = r.$$

Corollary 3.3 *Let $f \in T$ belong to the class $P_q(\lambda, 1 - 2\alpha, -1)$, $q \in (0, 1)$, $\lambda \in [0, 1]$, $\alpha \in [0, 1)$. Then,*

$$r - \frac{1 - \alpha}{\left([2]_q - \alpha\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{\left([2]_q - \alpha\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} r^2,$$

$$|z| = r, r \leq 1$$

such that equality is achieved for the function

$$f(z) = z - \frac{1 - \alpha}{\left([2]_q - \alpha\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} z^2, |z| = r.$$

Corollary 3.4 *Let $f \in T$ belong to the class $S_q^*(\alpha)$, $q \in (0, 1)$, $\alpha \in [0, 1)$. Then,*

$$r - \frac{1 - \alpha}{[2]_q - \alpha} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{[2]_q - \alpha} r^2, |z| = r, r \leq 1,$$

such that equality is achieved for the function

$$f(z) = z - \frac{1 - \alpha}{[2]_q - \alpha} z^2, |z| = r.$$

Corollary 3.5 *Let $f \in T$ belong to the class $C_q(\alpha)$, $q \in (0, 1)$, $\alpha \in [0, 1)$. Then,*

$$r - \frac{1 - \alpha}{\left([2]_q - \alpha\right) [2]_q} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{\left([2]_q - \alpha\right) [2]_q} r^2, |z| = r, r \leq 1,$$

such that equality is achieved for the function

$$f(z) = z - \frac{1 - \alpha}{\left([2]_q - \alpha\right) [2]_q} z^2, |z| = r.$$

Remark 3.6 Numerous consequences of the results obtained in the above corollaries can be given for different values of parameters λ and α .

The related results obtained in Corollaries 3.4 and 3.5 for the classes $S^*(\alpha)$ and $C(\alpha)$, respectively, were obtained by Silverman [25] as follows.

Theorem 3.7 (see [25], Theorem 4) If $f \in S^*(\alpha)$, $\alpha \in [0, 1)$, then

$$r - \frac{1 - \alpha}{2 - \alpha} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{2 - \alpha} r^2, |z| = r, r \leq 1$$

such that equality is achieved for the function

$$f(z) = z - \frac{1 - \alpha}{2 - \alpha} z^2, |z| = r.$$

Corollary 3.8 (see [25], Corollary of the Theorem 4) If $f \in C(\alpha)$, $\alpha \in [0, 1)$, then

$$r - \frac{1 - \alpha}{2(2 - \alpha)} r^2 \leq |f(z)| \leq r + \frac{1 - \alpha}{2(2 - \alpha)} r^2, |z| = r, r \leq 1$$

such that equality is achieved for the function

$$f(z) = z - \frac{1 - \alpha}{2(2 - \alpha)} z^2, |z| = r.$$

The following theorem is the growth theorem for the functions belonging to class $P_q(\lambda, A, B)$.

Theorem 3.9 Let $f \in T$ belong to the class $P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$. Then,

$$1 - \frac{(1 - \beta) [2]_q}{\left([2]_q - \beta\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} r \leq |D_q f(z)| \leq 1 + \frac{(1 - \beta) [2]_q}{\left([2]_q - \beta\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} r,$$

$$|z| = r, r \leq 1,$$

where $\beta = \frac{1-A}{1-B}$, $-1 \leq B < A \leq 1$, such that equality is achieved for the function

$$f(z) = z - \frac{(1 - \beta) [2]_q}{\left([2]_q - \beta\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} z^2, |z| = r. \tag{3.4}$$

Proof Let $f \in P_q(\lambda, A, B)$, $q \in (0, 1)$, $\lambda \in [0, 1]$. By simple computation, we obtain

$$|D_q f(z)| \leq 1 + \sum_{n=2}^{\infty} [n]_q a_n |z^{n-1}| \leq 1 + r \sum_{n=2}^{\infty} [n]_q a_n.$$

Using the second assertion of the Theorem 2.8, we obtain

$$|D_q f(z)| \leq 1 + \frac{(1 - \beta) [2]_q}{\left([2]_q - \beta\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} r.$$

Similarly, we can easily show that

$$|D_q f(z)| \geq 1 - \sum_{n=2}^{\infty} [n]_q a_n |z^{n-1}| \geq 1 - \frac{(1-\beta) [2]_q}{\left([2]_q - \beta\right) \left[1 + \left([2]_q - 1\right) \lambda\right]} r.$$

Thus, the proof of the inequalities is completed.

We can easily show that equalities are attained in the inequalities when the function $f(z)$ is chosen as in (3.4).

With this, the proof of Theorem 3.9 is completed. \square

Remark 3.10 Numerous consequences of the results obtained in Theorem 3.9 can be given for different values of parameters A, B and λ .

References

- [1] Adams CR. On the linear ordinary q -difference equation. *Annals of Mathematics* 1929; 30: 195-205.
- [2] Agrawal S, Sahoo SK. Geometric properties of basic hypergeometric functions. *Journal of Difference Equations and Applications* 2014; 20 (11): 1502-1522.
- [3] Ali A, Viyaj G. The q -derivative and applications to q -Szász Mirakyan operator. *Calcolo* 2006; 43 (3): 151-170.
- [4] Altıntas O. On a subclass of certain starlike functions with negative coefficients. *Mathematica Japonica* 1983; 36 (3): 489-495.
- [5] Altıntas O. Certain application of subordination associated with neighborhoods. *Haceteppe Journal of Mathematics and Statistics* 2010; 39 (4): 527-534.
- [6] Altıntas O, Irmak H, Srivastava HM. Fractional calculus and certain starlike functions with negative coefficients. *Computers & Mathematics with Applications* 1995; 30 (2): 9-15
- [7] Altıntas O, Özkan Ö, Srivastava HM. Neighbourhoods of certain family of multivalent functions with negative coefficients. *Computers & Mathematics with Applications* 2004; 47: 1667-1672.
- [8] Baricz A, Swaminathan A. Mapping properties of basic hypergeometric functions. *Journal of Classical Analysis* 2014; 5 (2): 115-128.
- [9] Bangerezako G. An Introduction to q -Difference Equations. University of Burundi, Bujumbura, 2007, preprint.
- [10] Carmichael RD. The general theory of linear q -difference equations. *American Journal of Mathematics* 1912; 34: 147-168.
- [11] Duren PL. *Univalent Functions*. Grundlehren der Mathematischen Wissenschaften. New York, NY, USA: Springer, 1983.
- [12] Exton H. *q -Hypergeometric Functions and Applications*. Chichester, UK: Ellis Horwood Limited, 1983.
- [13] Gasper G, Rahman M. *Basic hypergeometric series*. Cambridge, UK: Cambridge University Press, 1990.
- [14] Goodman AW. *Univalent Functions*. Volume I. Washington, USA: Polygonal, 1983.
- [15] Ismail MEH, Merkes E, Styer D. A generalization of starlike functions. *Complex Variables* 1990; 14: 77-84.
- [16] Jackson FH. On q -functions and a certain difference operator. *Transactions of the Royal Society of Edinburgh* 1908; 46: 253-281.
- [17] Jackson FH. On q -definite integrals. *The Quarterly Journal of Pure and Applied Mathematics* 1910; 41: 193-203.
- [18] Jackson FH. On basic double hypergeometric functions. *The Quarterly Journal of Mathematics* 1942; 13: 69-82.

- [19] Mason TE. On properties of the solution of linear q -difference equations with entire function coefficients. *American Journal of Mathematics* 1915; 37: 439-444.
- [20] Polatoğlu Y. Growth and distortion theorems for generalized q -starlike functions. *Advances in Mathematics Scientific Journal* 2016; 5 (1): 7-12.
- [21] Raghavender K, Swaminathan A. Close-to-convexity of basic hypergeometric functions using their Taylor coefficients. *Journal of Mathematical Analysis and Applications* 2012; 35: 111-125.
- [22] Ramachandran C, Kavitha D, Soupramanien T. Certain Bound for q -Starlike and q -Convex Functions with Respect to Symmetric Points. *International Journal of Mathematics and Mathematical Sciences* 2015; Article ID 205682, 7 pages.
- [23] Ronning F. Szegő quadrature formula arising from q -starlike functions: In continued fractions and orthogonal functions (Leon,1992), 345-352, Dekker, New York.
- [24] Sahoo SK, Sharma NL. On a generalization of close-to-convex functions. *Annales Polonici Mathematici* 2015; 113 (1): 93-108.
- [25] Silverman H. Univalent functions with negative coefficients. *Proceedings of the American Mathematical Society* 1975; 51 (1): 109-116.
- [26] Srivastava HM, Altıntaş O, Serenbay SK. Coefficient bounds for certain subclasses of starlike functions of complex order. *Applied Mathematics Letters* 2011; 24: 1359-1363.
- [27] Srivastava HM, Owa S. (editors) *Current topics in analytic function theory*. Singapore: World Scientific, 1992.
- [28] Trijitzinsky WJ. Analytic theory of linear q -difference equations. *Acta Mathematica* 1933; 61 (1): 1-38.
- [29] Uçar HEÖ. Coefficient inequality for q -starlike functions. *Applied Mathematics and Computation* 2016; 276: 122-126.
- [30] Uçar HEÖ, Mert O, Polatoğlu Y. Some properties of q -close-to-convex functions. *Hacettepe Journal of Mathematics and Statistics* 2017; 46 (6): 1105-1112.