

1-1-2019

## A Neutral relation between metallic structure and almost quadratic $\phi$ -structure

SİNEM GÖNÜL

İREM KÜPELİ ERKEN

AZİZ YAZLA

CENGİZHAN MURATHAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

GÖNÜL, SİNEM; ERKEN, İREM KÜPELİ; YAZLA, AZİZ; and MURATHAN, CENGİZHAN (2019) "A Neutral relation between metallic structure and almost quadratic  $\phi$ -structure," *Turkish Journal of Mathematics*: Vol. 43: No. 1, Article 21. <https://doi.org/10.3906/mat-1807-72>  
Available at: <https://journals.tubitak.gov.tr/math/vol43/iss1/21>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## A Neutral relation between metallic structure and almost quadratic $\phi$ -structure

Sinem GÖNÜL<sup>1</sup> , İrem KÜPELİ ERKEN<sup>2\*</sup> , Aziz YAZLA<sup>3</sup> , Cengizhan MURATHAN<sup>4</sup> 

<sup>1</sup>Science Institute, Uludag University, Gorukle, Bursa, Turkey

<sup>2</sup>Department of Mathematics, Faculty of Engineering and Natural Sciences, Bursa Technical University, Bursa, Turkey

<sup>3</sup>Department of Mathematics, Faculty of Sciences, Selcuk University, Konya, Turkey

<sup>4</sup>Department of Mathematics, Faculty of Art and Science, Uludag University, Bursa, Turkey

Received: 09.07.2018

Accepted/Published Online: 29.11.2018

Final Version: 18.01.2019

**Abstract:** In this paper, we give a neutral relation between metallic structure and almost quadratic metric  $\phi$ -structure. Considering  $N$  as a metallic Riemannian manifold, we show that the warped product manifold  $\mathbb{R} \times_f N$  has an almost quadratic metric  $\phi$ -structure. We define Kenmotsu quadratic metric manifolds, which include cosymplectic quadratic manifolds when  $\beta = 0$ . Then we give nice almost quadratic metric  $\phi$ -structure examples. In the last section, we construct a quadratic  $\phi$ -structure on the hypersurface  $M^n$  of a locally metallic Riemannian manifold  $\tilde{M}^{n+1}$ .

**Key words:** Polynomial structure, golden structure, metallic structure, almost quadratic  $\phi$ -structure

### 1. Introduction

In [10] and [9], Goldberg and Yano and Goldberg and Petridis respectively defined a new type of structure called a polynomial structure on an  $n$ -dimensional differentiable manifold  $M$ . The polynomial structure of degree 2 can be given by

$$J^2 = pJ + qI, \quad (1.1)$$

where  $J$  is a  $(1,1)$  tensor field on  $M$ ,  $I$  is the identity operator on the Lie algebra  $\Gamma(TM)$  of vector fields on  $M$ , and  $p, q$  are real numbers. This structure can be also viewed as a generalization of the following well known structures:

· If  $p = 0$ ,  $q = 1$ , then  $J$  is called an almost product or almost para complex structure and denoted by  $F$  [12, 16];

· If  $p = 0$ ,  $q = -1$ , then  $J$  is called an almost complex structure [18];

· If  $p = 1$ ,  $q = 1$ , then  $J$  is called a golden structure [6, 7];

· If  $p \in \mathbb{R} - (-2, 2)$  and  $q = -1$ , then  $J$  is called a poly-Norden structure [17];

· If  $p = -1$ ,  $q = \frac{3}{2}$ , then  $J$  is called an almost complex golden structure [1];

· If  $p$  and  $q$  are positive integers, then  $J$  is called a metallic structure [11].

If a differentiable manifold is endowed with a metallic structure  $J$  then the pair  $(M, J)$  is called a metallic

\*Correspondence: irem.erken@btu.edu.tr

2010 AMS Mathematics Subject Classification: Primary 53B25, 53B35, 53C15, 53C55; Secondary 53D15

manifold. Any metallic structure  $J$  on  $M$  induces two almost product structures on  $M$ :

$$F_{\pm} = \pm \left( \frac{2}{2\sigma_{p,q} - p} J - \frac{p}{2\sigma_{p,q} - p} I \right),$$

where  $\sigma_{p,q} = \frac{p + \sqrt{p^2 + 4q}}{2}$  is the metallic number, which is the positive solution of the equation  $x^2 - px - q = 0$  for  $p$  and  $q$  nonzero natural numbers. Conversely, any almost product structure  $F$  on  $M$  induces two metallic structures on  $M$ :

$$J_{\pm} = \pm \frac{2\sigma_{p,q} - p}{2} F + \frac{p}{2} I.$$

If  $M$  is Riemannian, the metric  $g$  is said to be compatible with the polynomial structure  $J$  if

$$g(JX, Y) = g(X, JY) \tag{1.2}$$

for  $X, Y \in \Gamma(TM)$ . In this case,  $(g, J)$  is called a metallic Riemannian structure and  $(M, g, J)$  a metallic Riemannian manifold [8]. By (1.1) and (1.2), one can get

$$g(JX, JY) = pg(JX, Y) + qg(X, Y),$$

for  $X, Y \in \Gamma(TM)$ . The Nijenhuis torsion  $N_K$  for arbitrary tensor field  $K$  of type  $(1, 1)$  on  $M$  is a tensor field of type  $(1, 2)$  defined by

$$N_K(X, Y) = K^2[X, Y] + [KX, KY] - K[KX, Y] - K[X, KY], \tag{1.3}$$

where  $[X, Y]$  is the commutator for arbitrary differentiable vector fields  $X, Y \in \Gamma(TM)$ . The polynomial structure  $J$  is said to be integrable if  $N_J \equiv 0$ . A metallic Riemannian structure  $J$  is said to be locally metallic if  $\nabla J = 0$ , where  $\nabla$  is the Levi-Civita connection with respect to  $g$ . Thus, one can deduce that a locally metallic Riemannian manifold is always integrable.

On the other hand, Debnath and Konar [8] recently introduced a new type of structure named the almost quadratic  $\phi$ -structure  $(\phi, \eta, \xi)$  on an  $n$ -dimensional differentiable manifold  $M$ , determined by a  $(1, 1)$ -tensor field  $\phi$ , a unit vector field  $\xi$ , and a 1-form  $\eta$ , which satisfy the following relations:

$$\phi\xi = 0,$$

$$\phi^2 = a\phi + b(I - \eta \otimes \xi); \quad a^2 + 4b \neq 0, \tag{1.4}$$

where  $a$  is an arbitrary constant and  $b$  is a nonzero constant. If  $M$  is a Riemannian manifold the Riemannian metric  $g$  is said to be compatible with the polynomial structure  $\phi$  if

$$g(\phi X, Y) = g(X, \phi Y),$$

which is equivalent to

$$g(\phi X, \phi Y) = ag(\phi X, Y) + b(g(X, Y) - \eta(X)\eta(Y)). \tag{1.5}$$

In this case,  $(g, \phi, \eta, \xi)$  is called an almost quadratic metric  $\phi$ -structure. The manifold  $M$  is said to be an almost quadratic metric  $\phi$ -manifold if it is endowed with an almost quadratic metric  $\phi$ -structure [8]. They

proved the necessary and sufficient conditions for an almost quadratic  $\phi$ -manifold to induce an almost contact or almost paracontact manifold.

Recently, Blaga and Hretcanu [3] characterized the metallic structure on the product of two metallic manifolds in terms of metallic maps and provided a necessary and sufficient condition for the warped product of two locally metallic Riemannian manifolds to be locally metallic. Moreover, Özkan and F. Yılmaz [15] investigated integrability and parallelism conditions for the metallic structure on a differentiable manifold.

This paper is organized in the following way.

Section 2 is the preliminaries section, where we recall some properties of an almost quadratic metric  $\phi$ -structure and warped product manifolds. In Section 3, we define the  $(\beta, \phi)$ -Kenmotsu quadratic metric manifold and cosymplectic quadratic metric manifold. We mainly prove that if  $(N, g, \nabla, J)$  is a locally metallic Riemannian manifold, then  $\mathbb{R} \times_f N$  is a  $(-\frac{f'}{f}, \phi)$ -Kenmotsu quadratic metric manifold, and we show that every differentiable manifold  $M$  endowed with an almost quadratic  $\phi$ -structure  $(\phi, \eta, \xi)$  admits an associated Riemannian metric. We prove that on a  $(\beta, \phi)$ -Kenmotsu quadratic metric manifold the Nijenhuis tensor  $N_\phi \equiv 0$ . We also give examples of  $(\beta, \phi)$ -Kenmotsu quadratic metric manifolds. Section 4 is devoted to quadratic  $\phi$ -hypersurfaces of metallic Riemannian manifolds. We show that there are almost quadratic  $\phi$ -structures on hypersurfaces of metallic Riemannian manifolds. Then we give the necessary and sufficient condition for the characteristic vector field  $\xi$  to be Killing in a quadratic metric  $\phi$ -hypersurface. Furthermore, we obtain the Riemannian curvature tensor of a quadratic metric  $\phi$ -hypersurface.

## 2. Preliminaries

Let  $M^n$  be an almost quadratic  $\phi$ -manifold. As in almost contact manifolds, Debmah and Konar [8] proved that  $\eta \circ \phi = 0, \eta(\xi) = 1$ , and  $rank \phi = n - 1$ . They also showed that the eigenvalues of the structure tensor  $\phi$  are  $\frac{a+\sqrt{a^2+4b}}{2}, \frac{a-\sqrt{a^2+4b}}{2}$ , and 0. If  $\lambda_i, \sigma_j$ , and  $\xi$  are eigenvectors corresponding to the eigenvalues  $\frac{a+\sqrt{a^2+4b}}{2}, \frac{a-\sqrt{a^2+4b}}{2}$ , and 0 of  $\phi$ , respectively, then  $\lambda_i, \sigma_j$ , and  $\xi$  are linearly independent. Denote the following distributions:

$$\begin{aligned} \cdot \Pi_r &= \{X \in \Gamma(TM) : \alpha LX = -\phi^2 X - (\frac{\sqrt{a^2+4b}-a}{2})\phi, \alpha = -2b - \frac{a^2+a\sqrt{a^2+4b}}{2}\}; \dim \Pi_r = r, \\ \cdot \Pi_s &= \{X \in \Gamma(TM) : \beta QX = -\phi^2 X + (\frac{\sqrt{a^2+4b}+a}{2})\phi X, \beta = -2b - \frac{a^2-a\sqrt{a^2+4b}}{2}\}; \dim \Pi_s = s, \\ \cdot \Pi_1 &= \{X \in \Gamma(TM) : bRX = \phi^2 X - a\phi X - bX = -b\eta(X)\xi\}; \dim \Pi_1 = 1. \end{aligned}$$

By the above notations, Debmah and Konar proved following theorem.

**Theorem 2.1 ([8])** *The necessary and sufficient condition that a manifold  $M^n$  will be an almost quadratic  $\phi$ -manifold is that at each point of the manifold  $M^n$  it contains distributions  $\Pi_r, \Pi_s$ , and  $\Pi_1$  such that  $\Pi_r \cap \Pi_s = \{\emptyset\}, \Pi_r \cap \Pi_1 = \{\emptyset\}, \Pi_s \cap \Pi_1 = \{\emptyset\}$ , and  $\Pi_r \cup \Pi_s \cup \Pi_1 = TM$ .*

Let  $(M^m, g_M)$  and  $(N^n, g_N)$  be two Riemannian manifolds and  $\tilde{M} = M \times N$ . The warped product metric  $\langle, \rangle$  on  $\tilde{M}$  is given by

$$\langle \tilde{X}, \tilde{Y} \rangle = g_M(\pi_* \tilde{X}, \pi_* \tilde{Y}) + (f \circ \pi)^2 g_N(\sigma_* \tilde{X}, \sigma_* \tilde{Y})$$

for every  $\tilde{X}$  and  $\tilde{Y} \in \Gamma(T\tilde{M})$  where  $f : M \xrightarrow{C^\infty} \mathbb{R}^+$  and  $\pi : M \times N \rightarrow M, \sigma : M \times N \rightarrow N$  the canonical projections (see [2]). The warped product manifolds are denoted by  $\tilde{M} = (M \times_f N, \langle, \rangle)$ . The function  $f$  is

called the warping function of the warped product. If the warping function  $f$  is 1, then  $\tilde{M} = (M \times_f N, \langle, \rangle)$  reduces the Riemannian product manifold. The manifolds  $M$  and  $N$  are called the base and the fiber of  $\tilde{M}$ , respectively. For a point  $(p, q) \in M \times N$ , the tangent space  $T_{(p,q)}(M \times N)$  is isomorphic to the direct sum  $T_{(p,q)}(M \times N) \oplus T_{(p,q)}(p \times N) \equiv T_p M \oplus T_q N$ . Let  $\mathcal{L}_{\mathcal{H}}(M)$  (resp.  $\mathcal{L}_{\mathcal{V}}(N)$ ) be the set of all vector fields on  $M \times N$ , which is the horizontal lift (resp. the vertical lift) of a vector field on  $M$  (a vector field on  $N$ ). Thus, a vector field on  $M \times N$  can be written as  $\bar{E} = \bar{X} + \bar{U}$ , with  $\bar{X} \in \mathcal{L}_{\mathcal{H}}(M)$  and  $\bar{U} \in \mathcal{L}_{\mathcal{V}}(N)$ . One can see that

$$\pi_*(\mathcal{L}_{\mathcal{H}}(M)) = \Gamma(TM), \sigma_*(\mathcal{L}_{\mathcal{V}}(N)) = \Gamma(TN)$$

and so  $\pi_*(\bar{X}) = X \in \Gamma(TM)$  and  $\sigma_*(\bar{U}) = U \in \Gamma(TN)$ . If  $\bar{X}, \bar{Y} \in \mathcal{L}_{\mathcal{H}}(M)$ , then  $[\bar{X}, \bar{Y}] = [\bar{X}, \bar{Y}] \in \mathcal{L}_{\mathcal{H}}(M)$  and similarly for  $\mathcal{L}_{\mathcal{V}}(N)$ , and also if  $\bar{X} \in \mathcal{L}_{\mathcal{H}}(M), \bar{U} \in \mathcal{L}_{\mathcal{V}}(N)$  then  $[\bar{X}, \bar{U}] = 0$  [13].

The Levi-Civita connection  $\bar{\nabla}$  of  $M \times_f N$  is related to the Levi-Civita connections of  $M$  and  $N$  as follows:

**Proposition 2.2** ([13]) *For  $\bar{X}, \bar{Y} \in \mathcal{L}_{\mathcal{H}}(M)$  and  $\bar{U}, \bar{V} \in \mathcal{L}_{\mathcal{V}}(N)$ ,*

(a)  $\bar{\nabla}_{\bar{X}} \bar{Y} \in \mathcal{L}_{\mathcal{H}}(M)$  is the lift of  ${}^M \nabla_X Y$ , that is,  $\pi_*(\bar{\nabla}_{\bar{X}} \bar{Y}) = {}^M \nabla_X Y$ ;

(b)  $\bar{\nabla}_{\bar{X}} \bar{U} = \bar{\nabla}_{\bar{U}} \bar{X} = \frac{X(f)}{f} \bar{U}$ ;

(c)  $\bar{\nabla}_{\bar{U}} \bar{V} = {}^N \nabla_U V - \frac{\langle U, V \rangle}{f} \text{grad} f$ , where  $\sigma_*(\bar{\nabla}_{\bar{U}} \bar{V}) = {}^N \nabla_U V$ .

Here the notation is simplified by writing  $f$  for  $f \circ \pi$  and  $\text{grad} f$  for  $\text{grad}(f \circ \pi)$ .

Now we consider the special warped product manifold

$$\tilde{M} = I \times_f N, \langle, \rangle = dt^2 + f^2(t)g_N.$$

In practice,  $(-)$  is omitted from lifts. In this case,

$$\tilde{\nabla}_{\partial_t} \partial_t = 0, \tilde{\nabla}_{\partial_t} X = \tilde{\nabla}_X \partial_t = \frac{f'(t)}{f(t)} X \text{ and } \tilde{\nabla}_X Y = {}^N \nabla_X Y - \frac{\langle X, Y \rangle}{f(t)} f'(t) \partial_t. \tag{2.1}$$

### 3. Almost quadratic metric $\phi$ -structure

Let  $(N, g, J)$  be a metallic Riemannian manifold with metallic structure  $J$ . By (1.1) and (1.2) we have

$$g(JX, JY) = pg(X, JY) + qg(X, Y).$$

Let us consider the warped product  $\tilde{M} = \mathbb{R} \times_f N$ , with warping function  $f > 0$ , endowed with the Riemannian metric

$$\langle, \rangle = dt^2 + f^2 g.$$

Now we will define an almost quadratic metric  $\phi$ -structure on  $(\tilde{M}, \tilde{g})$  by using a method similar to that in [5]. Denote arbitrarily any vector field on  $\tilde{M}$  by  $\tilde{X} = \eta(\tilde{X})\xi + X$ , where  $X$  is any vector field on  $N$  and  $dt = \eta$ . By the help of tensor field  $J$ , a new tensor field  $\phi$  of type (1, 1) on  $\tilde{M}$  can be given by

$$\phi \tilde{X} = JX, \quad X \in \Gamma(TN), \tag{3.1}$$

for  $\tilde{X} \in \Gamma(T\tilde{M})$ . Thus, we get  $\phi\xi = \phi(\xi + 0) = J0 = 0$  and  $\eta(\phi\tilde{X}) = 0$ , for any vector field  $\tilde{X}$  on  $\tilde{M}$ . Hence, we obtain

$$\phi^2\tilde{X} = p\phi\tilde{X} + q(\tilde{X} - \eta(\tilde{X})\xi) \tag{3.2}$$

and arrive at

$$\begin{aligned} &< \phi\tilde{X}, \tilde{Y} > = f^2g(JX, Y) \\ &= f^2g(X, JY) \\ &= < \tilde{X}, \phi\tilde{Y} >, \end{aligned}$$

for  $\tilde{X}, \tilde{Y} \in \Gamma(T\tilde{M})$ . Moreover, we get

$$\begin{aligned} &< \phi\tilde{X}, \phi\tilde{Y} > = f^2g(JX, JY) \\ &= f^2(pg(X, JY) + qg(X, Y)) \\ &= p < \tilde{X} - \eta(\tilde{X})\xi, \phi\tilde{Y} > + q(< \tilde{X}, \tilde{Y} > - \eta(\tilde{X})\eta(\tilde{Y})) \\ &= p < \tilde{X}, \phi\tilde{Y} > + q(< \tilde{X}, \tilde{Y} > - \eta(\tilde{X})\eta(\tilde{Y})). \end{aligned}$$

Thus, we have proved the following proposition.

**Proposition 3.1** *If  $(N, g, J)$  is a metallic Riemannian manifold, then there is an almost quadratic metric  $\phi$ -structure on warped product manifold  $(\tilde{M} = \mathbb{R} \times_f N, <, > = dt^2 + f^2g)$ .*

An almost quadratic metric  $\phi$ -manifold  $(M, g, \nabla, \phi, \xi, \eta)$  is called a  $(\beta, \phi)$ -Kenmotsu quadratic metric manifold if

$$(\nabla_X \phi)Y = \beta\{g(X, \phi Y)\xi + \eta(Y)\phi X\}, \beta \in C^\infty(M). \tag{3.3}$$

Taking  $Y = \xi$  in (3.3) and using (1.4), we obtain

$$\nabla_X \xi = -\beta(X - \eta(X)\xi). \tag{3.4}$$

Moreover, by (3.4) we get  $d\eta = 0$ . If  $\beta = 0$ , then this kind of manifold is called a cosymplectic quadratic manifold.

**Theorem 3.2** *If  $(N, g, \nabla, J)$  is a locally metallic Riemannian manifold, then  $\mathbb{R} \times_f N$  is a  $(-\frac{f'}{f}, \phi)$ -Kenmotsu quadratic metric manifold.*

**Proof** We consider  $\tilde{X} = \eta(\tilde{X})\xi + X$  and  $\tilde{Y} = \eta(\tilde{Y})\xi + Y$  vector fields on  $\mathbb{R} \times_f N$ , where  $X, Y \in \Gamma(TN)$  and  $\xi = \frac{\partial}{\partial t} \in \Gamma(\mathbb{R})$ . By help of (3.1), we have

$$\begin{aligned} (\tilde{\nabla}_{\tilde{X}} \phi)\tilde{Y} &= \tilde{\nabla}_{\tilde{X}} \phi\tilde{Y} - \phi\tilde{\nabla}_{\tilde{X}} \tilde{Y} \\ &= \tilde{\nabla}_X JY + \eta(\tilde{X})\tilde{\nabla}_\xi JY - \phi(\tilde{\nabla}_X \tilde{Y} + \eta(\tilde{X})\tilde{\nabla}_\xi \tilde{Y}) \\ &= \tilde{\nabla}_X JY + \eta(\tilde{X})\tilde{\nabla}_\xi JY - \phi(\tilde{\nabla}_X Y + X(\eta(\tilde{Y}))\xi + \eta(\tilde{Y})\tilde{\nabla}_X \xi \\ &\quad + \eta(\tilde{X})\tilde{\nabla}_\xi Y + \xi(\eta(\tilde{Y}))\eta(\tilde{X})\xi). \end{aligned} \tag{3.5}$$

Using (2.1) in (3.5), we get

$$\begin{aligned} (\tilde{\nabla}_{\tilde{X}}\phi)\tilde{Y} &= (\nabla_X J)Y - \frac{f'}{f} \langle X, JY \rangle \xi + \eta(\tilde{X}) \frac{f'}{f} JY - \phi(\eta(\tilde{Y})) \frac{f'}{f} X + \eta(\tilde{X}) \frac{f'}{f} Y \\ &= (\nabla_X J)Y - \frac{f'}{f} (\langle \tilde{X}, \phi\tilde{Y} \rangle \xi + \eta(\tilde{Y})\phi\tilde{X}). \end{aligned}$$

Since  $\nabla J = 0$ , the last equation is reduced to

$$(\tilde{\nabla}_{\tilde{X}}\phi)\tilde{Y} = -\frac{f'}{f} (\langle \tilde{X}, \phi\tilde{Y} \rangle \xi + \eta(\tilde{Y})\phi\tilde{X}). \tag{3.6}$$

Using  $\tilde{\nabla}_X \xi = \frac{f'}{f} X$ , we have

$$\tilde{\nabla}_{\tilde{X}} \xi = \frac{f'}{f} (\tilde{X} - \eta(\tilde{X})\xi).$$

Thus,  $\mathbb{R} \times_f N$  is a  $(-\frac{f'}{f}, \phi)$ -Kenmotsu quadratic metric manifold. □

**Corollary 3.3** *Let  $(N, g, \nabla, J)$  be a locally metallic Riemannian manifold. Then product manifold  $\mathbb{R} \times N$  is a cosymplectic quadratic metric manifold.*

**Example 3.4** *Blaga and Hretcanu [3] constructed a metallic structure on  $\mathbb{R}^{n+m}$  in the following manner:*

$$J(x_1, \dots, x_n, y_1, \dots, y_m) = (\sigma x_1, \dots, \sigma x_n, \bar{\sigma} y_1, \dots, \bar{\sigma} y_m),$$

where  $\sigma = \sigma_{p,q} = \frac{p+\sqrt{p^2+4pq}}{2}$  and  $\bar{\sigma} = \bar{\sigma}_{p,q} = \frac{p-\sqrt{p^2+4pq}}{2}$  for  $p, q$  positive integers. By Theorem 3.2  $H^{n+m+1} = \mathbb{R} \times_{e^t} \mathbb{R}^{n+m}$  is a  $(-1, \phi)$ -Kenmotsu quadratic metric manifold.

$M$  is said to be metallic shaped hypersurface in a space form  $N^{n+1}(c)$  if the shape operator  $A$  of  $M$  is a metallic structure (see [14]).

**Example 3.5** *In [14], Özgür and Yılmaz Özgür proved that an  $S^n(\frac{2}{p+\sqrt{p^2+4pq}})$  sphere is a locally metallic shaped hypersurfaces in  $\mathbb{R}^{n+1}$ . Using Theorem 3.2, we have*

$$H^{n+1} = \mathbb{R} \times_{\cosh(t)} S^n\left(\frac{2}{p + \sqrt{p^2 + 4q}}\right),$$

a  $(-\tanh t, \phi)$ -Kenmotsu quadratic metric manifold.

**Example 3.6** *Debnath and Konar [8] gave an example of an almost quadratic  $\phi$ -structure on  $\mathbb{R}^4$  as follows:*

*If the  $(1, 1)$  tensor field  $\phi$ , 1-form  $\eta$ , and vector field  $\xi$  are defined as*

$$\phi = \begin{bmatrix} 2 & 1 & 0 & 0 \\ 9 & 2 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \eta = [0 \ 0 \ 0 \ 1], \xi = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix},$$

then

$$\phi^2 = 4\phi + 5(I_4 - \eta \otimes \xi).$$

Thus,  $\mathbb{R}^4$  has an almost quadratic  $\phi$ -structure.

**Theorem 3.7** Every differentiable manifold  $M$  endowed with an almost quadratic  $\phi$ -structure  $(\phi, \eta, \xi)$  admits an associated Riemannian metric.

**Proof** Let  $\tilde{h}$  be any Riemannian metric. Putting

$$h(X, Y) = \tilde{h}(\phi^2 X, \phi^2 Y) + \eta(X)\eta(Y),$$

we have  $\eta(X) = h(X, \xi)$ . We now define  $g$  by

$$g(X, Y) = \frac{1}{\alpha + \delta} [\alpha h(X, Y) + \beta h(\phi X, \phi Y) + \frac{\gamma}{2} (h(\phi X, Y) + h(X, \phi Y)) + \delta \eta(X)\eta(Y)],$$

where  $\alpha, \beta, \gamma, \delta, q$  are nonzero constants satisfying  $\beta q = p\frac{\gamma}{2} + \alpha$ ,  $\alpha + \delta \neq 0$ . It is clearly seen that

$$g(\phi X, \phi Y) = pg(\phi X, Y) + q(g(X, Y) - \eta(X)\eta(Y))$$

for any  $X, Y \in \Gamma(TM)$ . □

**Remark 3.8** If we choose  $\alpha = \delta = q, \beta = \gamma = 1$ , then we have  $p = 0$ . In this case, we obtain Theorem 4.1 of [8].

**Proposition 3.9** Let  $(M, g, \nabla, \phi, \xi, \eta)$  be a  $(\beta, \phi)$ -Kenmotsu quadratic metric manifold. Then quadratic structure  $\phi$  is integrable; that is, the Nijenhuis tensor  $N_\phi \equiv 0$ .

**Proof** Using (3.2) in (1.3), we have

$$\begin{aligned} N_\phi(X, Y) &= \phi^2[X, Y] + [\phi X, \phi Y] - \phi[\phi X, Y] - \phi[X, \phi Y] \\ &= p\phi[X, Y] + q([X, Y] - \eta([X, Y])\xi) + \tilde{\nabla}_{\phi X}\phi Y \\ &\quad - \nabla_{\phi Y}\phi X - \phi(\nabla_{\phi X}Y - \nabla_Y\phi X) - \phi(\nabla_X\phi Y - \nabla_{\phi Y}X) \\ &= p\phi\nabla_X Y - p\phi\nabla_Y X + q\nabla_X Y - q\nabla_Y X - q\eta([X, Y])\xi \\ &\quad + (\nabla_{\phi X}\phi)Y - (\nabla_{\phi Y}\phi)X + \phi\nabla_Y\phi X - \phi\nabla_X\phi Y \end{aligned} \tag{3.7}$$

for  $X, Y \in \Gamma(TM)$ . By using (3.2), we have

$$\begin{aligned} p\phi\nabla_X Y - \phi\nabla_X\phi Y &= p\phi\nabla_X Y + (\nabla_X\phi)\phi Y - \nabla_X\phi^2 Y \\ &= -p(\nabla_X\phi)Y + (\nabla_X\phi)\phi Y - q\nabla_X Y; \\ &\quad + qX(\eta(Y))\xi + q(\eta(Y))\nabla_X\xi. \end{aligned}$$

If we write the last equation in (3.7), we get

$$\begin{aligned} N_\phi(X, Y) &= -p(\nabla_X\phi)Y + p(\nabla_Y\phi)X + (\nabla_X\phi)\phi Y - (\nabla_Y\phi)\phi X \\ &\quad + (\nabla_{\phi X}\phi)Y - (\nabla_{\phi Y}\phi)X + q(X\eta(Y)\xi - Y\eta(X)\xi - \eta([X, Y])\xi) \\ &\quad + q(\eta(Y)\nabla_X\xi - \eta(X)\nabla_Y\xi). \end{aligned} \tag{3.8}$$



Employing (3.6) and (3.2) in (3.8), we deduce that

$$\begin{aligned} N_\phi(X, Y) &= q(X\eta(Y)\xi - Y\eta(X)\xi - \eta([X, Y])\xi) \\ &= 0. \end{aligned}$$

This completes the proof of the theorem. □

#### 4. Quadratic metric $\phi$ -hypersurfaces of metallic Riemannian manifolds

**Theorem 4.1** *Let  $\tilde{M}^{n+1}$  be a differentiable manifold with metallic structure  $J$  and  $M^n$  be a hypersurface of  $\tilde{M}^{n+1}$ . Then there is an almost quadratic  $\phi$ -structure  $(\phi, \eta, \xi)$  on  $M^n$ .*

**Proof** Denote by  $\nu$  a unit normal vector field of  $M^n$ . For any vector field  $X$  tangent to  $M^n$ , we put

$$JX = \phi X + \eta(X)\nu, \tag{4.1}$$

$$J\nu = q\xi + p\nu, \tag{4.2}$$

$$J\xi = \nu, \tag{4.3}$$

where  $\phi$  is a (1,1) tensor field on  $M^n$ ,  $\xi \in \Gamma(TM)$  and  $\eta$  is a 1-form such that  $\eta(\xi) = 1$  and  $\eta \circ \phi = 0$ . On applying operator  $J$  to the above equality (4.1) and using (4.2), we have

$$\begin{aligned} J^2X &= J(\phi X) + \eta(X)J\nu \\ &= \phi^2X + \eta(X)(q\xi + p\nu). \end{aligned} \tag{4.4}$$

Using (1.1) in (4.4),

$$p\phi X + p\eta(X)\nu + qX = \phi^2X + \eta(X)(q\xi + p\nu).$$

Hence, we are led to the conclusion:

$$\phi^2X = p\phi X + q(X - \eta(X)\xi). \tag{4.5}$$

□

Let  $M^n$  be a hypersurface of an  $n + 1$ -dimensional metallic Riemannian manifold  $\tilde{M}^{n+1}$  and let  $\nu$  be a globally unit normal vector field on  $M^n$ . Denote  $\tilde{\nabla}$  the Levi-Civita connection with respect to the Riemannian metric  $\tilde{g}$  of  $\tilde{M}^{n+1}$ . Then the Gauss and Weingarten formulas are given respectively by

$$\tilde{\nabla}_X Y = \nabla_X Y + g(AX, Y)\nu,$$

$$\tilde{\nabla}_X \nu = -AX$$

for any  $X, Y \in \Gamma(TM)$ , where  $g$  denotes the Riemannian metric of  $M^n$  induced from  $\tilde{g}$  and  $A$  is the shape operator of  $M^n$ .

**Proposition 4.2** *Let  $(\tilde{M}^{n+1}, \langle, \rangle, \tilde{\nabla}, J)$  be a locally metallic Riemannian manifold. If  $(M^n, g, \nabla, \phi)$  is a quadratic metric  $\phi$ -hypersurface of  $\tilde{M}^{n+1}$ , then*

$$(\nabla_X \phi)Y = \eta(Y)AX + g(AX, Y)\xi, \tag{4.6}$$

$$\nabla_X \xi = pAX - \phi AX, \quad A\xi = 0, \tag{4.7}$$

and

$$(\nabla_X \eta)Y = pg(AX, Y) - g(AX, \phi Y). \tag{4.8}$$

**Proof** If we take the covariant derivatives of the metallic structure tensor  $J$  according to  $X$  by (4.1)–(4.3), the Gauss and Weingarten formulas, we get

$$\begin{aligned} 0 &= (\nabla_X \phi)Y - \eta(Y)AX - qg(AX, Y)\xi \\ &\quad + (g(AX, \phi Y) + X(\eta(Y)) - \eta(\nabla_X Y) - pg(AX, Y))\nu. \end{aligned} \tag{4.9}$$

If we identify the tangential components and the normal components of the equation (4.9), respectively, we have

$$(\nabla_X \phi)Y - \eta(Y)AX - qg(AX, Y)\xi = 0. \tag{4.10}$$

$$g(AX, \phi Y) + X(\eta(Y)) - \eta(\nabla_X Y) - pg(AX, Y) = 0.$$

Using the compatible condition of  $J$  and (4.1), we get

$$\begin{aligned} g(JX, JY) &= pg(X, JY) + qg(X, Y) \\ &= pg(X, \phi Y) + qg(X, Y). \end{aligned} \tag{4.11}$$

Expressed in another way, by help of (1.5) and (4.1), we obtain

$$\begin{aligned} g(JX, JY) &= g(\phi X, \phi Y) + \eta(X)\eta(Y) \\ &= pg(X, \phi Y) + q(g(X, Y) - \eta(X)\eta(Y)) + \eta(X)\eta(Y) \\ &= pg(X, \phi Y) + qg(X, Y) + (1 - q)\eta(X)\eta(Y). \end{aligned} \tag{4.12}$$

Considering (4.11) and (4.12), we get  $q = 1$ . By (4.10) we arrive at (4.6). If we put  $Y = \xi$  in (4.10) we get

$$\phi \nabla_X \xi = -AX - g(AX, \xi)\xi. \tag{4.13}$$

If we apply  $\xi$  on both sides of (4.13), we have  $A\xi = 0$ .

Applying  $\phi$  on both sides of the equation (4.13) and using  $A\xi = 0$ ,

$$\begin{aligned} -\phi AX &= p\phi \nabla_X \xi + (\nabla_X \xi - \eta(\nabla_X \xi)\xi) \\ &= -pAX + \nabla_X \xi. \end{aligned}$$

Hence, we arrive at the first equation of (4.7). By help of (4.7), we readily obtain (4.8). This completes the proof.  $\square$

**Proposition 4.3** ([4]) *Let  $(M, g)$  be a Riemannian manifold and let  $\nabla$  be the Levi-Civita connection on  $M$  induced by  $g$ . For every vector field  $X$  on  $M$ , the following conditions are equivalent:*

- (1)  $X$  is a Killing vector field; that is,  $L_X g = 0$ .
- (2)  $g(\nabla_Y X, Z) + g(\nabla_Z X, Y) = 0$  for all  $Y, Z \in \chi(M)$ .

**Proposition 4.4** *Let  $(M^n, g, \nabla, \phi, \eta, \xi)$  be a quadratic metric  $\phi$ -hypersurface of a locally metallic Riemannian manifold  $(\tilde{M}^{n+1}, \tilde{g}, \tilde{\nabla}, J)$ . The characteristic vector field  $\xi$  is a Killing vector field if and only if  $\phi A + A\phi = 2pA$ .*

**Proof** From Proposition 4.3, we have

$$g(\nabla_X \xi, Y) + g(\nabla_Y \xi, X) = 0.$$

Making use of (4.7) in the last equation, we get

$$pg(AX, Y) - g(\phi AX, Y) + pg(AY, X) - g(\phi AY, X) = 0.$$

Using the symmetric property of  $A$  and  $\phi$ , we obtain

$$2pg(AX, Y) = g(\phi AX, Y) + g(A\phi X, Y). \quad (4.14)$$

We arrive at the desired equation from (4.14).  $\square$

**Proposition 4.5** *If  $(M^n, g, \nabla, \phi, \xi)$  is a  $(\beta, \phi)$ -Kenmotsu quadratic hypersurface of a locally metallic Riemannian manifold on  $(\tilde{M}^{n+1}, \tilde{g}, \tilde{\nabla}, J)$ , then  $\phi A = A\phi$  and  $A^2 = \beta pA + \beta^2(I - \eta \otimes \xi)$ .*

**Proof** Since  $d\eta = 0$ , using (4.7), we have

$$\begin{aligned} 0 &= g(Y, \nabla_X \xi) - g(X, \nabla_Y \xi) \\ &= pg(Y, AX) - g(Y, \phi AX) - pg(X, AY) + g(X, \phi AY) \\ &= g(A\phi X - \phi AX, Y). \end{aligned}$$

Thus, we get  $\phi A = A\phi$ . By (3.3) and (4.6), we get

$$\beta(g(X, \phi Y)\xi + \eta(Y)\phi X) = \eta(Y)AX + g(AX, Y)\xi.$$

If we apply  $\xi$  on both sides of the last equation, we obtain

$$\beta g(X, \phi Y) = g(AX, Y).$$

Namely,

$$\beta \phi X = AX. \quad (4.15)$$

Putting  $AX$  instead of  $X$  and using (4.5) in (4.15), we get  $A^2 X = \beta pAX + \beta^2(X - \eta(X)\xi)$ . This completes the proof.  $\square$

By help of (4.15) we obtain the following:

**Corollary 4.6** *Let  $(M^n, g, \nabla, \phi, \xi)$  be a cosymplectic quadratic metric  $\phi$ -hypersurface of a locally metallic Riemannian manifold. Then  $M$  is totally geodesic.*

**Remark 4.7** *Hretcanu and Crasmareanu [11] investigated some properties of the induced structure on a hypersurface in a metallic Riemannian manifold, but the argument in Proposition 4.2 is to get the quadratic  $\phi$ -hypersurface of a metallic Riemannian manifold. In the same paper, they proved that the induced structure on  $M$  is parallel to the induced Levi-Civita connection if and only if  $M$  is totally geodesic.*

By Proposition 4.2, we have the following.

**Proposition 4.8** *Let  $(M^n, g, \nabla, \phi, \xi)$  be a quadratic metric  $\phi$ -hypersurface of a locally metallic Riemannian manifold. Then*

$$R(X, Y)\xi = p((\nabla_X A)Y - (\nabla_Y A)X) - \phi((\nabla_X A)Y - (\nabla_Y A)X),$$

for any  $X, Y \in \Gamma(TM)$ .

**Corollary 4.9** *Let  $(M^n, g, \nabla, \phi, \xi)$  be a quadratic metric  $\phi$ -hypersurface of a locally metallic Riemannian manifold. If the second fundamental form is parallel, then  $R(X, Y)\xi = 0$ .*

### References

- [1] Bilen L, Turanlı S, Gezer A. On Kaehler–Norden–Codazzi golden structures on pseudo-Riemannian manifolds. *Int J Geom Methods Mod Phys* 2018; 15: 1850080.
- [2] Bishop RL, O’Neill B. Manifolds of negative curvature. *T Am Math Soc* 1969; 145: 1-49.
- [3] Blaga AM, Hretcanu CE. Invariant, anti-invariant and slant submanifolds of a metallic Riemannian manifold. *arXiv: 1803.01415*.
- [4] Blair DE. *Riemannian Geometry of Contact and Symplectic Manifolds*. Progress in Mathematics, 203. Boston, MA, USA: Birkhäuser, 2002.
- [5] Carriazo A, Pérez-García MJ. Slant submanifolds in neutral almost contact pseudo-metric manifolds. *Differential Geom Appl* 2017; 54: 71-80.
- [6] Crasmareanu M, Hretcanu CE. Golden differential geometry. *Chaos Solitons Fractals* 2008; 38: 1229-1238.
- [7] Crasmareanu M, Hretcanu CE, Munteanu MI. Golden- and product-shaped hypersurfaces in real space forms. *Int J Geom Methods Mod Phys* 2013; 10: 1320006.
- [8] Debmath P, Konar A. A new type of structure on differentiable manifold. *Int Electron J Geom* 2011; 4: 102-114.
- [9] Goldberg SI, Petridis NC. Differentiable solutions of algebraic equations on manifolds. *Kodai Math Sem Rep* 1973; 25: 111-128.
- [10] Goldberg SI, Yano K. Polynomial structures on manifolds. *Kodai Math Sem Rep* 1970; 22: 199-218.
- [11] Hretcanu CE, Crasmareanu M. Metallic structures on Riemannian manifolds. *Revista de Launi on Matematica Argentina* 2013; 54: 15-27.
- [12] Naveira A. A classification of Riemannian almost-product manifolds. *Rend Di Mat Di Roma* 1983; 3: 577-592.
- [13] O’Neill B. *Semi-Riemannian Geometry*. New York, NY, USA: Academic Press, 1983.
- [14] Özgür C, Yılmaz Özgür N. Classification of metallic shaped hypersurfaces in real space forms. *Turk J Math* 2015; 39: 784-794.
- [15] Özkan M, Yılmaz F. Metallic structures on differentiable manifolds. *Journal of Science and Arts* 2018; 44: 645-660.
- [16] Pitis G. On some submanifolds of a locally product manifold. *Kodai Math J* 1986; 9: 327-333.
- [17] Şahin B. Almost poly-Norden manifolds. *International Journal of Maps in Mathematics* 2018; 1: 68-79.
- [18] Yano K, Kon M. *Structures on Manifolds*. Series in Pure Mathematics. Singapore: World Scientific, 1984.