

1-1-2014


Micropropagation of *Erodium sibthorpiatum* subsp. *sibthorpiatum*, an endemic threatened species of Uludağ Mountain (Bursa-Turkey)

BETÜL AKIN

İSMAİL KOCAÇALIŞKAN

GÜRCAN GÜLERYÜZ

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>

 Part of the [Botany Commons](#)

Recommended Citation

AKIN, BETÜL; KOCAÇALIŞKAN, İSMAİL; and GÜLERYÜZ, GÜRCAN (2014) "Micropropagation of *Erodium sibthorpiatum* subsp. *sibthorpiatum*, an endemic threatened species of Uludağ Mountain (Bursa-Turkey)," *Turkish Journal of Botany*. Vol. 38: No. 1, Article 15. <https://doi.org/10.3906/bot-1304-24>
Available at: <https://journals.tubitak.gov.tr/botany/vol38/iss1/15>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Micropropagation of *Erodium sibthorpiatum* subsp. *sibthorpiatum*, an endemic threatened species of Uludağ Mountain (Bursa-Turkey)

Betül AKIN^{1*}, İsmail KOCAÇALIŞKAN², Gürcan GÜLERYÜZ³

¹Department of Biology, Faculty of Science and Arts, Dumlupınar University, Kütahya, Turkey

²Department of Molecular Biology and Genetics, Faculty of Science and Arts, Yıldız Technical University, İstanbul, Turkey

³Department of Biology, Faculty of Science and Arts, Uludağ University, Bursa, Turkey

Received: 09.04.2013

Accepted: 23.09.2013

Published Online: 02.01.2014

Printed: 15.01.2014

Abstract: Uludağ Mountain is one of the Important Plant Areas of Turkey. Its native plant species are threatened by various human and environmental factors. In this study, 2 different micropropagation techniques, with Murashige Skoog (MS) and modified MS (ModMS) media, were used for *Erodium sibthorpiatum* Boiss. subsp. *sibthorpiatum*. Mineral composition was developed according to the plant requirements. Shoot tips were cultured on these media. Seedlings, obtained from in vitro germinated seeds, were used as explant sources in micropropagation studies. The highest percentage of shoot induction (11.33 shoots/explant) was determined in 1 ppm 6-benzyl amino purine and plant growth regulator-free MS medium (35.97 mm shoot length). Elongated shoots were successfully rooted in the ModMS medium with different naphthalene acetic acid concentrations. Shoots with regenerated roots were transferred to the greenhouse. This protocol could be used for conservation studies of endemic and endangered plant species.

Key words: *Erodium sibthorpiatum*, endangered, endemic, micropropagation

1. Introduction

Native plants are important components of the global biological diversity and are often highly adapted to specific environmental and climatic conditions (Sudharsan et al., 2003). Thus, many endemic plant species are becoming endangered because of the increasing human activity and consequent habitat deterioration (Güleryüz et al., 2011). Hence, there is a need to develop efficient techniques that allow large-scale multiplication and preservation of rare endemic plant species (Kapai et al., 2010). Besides the conventional propagation methods, in vitro methods provide an important opportunity for the propagation and preservation of endemic and endangered plant species (Fay, 1992; Rao, 2004).

Erodium sibthorpiatum Boiss. subsp. *sibthorpiatum* (Geraniaceae) is an endemic species of Uludağ Mountain in Turkey, the highest mountain in the Marmara region, including Thrace and the northwestern side of the Anatolian peninsula. It has a geomorphologically interesting structure with steep southern slopes of calcareous rocks (Ketin, 1983). The climate of the mountain varies with elevation, resulting in high biological diversity. This high plant diversity has led to the designation of Uludağ Mountain as one of the Important Plant Areas

(IPAs) of Turkey (Güleryüz et al., 2010). *Erodium sibthorpiatum* grows on limestone rocks in poor soils and prefers calcareous soils. The species is spread among the alpine communities (*Acantholimon ulucinum* Boiss. and *Festuca punctoria* Sm. communities) at upper altitudes of ridges and slopes above 2200 m. It is a densely caespitose perennial, 5–12 cm, forming hard wide hummocks; basal leaves 2-pinnatisect, canescent, blade ovate, laciniae shortly oblong, acute; stems erect or ascending and flowers pale lilac. *Erodium sibthorpiatum* shares its habitats with other endemic species including *Veronica caespitose* Boiss. var. *caespitose* Boiss., *Astragalus hirsutus* Vahl., *A. sibthorpiatus* Boiss., *Arabis drabiformis* Boiss., *Festuca punctoria* Sm., *Linum olympicum* Boiss., *Muscari bourgaei* Baker, *Galium olympicum* Boiss. Flowering time is July to August (Güleryüz, 2000).

The habitat of this species has been damaged by heavy recreational and winter sports activities. In alpine environments, natural or human-induced disturbances are fairly common. Damage to plant diversity on Uludağ Mountain has been primarily caused by the construction of hotels and ski lift areas, recreational activities, and the use of artificial snow (Pintar et al., 2009; Güleryüz et al., 2011). *Erodium sibthorpiatum* is classified as an Endangered

* Correspondence: bortaca@yahoo.com

(EN) species according to the IUCN classification and it is considered under threat of extinction according to the IPA Project (Ekim et al., 2000; Güler, 2010).

This species occupies less than 10 km² and its habitat is severely fragmented. An inferred decline in the area and/or quality of habitat leads to the assumption that the population size of approximately <250 mature individuals will decline. It is estimated that it will decline by at least 25% within 3 years. For this reason, the development of a viable micropropagation protocol is essential. It must be an effective and rapid method for clonal propagation and could prove useful for the conservation of many rare and endemic species (Sudha et al., 1998).

In vitro multiplication efficiency is widely dependent on culture medium, growth regulators, and genotypes (Ansari et al., 2009; Farsi et al., 2012). The deficiency of minerals in plants can cause biochemical, physiological, and morphological changes according to nutrients and the level of deficiency (Monteiro et al., 2000). The interaction of nutrient salts and plant growth regulators (PGRs) in the culture medium is also important and has long been recognized (Murashige and Skoog, 1962). Optimizing the culture medium is difficult because of interactions between different medium components, leading to the creation of various tissue culture media (Nas and Read, 2004).

Until now no study has been initiated on in vitro propagation of *Erodium sibthorpiatum*. It is hypothesized that developing a medium according to the results of soil analyses of the plant habitat will result in the best nutritional conditions for the growth of shoot tips in the culture medium.

2. Materials and methods

2.1. Plant material and sterilization

Mature seeds of the *Erodium sibthorpiatum* were collected from the alpine belt between 2000 and 2400 m on Uludağ Mountain during July–August 2009. Seeds were dried and cleaned in the laboratory before the experiment. Surface sterilization of the seeds was carried out with 0.5%, 1%, and 2% sodium hypochlorite (NaOCl) solution containing a few drops of surfactant Tween 20 for 10 min. This was followed by 3 rinsing steps with sterile distilled water.

2.2. Culture media and isolation of explants

Throughout the study, different strengths of Murashige Skoog (MS) medium were used for experiments. Full-strength MS basal medium with different concentrations of 6-benzyl amino purine (BAP), kinetin (Ki), and modified MS (ModMS) medium with different combinations of BAP/naphthalene acetic acid (NAA) and auxins [indole-3-butyric acid (IBA), indole-3-acetic acid (IAA), and NAA] were used as the initiation media for in vitro multiplication responses. MS medium was modified according to the soil analyses of the plant habitat (Table 1). The amounts

of NH₄⁺, NO₃⁻, Ca⁺⁺, and Cl⁻ ions and vitamins, including nicotinic acid and thiamine-HCl, were changed (Table 2). Sucrose (30%) was used as the carbon source in all the combinations. All media combinations were solidified by adding 0.7% agar.

Seeds were germinated in a growth chamber at 21/16 °C for a 16/8 h photoperiod, respectively. The pH of the nutrient medium was adjusted to 5.8 before autoclaving the medium at 121 °C and 1.1 atm for 20 min. Seeds were aseptically transferred into Magenta vessels containing medium with 0 (control) and 100 ppm of gibberellic acid (GA₃) (filter-sterilized and added to the growth media after autoclaving) (Kocaalişkan, 2008). Each trial consisted of 5 magenta vessels containing 10 seeds. Three replicates were used for each treatment.

At the end of 30–40 days (Figure 1), *Erodium sibthorpiatum* shoot tips were placed on MS media with various concentrations of BAP (0, 1, 2, 4, 6, and 8 ppm), Ki (0.5, 1, 2, and 3 ppm), and ModMS with various concentrations of BAP/NAA (5/1, 2/0.04, 2/0.1, 4/0.1, and 2/2 ppm) for 8 weeks. Shoots were subcultured 3 times for each application.

The regenerated shoots (2 to 3 cm) were excised and individually transferred to MS medium without plant growth regulators (control), with various concentrations of IBA and IAA (0.5, 1, 1.5, and 2 ppm) and ModMS medium control, and with various concentrations of NAA (0.1, 0.5, 1, and 2 ppm) in 50-mL Magenta vessels to test the rooting potential. The number of roots per shoot, root lengths, and rooting percentages were determined after 8 weeks from the culture initiation. Rooted plantlets were acclimatized in a growth chamber for several days. After that, plantlets were transferred to 16-cm pots containing 2:1 turf and perlite and were grown under greenhouse conditions.

2.3. Statistical analysis

Each treatment had 3 replicates containing 5 explants in each culture vessel. Collected data were subjected to t-tests and one-way analysis of variance (ANOVA, SPSS for Windows 18.0), and post-hoc analysis was performed using Duncan's multiple range test. Statistical analysis was performed separately for MS and ModMS media.

3. Results

In this study, 0.5%, 1%, and 2% NaOCl were used to identify the effect of sterilization on the decontamination and germination of *Erodium sibthorpiatum* seeds. When 3 different sterilization treatments were compared, a high level of contamination (45%) was observed at the lower percentage of NaOCl (0.5%). According to this result, 1% sodium hypochlorite treatment (67.33%) was selected as the optimum concentration for the decontamination and germination of *Erodium sibthorpiatum* seeds (Table 3).

The highest number of shoots per explant was obtained with 1 ppm BAP (11.33; Table 4), but the highest shoot

Table 1. Nutritional composition of MS basal and modified medium.

Elements	Basal MS medium (mg/L)	Modified MS medium (mg/L)
NH ₄ NO ₃	1650	2000
CaCl ₂ .2H ₂ O	440	660
MgSO ₄ .7H ₂ O	370	370
KH ₂ PO ₄	170	170
KNO ₃	1900	1900
H ₃ BO ₃	6.2	6.2
CoCl ₂ .6H ₂ O	0.025	0.025
CuSO ₄ .5H ₂ O	0.025	0.025
MnSO ₄ .4H ₂ O	22.3	22.3
KI	0.83	0.83
Na ₂ MoO ₄ .2H ₂ O	0.25	0.25
ZnSO ₄ .4H ₂ O	8.6	8.6
Na ₂ EDTA	37.3	37.3
FeSO ₄ .7H ₂ O	27.8	27.8
Vitamins		
Glycine	2.0	2.0
myo-Inositol	100	100
Nicotinic acid	0.5	0.025
Pyridoxine-HCl	0.5	0.5
Thiamine-HCl	0.1	0.5
Saccharose	30 g/L	30 g/L

length was determined in PGR-free MS medium (Table 4; Figure 1). The shoot explants of *Erodium sibthorpiatum* cultures were also successfully established in modified MS medium. The combination of cytokinin with auxin (4 BAP/0.1 NAA) in modified MS medium showed a higher response in shoot length (29.40 mm; Table 4). According to the results given in Table 5, while MS medium had significant impact on stem length and on number of leaves

Table 2. Alterations in MS basal medium for preparing ModMS nutritional composition.

Ions	Modifications
Ammonium (NH ₄ ⁺)	+4.38 mmol
Nitrate (NO ₃ ⁻)	+4.38 mmol
Calcium (Ca ⁺⁺)	+1.98 mmol
Chloride (Cl ⁻)	+3.96 mmol
Vitamins	
Nicotinic acid	-0.475 mg
Thiamine-HCl	+0.4 mg

+: Increased values; -: decreased values.

and shoots, ModMS medium had an impact on rooting parameters of *Erodium sibthorpiatum*.

In vitro regenerated shoots of *Erodium sibthorpiatum* showed different behaviors during the processes of rooting and acclimatization. According to the results of this study, the presence of auxins, including IBA, IAA, and NAA, in the culture medium is generally necessary to promote rooting of *Erodium sibthorpiatum* in vitro cultures. The number of roots per shoot (10.57 roots/shoot), root length (16.40 mm), and the percentage of rooting (84.17%) changed significantly with a concentration of 0.5 ppm IBA (Table 6). The addition of 0.5 ppm IAA to the MS culture medium increased the number of roots (8.47 roots/shoot), root length (18.87 mm), and rooting percentage (80.37%), but not as much as IBA. We compared the effect of MS and ModMS media on the rooting of seedlings and the results showed that ModMS medium with 0.1 ppm NAA was superior compared to MS medium with IBA and IAA (Table 6; Figure 1).

Well-rooted *Erodium sibthorpiatum* shoots were rinsed with sterilized water to remove residual rooting medium, were transplanted to pots containing a 2:1 mixture of turf and perlite, and were grown in a growth chamber. After that, regenerated *Erodium sibthorpiatum* plantlets were transferred to the greenhouse (Figure 1).



Figure 1. Seed germination, shoot regeneration, rooting, and acclimatization of *Erodium sibthorpiatum* subsp. *sibthorpiatum*. **a-** Seedlings of 30–40 days old cultured on MS medium, **b-** in vitro shoot regeneration from a shoot tip explant cultured on MS medium with 1 ppm BAP, **c-** microshoots rooted on ModMS medium with 0.1 ppm NAA, **d-** acclimatized plantlets grown in a greenhouse.

4. Discussion

The main objective of this study was to establish a rapid and proper micropropagation system for *Erodium sibthorpiatum*, which is a rare and threatened endemic plant of Uludağ-Bursa. The most important and rather difficult aspect of the in vitro techniques is the requirement to carry out various operations under aseptic conditions. Contamination with microorganisms, including viruses, bacteria, yeast, and fungi, is considered the most important reason for plant

losses during in vitro culturing (Sarasan et al., 2006; Omamor et al., 2007). During seed formation, embryos and associated tissues tend to exclude pathogens and foreign materials. The contents of the seed are essentially aseptic and the resultant seedlings can be maintained in aseptic conditions if the outer surface of the seed (seed coat) is sterilized with sodium hypochlorite (or other surface sterilants) (Mineo, 1990).

In our study, different concentrations of NaOCl were tested; 1% NaOCl showed significant reduction

Table 3. Effects of different concentrations of NaOCl on decontamination and germination of *Erodium sibthorpiatum* subsp. *sibthorpiatum* seeds.

Treatments	% of seeds germinated	% of seeds contaminated
0.5% NaOCl × 10 min	51.00 ± 2.08 b	45.00 ± 2.89 a
1.0% NaOCl × 10 min	67.33 ± 0.92 a	34.33 ± 1.48 b
2.0% NaOCl × 10 min	28.00 ± 1.04 c	33.00 ± 0.29 b

Means within a column followed by the same letter are not significantly different by Duncan's multiple range test ($P < 0.05$); ±SE.

Table 4. Effect of different culture media (MS and ModMS) and growth regulators (BAP, Ki, BAP, and NAA) on shoot formation of *Erodium sibthorpiatum* subsp. *sibthorpiatum*.

Medium	BAP	Ki	Number of	Shoot length	Shoot	Callus-forming
	(ppm)	(ppm)	shoots/explant	(mm)	explant %	explant %
MS	0	0	5.27 ± 0.37 de	35.97 ± 0.55 a	100.00 ± 0.0 a	0
	1	0	11.33 ± 0.44 a	27.87 ± 0.45 b	100.00 ± 0.0 a	100.00 ± 0.0 a
	2	0	7.83 ± 0.44 b	25.37 ± 0.32 c	100.00 ± 0.0 a	100.00 ± 0.0 a
	4	0	6.80 ± 0.42 bc	16.73 ± 0.37 ef	100.00 ± 0.0 a	100.00 ± 0.0 a
	6	0	4.70 ± 0.35 ef	14.77 ± 0.39 g	87.03 ± 0.58 c	100.00 ± 0.0 a
	8	0	7.47 ± 0.29 b	20.00 ± 0.58 d	93.73 ± 0.67 b	100.00 ± 0.0 a
	0	0.5	1.43 ± 0.23 g	9.8 ± 0.42 h	73.17 ± 0.44 d	100.00 ± 0.0 a
	0	1	4.17 ± 0.20 f	20.60 ± 0.38 d	86.90 ± 0.23 c	100.00 ± 0.0 a
	0	2	5.77 ± 0.39 cde	18.00 ± 0.58 e	100.00 ± 0.0 a	100.00 ± 0.0 a
	0	3	6.17 ± 0.18 cd	16.57 ± 0.38 f	100.00 ± 0.0 a	100.00 ± 0.0 a
Medium	BAP	NAA	Number of	Shoot length	Shoot	Callus-forming
	(ppm)	(ppm)	shoots/explant	(mm)	explant %	explant %
ModMS	5	1	6.17 ± 0.20 b	22.90 ± 0.29 b	75.00 ± 0.58 c	100.00 ± 0.0 a
	2	0.04	3.93 ± 0.23 c	19.03 ± 0.20 d	78.67 ± 0.88 b	100.00 ± 0.0 a
	2	0.1	3.72 ± 0.25 c	22.77 ± 0.15 b	81.33 ± 0.67 a	100.00 ± 0.0 a
	4	0.1	6.88 ± 0.20 a	29.40 ± 0.26 a	79.33 ± 0.33 b	0
	2	2	5.90 ± 0.26 b	21.97 ± 0.26 c	69.67 ± 0.67 d	0

Means within a column followed by the same letter are not significantly different by Duncan's multiple range test ($P < 0.05$); \pm SE.

in contamination of *Erodium sibthorpiatum* seeds and healthy plants were obtained. These results are very much in conformity with other previous studies on various rare, endangered, and endemic plants including *Gentiana cerina* Hook.f., *G. corymbifera* Kirk, *Centaurea tchihatcheffii* Fisch. et Mey., *Linum usitatissimum* L., and *Arabis drabiformis* Boiss. (Morgan et al., 1997; Özel et al., 2006; Yıldız et al., 2010; Akin and Kocaçalışkan, 2011).

For multiplication, many plant species require specific in vitro conditions. A range of cytokinins (kinetin, BAP, 2-iP, and zeatin) have been used in micropropagation

Table 5. Comparison of MS and ModMS medium for seedling growth.

Measured parameters	MS	ModMS
Stem length (mm)	28.93 ± 0.23*	25.60 ± 0.21
Root length (mm)	13.37 ± 0.27	39.17 ± 0.28*
Number of leaves/seedling	11.03 ± 0.20*	5.67 ± 0.35
Number of shoots/seedling	5.67 ± 0.28*	2.77 ± 0.19
Number of roots/seedling	3.40 ± 0.23	7.40 ± 0.21*

* t ($P < 0.05$); \pm SE.

studies (Babaoğlu et al., 2001; Sarasan et al., 2006; Bektaş et al., 2013). If in vitro grown plantlets are to be used as a source of explants for the induction of organogenesis or embryogenesis, well-developed plantlets may also be of significant importance (Monteiro et al., 2000). The nutrient composition of in vitro culture medium may play a decisive role in research of in vitro culturing of new species and development of industrial applications (production of secondary metabolites) and horticulture (micropropagation). The formulation of in vitro culture media can be adjusted to meet the various requirements of plant species (Nas and Read, 2004). In this study, shoot tips were tested for shoot organogenesis. The highest multiplication rate of *Erodium sibthorpiatum* shoot tips was achieved on MS medium with 1 ppm BAP, but the highest shoot length was obtained in PGR-free medium. BAP also significantly induced the auxiliary shoots when it was compared to Ki. Among the cytokinins, the superior effect of BAP over Ki on multiplication was reported by Sudha et al. (1998), Gümüşçü et al. (2008), and Akin and Kocaçalışkan (2011). However, as shown in different studies, high BAP levels caused necrosis in various plant species (Bhatia et al., 2002; Gümüşçü et al., 2008). In our research, MS medium proved to be superior to ModMS medium for shoot multiplication.

Table 6. Effect of different culture media (MS & ModMS) and growth regulators (IBA, IAA & NAA) on rooting of in vitro regenerated shoots after 8 weeks of rooting treatments.

Medium	IBA (ppm)	IAA (ppm)	Number of roots/shoot	Root length (mm)	Rooting (%)
MS	Control		3.10 ± 0.31 e	12.83 ± 0.44 d	32.13 ± 0.41 i
	0.5	0.0	10.57 ± 0.30 a	16.40 ± 0.46 b	84.17 ± 0.60 a
	1.0	0.0	8.97 ± 0.26 b	13.27 ± 0.49 d	77.83 ± 0.38 c
	1.5	0.0	8.57 ± 0.35 b	10.07 ± 0.41 e	72.70 ± 0.35 d
	2.0	0.0	4.37 ± 0.35 d	6.33 ± 0.28 f	65.40 ± 0.61 h
	0.0	0.5	8.47 ± 0.32 b	18.87 ± 0.49 a	80.37 ± 0.69 b
	0.0	1.0	6.30 ± 0.44 c	14.90 ± 0.46 c	70.93 ± 0.47 e
	0.0	1.5	5.47 ± 0.38 c	10.70 ± 0.35 e	69.33 ± 0.35 f
	0.0	2.0	4.00 ± 0.10 de	10.30 ± 0.15 e	67.63 ± 0.47 g
Medium	NAA (ppm)		Number of roots/shoot	Root length (mm)	Rooting (%)
ModMS	Control		7.87 ± 0.20 d	39.17 ± 0.41 b	43.10 ± 0.55 d
	0.1		21.17 ± 0.43 a	44.03 ± 0.58 a	100.00 ± 0.00 a
	0.5		20.80 ± 0.40 ab	40.77 ± 0.43 b	100.00 ± 0.00 a
	1		17.00 ± 0.29 c	22.83 ± 0.52 c	90.03 ± 0.58 c
	2		19.83 ± 0.44 b	23.33 ± 0.85 c	92.17 ± 0.62 b

Means within a column followed by the same letter are not significantly different by Duncan's multiple range test ($P < 0.05$); \pm SE.

In vitro shoot formation may have resulted in different success levels depending upon the explant types used (Zobayed and Saxena, 2003). Our results confirm those of Pevalék-Kozlina et al. (1999), Makowczynska and Andrzejewska-Golec (2003), and Gümüşçü et al. (2008) showing the achievement of shoot induction from shoot tips, which is the best source for the highest shoot induction.

The rooting parameters were changed significantly with different concentrations of IBA. The best rooting was achieved in the MS medium containing 0.5 ppm IBA. IBA is by far the most commonly used auxin to obtain root initiation. Similarly, IBA, as a synthetic auxin, was found to be the most favorable root inducer and was preferred based on the results of prior studies (Mikulík, 1999; Pretto and Santarém, 2000; Prasad et al., 2004; Akbar and Roy, 2006; George et al., 2008; Gümüşçü et al., 2008; Yasodha et al., 2008; Akin and Kocaalışkan, 2011; Okumuş et al., 2011; Chalageri and Babu, 2012). However, when we compared the effects of MS and ModMS media on the rooting of seedlings, ModMS medium was superior to MS medium. The best rooting was obtained when ModMS with 0.1 ppm NAA was added to culture medium. ModMS medium contains a higher level of nitrogen and calcium. Because nitrogen affects the rate of plant growth and is an essential element in the molecular make-up of nucleic

acids, proteins, chlorophyll, amino acids, alkaloids, and some plant hormones, it has a great influence on the rooting of shoot explants in vitro. Additionally, calcium is an integral component of plant cell walls and helps in the formation of pectin, a substance that bonds cell walls together. It also promotes root development (Wu et al., 2005; Sathyanarayana and Varghese, 2007). As a result, rooted plants were transplanted into pots in a greenhouse for further acclimatization studies.

The in vitro techniques reported here offer a powerful tool for mass multiplication of this threatened endemic plant species. The possibilities of plantlet regeneration through tissue culture technology were explored in this study. Several plantlet regeneration protocols were developed for the first time for this species. The regeneration system described here can be used in studies dealing with in vitro conservation of *Erodium sibthorpiatum*. It is concluded that mineral salt of media and PGRs are very important in the success of plant tissue cultures.

Acknowledgments

We gratefully acknowledge the Dumlupınar University Research Fund (Project No: 2010 / 7) for supporting this research financially. We also thank James Innes (Royal Botanic Gardens, Kew, UK) for linguistic and editing help.

References

- Akbar MA, Roy SK (2006). Effects of liquid medium on rooting and acclimation of regenerated microshoots of Banana (*Musa sapientum* L.) cv. Sagar. *Plant Tissue Cult Biotech* 16: 11–18.
- Akın B, Kocaçalışkan İ (2011). In vitro propagation of *Arabis drabiformis* Boiss. (Brassicaceae) an endemic rare species of Uludağ mountain (Bursa-Turkey). *Afr J Biotechnol* 10: 18356–18361.
- Ansar A, Touqeer A, Abbasi NA, Hafız IA (2009). Effect of different media and growth regulators on in vitro shoot proliferation of olive cultivar 'Moraiolo'. *Pak J Bot* 41: 783–795.
- Avilla A de L, Pereyra SM, Argüello JA (1998). Nitrogen concentration and proportion of NH_4^+ -N affect potato cultivar response in solid and liquid media. *HortScience* 33: 336–338.
- Babaoğlu M, Gürel E, Özcan S (2001). *Bitki Biyoteknolojisi I: Doku Kültürü ve Uygulamaları*. Konya, Turkey: Selçuk University Press (in Turkish).
- Bektaş E, Cüce M, Sökmen A (2013). In vitro germination, protocorm formation, and plantlet development of *Orchis coriophora* (Orchidaceae), a naturally growing orchid species in Turkey. *Turk J Bot* 37: 336–342.
- Bhatia P, Bhatia NP, Ashwath N (2002). In vitro propagation of *Stackhousia tryonii* Bailey (Stackhousiaceae): a rare and serpentine-endemic species of central Queensland, Australia. *Biodivers Conserv* 11: 1469–1477.
- Chalageri G, Babu UV (2012). In vitro plant regeneration via petiole callus of *Viola patrinii* and genetic fidelity assessment using RAPD markers. *Turk J Bot* 36: 358–368.
- Ekim T, Koyuncu M, Vural M, Duman H, Aytaç Z, Adıgüzel N (2000). *Türkiye Bitkileri Kırmızı Kitabı (Eğrelti ve Tohumlu Bitkiler)*. Ankara, Turkey: Türkiye Tabiatını Koruma Derneği Yayını (in Turkish).
- Farsi M, Taghavizadeh Yazdi ME, Qasemionran V (2012). Micropropagation of *Anthurium andreaeanum* cv. Terra. *Afr J Biotechnol* 11: 13162–13166.
- Fay MF (1992). Conservation of rare and endangered plants using in vitro methods. *In Vitro Cell Dev B* 28: 1–4.
- George EF, Hall MA, Klerk GJ (2008). Plant growth regulators I: Introduction; auxins, their analogues and inhibitors. In: George EF, editor. *Plant Propagation by Tissue Culture*. 3rd ed. Basingstoke, UK: Exegetics, pp. 175–204.
- Gülyüz G (2000). *Uludağ Alpin Çiçekleri*. Bursa, Turkey: Bursa İl Turizm Müdürlüğü ve Uludağ Turizmini Geliştirme Derneği Yayını (in Turkish).
- Gülyüz G, Kırmızı S, Arslan H, Kondu Yakut E (2011). Alterations of the nitrogen mineralization rates in soils of forest community depending on the ski run constructions (Mount Uludağ, Bursa, Turkey). *J Mt Sci* 8: 53–61.
- Gülyüz G, Malyer H, Kaynak G, Özhatay N (2010). Uludağ A2 (A) Bursa. In: Byfield A, Atay S, Özhatay N, editors. *Türkiye'nin 122 Önemli Bitki Alanı*. İstanbul, Turkey: WWF Türkiye (book chapter in Turkish).
- Gümüşçü A, Çöçü S, Uranbey S, İpek A, Çalışkan M, Arslan N (2008). In vitro micro-propagation of endangered ornamental plant-*Neotchihatchewia isatidea* (Boiss.) Rauschert. *Afr J Biotechnol* 7: 234–238.
- Kapai VY, Kapoor P, Rao IU (2010). In vitro propagation for conservation of rare and threatened plants of India-a review. *Int J Biol Tech* 1: 1–14.
- Ketin İ (1983). *Türkiye Jeolojisine Genel Bir Bakış*. İstanbul, Turkey: İstanbul Teknik Üniversitesi Yayını (in Turkish).
- Kocaçalışkan İ (2008). *Bitki Doku Kültürleri (Organ, Doku ve Hücre)*. Ankara, Turkey: Bizim Büro Basımevi (in Turkish).
- Makowczynska J, Andrzejewska-Golec E (2003). Micropropagation of *Plantago asiatica* L. through culture of shoot-tips. *Acta Soc Bot Pol* 72: 191–194.
- Mikulík J (1999). Propagation of endangered plant species by tissue cultures. *Acta Univ Palack Olomuc Fac Rerum Natur Biol* 37: 27–33.
- Mineo L (1990). Plant tissue culture techniques. In: Goldman CA, editor. *Tested Studies for Laboratory Teaching. Proceedings of the Eleventh Workshop / Conference of the Association for Biology Laboratory Education (ABLE)*, pp. 151–174.
- Monteiro ACB de A, Higashi EN, Gonçalves AN, Rodriguez APM (2000). A novel approach for the definition of the inorganic medium components for micropropagation of yellow passionfruit (*Passiflora edulis* Sims. f. *Flavicarpa* Deg.). *In Vitro Cell Dev-Pl* 36: 527–531.
- Morgan ER, Butler RM, Bicknell RA (1997). In vitro propagation of *Gentiana cerina* and *Gentiana corymbifera*. *New Zeal J Crop Hort* 25: 1–8.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plantarum* 15: 473–497.
- Nas MN, Read PE (2004). A hypothesis for the development of a defined tissue culture medium of higher plants and micropropagation of hazelnuts. *Sci Hortic-Amsterdam* 101: 189–200.
- Okumuş V, Pirinç V, Onay A, Başaran D (2011). In vitro propagation of Diyarbakır watermelons and comparison of direct-seeded and transplanted watermelon. *Turk J Biol* 35: 601–610.
- Omamor IB, Asemota AO, Eke CR, Eziashi EI (2007). Fungal contaminants of the oil palm tissue culture in Nigerian institute for oil palm research (NIFOR). *Afr J Agr Res* 2: 534–537.
- Özel ÇA, Khawar KM, Mirici S, Özcan S, Arslan O (2006). Factors affecting in vitro plant regeneration of critically endangered Mediterranean knapweed (*Centaurea tchihatcheffii* Fisch et. Mey). *Naturwissenschaften* 93: 511–517.
- Pevalek-Kozlina B, Pavlica M, Vujević M (1999). Micropropagation of *Degenia velebitica* (Deg.) Hay. *Phyton-Ann Rei Bota* 39: 293–296.

- Pintar M, Mali B, Kraigher H (2009). The impact of ski slopes management on Kravvec ski resort (Slovenia) on hydrological function of soils. *Biologia* 64: 639–642.
- Prasad PJN, Chakradhar T, Pullaiah T (2004). Micropropagation of *Cryptolepis buchanani* Roem. & Schult. *Taiwania* 49: 57–65.
- Pretto FR, Santarém ER (2000). Callus formation and plant regeneration from *Hypericum perforatum* leaves. *Plant Cell Tiss Org* 62: 107–113.
- Rao NK (2004) Plant genetic resources: advancing conservation and use through biotechnology. *Afr J Biotechnol* 3: 136–145.
- Sarasan V, Cripps R, Ramsay MM, Atherton C, McMichen M, Prendergast G, Rowntree JK (2006). Conservation in vitro of threatened plants-progress in the past decade. *In Vitro Cell Dev-Pl* 42: 206–214.
- Sathyanarayana BN, Varghese Dalia B (2007). *Plant Tissue Culture: Practices and New Experimental Protocols*. New Delhi, India: IK International Publishing House.
- Sudha CG, Krishnan PN, Pushpangadan P (1998). In vitro propagation of *Holostemma annulare* (Roxb.) K. Schum., a rare medicinal plant. *In Vitro Cell Dev-Pl* 34: 57–63.
- Sudharsan C, AboEl-Nil M, Hussain J (2003). Tissue culture technology for the conservation and propagation of certain native plants. *J Arid Environ* 54: 133–147.
- Wu Y, Yi G, Yang H, Zhou B, Zeng J (2005). Basal medium with modified nitrogen source and other factors influence the rooting of banana. *HortScience* 40: 428–430.
- Yasodha R, Kamala S, Anand Kumar SP, Durai Kumar P, Kalaiarasi K (2008). Effect of glucose on in vitro of mature plants of *Bambusa nutans*. *Sci Hortic-Amsterdam* 116: 113–116.
- Yıldız M, Sağlık Ç, Telci C, Erkök EG (2010). The effect of in vitro competition on shoot regeneration from hypocotyl explants of *Linum usitatissimum*. *Turk J Bot* 35: 211–218.
- Zobayed SMA, Saxena PK (2003). In vitro-grown roots: a superior explant for prolific shoot regeneration of St. John's wort (*Hypericum perforatum* L. cv 'New Stem') in a temporary immersion bioreactor. *Plant Sci* 165: 463–470.