

1-1-2016

The determination of in vitro gas production and metabolizable energy value of rice straw treated with exogenous fibrolytic enzymes

ZEHRA SELÇUK

NURCAN ÇETİNKAYA

MUSTAFA SALMAN

BUĞRA GENÇ

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

Recommended Citation

SELÇUK, ZEHRA; ÇETİNKAYA, NURCAN; SALMAN, MUSTAFA; and GENÇ, BUĞRA (2016) "The determination of in vitro gas production and metabolizable energy value of rice straw treated with exogenous fibrolytic enzymes," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 40: No. 6, Article 5. <https://doi.org/10.3906/vet-1601-113>

Available at: <https://journals.tubitak.gov.tr/veterinary/vol40/iss6/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The determination of in vitro gas production and metabolizable energy value of rice straw treated with exogenous fibrolytic enzymes

Zehra SELÇUK^{1*}, Nurcan ÇETİNKAYA¹, Mustafa SALMAN¹, Buğra GENÇ²

¹Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine, Ondokuz Mayıs University, Kurupelit Campus, Atakum, Samsun, Turkey

²Department of Laboratory Animals, Faculty of Veterinary Medicine, Ondokuz Mayıs University, Kurupelit Campus, Atakum, Samsun, Turkey

Received: 29.01.2016 • Accepted/Published Online: 09.05.2016 • Final Version: 15.12.2016

Abstract: This study was conducted to determine the effects of cellulase and xylanase treatments on in vitro 24-h gas production (GP), organic matter digestibility (OMD), and metabolizable energy (ME) values of rice straw (RS). Rice straws were treated with cellulase and xylanase at the levels of 0% (C1), 0.5% (RS+CEL0.5), 1% (RS+CEL1), 1.5% (RS+CEL1.5), and 2% (RS+CEL2) and 0% (C2), 0.5% (RS+XYL0.5), 1% (RS+XYL1), 1.5% (RS+XYL1.5), and 2% (RS+XYL2), respectively. Rice straws treated with each level of cellulase and xylanase were ensiled in 6 glass jars. For each level of cellulase and xylanase treatment, 3 glass jars were incubated at room temperature, while 3 glass jars were incubated at 40 ± 0.2 °C in an incubator for 30 days. In vitro GP for 24 h, OMD, and ME values in RS+CEL2 were higher ($P < 0.05$) than those of C1 and the other groups treated with cellulase, while significant increases ($P < 0.05$) for the same parameters were obtained from RS+XYL1 incubated at 40 ± 0.2 °C, RS+XYL1.5, and RS+XYL2 compared to C2, RS+XYL0.5, and RS+XYL1 incubated at room temperature. In conclusion, in vitro 24-h GP, OMD, and ME values were the highest ($P < 0.05$) in the groups treated with the highest level of cellulase and xylanase incubated at 40 ± 0.2 °C.

Key words: Fibrolytic enzyme, in vitro gas production, metabolizable energy, rice straw

1. Introduction

Rice straw (RS) is a crop residue of rice production and can be used as a feed for ruminants because it is easily and cheaply available in many Asian countries. Kadam et al. (1) stated that about 1.35 metric tons of RS are left for each metric ton of harvested rice. Rice straw, such as other cereal straws, does not provide enough nutrients to meet the requirement of ruminants due to its high lignocellulosic structure, relatively low digestibility, and poor palatability (2). Average dry matter (DM), organic matter (OM), crude protein (CP), neutral detergent fiber (NDF), acid detergent fiber (ADF), and lignin contents of different varieties of RS range from 92.21% to 93.05%, 81.21% to 86.24%, 3.49% to 5.10%, 72.16% to 77.57%, 41.38% to 46.32%, and 4.3% to 6.97%, respectively (3).

Low quality forages (straws) can be treated with different physical methods (grinding, pelleting/chopping for reducing particle size, soaking, steam treatments, cooking under pressure, etc.) to improve their feeding value. Likewise, chemical treatment of low-quality forages improves their degradability (4); however, the chemicals may result in environmental pollution. Thus, the interest

in treatment of straws is focused on biological methods owing to the ongoing concerns about pollution and food safety. Exogenous enzymes obtained from fungi and bacteria can be used in biological treatment to improve the digestibility of low-quality forages due to being able to break down lignocellulosic complexes (5). Improving the availability of the nutrients in the cell wall structure of fibrous feeds can be possible by using exogenous fibrolytic enzymes. These enzymes can be directly added to feeds before feeding or during ensiling (6). It was reported that fibrolytic enzymes have effects on the degradation of cereal straw cell wall and improvement of the nutritive value of cereal straws (7,8).

The aim of the present study was to determine the effects of different levels of cellulase and xylanase treatments of RS on in vitro 24-h gas production (GP), organic matter digestibility (OMD), and metabolizable energy (ME) values.

2. Materials and methods

Chemical analyses and in vitro gas production techniques were conducted in the Ruminant Feed Evaluation

* Correspondence: zselcuk@omu.edu.tr

Laboratory of the Department of Animal Nutrition and Nutritional Diseases, Faculty of Veterinary Medicine of Ondokuz Mayıs University, Samsun, Turkey.

2.1. Enzyme treatments

Rice straw was chopped to 5 cm in length and treated with the cellulase (from *Trichoderma reesei* with activity ≥ 5000 U/kg; 1 unit of cellulase activity is described as the enzyme required to release 1 μg of reducing sugar from 4 mg/mL sodium cellulose glycolate in 1 min at 37 °C and pH 5.5) and xylanase (from *Bacillus subtilis* with activity $\geq 100,000$ U/kg; 1 unit of xylanase activity is described as the enzyme required to release 1 μmol xylan from xylan solution in 1 min at 37 °C and pH 5.5) at the levels of 0% (C1), 0.5% (RS+CEL0.5), 1% (RS+CEL1), 1.5% (RS+CEL1.5), and 2.0% (RS+CEL2) and 0% (C2), 0.5% (RS+XYL0.5), 1% (RS+XYL1), 1.5% (RS+XYL1.5), and 2.0% (RS+XYL2) of DM basis of RS according to the method reported by Nakashima et al. (9), respectively. The amount of water added to RS for cellulase and xylanase treatments was calculated to provide approximately 40% DM of RS. For ensuring homogeneous distribution of cellulase and xylanase enzymes in RS, the enzymes were dissolved in water and then were immediately sprayed on RS and mixed for absorption by the RS. After mixing, 6 glass jars with a volume of 1 L were filled completely with RS treated with cellulase and xylanase, for each level of cellulase and xylanase treatment. The perimeters of the jar lids were coated with silicone to prevent air intake. For each level of cellulase and xylanase treatments, 3 glass jars were incubated at room temperature (22 °C) in the dark for 30 days and 3 glass jars were incubated at 40 ± 0.2 °C in an incubator for 30 days. After incubation of the jars containing RS treated with cellulase and xylanase, all jars were opened and dried at 100 ± 0.2 °C in an incubator to stop the enzyme activities. Dried RS samples were ground for the in vitro gas production technique and chemical analysis.

2.2. Ruminant fluid collection

Ruminal fluid was obtained from different sites within the rumen of three ruminally cannulated Karayaka rams (average weight 50 ± 5 kg; 2 years old) fed twice daily at 0800 and 1700 hours with a diet containing 650 g of alfalfa hay and 350 g of concentrate estimated to the level of 1.25 \times maintenance requirements according to the NRC (10). Fresh drinking water was freely available. Ruminal fluid was collected at 3 h after the morning feeding in thermos flasks and transported directly to the laboratory. Ruminal fluid was strained through four layers of cheesecloth and held at 39 °C under CO₂ atmosphere.

2.3. Chemical analysis

Rice straw samples were milled for passing through a 1 mm sieve and then DM, ash, ether extract (EE), and CP of RS samples were analyzed according to AOAC methods

(11). ADF, NDF (with sodium sulfite), and ADL contents of RS samples were determined by the method of Van Soest et al. (12).

2.4. In vitro gas production

In vitro gas production was measured using the ANKOM^{RF} gas production system (13). Each unit of the system consisted of a 250 mL glass bottle equipped with an attached module. There was wireless communication between the attached module and computer. When the pressure inside of each unit reached 1.5 kPa above ambient pressure, a valve of the attached module automatically released the gas accumulated in the headspace of the glass bottles. Pressure inside of the glass bottles was measured every 10 min. The buffer solution was prepared according to Menke and Steingass (13), preheated, and mixed with rumen fluid at 4:1, and then 100 mL of buffered rumen fluid was added to each preheated glass bottle containing 1 g of the milled feed samples. The incubation media preparation was carried out under anaerobic conditions purged with CO₂, at approximately 39 °C and pH of about 6.5–6.8. After filling, all glass bottles were closed and placed into an incubator at 39.0 ± 0.5 °C. The incubation procedure was conducted in triplicate for each treatment group. All glass bottles containing incubation medium and feed samples were incubated for 0, 3, 6, 12, 24, 48, 72, and 96 h. The average cumulative pressure measured for each incubation time of the feed samples and ME (MJ/kg DM) and OMD (%) values of RS samples were estimated from the measured pressure by in vitro gas production method at 24 h by using the following equations (14).

$$\text{ME (MJ/kg DM)} = 2.20 + 0.1136\text{GP} + 0.0057\text{CP} + 0.00029\text{EE}^2$$

$$\text{OMD (\%)} = 57.2 + 0.365\text{GP} + 0.304\text{CP} - 1.98\text{ADL}$$

2.5. Statistical analysis

Data were designed for appropriate one-way classification with a 2 \times 5 factorial experimental design (full factorial design model) and analyzed with the GLM procedure. Data were fitted to the statistical model with a full factorial design including main effects and all possible interaction effects of the factor combinations (enzyme treatments), temperature, and levels of the traits due to selecting for the reducing model. Therefore, a mathematical model including three main effects and only traits \times levels interaction effects was fitted to data due to the fact that the other interaction effects were insignificant. Main effects were compared with Duncan's multiple range test. Mean differences of interaction effects were compared by Tukey test. All analyses and calculations were performed with SAS (15).

3. Results

Dry matter, ash, CP, ADF, NDF, and ADL contents of rice straw used in the study were 93.53%, 16.68%, 4.15%,

41.15%, 65.72%, and 7.85%, respectively. Cumulative GP for 24 h, OMD%, and ME (MJ/kg DM) values of RS untreated and treated with different levels of cellulase and xylanase enzymes and incubated at room temperature (22 °C) and 40 ± 0.2 °C for 30 days are given in Tables 1 and 2, respectively. Least square of means and standard errors of means for interaction effects on the traits × levels are presented in Table 3. There were differences ($P < 0.05$) in cumulative GP for 24 h, OMD%, and ME (MJ/kg DM) values of groups treated with the increasing level of cellulase enzyme and incubated at 40 ± 0.2 °C for 30 days (Table 1). Cumulative GP for 24 h, OMD%, and ME (MJ/kg DM) values of RS+XYL1, RS+XYL1.5, and RS+XYL2 incubated at 40 ± 0.2 °C were higher ($P < 0.05$) than those of C2 and RS+XYL0.5 incubated at the same incubation temperature (Tables 2 and 3).

Cumulative gas production values for 3, 6, 12, 24, 48, 72, and 96 h of RS untreated and treated with different levels of cellulase and xylanase enzymes and incubated at room temperature (22 °C) and 40 ± 0.2 °C for 30 days are given in Table 4.

4. Discussion

Plant cell wall components are broken down with fibrolytic enzymes such as cellulase and xylanase produced by microorganisms in the reticulorumen of ruminants. However, many forages like cereal straws are of low quality due to poor digestibility and limited energy value for ruminants (16) because of insufficient degradation of the cell wall in the rumen. The use of exogenous enzymes

may be an option to increase the degradability of cell wall fraction (17). Fibrolytic enzymes as feed additives, added in liquid or granular form to diets, hay, silages, etc. to increase the availability of nutrients in the plant cell wall may be obtained from microbial fermentation of bacteria or fungi, and they show specific activity. Exogenous fibrolytic enzymes applied both during feed ensiling and directly during feeding increased in vitro GP and improved the availability of nutrients of fibrous feeds. These supplementations may allow the attachment of microorganisms to feeds (18) and/or the formation of enzyme–feed complexes (19). Beauchemin et al. (16) stated that exogenous enzyme treatments in a liquid form of dried feeds were important for adsorption of enzymes to provide suitable attachment to feeds before consumption. In the present study, 24-h GP, OMD%, and ME values of the RS+CEL1, RS+CEL1.5, RS+CEL2, RS+XYL1.5, and RS+XYL2 groups treated with cellulase and xylanase dissolved in water and sprayed on rice straw and incubated at room temperature (22 °C) for 30 days improved compared to those of controls and other treatment groups, while the same parameters increased in RS+CEL0.5, RS+CEL1, RS+CEL1.5, RS+CEL2, RS+XYL1, RS+XYL1.5, and RS+XYL2 incubated at 40 ± 0.2 °C for 30 days compared to those of controls.

Kumar et al. (20) stated that some commercial fibrolytic enzymes had effects on increasing of total GP of feeds. Akinfemi and Ogunwole (21) reported that 24-h GP, ME, and OMD values increased in RS treated with different mushroom species. In the present study, treatments of RS

Table 1. Cumulative gas production for 24 h, the percentage of OMD, and ME values of rice straw untreated and treated with cellulase and incubated at room temperature and 40 °C for 30 days.

Treatments	Incubations	GP, mL Mean ± SE	OMD% Mean ± SE	ME, MJ/kg DM Mean ± SE
C1	22 °C	28.32 ± 0.48 ^f	53.25 ± 0.17 ^f	6.29 ± 0.06 ^f
	40 °C	28.60 ± 0.31 ^f	53.36 ± 0.31 ^f	6.33 ± 0.04 ^f
RS+CEL0.5	22 °C	29.32 ± 0.28 ^f	53.62 ± 0.10 ^f	6.42 ± 0.03 ^f
	40 °C	30.63 ± 0.43 ^e	54.09 ± 0.15 ^e	6.60 ± 0.05 ^e
RS+CEL1	22 °C	31.53 ± 0.48 ^{de}	54.42 ± 0.17 ^{de}	6.72 ± 0.06 ^{de}
	40 °C	32.11 ± 0.08 ^{cd}	54.63 ± 0.03 ^{cd}	6.80 ± 0.01 ^{cd}
RS+CEL1.5	22 °C	31.73 ± 0.21 ^{cd}	54.50 ± 0.07 ^{cd}	6.75 ± 0.02 ^{cd}
	40 °C	32.68 ± 0.29 ^{bc}	54.84 ± 0.10 ^{bc}	6.88 ± 0.03 ^{bc}
RS+CEL2	22 °C	33.20 ± 0.20 ^b	55.03 ± 0.07 ^b	6.95 ± 0.02 ^b
	40 °C	34.56 ± 0.29 ^a	55.53 ± 0.10 ^a	7.14 ± 0.04 ^{ba}

Different letters within the same column indicate differences among groups ($P < 0.05$).

GP: Gas production; OMD: organic matter digestibility; ME: metabolizable energy; DM: dry matter; C1: control; RS+CEL0.5: rice straw+cellulase 0.5%; RS+CEL1: rice straw+cellulase 1%; RS+CEL1.5: rice straw+cellulase 1.5%; RS+CEL2: rice straw+cellulase 2%.

Table 2. Cumulative gas production for 24 h, the percentage of OMD, and ME values of rice straw untreated and treated with xylanase and incubated at room temperature and 40 °C for 30 days.

Treatments	Incubations	GP, mL Mean ± SE	OMD% Mean ± SE	ME, MJ/kg DM Mean ± SE
C2	22 °C	28.32 ± 0.48 ^d	53.25 ± 0.17 ^d	6.29 ± 0.06 ^d
	40 °C	28.60 ± 0.31 ^d	53.36 ± 0.31 ^d	6.33 ± 0.04 ^d
RS+XYL0.5	22 °C	28.85 ± 0.31 ^d	53.44 ± 0.11 ^d	6.36 ± 0.04 ^d
	40 °C	29.00 ± 0.20 ^d	53.50 ± 0.07 ^d	6.38 ± 0.02 ^d
RS+XYL1	22 °C	29.29 ± 0.45 ^d	53.60 ± 0.16 ^d	6.42 ± 0.06 ^d
	40 °C	30.69 ± 0.35 ^c	54.12 ± 0.13 ^c	6.61 ± 0.04 ^c
RS+XYL1.5	22 °C	31.39 ± 0.12 ^c	54.37 ± 0.04 ^c	6.71 ± 0.01 ^c
	40 °C	33.04 ± 0.26 ^b	54.97 ± 0.09 ^b	6.93 ± 0.03 ^b
RS+XYL2	22 °C	33.65 ± 0.25 ^{ab}	55.20 ± 0.09 ^{ab}	7.01 ± 0.03 ^{ab}
	40 °C	34.18 ± 0.16 ^a	55.39 ± 0.06 ^a	7.08 ± 0.02 ^a

Different letters within the same column indicate differences among groups ($P < 0.05$).

GP: Gas production; OMD: organic matter digestibility; ME: metabolizable energy; DM: dry matter; C2: control, RS+XYL0.5: rice straw+xylanase 0.5%; RS+XYL1: rice straw+xylanase 1%; RS+XYL1.5: rice straw+xylanase 1.5%; RS+XYL2: rice straw+xylanase 2%.

Table 3. Least square of means and standard error of means for interaction effects on traits × levels.

Factors	Levels	Traits		
		GP 24-h, mL	OMD, %	ME, MJ/kg DM
C	0	28.46 ± 0.23a	53.30 ± 0.07a	6.31 ± 0.031a
RS+CEL	0.5	29.97 ± 0.23b	53.86 ± 0.10a	6.51 ± 0.031b
	1	31.82 ± 0.23c	54.53 ± 0.10b	6.76 ± 0.031b
	1.5	32.21 ± 0.23c	54.67 ± 0.10b	6.82 ± 0.031b
	2.0	33.88 ± 0.23d	55.28 ± 0.10c	7.04 ± 0.031d
RS+XYL	0.5	28.92 ± 0.23a	53.47 ± 0.10a	6.37 ± 0.031a
	1	29.99 ± 0.23b	53.87 ± 0.10a	6.51 ± 0.031b
	1.5	32.21 ± 0.23c	54.67 ± 0.10b	6.82 ± 0.031c
	2.0	33.91 ± 0.23d	55.29 ± 0.10bc	7.05 ± 0.031d
P-value		0.0003	0.006	0.0003

GP 24-h: Gas production for 24 h; OMD: organic matter digestibility; ME: metabolizable energy; DM: dry matter; C: control, RS+CEL: rice straw+cellulase; RS+XYL: rice straw+xylanase.

with increasing levels of cellulase significantly increased 24-h GP, ME, and OMD of the RS+CEL1, RS+CEL1.5 and RS+CEL2 groups incubated at two different incubation temperatures for 30 days compared to those of control and RS+CEL0.5 groups incubated at room temperature. Similar increases in the same parameters were obtained from RS+XYL1 incubated at 40 ± 0.2 °C and RS+XYL1.5 and RS+XYL2 groups incubated at two different temperatures for 30 days.

Tang et al. (22) reported that fibrolytic enzyme supplementations improved the in vitro OMD of RS. OMDs of treated rice straws varied from 53.62 ± 0.10% to 55.53 ± 0.10% for cellulase treatments and from 53.44 ± 0.11% to 55.39 ± 0.06% for xylanase treatments in the present study.

Although gas production, basically the result of rumen fermentation, seems like a loss of energy in terms of nutrition, it has been used to predict OMD and ME

Table 4. Cumulative gas production values (mean \pm SE) of rice straw untreated and treated with cellulase or xylanase.

Cumulative gas production, mL									
Treatments		Levels, %	3 h	6 h	12 h	24 h	48 h	72 h	96 h
C	22 °C	0	3.42 \pm 0.14	7.76 \pm 0.22	11.39 \pm 0.14	28.32 \pm 0.48	31.99 \pm 0.35	39.37 \pm 0.2	41.64 \pm 0.10
	40 °C	0	3.38 \pm 0.14	7.84 \pm 0.10	11.68 \pm 0.21	28.60 \pm 0.31	32.11 \pm 0.43	39.54 \pm 0.21	42.16 \pm 0.10
RS+CEL	22 °C	0.5	3.21 \pm 0.07	7.92 \pm 0.07	11.51 \pm 0.14	29.32 \pm 0.28	33.35 \pm 0.10	39.62 \pm 0.21	41.51 \pm 0.25
		1.0	3.59 \pm 0.07	11.59 \pm 0.27	25.38 \pm 0.53	31.53 \pm 0.48	35.70 \pm 0.25	43.09 \pm 0.18	44.87 \pm 0.17
		1.5	4.76 \pm 0.09	13.99 \pm 0.14	28.81 \pm 0.17	31.73 \pm 0.21	37.23 \pm 0.14	43.09 \pm 0.07	45.52 \pm 0.22
		2.0	5.65 \pm 0.10	14.86 \pm 0.28	27.24 \pm 0.58	33.20 \pm 0.20	43.09 \pm 0.18	50.60 \pm 0.50	53.04 \pm 0.21
	40 °C	0.5	3.46 \pm 0.14	8.50 \pm 0.17	11.88 \pm 0.14	30.63 \pm 0.43	33.18 \pm 0.37	39.70 \pm 0.17	42.39 \pm 0.10
		1.0	4.25 \pm 0.10	13.08 \pm 0.25	27.73 \pm 0.35	32.11 \pm 0.08	36.86 \pm 0.14	43.83 \pm 0.14	45.68 \pm 0.25
		1.5	5.40 \pm 0.10	14.98 \pm 0.21	30.15 \pm 0.10	32.68 \pm 0.29	39.03 \pm 0.14	44.95 \pm 0.32	47.01 \pm 0.10
		2.0	6.48 \pm 0.21	16.88 \pm 0.10	28.77 \pm 0.32	34.56 \pm 0.29	44.16 \pm 0.14	52.42 \pm 0.17	54.85 \pm 0.21
RS+XYL	22 °C	0.5	3.62 \pm 0.21	9.12 \pm 0.10	14.73 \pm 0.14	28.85 \pm 0.31	32.27 \pm 0.32	39.04 \pm 0.35	41.25 \pm 0.18
		1.0	3.54 \pm 0.17	11.66 \pm 0.18	23.52 \pm 0.28	29.29 \pm 0.45	32.77 \pm 0.21	40.03 \pm 0.17	41.89 \pm 0.28
		1.5	4.62 \pm 0.14	13.12 \pm 0.28	28.85 \pm 0.25	31.39 \pm 0.12	35.12 \pm 0.17	41.15 \pm 0.10	43.09 \pm 0.14
		2.0	5.64 \pm 0.17	13.47 \pm 0.16	25.26 \pm 0.32	33.65 \pm 0.25	42.47 \pm 0.28	51.39 \pm 0.32	53.24 \pm 0.14
	40 °C	0.5	4.08 \pm 0.13	9.32 \pm 0.10	15.14 \pm 0.14	29.00 \pm 0.20	33.22 \pm 0.14	40.08 \pm 0.10	42.23 \pm 0.14
		1.0	4.21 \pm 0.14	12.49 \pm 0.20	24.51 \pm 0.21	30.69 \pm 0.35	33.53 \pm 0.07	40.47 \pm 0.16	42.59 \pm 0.07
		1.5	5.77 \pm 0.17	14.69 \pm 0.17	29.88 \pm 0.21	33.04 \pm 0.27	35.87 \pm 0.10	41.89 \pm 0.21	43.87 \pm 0.14
		2.0	6.76 \pm 0.17	14.90 \pm 0.28	27.03 \pm 0.14	34.18 \pm 0.16	44.45 \pm 0.25	53.08 \pm 0.21	55.14 \pm 0.14

C: Control; RS+CEL; rice straw+cellulase; RS+XYL: rice straw+xylanase.

values of feeds. Therefore, the increase in the amount of gas production due to the increase in the activity of rumen microorganisms causes an increase in both OMD and ME values. In this study, digestible organic matter and ME values of RS+CEL2 and RS+XYL2 were higher than those of control. Sommart et al. (23) and Nitipot and Sommart (24) stated that there was a positive correlation between in vitro OMD and the volume of gas released during fermentation.

Jalc (5) reported that usage of exogenous enzymes obtained from fungi, white rot fungi, and mushroom has a potential effect on improving the digestibility of low-quality forages due to being able to break down lignocellulosic complex. Some toxic metabolites may be produced by the fungi during the treatment for increasing feeding value of rice straw. However, treatment of the straws with fungi is difficult in terms of ensuring optimal conditions (pH, temperature, CO₂ concentration, etc.) for fungal growth. Hence, commercial products will be important in the future for ruminant nutrition (16). Fazaeli et al. (25) and

Rodrigues et al. (26) reported that enzyme treatment or its combination with other treatments has effects on increasing the digestibility of cereal straws for ruminants.

It was stated that treatment with cellulase (27) and cellulase and xylanase (28) improved the digestibility of RS. In the present study, treatments of rice straw with increasing levels of cellulase and xylanase significantly improved OMD compared to untreated rice straw. This result was consistent with the statements of some other researchers (25,26,28). OMD was the highest in RS+CEL2 and RS+XYL incubated at 40 \pm 0.2 °C for 30 days. This result can be attributed to the cellulase and xylanase levels and their activity resulting in changes in cell wall structure in incubation.

In conclusion, cellulase and xylanase treatments of rice straw had higher 24-h GP, OMD, and ME values compared to untreated rice straw. The best result was obtained with the highest level of cellulase (RS+CEL2) and xylanase (RS+XYL2) treatments at 40 \pm 0.2 °C when in vitro 24-h GP, OMD, and ME values were considered.

Although fibrolytic enzyme treatments improved the 24-h GP, OMD, and ME values, further low-quality forage treatment experiments with exogenous enzymes alone and in combinations are required to improve digestibility. The findings of in vitro studies should also be supported by the results of in vivo studies.

References

- Kadam KL, Forrest LH, Jacobson WA. Rice straw as a lignocellulosic resource: collection, processing, transportation, and environmental aspects. *Biomass Bioenergy* 2000; 18: 369-389.
- Widyastuti Y, Terada F, Kajikawa H, Abe A. Digestion of rice straw cell wall constituents in various rumen conditions. *Jpn Agric Res Quart* 1987; 21: 59-64.
- Rahman MM, Alam MR, Amin MR, Das NG. Comparative study of the nutritive values of the different varieties of rice straw. *Bangladesh Journal of Animal Science* 2010; 39: 75-82.
- Vadiveloo J. Nutritional properties of the leaf and stem of rice straw. *Anim Feed Sci Technol* 2000; 83: 57-65.
- Jalc D. Straw enrichment for fodder production by fungi. In: Kempken F, editor. *The Mycota. XI. Agricultural Applications*. Berlin, Germany: Springer-Verlag; 2002. pp. 19-38.
- Khattab HM, Gado HM, Kholif AE, Mansour AM, Kholif AM. The potential of feeding goats sun dried rumen contents with or without bacterial inoculums as replacement for berseem clover and the effects on milk production and animal health. *Int J Dairy Sci* 2011; 6: 267-277.
- Yang HE, Son YS, Beauchemin KA. Effects of exogenous enzymes on ruminal fermentation and degradability of alfalfa hay and rice straw. *Asian-Aust J Anim Sci* 2011; 24: 56-64.
- Wang Y, Ramirez-Bribiesca JE, Yanke LJ, Tsang A, McAllister TA. Effect of exogenous fibrolytic enzyme application on the microbial attachment and digestion of barley straw in vitro. *Asian-Aust J Anim Sci* 2012; 25: 66-74.
- Nakashima Y, Ørskov ER, Adebawale EA, Ambo K. Enzymatic manipulation of straw quality: experience on straw upgrading. In: *Proceedings of International Conference on Increasing Livestock Production through Utilization of Local Resources*; Beijing, China; 1993. pp. 139-152.
- NRC. *Nutrient Requirements of Sheep*. 6th ed. Washington, DC, USA: National Academy Press; 1985.
- AOAC. *Official Methods of Analysis*. 18th ed. Washington, DC, USA: Association of Official Analytical Chemists; 2006.
- Van Soest PJ, Robertson JD, Lewis BA. Methods for dietary fibre, neutral detergent fibre and non-starch polysaccharides in relation to animal nutrition. *J Dairy Sci* 1991; 74: 3583-3597.
- ANKOM. *ANKOM^{RF} Gas Production System Operator's Manual*. Macedon, NY, USA: ANKOM; 2011.
- Menke KH, Steingass H. Estimation of the energetic feed value obtained from chemical analysis and in vitro gas production using rumen fluid. *Anim Res Dev* 1988; 28: 7-55.
- SAS. *SAS Statistic Software*. Cary, NC, USA: SAS; 2007.
- Beauchemin KA, Colombatto D, Morgavi DP. A rationale for the development of feed enzyme products for ruminants. *Can J Anim Sci* 2004; 84: 23-36.
- Avellaneda JH, Pinos-Rodríguez JM, González SS, Bárcena R, Hernández A, Cobos M, Hernández D, Montañez O. Effects of exogenous fibrolytic enzymes on ruminal fermentation and digestion of Guinea grass hay. *Anim Feed Sci Technol* 2009; 149: 70-77.
- Nsereko VL, Beauchemin KA, Morgavi DP, Rode LM, Furtado AF, McAllister TA, Iwaasa AD, Yang WZ, Wang Y. Effect of a fibrolytic enzyme preparation from *Trichoderma longibrachiatum* on the rumen microbial population of dairy cows. *Can J Microbiol* 2002; 48: 14-20.
- Kung L Jr, Treacher RJ, Nauman GA, Smagala AM, Endres KM, Cohen MA. The effect of treating forages with fibrolytic enzymes on its nutritive value and lactation performance of dairy cows. *J Dairy Sci* 2000; 83: 115-122.
- Kumar S, Dagar SS, Sirohi SK, Upadhyay RC, Puniya AK. Microbial profiles, in vitro gas production and dry matter digestibility based on various ratios of roughage to concentrate. *Ann Microbiol* 2013; 63: 541-545.
- Akinfemi A, Ogunwale OA. Chemical composition and in vitro digestibility of rice straw treated with *Pleurotus ostreatus*, *Pleurotus pulmonarius* and *Pleurotus tuber-regium*. *Slovak J Anim Sci* 2012; 45: 14-20.
- Tang SX, Tayo GO, Tan ZL, Sun ZH, Shen LX, Zhou CS, Xiao WJ, Ren GP, Han XF, Shen SB. Effects of yeast culture and fibrolytic enzyme supplementation on in vitro fermentation characteristics of low-quality cereal straws. *J Anim Sci* 2008; 86: 1164-1172.
- Sommart K, Parker DS, Rowlinson P, Wanapat M. Fermentation characteristics and microbial protein synthesis in an in vitro system using cassava, rice straw and dried ruzi grass as substrate. *Asian-Austral J Anim Sci* 2000; 13: 1084-1093.
- Nitipot P, Sommart K. Evaluation of ruminant nutritive value of cassava starch industry by using in vitro gas production technique. In: *Proceedings of Annual Agricultural Seminar for Year 2003*; Khon Kaen, Thailand; 2003. pp. 179-190.
- Fazaeli H, Azizi A, Amile M. Nutritive value index of treated wheat straw with *Pleurotus* fungi fed to sheep. *Pak J Biol Sci* 2006; 9: 2444-2449.

Acknowledgments

This study was supported by the Ondokuz Mayıs University Research Fund (Project number: PYO.VET.1901.12.009). The authors would like to thank FARMAVET International and Kurtbey Türk for supplying fibrolytic enzymes and rice straw, respectively.

26. Rodrigues MAM, Pinto P, Bezerra RMF, Dias AA, Guedes CVM, Gardoso VMG, Cone JW, Ferreira LMM, Colaco J, Sequeira CA. Effect of enzyme extracts isolated from white-rot fungi on chemical composition and in vitro digestibility of wheat straw. *Anim Feed Sci Technol* 2008; 141: 326-338.
27. Liu JX, Ørskov ER. Cellulase treatment of untreated and steam pre-treated rice straw-effect on in vitro fermentation characteristics. *Anim Feed Sci Technol* 2000; 88: 189-200.
28. Eun JS, Beauchemin KA, Hong SH, Bauer MW. Exogenous enzymes added to untreated or ammoniated rice straw: effects on in vitro fermentation characteristics and degradability. *Anim Feed Sci Technol* 2006; 131: 86-101.