

1-1-2018

Two asymptotic results of solutions for nabla fractional (q,h) -difference equations

FEIFEI DU

LYNN ERBE

BAOGUO JIA

ALLAN PETERSON

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

DU, FEIFEI; ERBE, LYNN; JIA, BAOGUO; and PETERSON, ALLAN (2018) "Two asymptotic results of solutions for nabla fractional (q,h) -difference equations," *Turkish Journal of Mathematics*: Vol. 42: No. 5, Article 11. <https://doi.org/10.3906/mat-1802-49>

Available at: <https://journals.tubitak.gov.tr/math/vol42/iss5/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Two asymptotic results of solutions for nabla fractional (q, h) -difference equations

Feifei DU^{1,*}, Lynn ERBE², Baoguo JIA¹, Allan PETERSON²

¹School of Mathematics, Sun Yat-Sen University, Guangzhou, P.R. China

²Department of Mathematics, University of Nebraska-Lincoln, Lincoln, Nebraska, USA

Received: 13.02.2018

Accepted/Published Online: 09.06.2018

Final Version: 27.09.2018

Abstract: In this paper we study the Caputo and Riemann–Liouville nabla (q, h) -fractional difference equation and obtain the following two main results:

Theorem A Assume $0 < \alpha < 1$ and there is a constant b such that $c(t) \leq b < 0$, for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. Then any solution, $x(t)$, of the nabla Caputo (q, h) -fractional difference equation

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \quad (0.1)$$

with $x(a) > 0$ satisfies

$$\lim_{t \rightarrow \infty} x(t) = 0.$$

Theorem B Assume $0 < \alpha < 1$, $c(t) \leq 0$, $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$, and $x(t)$ is a solution of the equation

$${}_a \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}, \quad (0.2)$$

satisfying $x(\sigma(a)) > 0$. Then $x(t) > 0$, $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$ and

$$\lim_{t \rightarrow \infty} x(t) = 0.$$

Theorem A and Theorem B extend the results in other recent works of the authors.

Key words: Nabla fractional difference, (q, h) -calculus, monotonicity, asymptotic behavior

1. Introduction

In recent years, fractional calculus has attracted increasing interest. Although several results of fractional differential/difference equations are already published [1–6, 13–16, 22, 23, 25–30, 32, 33, 35, 40–44, 46, 47], the development of a qualitative theory for fractional difference equations is still in its beginning due to the memory effects of fractional operators.

*Correspondence: qinjin65@126.com

2010 AMS Mathematics Subject Classification: 39A12, 39A70

This work was supported by the National Natural Science Foundation of China (No. 11271380) and the Guangdong Province Key Laboratory of Computational Science.

The extension of the basic notions of discrete fractional calculus to the (q, h) -calculus setting appear in [10, 12]. The (q, h) -calculus reduces to discrete nabla-calculus (see [18, Chapter 3]) when $q = 1 = h$ and q -difference calculus (quantum-calculus) (see [18, Chapter 4]) when $h = 0$. Some interesting results concerning fractional (q, h) -calculus can be found in [36–38].

In 1996, Matignon [34] gave a classical stability condition for homogeneous linear fractional differential equations. This criterion was developed by several authors. In [31], Chen et al. proposed a fractional Lyapunov direct method. Later, Chen et al. [13] used the Schauder fixed point theorem in the asymptotic stability of nonlinear fractional difference equations. In 2013, Cermak et al. [11] investigated the stability regions for linear fractional difference systems using Laplace transform. In 2015, Jia et al. [25] established comparison theorems to extend the corresponding asymptotic result in [4]. In 2016, Mozyrska et al. [35] presented linearization to decide the stability of fractional difference systems. In 2017, Jia et al. [20] constructed a Liapunov functional to decide the stability of fractional (q, h) -difference equations.

Some efforts have been made in stability of nabla fractional q -difference equations [19, 25] and nabla fractional h -difference equations (see [11, 21], [39, Chapter 6]). Other efforts have been made in stability of delta fractional difference equations ($h = 1$) [9]. However, [20] is the only result on the asymptotic behavior of nabla fractional (q, h) -difference equations. The formal solutions (when $c(t) = \lambda$ is a constant) of (0.1) and (0.2) are given through (q, h) -Mittag-Leffler function (see details in [10, 38]),

$$E_{\alpha, \beta}^{s, \lambda}(t) = \sum_{k=0}^{\infty} \lambda^k \hat{h}_{\alpha k + \beta - 1}(t, s),$$

which converges (absolutely) if $|\lambda| < (\nu(t))^{-\alpha}$, which means λ only can be zero when $q > 1$ and $h > 0$ (since $\lim_{t \rightarrow \infty} \nu(t) = \infty, t \in \mathbb{T}_{(q, h)}^{t_0}$). This illustrates that there are substantial differences between the h -time scale and q -time scale (or (q, h) -time scale). Thus, we cannot directly use (q, h) -Mittag-Leffler function to study their asymptotic behavior.

The aim of this paper is to further investigate asymptotic stability of the nabla fractional (q, h) -difference equations. Theorem A is motivated by Jia et al. [23], who obtained asymptotic results for the nabla Caputo fractional difference equation

$$\begin{aligned} \nabla_{a*}^{\nu} x(t) &= c(t)x(t), \quad t \in \mathbb{N}_{a+1}, \\ x(a) &> 0, \end{aligned} \tag{1.1}$$

where $0 < \nu < 1, c(t) \leq b_1 < 0$, and motivated by Baleanu et al. [6], who obtained the solution’s monotonicity for $\lambda < 0$ and asymptotic stability of fractional Caputo-like h -difference equation

$$\begin{aligned} {}_h^C \Delta_a^{\nu} u(t) &= \lambda u(t + \nu h), \quad 0 < \nu \leq 1, t \in (h\mathbb{N})_{a+(1-\nu)h}, \\ u(a) &> 0. \end{aligned} \tag{1.2}$$

Theorem B is motivated by Jia et al. [25] and Goodrich and Peterson [18], who obtained asymptotic results for the nabla fractional q -difference equation

$$\begin{aligned} \nabla_{q,p}^{\alpha} x(t) &= c(t)x(t), \quad t \in q^{\mathbb{N}_1}, \\ x(1) &> 0, \end{aligned} \tag{1.3}$$

where $p = q^{-1}, c(t) \leq 0$.

This paper is organized as follows. In Section 2, some preliminaries about (q, h) -discrete time scales are introduced. In Section 3, Theorem A is proved and we show that the solution of the nabla Caputo (q, h) -fractional difference equation

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad c(t) \leq b < 0, \quad 0 < \alpha < 1,$$

has similar asymptotic behavior with the solution of the ordinary nabla Caputo difference equation

$${}_a^C \nabla_{(1,1)}^\alpha x(t) = c(t)x(t), \quad c(t) \leq b < 0, \quad 0 < \alpha < 1.$$

In Section 4, Theorem B is given and we show that the solution of nabla (q, h) -fractional difference equation ${}_a \nabla_{(q,h)}^\alpha x(t) = cx(t)$, $c < 0$, $0 < \alpha < 1$, has similar asymptotic behavior with the solution of the first-order nabla q -difference equation $\nabla_q x(t) = cx(t)$, $c < 0$, $t \in \mathbb{N}_1$. Four numerical examples are given in Section 5 and asymptotic stability of numerical solution is shown to support our the qualitative analysis.

2. Preliminaries

First we introduce some notation used in (q, h) -calculus (see [10] and [12]). For any real number α and $q > 0, q \neq 1$, we set $[\alpha]_q := \frac{q^\alpha - 1}{q - 1}$. Then we have the q -analogy of $n!$ in the form $[n]_q! := [n]_q [n - 1]_q \cdots [1]_q$ for $n = 1, 2, \dots$, whereas for $n = 0$ we put $[0]_q! := 1$. If $q = 1$, then $[\alpha]_1 := \alpha$ and $[n]_1!$ becomes the standard factorial. Further, the q -binomial coefficient is introduced by use of relations

$$\begin{bmatrix} \alpha \\ 0 \end{bmatrix}_{\tilde{q}} := 1,$$

$$\begin{bmatrix} \alpha \\ n \end{bmatrix}_{\tilde{q}} := \frac{[\alpha]_{\tilde{q}} [\alpha - 1]_{\tilde{q}} \cdots [\alpha - n + 1]_{\tilde{q}}}{[n]_{\tilde{q}}!}, \tag{2.1}$$

where $\alpha \in \mathbb{R}$ and $n \in \mathbb{N}_0$. The extension of the q -binomial coefficient to noninteger value n is allowed via the $\Gamma_{\tilde{q}}(t)$ function defined for $0 < \tilde{q} < 1$ as

$$\Gamma_{\tilde{q}}(t) := \frac{(\tilde{q}, \tilde{q})_\infty (1 - \tilde{q})^{1-t}}{(\tilde{q}^t, \tilde{q})_\infty}, \quad 0 < \tilde{q} < 1, \tag{2.2}$$

where $(a, \tilde{q})_\infty = \prod_{j=0}^\infty (1 - a\tilde{q}^j)$ and $t \in \mathbb{R} \setminus \{0, -1, -2, \dots\}$. It is easy to check that $\Gamma_{\tilde{q}}$ satisfies the functional relation $\Gamma_{\tilde{q}}(t + 1) = [t]_{\tilde{q}} \Gamma_{\tilde{q}}(t)$ and $\Gamma_{\tilde{q}}(1) = 1$. The q -binomial coefficient was introduced as

$$\begin{bmatrix} x \\ k \end{bmatrix}_{\tilde{q}} = \frac{\Gamma_{\tilde{q}}(x + 1)}{\Gamma_{\tilde{q}}(k + 1) \Gamma_{\tilde{q}}(x - k + 1)}, \quad x \in \mathbb{R}, \quad k \in \mathbb{Z}. \tag{2.3}$$

From (2.1), it is easy to derive the recursion formula

$$\begin{bmatrix} x \\ k \end{bmatrix}_{\tilde{q}} = \frac{1 - \tilde{q}^{x+1-k}}{1 - \tilde{q}^k} \begin{bmatrix} x \\ k - 1 \end{bmatrix}_{\tilde{q}}, \quad \begin{bmatrix} x \\ 0 \end{bmatrix}_{\tilde{q}} = 1. \tag{2.4}$$

The q -analogue of the power function is introduced as

$$(t - s)_{\tilde{q}}^{(\alpha)} = t^\alpha \frac{(s/t, \tilde{q})_\infty}{(\tilde{q}^\alpha s/t, \tilde{q})_\infty}, \quad t \neq 0, 0 < \tilde{q} < 1, \alpha \in \mathbb{R}.$$

For $\alpha = n$, a positive integer, this expression reduces to

$$(t - s)_{\tilde{q}}^{(n)} = t^n \prod_{j=0}^{n-1} (1 - \tilde{q}^j \frac{s}{t}).$$

We consider here the (q, h) -time scale (for details, see [10] and [12]):

$$\mathbb{T}_{(q,h)}^{t_0} = \{t_0 q^k + [k]_q h, k \in \mathbb{Z}\} \cup \left\{ \frac{h}{1-q} \right\}, t_0 > 0, q \geq 1, h \geq 0, q + h > 1.$$

Note that if $q = 1$ then the cluster point $h/(1 - q) = -\infty$ is not involved in $\mathbb{T}_{(q,h)}^{t_0}$. The forward and backward jump operator is the linear function $\sigma(t) = qt + h$ and $\rho(t) = q^{-1}(t - h)$, respectively. Similarly, the forward and backward graininess is given by $\mu(t) = (q - 1)t + h$ and $\nu(t) = q^{-1}\mu(t)$, respectively. We use the standard notation

$$\sigma^{k+1}(t) := \sigma^k(\sigma(t)), \quad \rho^{k+1}(t) := \rho^k(\rho(t)), \quad k = 0, 1, \dots .$$

It is easy to show that

$$\sigma^k(t) = q^k t + [k]_q h \quad \text{and} \quad \rho^k(t) = q^{-k}(t - [k]_q h).$$

The relation

$$\nu(\rho^k(t)) = q^{-k}\nu(t)$$

holds for $t \in \mathbb{T}_{(q,h)}^{t_0}$.

Let $a \in \mathbb{T}_{(q,h)}^{t_0}, a > h/(1 - q)$ be fixed. Then we introduce restrictions of the time scale $\mathbb{T}_{(q,h)}^{t_0}$ by relation

$$\tilde{\mathbb{T}}_{(q,h)}^{\sigma^i(a)} = \{t \in \mathbb{T}_{(q,h)}^{t_0}, t \geq \sigma^i(a)\}, \quad i = 0, 1, \dots ,$$

where the symbol σ^i stands for the i th iterate of σ (analogously, we use the symbol ρ^i). To simplify the notation, we put $\tilde{q} = 1/q$ whenever considering the time scale $\mathbb{T}_{(q,h)}^{t_0}$ or $\tilde{\mathbb{T}}_{(q,h)}^{\sigma^i(a)}$.

The nabla (q, h) -derivative of the function $f : \mathbb{T}_{(q,h)}^{t_0} \rightarrow \mathbb{R}$ is defined by

$$\nabla_{(q,h)} f(t) := \frac{f(t) - f(\rho(t))}{\nu(t)} = \frac{f(t) - f(\tilde{q}(t - h))}{(1 - \tilde{q})t + \tilde{q}h},$$

where $\tilde{q} = q^{-1}$.

The power functions and the nabla fractional (q, h) -Taylor monomial of degree α on $\mathbb{T}_{(q,h)}^{t_0}$ (see [10]) are defined by

$$(t - s)_{(\tilde{q},h)}^{(\alpha)} := ([t] - [s])_{\tilde{q}}^{(\alpha)}, \tag{2.5}$$

$$\hat{h}_\alpha(t, s) := \frac{(t - s)_{(\bar{q}, h)}^{(\alpha)}}{\Gamma_{\bar{q}}(\alpha + 1)}, \quad \alpha \in \mathbb{R}, \tag{2.6}$$

respectively, where the square bracket of t , $[t]$ is defined by

$$[t] := t + h\bar{q}/(1 - \bar{q}). \tag{2.7}$$

It follows that

$$\nu(t) := [t](1 - \bar{q}). \tag{2.8}$$

The following definitions appear in [10].

Definition 2.1 Assume $x : \mathbb{T}_{(q, h)}^{\sigma(a)} \rightarrow \mathbb{R}$ and $t = \sigma^n(a)$, $n \geq 1$. Then the nabla (q, h) -integral of x from a to t is defined by

$$\int_a^t x(\tau) \nabla \tau := \sum_{i=1}^n x(\sigma^i(a)) \nu(\sigma^i(a))$$

with the standard convention that $\int_a^a x(\tau) \nabla \tau = 0$. We also define the nabla (q, h) -fractional integral of order $\alpha > 0$ by

$${}_a \nabla_{(q, h)}^{-\alpha} x(t) := \int_a^t \hat{h}_{\alpha-1}(t, \rho(\tau)) x(\tau) \nabla \tau. \tag{2.9}$$

Definition 2.2 The nabla (q, h) -fractional difference (in the Riemann–Liouville sense) of order $\alpha > 0$ is defined by

$${}_a \nabla_{(q, h)}^\alpha x(t) := \nabla_{(q, h)}^m {}_a \nabla_{(q, h)}^{-(m-\alpha)} x(t),$$

where $m = \lceil \alpha \rceil$; that is, m is the ceiling of α .

Definition 2.3 The nabla (q, h) -fractional difference (in the sense of Caputo) of order $\alpha > 0$ is defined by

$${}_a^C \nabla_{(q, h)}^\alpha x(t) := {}_a \nabla_{(q, h)}^{-(m-\alpha)} \nabla_{(q, h)}^m x(t).$$

3. Asymptotic behavior for Caputo (q, h) -fractional difference equation

From the definition of the q -gamma function (2.2), it is easy to prove the following lemma.

Lemma 3.1 (Sign of q -gamma function)

1. If $x > 0$ or $x \in (-2n - 2, -2n - 1)$, $n \in \mathbb{N}_0$, then $\Gamma_{\bar{q}}(x) > 0$.
2. If $x \in (-2n - 1, -2n)$, $n \in \mathbb{N}_0$, then $\Gamma_{\bar{q}}(x) < 0$.

The following Figure 1 illustrates the above lemma.

The following lemma gives a very nice form of the Taylor monomials on $\mathbb{T}_{(q, h)}^{t_0}$ (see [10]).




Figure 1. The q -gamma function on interval $(-3, 3)$, $\tilde{q} = 0.5$.

Lemma 3.2 Let $\alpha \in \mathbb{R}, s, t \in \mathbb{T}_{(q,h)}^{t_0}$ and $n \in \mathbb{N}_0$ such that $t = \sigma^n(s)$ Then:

$$\hat{h}_\alpha(t, s) = (\nu(t))^\alpha \begin{bmatrix} \alpha + n - 1 \\ n - 1 \end{bmatrix}_{\tilde{q}}. \tag{3.1}$$

Lemma 3.3 Let $\alpha \in \mathbb{R}, s, t \in \mathbb{T}_{(q,h)}^{t_0}$ and $n \in \mathbb{N}_0$ such that $t = \sigma^n(s)$. Then:

- (i) $\hat{h}_\alpha(t, t) = 0, \alpha \neq 0$.
- (ii) $\hat{h}_0(t, s) = 1, \text{ for } n \in \mathbb{N}_0$.
- (iii) for $k \in \mathbb{N}_1$,

$$\hat{h}_{-k}(t, s) = \begin{cases} (\nu(t))^{-k}, & n = 1, \\ (\nu(t))^{-k} \frac{[n - 1 - k]_{\tilde{q}} [n - 2 - k]_{\tilde{q}} \cdots [1 - k]_{\tilde{q}}}{\Gamma_{\tilde{q}}(n)}, & n \in \mathbb{N}_2, \end{cases} \tag{3.2}$$

when $1 \leq k \leq n - 1$ and $n \in \mathbb{N}_2, \hat{h}_{-k}(t, s) = 0$.

Proof (i) If $\alpha \neq 0, -1, -2, \dots$, then

$$\begin{aligned} \hat{h}_\alpha(t, t) &= (\nu(t))^\alpha \begin{bmatrix} \alpha + 0 - 1 \\ 0 - 1 \end{bmatrix}_{\tilde{q}} \\ &= (\nu(t))^\alpha \frac{\Gamma_{\tilde{q}}(\alpha + 0)}{\Gamma_{\tilde{q}}(0)\Gamma_{\tilde{q}}(\alpha + 1)} \\ &= (\nu(t))^\alpha \frac{1}{\Gamma_{\tilde{q}}(0)[\alpha]_{\tilde{q}}} \\ &= 0. \end{aligned}$$

If $\alpha = -1, -2, \dots$, then

$$\begin{aligned} \hat{h}_\alpha(t, t) &= \lim_{\delta \rightarrow \alpha} (\nu(t))^\delta \begin{bmatrix} \delta + 0 - 1 \\ 0 - 1 \end{bmatrix}_{\bar{q}} \\ &= \lim_{\delta \rightarrow \alpha} (\nu(t))^\delta \frac{\Gamma_{\bar{q}}(\delta + 0)}{\Gamma_{\bar{q}}(0)\Gamma_{\bar{q}}(\delta + 1)} \\ &= \lim_{\delta \rightarrow \alpha} (\nu(t))^\delta \frac{1}{\Gamma_{\bar{q}}(0)[\delta]_{\bar{q}}} \\ &= \lim_{\delta \rightarrow \alpha} 0 = 0. \end{aligned}$$

(ii) If $n = 0$, then

$$\begin{aligned} \hat{h}(t, t) &= (\nu(t))^0 \begin{bmatrix} -1 \\ -1 \end{bmatrix}_{\bar{q}} \\ &= \lim_{\gamma \rightarrow -1} (\nu(t))^0 \begin{bmatrix} \gamma \\ \gamma \end{bmatrix}_{\bar{q}} \\ &= 1. \end{aligned}$$

If $n \in \mathbb{N}_1$, then

$$\begin{aligned} \hat{h}_0(t, s) &= (\nu(t))^0 \begin{bmatrix} 0 + n - 1 \\ n - 1 \end{bmatrix}_{\bar{q}} \\ &= (\nu(t))^0 \frac{\Gamma_{\bar{q}}(0 + n)}{\Gamma_{\bar{q}}(n)\Gamma_{\bar{q}}(1)} \\ &= 1. \end{aligned}$$

(iii) If $n = 1$, then

$$\begin{aligned} \hat{h}_{-k}(t, s) &= \lim_{\alpha \rightarrow k} \hat{h}_{-\alpha}(t, s) \\ &= \lim_{\alpha \rightarrow k} (\nu(t))^{-\alpha} \begin{bmatrix} -\alpha + n - 1 \\ n - 1 \end{bmatrix}_{\bar{q}} \\ &= \lim_{\alpha \rightarrow k} (\nu(t))^{-\alpha} \frac{\Gamma_{\bar{q}}(-\alpha + n)}{\Gamma_{\bar{q}}(n)\Gamma_{\bar{q}}(1 - \alpha)} \\ &= \lim_{\alpha \rightarrow k} (\nu(t))^{-\alpha} \\ &= (\nu(t))^{-k}. \end{aligned}$$

If $n \in \mathbb{N}_2$, then

$$\begin{aligned} \hat{h}_{-k}(t, s) &= \lim_{\alpha \rightarrow k} \hat{h}_{-\alpha}(t, s) \\ &= \lim_{\alpha \rightarrow k} (\nu(t))^{-\alpha} \begin{bmatrix} -\alpha + n - 1 \\ n - 1 \end{bmatrix}_{\tilde{q}} \\ &= \lim_{\alpha \rightarrow k} (\nu(t))^{-\alpha} \frac{\Gamma_{\tilde{q}}(-\alpha + n)}{\Gamma_{\tilde{q}}(n)\Gamma_{\tilde{q}}(1 - \alpha)} \\ &= \lim_{\alpha \rightarrow k} (\nu(t))^{-\alpha} \frac{[n - 1 - \alpha]_{\tilde{q}}[n - 2 - \alpha]_{\tilde{q}} \cdots [1 - \alpha]_{\tilde{q}}}{\Gamma_{\tilde{q}}(n)} \\ &= (\nu(t))^{-k} \frac{[n - 1 - k]_{\tilde{q}}[n - 2 - k]_{\tilde{q}} \cdots [1 - k]_{\tilde{q}}}{\Gamma_{\tilde{q}}(n)}. \end{aligned}$$

Note that if $1 \leq k \leq n - 1$ and $n \in \mathbb{N}_2$, then $\hat{h}_{-k}(t, s) = 0$. □

Next we give nabla power rule formulas as an extension of results that appear in [24] and [25].

Lemma 3.4 Assume $q > 1$, $\tilde{q} = q^{-1}$, $\alpha \in \mathbb{R}$, and let $t = \sigma^n(s)$, $n \in \mathbb{N}_0$, where $s \in \mathbb{T}_{(q,h)}^{t_0}$. Then:

1. The nabla (q, h) -difference of the function $\hat{h}_{-\alpha}(t, s)$ with respect to t is given by

$${}_t \nabla_{(q,h)} \hat{h}_{-\alpha}(t, s) = \hat{h}_{-\alpha-1}(t, s).$$

2. The nabla (q, h) -difference of the function $\hat{h}_{-\alpha}(t, s)$ with respect to s is given by

$${}_s \nabla_{(q,h)} \hat{h}_{-\alpha}(t, s) = -\hat{h}_{-\alpha-1}(t, \rho(s)).$$

Proof These results follow easily from Lemma 2.2 in [17]. □

Lemma 3.5 Assume $c(t) < (\nu(t))^{-\alpha}$, $0 < \alpha < 1$. Then any solution of

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \tag{3.3}$$

satisfying $x(a) > 0$ is positive on $\tilde{\mathbb{T}}_{(q,h)}^a$.

Proof Using integration by parts and

$${}_s \nabla_{(q,h)} \hat{h}_{-\alpha}(t, s) = -\hat{h}_{-\alpha-1}(t, \rho(s)),$$

we have

$$\begin{aligned}
 {}_a^C \nabla_{(q,h)}^\alpha x(t) &\stackrel{\text{Def.2.3}}{=} {}_a \nabla_{(q,h)}^{-(1-\alpha)} \nabla_{(q,h)} x(t) \\
 &\stackrel{\text{Def.2.1}}{=} \int_a^t \hat{h}_{-\alpha}(t, \rho(s)) \nabla_{(q,h)} x(s) \nabla s \\
 &= \hat{h}_{-\alpha}(t, s) x(s) \Big|_{s=a}^t + \int_a^t \hat{h}_{-\alpha-1}(t, \rho(s)) x(s) \nabla s \\
 &= -\hat{h}_{-\alpha}(t, a) x(a) + \sum_{i=1}^k \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) x(\sigma^i(a)) \nu(\sigma^i(a)).
 \end{aligned}$$

Taking $t = \sigma^k(a), k \geq 1$, we have

$$\begin{aligned}
 {}_a^C \nabla_{(q,h)}^\alpha x(t) &= {}_a^C \nabla_{(q,h)}^\alpha x(\sigma^k(a)) \\
 &= -\hat{h}_{-\alpha}(t, a) x(a) + \hat{h}_{-\alpha-1}(\sigma^k(a), \sigma^{k-1}(a)) x(\sigma^k(a)) \nu(\sigma^k(a)) \\
 &\quad + \sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) x(\sigma^i(a)) \nu(\sigma^i(a)) \\
 &= -\hat{h}_{-\alpha}(t, a) x(a) + x(\sigma^k(a)) (\nu(\sigma^k(a)))^{-\alpha} \\
 &\quad + \sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) x(\sigma^i(a)) \nu(\sigma^i(a)).
 \end{aligned}$$

Using $x(t)$ as a solution of (3.3), we get

$$\begin{aligned}
 x(\sigma^k(a)) &= \frac{1}{(\nu(\sigma^k(a)))^{-\alpha} - c(\sigma^k(a))} \left\{ \hat{h}_{-\alpha}(t, a) x(a) \right. \\
 &\quad \left. - \sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) x(\sigma^i(a)) \nu(\sigma^i(a)) \right\}.
 \end{aligned}$$

Note that

$$\begin{aligned}
 \hat{h}_{-\alpha}(t, a) &= (\nu(t))^{-\alpha} \begin{bmatrix} -\alpha + k - 1 \\ k - 1 \end{bmatrix}_{\bar{q}} \\
 &= (\nu(t))^{-\alpha} \frac{\Gamma_{\bar{q}}(-\alpha + k)}{\Gamma_{\bar{q}}(k) \Gamma_{\bar{q}}(1 - \alpha)} \\
 &> 0,
 \end{aligned} \tag{3.4}$$

for $k \geq 1$.

$$\begin{aligned}
 \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) &= (\nu(t))^{-\alpha-1} \begin{bmatrix} -\alpha - 1 + k - i \\ k - i \end{bmatrix}_{\bar{q}} \\
 &= (\nu(t))^{-\alpha-1} \frac{\Gamma_{\bar{q}}(-\alpha + k - i)}{\Gamma_{\bar{q}}(k - i + 1) \Gamma_{\bar{q}}(-\alpha)} \\
 &< 0,
 \end{aligned} \tag{3.5}$$

for $1 \leq i \leq k - 1$.

Using the strong induction principle, (3.4), (3.5), and $x(a) > 0$, it is easy to see that $x(\sigma^i(a)) > 0$ for $i \in \mathbb{N}_0$, where we use the convention $\sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a))x(\sigma^i(a))\nu(\sigma^i(a)) = 0$ for $k < 2$. \square

The following comparison theorem plays an important role in proving our main results.

Theorem 3.6 Assume $c_2(t) \leq c_1(t) < (\nu(t))^{-\alpha}$, $0 < \alpha < 1$, and $x(t), y(t)$ are solutions of the equations

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = c_1(t)x(t), \tag{3.6}$$

and

$${}_a^C \nabla_{(q,h)}^\alpha y(t) = c_2(t)y(t), \tag{3.7}$$

respectively, for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$ satisfying $x(a) \geq y(a) > 0$. Then

$$x(t) \geq y(t),$$

for $t \in \tilde{\mathbb{T}}_{(q,h)}^a$.

Proof Similar to the first part of the proof of Lemma 3.5, taking $t = \sigma^k(a), k \geq 1$, we have

$$x(\sigma^k(a)) = \frac{1}{(\nu(\sigma^k(a)))^{-\alpha} - c_1(\sigma^k(a))} \left\{ \hat{h}_{-\alpha}(t, a)x(a) - \sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a))x(\sigma^i(a))\nu(\sigma^i(a)) \right\}. \tag{3.8}$$

$$y(\sigma^k(a)) = \frac{1}{(\nu(\sigma^k(a)))^{-\alpha} - c_2(\sigma^k(a))} \left\{ \hat{h}_{-\alpha}(t, a)y(a) - \sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a))y(\sigma^i(a))\nu(\sigma^i(a)) \right\}. \tag{3.9}$$

We will prove $x(\sigma^k(a)) \geq y(\sigma^k(a))$ for $k \in \mathbb{N}_0$ by using the principle of strong induction. By assumption $x(a) \geq y(a) > 0$, so the base case holds. Now assume that $x(\sigma^i(a)) \geq y(\sigma^i(a))$, for $i = 0, 1 \dots, k - 1$.

Since $c_2(t) \leq c_1(t) < (\nu(t))^{-\alpha}$,

$$\hat{h}_{-\alpha}(t, a) > 0$$

and

$$\hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) < 0, \quad \text{for } 1 \leq i \leq k - 1.$$

From (3.8) and (3.9) we have

$$x(\sigma^k(a)) \geq y(\sigma^k(a)) > 0.$$

This completes the proof. \square

The following result appears in [38].

Lemma 3.7 ([38]) *Let $\alpha \in \mathbb{R}$ and $n \in \mathbb{Z}^+$ so that $n - 1 < \alpha \leq n$. Then*

$${}_a \nabla_{(q,h)a}^{-\alpha} {}^C \nabla_{(q,h)}^\alpha x(t) = x(t) - \sum_{k=0}^{n-1} \hat{h}_k(t, a) \nabla_{(q,h)}^k x(a). \tag{3.10}$$

Since $\hat{h}_0(t, a) = 1$ we get the following Lemma.

Lemma 3.8 *Assume that $0 < \alpha < 1$ and $x(t)$ is a solution of the fractional equation*

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \tag{3.11}$$

satisfying $x(a) > 0$. Then $x(t)$ satisfies the integral equation

$$x(t) = x(a) + \int_a^t \hat{h}_{\alpha-1}(t, \rho(s)) c(s) x(s) \nabla s. \tag{3.12}$$

Remark 3.9 *Letting $t = \sigma^k(a), k \in \mathbb{N}_0$, (3.12) can be rewritten explicitly as*

$$\begin{aligned} x(\sigma^k(a)) &= \frac{x(a) + \sum_{i=1}^{k-1} \hat{h}_{\alpha-1}(\sigma^k(a), \sigma^{i-1}(a)) c(\sigma^i(a)) \nu(\sigma^i(a)) x(\sigma^i(a))}{1 - (\nu(\sigma^k(a)))^\alpha c(\sigma^k(a))} \\ &= \frac{x(a) + \sum_{i=1}^{k-1} (\nu(\sigma^k(a)))^{\alpha-1} \begin{bmatrix} \alpha - 1 + k - i \\ k - i \end{bmatrix}_{\tilde{q}} c(\sigma^i(a)) \nu(\sigma^i(a)) x(\sigma^i(a))}{1 - (\nu(\sigma^k(a)))^\alpha c(\sigma^k(a))}, \end{aligned} \tag{3.13}$$

for $k \geq 1$, where we used formula (3.1).

Lemma 3.10

$$\lim_{n \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(n + \alpha)}{\Gamma_{\tilde{q}}(n)} = (1 - \tilde{q})^{-\alpha}, \quad 0 < \tilde{q} < 1, \alpha \in \mathbb{R}. \tag{3.14}$$

Proof From Lemma 3.1, we know that if $\alpha \in \mathbb{R}, a, t \in \mathbb{T}_{(q,h)}^{t_0}$ and $n \in \mathbb{N}_0$ such that $t = \sigma^n(a)$, then

$$\begin{aligned} \hat{h}_\alpha(t, a) &= (\nu(t))^\alpha \begin{bmatrix} \alpha + n - 1 \\ n - 1 \end{bmatrix}_{\tilde{q}} \\ &= (\nu(t))^\alpha \frac{\Gamma_{\tilde{q}}(\alpha + n)}{\Gamma_{\tilde{q}}(\alpha + 1) \Gamma_{\tilde{q}}(n)} \\ &= [t]^\alpha (1 - \tilde{q})^\alpha \frac{\Gamma_{\tilde{q}}(\alpha + n)}{\Gamma_{\tilde{q}}(\alpha + 1) \Gamma_{\tilde{q}}(n)}. \end{aligned}$$

On the other hand, from (2.6),

$$\begin{aligned} \hat{h}_\alpha(t, a) &= \frac{(t-a)_{(\tilde{q}, h)}^{(\alpha)}}{\Gamma_{\tilde{q}}(\alpha+1)} \\ &= \frac{([t]-[a])_{\tilde{q}}^{(\alpha)}}{\Gamma_{\tilde{q}}(\alpha+1)} \\ &= [t]^\alpha \frac{([a]/[t], \tilde{q})_\infty}{(\tilde{q}^\alpha [a]/[t], \tilde{q})_\infty \Gamma_{\tilde{q}}(\alpha+1)} \\ &= [t]^\alpha \frac{(\tilde{q}^n, \tilde{q})_\infty}{(\tilde{q}^{\alpha+n}, \tilde{q})_\infty \Gamma_{\tilde{q}}(\alpha+1)} \\ &= [t]^\alpha \frac{(1-\tilde{q}^{\alpha+1}) \cdots (1-\tilde{q}^{\alpha+n-1})}{(1-\tilde{q}) \cdots (1-\tilde{q}^{n-1}) \Gamma_{\tilde{q}}(\alpha+1)} \frac{\prod_{j=0}^\infty (1-\tilde{q}^{j+1})}{\prod_{j=0}^\infty (1-\tilde{q}^{j+\alpha+1})} \\ &= [t]^\alpha \frac{(1-\tilde{q}^{\alpha+1}) \cdots (1-\tilde{q}^{\alpha+n-1})}{(1-\tilde{q}) \cdots (1-\tilde{q}^{n-1}) \Gamma_{\tilde{q}}(\alpha+1)} \frac{\Gamma_{\tilde{q}}(\alpha+1)}{(1-\tilde{q})^{-\alpha}}, \end{aligned}$$

so

$$\frac{\Gamma_{\tilde{q}}(\alpha+n)}{\Gamma_{\tilde{q}}(n)} = \frac{(1-\tilde{q}^{\alpha+1}) \cdots (1-\tilde{q}^{\alpha+n-1})}{(1-\tilde{q}) \cdots (1-\tilde{q}^{n-1})} \Gamma_{\tilde{q}}(\alpha+1).$$

Letting n tend to infinity, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(\alpha+n)}{\Gamma_{\tilde{q}}(n)} &= \lim_{n \rightarrow \infty} \frac{(1-\tilde{q}^{\alpha+1}) \cdots (1-\tilde{q}^{\alpha+n-1})}{(1-\tilde{q}) \cdots (1-\tilde{q}^{n-1})} \Gamma_{\tilde{q}}(\alpha+1) \\ &= \frac{\prod_{j=0}^\infty (1-\tilde{q}^{j+\alpha+1})}{\prod_{j=0}^\infty (1-\tilde{q}^{j+1})} \Gamma_{\tilde{q}}(\alpha+1) \\ &= \frac{(\tilde{q}^{\alpha+1}, \tilde{q})_\infty}{(\tilde{q}, \tilde{q})_\infty} \Gamma_{\tilde{q}}(\alpha+1) \\ &= (1-\tilde{q})^{-\alpha}. \end{aligned}$$

□

Remark 3.11 Note that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(\alpha+n)}{\Gamma_{\tilde{q}}(n)} &= (1-\tilde{q})^{-\alpha} \\ &= \lim_{n \rightarrow \infty} \left(\frac{1-\tilde{q}^n}{1-\tilde{q}} \right)^\alpha \\ &= \lim_{n \rightarrow \infty} ([n]_{\tilde{q}})^\alpha. \end{aligned}$$

It follows that

$$\lim_{n \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(n+\alpha)}{\Gamma_{\tilde{q}}(n)[n]_{\tilde{q}}^\alpha} = 1,$$

so this limit can be regarded as the q -analogue of the Stirling formula (see [45])

$$\lim_{x \rightarrow \infty} \frac{\Gamma(x + \alpha)}{\Gamma(x)x^\alpha} = 1.$$

Considering the monotonicity of the solution for the following linear Caputo fractional difference equation,

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = cx(t), \quad c < 0, \tag{3.15}$$

we have the following result.

Lemma 3.12 *If $x(t)$ is a solution of equation (3.15), $x(a) > 0$ and $c < 0$, then $\nabla_{(q,h)}x(t) < 0$ for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$.*

Proof Letting $t = \sigma(a)$, from (3.12), we get

$$\begin{aligned} x(\sigma(a)) &= x(a) + \hat{h}_{\alpha-1}(\sigma(a), a)cx(\sigma(a))\nu(\sigma(a)) \\ &= x(a) + cx(\sigma(a))\nu^\alpha(\sigma(a)) \\ &< x(a), \end{aligned}$$

where we use $\hat{h}_{\alpha-1}(\sigma(a), a) = (\nu(\sigma(a)))^{\alpha-1}$, $x(\sigma(a)) > 0$ (from Lemma 3.5), and $c < 0$.

If $\nabla_{(q,h)}x(t) < 0$, for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$, this completes the proof.

If not, we assume that there exists two points $t_1 = \sigma^n(a), t_2 = \sigma^m(a), t_1 < t_2, 1 \leq n < m$ such that $\nabla_{(q,h)}x(\sigma^i(a)) < 0$ for $1 \leq i \leq n < m$ and $\nabla_{(q,h)}x(\sigma^i(a)) > 0$ for $n + 1 \leq i \leq m$.

$$\begin{aligned} &{}_a^C \nabla_{(q,h)}^\alpha x(t_1) - {}_a^C \nabla_{(q,h)}^\alpha x(t_2) \\ &= \int_a^{t_1} \hat{h}_{-\alpha}(t_1, \rho(s)) \nabla_{(q,h)}x(s) \nabla s - \int_a^{t_2} \hat{h}_{-\alpha}(t_2, \rho(s)) \nabla_{(q,h)}x(s) \nabla s \\ &= \int_a^{t_1} \left(\hat{h}_{-\alpha}(t_1, \rho(s)) - \hat{h}_{-\alpha}(t_2, \rho(s)) \right) \nabla_{(q,h)}x(s) \nabla s \\ &\quad - \int_{t_1}^{t_2} \hat{h}_{-\alpha}(t_2, \rho(s)) \nabla_{(q,h)}x(s) \nabla s \\ &= \sum_{i=1}^n \left(\hat{h}_{-\alpha}(\sigma^n(a), \sigma^{i-1}(a)) - \hat{h}_{-\alpha}(\sigma^m(a), \sigma^{i-1}(a)) \right) \nabla_{(q,h)}x(\sigma^i(a))\nu(\sigma^i(a)) \\ &\quad - \sum_{i=n+1}^m \hat{h}_{-\alpha}(\sigma^m(a), \sigma^{i-1}(a)) \nabla_{(q,h)}x(\sigma^i(a))\nu(\sigma^i(a)). \end{aligned}$$

Since

$$\begin{aligned} \hat{h}_{-\alpha}(\sigma^m(a), \sigma^{i-1}(a)) &= \left(\nu(\sigma^m(a)) \right)^{-\alpha} \left[\begin{matrix} -\alpha + m - i \\ m - i \end{matrix} \right]_{\bar{q}} \\ &= \left(\nu(\sigma^m(a)) \right)^{-\alpha} \frac{\Gamma_{\bar{q}}(-\alpha + m - i + 1)}{\Gamma_{\bar{q}}(m - i + 1)\Gamma_{\bar{q}}(1 - \alpha)} \\ &> 0, \end{aligned}$$

for $i \leq m$, similarly, we have

$$\hat{h}_{-\alpha}(\sigma^n(a), \sigma^{i-1}(a)) > 0, \quad \text{for } i \leq n.$$

Note that

$$\begin{aligned} & \frac{\hat{h}_{-\alpha}(\sigma^n(a), \sigma^{i-1}(a))}{\hat{h}_{-\alpha}(\sigma^m(a), \sigma^{i-1}(a))} \\ &= \frac{(\nu(t_1))^{-\alpha}}{(\nu(t_2))^{-\alpha}} \cdot \frac{\Gamma_{\tilde{q}}(-\alpha + n - i + 1)}{\Gamma_{\tilde{q}}(n - i + 1)\Gamma_{\tilde{q}}(1 - \alpha)} \cdot \frac{\Gamma_{\tilde{q}}(m - i + 1)\Gamma_{\tilde{q}}(1 - \alpha)}{\Gamma_{\tilde{q}}(-\alpha + m - i + 1)} \\ &= \frac{(\nu(t_1))^{-\alpha}}{(\nu(t_2))^{-\alpha}} \cdot \frac{\Gamma_{\tilde{q}}(-\alpha + n - i + 1)}{\Gamma_{\tilde{q}}(-\alpha + m - i + 1)} \cdot \frac{\Gamma_{\tilde{q}}(m - i + 1)}{\Gamma_{\tilde{q}}(n - i + 1)} \\ &= \left(\frac{\nu(t_1)}{\nu(t_2)}\right)^{-\alpha} \frac{[m - i]_{\tilde{q}}[m - i - 1]_{\tilde{q}} \cdots [n - i + 1]_{\tilde{q}}}{[-\alpha + m - i]_{\tilde{q}}[-\alpha + m - i - 1]_{\tilde{q}} \cdots [-\alpha + n - i + 1]_{\tilde{q}}}. \end{aligned}$$

Hence,

$$\begin{aligned} & \frac{\hat{h}_{-\alpha}(\sigma^n(a), \sigma^{i-1}(a))}{\hat{h}_{-\alpha}(\sigma^m(a), \sigma^{i-1}(a))} \\ &= \left(\frac{\nu(t_1)}{\nu(t_2)}\right)^{-\alpha} \cdot \frac{1 - \tilde{q}^{m-i}}{1 - \tilde{q}^{-\alpha+m-i}} \frac{1 - \tilde{q}^{m-i-1}}{1 - \tilde{q}^{-\alpha+m-i-1}} \cdots \frac{1 - \tilde{q}^{n-i+1}}{1 - \tilde{q}^{-\alpha+n-i+1}} \\ &= \left(\frac{(q-1)t_1 + h}{(q-1)t_2 + h}\right)^{-\alpha} \cdot \frac{1 - \tilde{q}^{m-i}}{1 - \tilde{q}^{-\alpha+m-i}} \frac{1 - \tilde{q}^{m-i-1}}{1 - \tilde{q}^{-\alpha+m-i-1}} \cdots \frac{1 - \tilde{q}^{n-i+1}}{1 - \tilde{q}^{-\alpha+n-i+1}} \\ &> 1, \end{aligned}$$

for $1 \leq i \leq n < m$.

Thus,

$$\hat{h}_{-\alpha}(\sigma^n(a), \sigma^{i-1}(a)) - \hat{h}_{-\alpha}(\sigma^m(a), \sigma^{i-1}(a)) > 0, \quad \text{for } 1 \leq i \leq n < m.$$

We can derive that

$${}_a^C \nabla_{(q,h)}^\alpha x(t_1) - {}_a^C \nabla_{(q,h)}^\alpha x(t_2) < 0.$$

On the other hand, due to $c < 0$ and $x(t_1) < x(t_2)$, we have

$${}_a^C \nabla_{(q,h)}^\alpha x(t_1) - {}_a^C \nabla_{(q,h)}^\alpha x(t_2) = c(x(t_1) - x(t_2)) > 0,$$

which is a contradiction. □

From Corollary 4.8 in [38], we know that

$$x(t) = x(a)E_{\alpha,1}^{a,c}(t)$$

is the solution of (3.15). As an application of Lemma 3.12, we have the following corollary.

Corollary 3.13 *The (q, h) -Mittag-Leffler function*

$$E_{\alpha,1}^{a,c}(t) = \sum_{k=0}^{\infty} c^k \hat{h}_{\alpha k}(t, a)$$

is monotonically decreasing if $c < 0$.

When $c(t) = c$ is a negative constant, we can give a simpler proof for Lemma 3.5.

Lemma 3.14 *Assume $c < 0$ and $x(t)$ is a solution of equation (3.15) satisfying $x(a) > 0$. Then $x(t) > 0$ for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$.*

Proof Let $t = \sigma^k(a), k \geq 1$. Note that

$$\begin{aligned} \hat{h}_{-\alpha}(\sigma^k(a), \sigma^{i-1}(a)) &= (\nu(\sigma^k(a)))^{-\alpha} \begin{bmatrix} -\alpha + k - i \\ k - i \end{bmatrix}_{\bar{q}} \\ &= (\nu(\sigma^k(a)))^{-\alpha} \frac{\Gamma_{\bar{q}}(-\alpha + k - i + 1)}{\Gamma_{\bar{q}}(k - i + 1)\Gamma_{\bar{q}}(1 - \alpha)} \\ &> 0, \end{aligned}$$

for $1 \leq i \leq k$. From Lemma 3.12, we have $\nabla_{(q,h)} x(\sigma^i(a)) < 0, i = 1, 2, \dots, k$. Hence,

$$\begin{aligned} {}_a^C \nabla_{(q,h)}^{\alpha} x(t) &= \int_a^t \hat{h}_{-\alpha}(t, \rho(s)) \nabla_{(q,h)} x(s) \nabla s \\ &= \sum_{i=1}^k \hat{h}_{-\alpha}(\sigma^k(a), \sigma^{i-1}(a)) \nabla_{(q,h)} x(\sigma^i(a)) \nu(\sigma^i(a)) \\ &< 0. \end{aligned}$$

As a result, $x(t)$ of the RHS of (3.15) should be positive when $c < 0$. □

Theorem 3.15 *If $c < 0$ and $x(t)$ is a solution of equation (3.15) satisfying $x(a) > 0$, then*

$$\lim_{t \rightarrow \infty} x(t) = 0.$$

Proof From Lemma 3.12 and Lemma 3.14, we know the limit exists. Arguing by contradiction, we assume $\lim_{t \rightarrow \infty} x(t) = A > 0$. By Lemma 3.12 we get that $x(t)$ is decreasing. Then, by equation (3.12), we have

$$\begin{aligned} x(t) - x(a) &= \int_a^t \hat{h}_{\alpha-1}(t, \rho(s)) cx(s) \nabla s \tag{3.16} \\ &< cx(t) \int_a^t \hat{h}_{\alpha-1}(t, \rho(s)) \nabla s \\ &= -cx(t) \hat{h}_{\alpha}(t, s) \Big|_{s=a}^t \\ &= cx(t) \hat{h}_{\alpha}(t, a). \end{aligned}$$

Letting t tend to infinity, the LHS of (3.16) becomes

$$\lim_{t \rightarrow \infty} x(t) - x(a) = A - x(a) < 0,$$

while the RHS gives for $t = \sigma^k(a)$, $k \geq 1$,

$$\begin{aligned} \lim_{t \rightarrow \infty} cx(t)\hat{h}_\alpha(t, a) &= cA \lim_{t \rightarrow \infty} \hat{h}_\alpha(t, a) \\ &= cA \lim_{t \rightarrow \infty} (\nu(t))^\alpha \lim_{t \rightarrow \infty} \left[\begin{matrix} \alpha + k - 1 \\ k - 1 \end{matrix} \right]_{\tilde{q}} \\ &= cA \lim_{t \rightarrow \infty} (\nu(t))^\alpha \lim_{k \rightarrow \infty} \left[\begin{matrix} \alpha + k - 1 \\ k - 1 \end{matrix} \right]_{\tilde{q}} \\ &\stackrel{(2.8)}{=} cA \lim_{t \rightarrow \infty} [\sigma^k(a)]^\alpha (1 - \tilde{q})^\alpha \lim_{k \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(\alpha + k)}{\Gamma_{\tilde{q}}(k)\Gamma_{\tilde{q}}(\alpha + 1)} \\ &= cA \cdot (+\infty) \frac{1}{\Gamma_{\tilde{q}}(\alpha + 1)} \\ &= -\infty, \end{aligned}$$

where we use

$$\begin{aligned} \lim_{k \rightarrow \infty} [\sigma^k(a)]^\alpha &= \lim_{k \rightarrow \infty} [q^k a + [k]_q h]^\alpha \\ &= \lim_{k \rightarrow \infty} \left(q^k a + [k]_q h + h \frac{\tilde{q}}{1 - \tilde{q}} \right)^\alpha \\ &= \lim_{k \rightarrow \infty} \left(q^k a + \frac{1 - q^k}{1 - q} h + \frac{h}{q - 1} \right)^\alpha \\ &= \lim_{k \rightarrow \infty} \left(q^k \left(a + \frac{h}{q - 1} \right) \right)^\alpha \\ &= +\infty \end{aligned}$$

and

$$\lim_{k \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(\alpha + k)}{\Gamma_{\tilde{q}}(k)} \stackrel{(3.14)}{=} (1 - \tilde{q})^{-\alpha}.$$

This yields a contradiction and we conclude that $\lim_{k \rightarrow \infty} x(t) = 0$. □

Theorem A. Assume $0 < \alpha < 1$ and there is a constant b such that $c(t) \leq b < 0$, for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. Then any solution, $x(t)$, of the nabla Caputo (q, h) -fractional difference equation

$${}_a^C \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \tag{3.17}$$

with $x(a) > 0$ satisfies

$$\lim_{t \rightarrow \infty} x(t) = 0.$$

Proof Assume $y(t)$ is the solution of Caputo nabla fractional equation

$${}_a^C \nabla_{(q,h)}^\alpha y(t) = by(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$$

satisfying $y(a) = 2x(a) > 0$. From Theorem 3.6, we have

$$0 < x(t) < y(t),$$

for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. From Theorem 3.15, we get that

$$0 \leq \lim_{t \rightarrow \infty} x(t) \leq \lim_{t \rightarrow \infty} y(t) = 0.$$

This completes the proof. □

Remark 3.16 (see Example 5.2) *If $c(t)$ in Theorem A is not a negative constant, then the solution of the equation (3.17) may not be monotonically decreasing.*

4. Asymptotic behavior for Riemann–Liouville (q, h) -fractional difference equation

From the Leibniz rule on time scales in [7, 8], it is easy to get the following nabla (q, h) -difference Leibniz rule.

Lemma 4.1 (Leibniz rule) *Assume $f : \tilde{\mathbb{T}}_{(q,h)}^a \times \tilde{\mathbb{T}}_{(q,h)}^a \rightarrow \mathbb{R}$. Then*

$$\nabla_{(q,h)} \int_a^t f(t, s) \nabla s = \int_a^t {}_t\nabla_{(q,h)} f(t, s) \nabla s + f(\rho(t), t)$$

for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$, where it is understood that the expression ${}_t\nabla_{(q,h)} f(t, s)$ in the last integral means the (partial) (q, h) -difference of f with respect to t .

Theorem 4.2 *Assume $0 < \alpha < 1$, $c(t) \leq 0$, and $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$. Then any solution of the equation*

$${}_a\nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)} \tag{4.1}$$

satisfying $x(\sigma(a)) > 0$ is positive on $\tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$.

Proof Since $m = \lceil \alpha \rceil = 1$, we have that

$$\begin{aligned} {}_a\nabla_{(q,h)}^\alpha x(t) &\stackrel{\text{Def. 2.2}}{=} \nabla_{(q,h)} {}_a\nabla_{(q,h)}^{-(1-\alpha)} x(t) \\ &\stackrel{(2.9)}{=} \nabla_{(q,h)} \int_a^t \hat{h}_{-\alpha}(t, \rho(\tau)) x(\tau) \nabla \tau \\ &\stackrel{\text{Lem. 4.1}}{=} \int_a^t \hat{h}_{-\alpha-1}(t, \rho(\tau)) x(\tau) \nabla \tau + \hat{h}_{-\alpha}(\rho(t), \rho(t)) x(t) \\ &= \int_a^t \hat{h}_{-\alpha-1}(t, \rho(\tau)) x(\tau) \nabla \tau. \end{aligned}$$

Letting $t = \sigma^k(a), k \geq 2$, we get that

$${}_a\nabla_{(q,h)}^\alpha x(t) = \sum_{i=1}^k \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a))x(\sigma^i(a))\nu(\sigma^i(a)).$$

Using that $x(t)$ is a solution of (4.1), we get that for $t = \sigma^k(a)$, $k \geq 2$,

$$\begin{aligned} & (\hat{h}_{-\alpha-1}(t, \sigma^{k-1}(a))\nu(t) - c(t))x(\sigma^k(a)) \\ &= - \sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(t\sigma^{i-1}(a))x(\sigma^i(a))\nu(\sigma^i(a)). \end{aligned} \tag{4.2}$$

We now show that since $0 < \alpha < 1$, $\hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) < 0$ for $1 \leq i \leq k - 1$. First note that

$$\begin{aligned} \hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) &\stackrel{(3.1)}{=} (\nu(t))^{-\alpha-1} \left[\begin{matrix} -\alpha - 1 + k - i \\ k - i \end{matrix} \right]_{\bar{q}} \\ &\stackrel{(2.3)}{=} (\nu(t))^{-\alpha-1} \frac{\Gamma_{\bar{q}}(-\alpha + k - i)}{\Gamma_{\bar{q}}(k - i + 1)\Gamma_{\bar{q}}(-\alpha)}. \end{aligned}$$

Also, since $\Gamma_{\bar{q}}(-\alpha + k - i) > 0, \Gamma_{\bar{q}}(k - i + 1) > 0$, for $k - i \geq 1$ and $\Gamma_{\bar{q}}(-\alpha) < 0$, we have

$$\hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) < 0,$$

for $1 \leq i \leq k - 1$.

Note that from Lemma 3.2 we have that

$$\hat{h}_{-\alpha-1}(t, \sigma^{k-1}(a)) > 0.$$

Since $c(t) \leq 0, \nu(t) > 0$ and $\hat{h}_{-\alpha-1}(t, \sigma^{k-1}(a)) > 0$, the coefficient of $x(t) = x(\sigma^k(a))$ on the left hand side of (4.2) is positive. Since $\hat{h}_{-\alpha-1}(t, \sigma^{i-1}(a)) < 0$ and $\nu(t) > 0$ for $1 \leq i \leq k - 1$, and $x(\sigma^i(a)) > 0$, it follows from (4.2) and the strong induction principle that $x(t) = x(\sigma^k(a)) > 0$, for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$.

This completes the proof. □

Remark 4.3 *In the above theorem we proved that if $x(t)$ is a solution of the IVP*

$${}_a\nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad x(\sigma(a)) = A, \tag{4.3}$$

then $x(t)$ is a solution of the summation equation (4.2) satisfying the initial condition $x(\sigma(a)) = A$. Since the steps in this calculation are reversible we have that if $x(t)$ is a solution of the summation equation (4.2) satisfying the initial condition $x(\sigma(a)) = A$, then $x(t)$ is a solution of the IVP (4.3). Hence, we say the IVP (4.3) is equivalent to the summation equation (4.2) along with the initial condition $x(\sigma(a)) = A$. Now solving the summation equation (4.2) for $x(\sigma^k(a))$, we obtain the summation equation

$$\begin{aligned} x(\sigma^k(a)) &= - \frac{\sum_{i=1}^{k-1} \hat{h}_{-\alpha-1}(\sigma^k(a), \sigma^{i-1}(a))x(\sigma^i(a))\nu(\sigma^i(a))}{\hat{h}_{-\alpha-1}(\sigma^k(a), \sigma^{k-1}(a))\nu(\sigma^k(a)) - c(\sigma^k(a))} \\ &= - \frac{\sum_{i=1}^{k-1} (\nu(\sigma^k(a)))^{-\alpha-1} \left[\begin{matrix} -\alpha - 1 + k - i \\ k - i \end{matrix} \right]_{\bar{q}} x(\sigma^i(a))\nu(\sigma^i(a))}{(\nu(\sigma^k(a)))^{-\alpha} - c(\sigma^k(a))}, \end{aligned} \tag{4.4}$$

for $k \geq 2$, where we used formula (3.1). Hence, the IVP (4.3) and the summation equation (4.4) with the initial condition $x(\sigma(a)) = A$ are equivalent. Note that if $x(\sigma(a)) = A$, then by the summation equation (4.4) with $k = 2$ we get that $x(\sigma^2(a))$ is uniquely determined, and then when $k = 3$ the values of $x(t)$ at $\sigma(a)$ and $\sigma^2(a)$ uniquely determine the value of $x(\sigma^3(a))$. Hence, it follows by strong mathematical induction that the solution of the IVP (4.3) exists and is unique on $\tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. Since the IVP (4.3) and the summation equation with the the initial condition $x(\sigma(a)) = A$ have a unique solution, we have that the IVP (4.3) has a unique solution and this solution exists on $\tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. Furthermore, it is easy to see that the nabla (q, h) -fractional equation has just one linearly independent solution $x_1(t)$ on $\tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$ and $x(t) := \alpha x_1(t)$ is a general solution of the nabla fractional equation (4.1).

Lemma 4.4 Assume that $\alpha \in \mathbb{R}$, $0 < \tilde{q} < 1$. Then for $[t] \neq 0$,

$$\nu(\sigma(a))\hat{h}_{-\alpha-1}(t, a) + \hat{h}_{-\alpha}(t, \sigma(a)) = \hat{h}_{-\alpha}(t, a). \tag{4.5}$$

Proof

On the one hand,

$${}_s\nabla_{(q,h)}\hat{h}_{-\alpha}(t, \sigma(a)) = \frac{\hat{h}_{-\alpha}(t, \sigma(a)) - \hat{h}_{-\alpha}(t, a)}{\nu(\sigma(a))}.$$

On the other hand, using Lemma 3.4 (2),

$${}_s\nabla_{(q,h)}\hat{h}_{-\alpha}(t, \sigma(a)) = -\hat{h}_{-\alpha-1}(t, a).$$

Thus,

$$\frac{\hat{h}_{-\alpha}(t, \sigma(a)) - \hat{h}_{-\alpha}(t, a)}{\nu(\sigma(a))} = -\hat{h}_{-\alpha-1}(t, a).$$

Hence, (4.5) holds. □

Lemma 4.5 Assume $f : \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \rightarrow \mathbb{R}$, $0 < \alpha < 1$, $q > 1$. Then

$$\sigma(a)\nabla_{(q,h)}^{-(1-\alpha)}\nabla_{(q,h)}f(t) = \nabla_{(q,h)}\sigma(a)\nabla_{(q,h)}^{-(1-\alpha)}f(t) - f(\sigma(a))\hat{h}_{-\alpha}(t, \sigma(a)) \tag{4.6}$$

$$= \sigma(a)\nabla_{(q,h)}^\alpha f(t) - f(\sigma(a))\hat{h}_{-\alpha}(t, \sigma(a)) \tag{4.7}$$

and

$$\sigma(a)\nabla_{(q,h)}^{-(1-\alpha)}\nabla_{(q,h)}f(t) = \nabla_{(q,h)}{}_a\nabla_{(q,h)}^{-(1-\alpha)}f(t) - f(\sigma(a))\hat{h}_{-\alpha}(t, a) \tag{4.8}$$

$$= {}_a\nabla_{(q,h)}^\alpha f(t) - f(\sigma(a))\hat{h}_{-\alpha}(t, a). \tag{4.9}$$

Proof Integrating by parts and using the Leibniz rule (Lemma 4.1), we have

$$\begin{aligned} & \sigma(a) \nabla_{(q,h)}^{-(1-\alpha)} \nabla_{(q,h)} f(t) \\ &= \int_{\sigma(a)}^t \hat{h}_{-\alpha}(t, \rho(\tau)) \nabla_{(q,h)} f(\tau) \nabla \tau \\ &= \left(\hat{h}_{-\alpha}(t, \tau) f(\tau) \right) \Big|_{\tau=\sigma(a)}^t + \int_{\sigma(a)}^t \hat{h}_{-\alpha-1}(t, \rho(\tau)) f(\tau) \nabla \tau \\ &= {}_{\sigma(a)} \nabla_{(q,h)}^{\alpha} f(t) - f(\sigma(a)) \hat{h}_{-\alpha}(t, \sigma(a)), \end{aligned}$$

and hence (4.7) holds. By the Leibniz rule (Lemma 4.1) we get that

$$\nabla_{(q,h)} \sigma(a) \nabla_{(q,h)}^{-(1-\alpha)} = {}_{\sigma(a)} \nabla_{(q,h)}^{\alpha} f(t). \tag{4.10}$$

Using (4.10) and (4.7), we see that (4.6) holds.

Next we will show that (4.8) holds.

$$\begin{aligned} & \nabla_{(q,h)a} \nabla_{(q,h)}^{-(1-\alpha)} f(t) \\ &= \nabla_{(q,h)} \int_a^t \hat{h}_{-\alpha}(t, \rho(s)) f(s) \nabla s \\ &= \int_a^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s + \hat{h}_{-\alpha}(\rho(t), \rho(t)) f(t) \\ &= \int_a^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s \\ &= \int_a^{\sigma(a)} \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s + \int_{\sigma(a)}^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s \\ &= \hat{h}_{-\alpha-1}(t, a) f(\sigma(a)) \nu(\sigma(a)) + \int_{\sigma(a)}^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s. \end{aligned}$$

Subtracting $f(\sigma(a)) \hat{h}_{-\alpha}(t, a)$ from both sides of this last equation we get

$$\begin{aligned} & \nabla_{(q,h)a} \nabla_{(q,h)}^{-(1-\alpha)} f(t) - f(\sigma(a)) \hat{h}_{-\alpha}(t, a) \\ &= \hat{h}_{-\alpha-1}(t, a) f(\sigma(a)) \nu(\sigma(a)) + \int_{\sigma(a)}^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s - f(\sigma(a)) \hat{h}_{-\alpha}(t, a) \\ &= \int_{\sigma(a)}^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s + f(\sigma(a)) \left(\nu(\sigma(a)) \hat{h}_{-\alpha-1}(t, a) - \hat{h}_{-\alpha}(t, a) \right) \\ &\stackrel{(4.5)}{=} \int_{\sigma(a)}^t \hat{h}_{-\alpha-1}(t, \rho(s)) f(s) \nabla s - f(\sigma(a)) \hat{h}_{-\alpha}(t, \sigma(a)) \\ &\stackrel{(4.7)}{=} {}_{\sigma(a)} \nabla_{(q,h)}^{-(1-\alpha)} \nabla_{(q,h)} f(t). \end{aligned}$$

Hence, (4.8) holds. Since by the Leibniz rule $\nabla_{(q,h)} a \nabla_{(q,h)}^{-(1-\alpha)} = a \nabla_{(q,h)}^\alpha f(t)$, we have that (4.8) implies (4.9) holds. \square

Replacing α by $1 - \alpha$ in Lemma 4.5, we get the following corollary, which will be used later.

Corollary 4.6 Assume $f : \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \rightarrow \mathbb{R}$, $0 < \alpha < 1$, $q > 1$. Then:

$$\sigma(a) \nabla_{(q,h)}^{-\alpha} \nabla_{(q,h)} f(t) = \nabla_{(q,h)} \sigma(a) \nabla_{(q,h)}^{-\alpha} f(t) - f(\sigma(a)) \hat{h}_{\alpha-1}(t, \sigma(a)), \tag{4.11}$$

$$\sigma(a) \nabla_{(q,h)}^{-\alpha} \nabla_{(q,h)} f(t) = \nabla_{(q,h)} a \nabla_{(q,h)}^{-\alpha} f(t) - f(\sigma(a)) \hat{h}_{\alpha-1}(t, a). \tag{4.12}$$

Lemma 4.7 Assume $f : \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \times \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)} \rightarrow \mathbb{R}$, $0 < \alpha < 1$. Then we have that

$$\int_a^t \int_{\rho(\xi)}^t f(\eta, \xi) \nabla \eta \nabla \xi = \int_a^t \int_a^\eta f(\eta, \xi) \nabla \xi \nabla \eta. \tag{4.13}$$

Proof Let

$$\begin{aligned} \phi(t) &:= \int_a^t \int_{\rho(\xi)}^t f(\eta, \xi) \nabla \eta \nabla \xi - \int_a^t \int_a^\eta f(\eta, \xi) \nabla \xi \nabla \eta, \\ F_1(t, \xi) &:= \int_{\rho(\xi)}^t f(\eta, \xi) \nabla \eta, \text{ and } F_2(t, \eta) := \int_a^\eta f(t, \eta) \nabla \xi. \end{aligned}$$

From Lemma 4.1, we have

$$\begin{aligned} \nabla_{(q,h)} \phi(t) &= \nabla_{(q,h)} \int_a^t F_1(t, \xi) \nabla \xi - \nabla_{(q,h)} \int_a^t F_2(t, \eta) \nabla \eta \\ &= \int_a^t \nabla_{(q,h)} F_1(t, \xi) \nabla \xi + F_1(\rho(t), t) - \int_a^t f(t, \xi) \nabla \xi \\ &= \int_a^t \nabla_{(q,h)} F_1(t, \xi) \nabla \xi + \int_{\rho(t)}^{\rho(t)} f(\eta, t) \nabla \eta - \int_a^t f(t, \xi) \nabla \xi \\ &= \int_a^t f(t, \xi) \nabla \xi - \int_a^t f(t, \xi) \nabla \xi = 0, \end{aligned}$$

and since $\phi(a) = 0$, we see that $\phi(t) \equiv 0$. This completes the proof. \square

The following lemma is from [10, page 14].

Lemma 4.8 Let $\alpha > 0$, $\beta \in \mathbb{R}$ and $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. Then

$$a \nabla_{(q,h)}^{-\alpha} \hat{h}_\beta(t, a) = \hat{h}_{\alpha+\beta}(t, a). \tag{4.14}$$

A special case of the composition rule in the next lemma is Theorem 3.107 [18]. Also, Cermak and Nechvatal [12] pointed out that this composition rule follows from a very general formula that they proved. With the aid of Lemma 4.6 and Lemma 4.7 we now give a straightforward proof of this result for the convenience of the reader.

Lemma 4.9 Let $\alpha, \beta \in \mathbb{R}^+$ and $f : \tilde{\mathbb{T}}_{(q,h)}^a \rightarrow \mathbb{R}$. Then

$${}_a\nabla_{(q,h)}^{-\alpha} {}_a\nabla_{(q,h)}^{-\beta} f(t) = {}_a\nabla_{(q,h)}^{-(\alpha+\beta)} f(t).$$

Proof For $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$ we have that

$$\begin{aligned} {}_a\nabla_{(q,h)}^{-\alpha} {}_a\nabla_{(q,h)}^{-\beta} f(t) &\stackrel{\text{Def. 2.1}}{=} \int_a^t \hat{h}_{\alpha-1}(t, \rho(s)) {}_a\nabla_{(q,h)}^{-\beta} f(s) \nabla s \\ &= \int_a^t \hat{h}_{\alpha-1}(t, \rho(s)) \int_a^s \hat{h}_{\beta-1}(s, \rho(\tau)) f(\tau) \nabla \tau \nabla s \\ &= \int_a^t \int_a^s \left(\hat{h}_{\alpha-1}(t, \rho(s)) \hat{h}_{\beta-1}(s, \rho(\tau)) f(\tau) \right) \nabla \tau \nabla s \\ &\stackrel{(4.13)}{=} \int_a^t \int_{\rho(\tau)}^t \left(\hat{h}_{\alpha-1}(t, \rho(s)) \hat{h}_{\beta-1}(s, \rho(\tau)) f(\tau) \right) \nabla s \nabla \tau \\ &= \int_a^t f(\tau) \int_{\rho(\tau)}^t \left(\hat{h}_{\alpha-1}(t, \rho(s)) \hat{h}_{\beta-1}(s, \rho(\tau)) \right) \nabla s \nabla \tau \\ &= \int_a^t f(\tau) \left(\rho(\tau) \nabla_{(q,h)}^{-\alpha} \hat{h}_{\beta-1}(t, \rho(\tau)) \right) \nabla \tau \\ &\stackrel{(4.14)}{=} \int_a^t f(\tau) \hat{h}_{\alpha+\beta-1}(t, \rho(\tau)) \nabla \tau \\ &= {}_a\nabla_{(q,h)}^{-(\alpha+\beta)} f(t). \end{aligned}$$

Also,

$${}_a\nabla_{(q,h)}^{-\alpha} {}_a\nabla_{(q,h)}^{-\beta} f(a) = {}_a\nabla_{(q,h)}^{-(\alpha+\beta)} f(a).$$

□

Theorem B. Assume $0 < \alpha < 1$, $c(t) \leq 0$, $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$, and $x(t)$ is a solution of the equation

$${}_a\nabla_{(q,h)}^{\alpha} x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)} \tag{4.15}$$

satisfying $x(\sigma(a)) > 0$. Then

$$\lim_{t \rightarrow \infty} x(t) = 0.$$

Proof Applying the operator ${}_{\sigma(a)}\nabla_{(q,h)}^{-\alpha}$ to each side of equation (4.15), we obtain

$${}_{\sigma(a)}\nabla_{(q,h)}^{-\alpha} {}_a\nabla_{(q,h)}^{\alpha} x(t) = {}_{\sigma(a)}\nabla_{(q,h)}^{-\alpha} c(t)x(t),$$

which can be written in the form

$${}_{\sigma(a)}\nabla_{(q,h)}^{-\alpha} \nabla_{(q,h)} {}_a\nabla_{(q,h)}^{-(1-\alpha)} x(t) = {}_{\sigma(a)}\nabla_{(q,h)}^{-\alpha} c(t)x(t).$$

Using (4.12), we get that

$$\begin{aligned} & \nabla_{(q,h)} a \nabla_{(q,h)}^{-\alpha} a \nabla_{(q,h)}^{-(1-\alpha)} x(t) - \hat{h}_{\alpha-1}(t, a) \left(a \nabla_{(q,h)}^{-(1-\alpha)} x(t) \Big|_{t=\sigma(a)} \right) \\ & = \sigma(a) \nabla_{(q,h)}^{-\alpha} c(t) x(t). \end{aligned}$$

Using

$$\begin{aligned} a \nabla_{(q,h)}^{-(1-\alpha)} x(t) \Big|_{t=\sigma(a)} & = \int_a^{\sigma(a)} \hat{h}_{-\alpha}(\sigma(a), \rho(s)) x(s) \nabla s \\ & = \hat{h}_{-\alpha}(\sigma(a), a) x(\sigma(a)) \nu(\sigma(a)) \\ & = (\nu(\sigma(a)))^{-\alpha} \begin{bmatrix} -\alpha \\ 0 \end{bmatrix}_{\tilde{q}} x(\sigma(a)) \nu(\sigma(a)) \\ & = [\sigma(a)]^{1-\alpha} (1 - \tilde{q})^{1-\alpha} x(\sigma(a)), \end{aligned}$$

we see that

$$\begin{aligned} \nabla_{(q,h)} a \nabla_{(q,h)}^{-\alpha} a \nabla_{(q,h)}^{-(1-\alpha)} x(t) & = \hat{h}_{\alpha-1}(t, a) [\sigma(a)]^{1-\alpha} (1 - \tilde{q})^{1-\alpha} x(\sigma(a)) \\ & \quad + \sigma(a) \nabla_{(q,h)}^{-\alpha} c(t) x(t). \end{aligned}$$

It is easy to see for $t \geq s$, letting $t = \sigma^n(s) = \sigma^{n+1}(\rho(s))$, for $n \geq 0$. Using the composition rule (Lemma 4.9), we have

$$a \nabla_{(q,h)}^{-\alpha} a \nabla_{(q,h)}^{-(1-\alpha)} x(t) = a \nabla_{(q,h)}^{-1} x(t)$$

and

$$\begin{aligned} \nabla_{(q,h)} a \nabla_{(q,h)}^{-1} x(t) & = \nabla_{(q,h)} \int_a^t \hat{h}_0(t, \rho(s)) x(s) \nabla s \\ & = \nabla_{(q,h)} \int_a^t x(s) \nabla s \\ & = x(t), \end{aligned}$$

where we used

$$\begin{aligned} \hat{h}_0(t, \rho(s)) & = (\nu(t))^0 \begin{bmatrix} 0 + n + 1 - 1 \\ n \end{bmatrix}_{\tilde{q}} \\ & = 1. \end{aligned}$$

Hence, we get that

$$\begin{aligned} x(t) & = \hat{h}_{\alpha-1}(t, a) [\sigma(a)]^{1-\alpha} (1 - \tilde{q})^{1-\alpha} x(\sigma(a)) \\ & \quad + \int_{\sigma(a)}^t \hat{h}_{\alpha-1}(t, \rho(s)) c(s) x(s) \nabla s. \end{aligned} \tag{4.16}$$

Since $x(\sigma(a)) > 0$, $0 < \alpha < 1$, $c(t) \leq 0$, and from Theorem 4.2, we have $x(t) > 0$, for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$. Then

$$\begin{aligned} \hat{h}_{\alpha-1}(t, \rho(s)) &= (\nu(t))^{\alpha-1} \left[\begin{matrix} \alpha - 1 + n \\ n \end{matrix} \right]_{\tilde{q}} \\ &= (\nu(t))^{\alpha-1} \frac{\Gamma_{\tilde{q}}(\alpha + n)}{\Gamma_{\tilde{q}}(n)\Gamma_{\tilde{q}}(\alpha)}. \end{aligned}$$

Note that $\Gamma_{\tilde{q}}(\alpha + n) > 0$, $\Gamma_{\tilde{q}}(\alpha) > 0$, and $\Gamma_{\tilde{q}}(n + 1) > 0$, so $\hat{h}_{\alpha-1}(t, \rho(s)) > 0$. It follows that

$$\int_{\sigma(a)}^t \hat{h}_{\alpha-1}(t, \rho(s))c(s)x(s)\nabla s < 0.$$

Since $c(s) \leq 0$ and from (4.16), we get that (taking $t = \sigma^k(a)$)

$$0 < x(\sigma^k(a)) \leq \hat{h}_{\alpha-1}(\sigma^k(a), a)[\sigma(a)]^{1-\alpha}(1 - \tilde{q})^{1-\alpha}x(\sigma(a)). \tag{4.17}$$

Note that

$$\begin{aligned} \hat{h}_{\alpha-1}(\sigma^k(a), a) &= (\nu(\sigma^k(a)))^{\alpha-1} \left[\begin{matrix} \alpha - 1 + k - 1 \\ k - 1 \end{matrix} \right]_{\tilde{q}} \\ &= (\nu(\sigma^k(a)))^{\alpha-1} \frac{\Gamma_{\tilde{q}}(\alpha - 1 + k)}{\Gamma_{\tilde{q}}(k)\Gamma_{\tilde{q}}(\alpha)} \\ &= [\sigma^k(a)]^{\alpha-1}(1 - \tilde{q})^{\alpha-1} \frac{\Gamma_{\tilde{q}}(\alpha - 1 + k)}{\Gamma_{\tilde{q}}(k)\Gamma_{\tilde{q}}(\alpha)} \\ &\rightarrow 0, \end{aligned} \tag{4.18}$$

where we used

$$\begin{aligned} \lim_{k \rightarrow \infty} [\sigma^k(a)]^{\alpha-1} &= \lim_{k \rightarrow \infty} [q^k a + [k]_q h]^{\alpha-1} \\ &= \lim_{k \rightarrow \infty} \left(q^k a + [k]_q h + h \frac{\tilde{q}}{1 - \tilde{q}} \right)^{\alpha-1} \\ &= \lim_{k \rightarrow \infty} \left(q^k a + \frac{1 - q^k}{1 - q} h + \frac{h}{q - 1} \right)^{\alpha-1} \\ &= \lim_{k \rightarrow \infty} \left(q^k \left(a + \frac{h}{q - 1} \right) \right)^{\alpha-1} \\ &= 0, \end{aligned}$$

and

$$\lim_{k \rightarrow \infty} \frac{\Gamma_{\tilde{q}}(\alpha - 1 + k)}{\Gamma_{\tilde{q}}(k)} = \left(\frac{1}{1 - \tilde{q}} \right)^{\alpha-1} \quad (\text{using (3.14)}).$$

From (4.17) and (4.18), we have the desired result:

$$\lim_{k \rightarrow \infty} x(\sigma^k(a)) = 0.$$

□

Remark 4.10 (see Example 5.4) If $c(t)$ in Theorem A is not a constant, then there are solutions of the equation (4.15) satisfying $x(\sigma(a)) > 0$ that are not monotonically decreasing.

We end this section by considering solutions $x(t)$ of the α th order nabla fractional (q, h) -difference equation

$${}_a \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$$

satisfying $x(\sigma(a)) < 0$. By making the transformation $x(t) = -y(t)$ and using Theorem A, we get the following theorem.

Theorem C. Assume $0 < \alpha < 1$, $c(t) \leq 0$, $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$, and $x(t)$ is a solution of the equation

$${}_a \nabla_{(q,h)}^\alpha x(t) = c(t)x(t), \quad t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma^2(a)}$$

satisfying $x(\sigma(a)) < 0$. Then $x(t) < 0$ for $t \in \tilde{\mathbb{T}}_{(q,h)}^{\sigma(a)}$ and

$$\lim_{t \rightarrow \infty} x(t) = 0.$$

5. Examples

In this section we give four examples to illustrate Lemma 3.5, Theorem 3.15, Theorem A, Remark 3.16, Theorem 4.2, Theorem B, and Remark 4.10.

Using the recursion formula (2.4), we can get the corresponding numerical solutions through formulae (3.13) and (4.4) using MATLAB. For the sake of convenience in plotting figures, letting $x_k = x(\sigma^k(a))$, we plot the points (k, x_k) instead of the points $(t, x(t))$.

Example 5.1 Consider the initial value problem in the form

$$\begin{aligned} {}_1^C \nabla_{(1.2,0.5)}^{0.3} x(t) &= -2x(t), \quad t \in \tilde{\mathbb{T}}_{(1.2,0.5)}^{\sigma(1)}, \\ x(1) &= 0.1. \end{aligned}$$

The solution $x(t)$ is given in Figure 2. Note that $x(t)$ monotonically tends to zero as $t \rightarrow \infty$.

Example 5.2 Consider the initial value problem in the form

$$\begin{aligned} {}_1^C \nabla_{(1.2,0.5)}^{0.3} x(t) &= -(0.1 + \sin^2 t) \cdot x(t), \quad t \in \tilde{\mathbb{T}}_{(1.2,0.5)}^{\sigma(1)}, \\ x(1) &= 0.1. \end{aligned}$$

There is a constant $b = -0.1$ such that

$$c(t) = -(0.1 + \sin^2 t) \leq b.$$

It is evident that the solution $x(t)$ is asymptotically stable from Theorem A.

We plot the solution $x(t)$ in Figure 3. Note that $x(t)$ tends to zero as $t \rightarrow \infty$, but is not monotonically decreasing.




Figure 2. Asymptotic behavior of $x(t)$ for $a = 1$, $\nu = 0.3$, $q = 1.2$, $h = 0.5$, $x(a) = 0.1$, $c(t) = -2$.




Figure 3. Asymptotic behavior of $x(t)$ for $a = 1$, $\nu = 0.3$, $q = 1.2$, $h = 0.5$, $x(a) = 0.1$, $c(t) = -(0.1 + \sin^2 t)$.

Example 5.3 Consider the initial value problem in the form

$$\begin{aligned}
 {}_1\nabla_{(1.2,0.5)}^{0.7}x(t) &= -0.00001x(t), & t \in \tilde{\mathbb{T}}_{(1.2,0.5)}^{\sigma^2(1)}, \\
 x(\sigma(1)) &= 0.1.
 \end{aligned}$$

We plot the solution $x(t)$ in Figure 4. Note that $x(t)$ monotonically tends to zero as $t \rightarrow \infty$.

Example 5.4 Consider the initial value problem in the form

$$\begin{aligned}
 {}_1\nabla_{(1.2,0.5)}^{0.9}x(t) &= -\frac{1}{100} \sin^{10} t \cdot x(t), & t \in \tilde{\mathbb{T}}_{(1.2,0.5)}^{\sigma^2(1)}, \\
 x(\sigma(1)) &= 0.1.
 \end{aligned}$$




Figure 4. Asymptotic behavior of $x(t)$ for $a = 1$, $\nu = 0.7$, $q = 1.2$, $h = 0.5$, $x(\sigma(1)) = 0.1$, $c(t) = -0.00001$.




Figure 5. Asymptotic behavior of $x(t)$ for $a = 1$, $\nu = 0.9$, $q = 1.2$, $h = 0.5$, $x(\sigma(1)) = 0.1$, $c(t) = -\frac{1}{100} \sin^{10} t$.

It is easy to check that

$$c(t) = -\frac{1}{100} \sin^{10} t \leq 0.$$

Hence, the solution $x(t)$ is asymptotically stable according to Theorem B.

We plot the solution $x(t)$ in Figure 5. Note that $x(t)$ tends to zero as $t \rightarrow \infty$, but is not monotonically decreasing.

6. Conclusions

In this paper, we present two asymptotic results on linear nabla fractional difference equations originating from the recent papers as well as their new extensions on the (q, h) -time scale. Furthermore, the fractional difference equations' numerical solutions are given to support the established theories. It remains to discuss the asymptotic stability of (0.1) and (0.2) with $c(t) > 0$, which is still an open problem.

References

- [1] Abadias L, Lizama C, Miana PJ, Velasco MP. On well-posedness of vector-valued fractional differential-difference equations. arXiv: 1606.05237.
- [2] Anastassiou G. Foundations of nabla fractional calculus on time scales and inequalities. COMPUT MATH APPL 2010; 59: 3750-3762.
- [3] Atici FM, Eloe PW. Eloe. Initial value problems in discrete fractional calculus. P AM MATH SOC 2009; 137: 981-989.
- [4] Atici FM, Eloe PW. Linear systems of fractional nabla difference equations. ROCKY MT J MATH 2011; 41: 353-370.
- [5] Bai Z, Xu R. The asymptotic behavior of solutions for a class of nonlinear fractional difference equations with damping term. DISCRETE DYN NAT SOC 2018; 2018: 5232147.
- [6] Baleanu D, Wu GC, Bai YR, Chen FL. Stability analysis of Caputo-like discrete fractional systems. COMMUN NONLINEAR SCI 2017; 48: 520-530.
- [7] Bohner M, Peterson A. Dynamic Equations on Time Scales: An Introduction with Applications. Boston, MA, USA: Birkhäuser, 2001.
- [8] Bohner M, Peterson A. Advances in Dynamic Equations on Time Scales. Boston, MA, USA: Birkhäuser, 2003.
- [9] Cermak J, Gyori I, Nechvatal L. On explicit stability conditions for a linear fractional difference system. FRACT CALC APPL ANAL 2015; 18: 652-672.
- [10] Cermak J, Kisela T, Nechvatal L. Discrete Mittag-Leffler functions in linear fractional difference equations. ABSTR APPL ANAL 2011; 2011: 565067.
- [11] Cermak J, Kisela T, Nechvatal L. Stability regions for linear fractional differential systems and their discretizations. APPL MATH COMPUT 2013; 219: 7012-7022.
- [12] Cermak J, Nechvatal L. On (q, h) -analogue of fractional calculus. J NONLINEAR MATH PHY 2010; 17: 51-68.
- [13] Chen F. Fixed points and asymptotic stability of nonlinear fractional difference equations. ELECTRON J QUAL THEO 2011; 2011: 39.
- [14] Dassios IK. A practical formula of solutions for a family of linear non-autonomous fractional nabla difference equations. J COMPUT APPL MATH 2017; 339: 317-328.
- [15] Dassios IK, Baleanu DI. Duality of singular linear systems of fractional nabla difference equations. APPL MATH MODEL 2015; 39: 4180-4195.

- [16] Dassios IK, Baleanu DI, Kalogeropoulos G. On non-homogeneous singular systems of fractional nabla difference equations. *APPL MATH COMPUT* 2014; 227: 112-131.
- [17] Du FF, Jia BG, Erbe L, Peterson A. Monotonicity and convexity for nabla fractional (q, h) -differences. *J DIFFER EQU APPL* 2016; 22: 1224-1243.
- [18] Goodrich C, Peterson A. *Discrete Fractional Calculus*. New York, NY, USA: Springer, 2015.
- [19] Jarad F, Abelfawad T, Baleanu D. Stability of q -fractional non-autonomous systems. *NONLINEAR ANAL-REAL* 2013; 14: 780-784.
- [20] Jia BG, Chen SY, Erbe L, Peterson A. Liapunov functional and stability of linear nabla (q, h) -fractional difference equations. *J DIFFER EQU APPL* 2017; 23: 1974-1985.
- [21] Jia BG, Du FF, Erbe L, Peterson A. Asymptotic behavior of half order fractional difference equation. *J Appl Anal Comput*, 2018.
- [22] Jia BG, Erbe L, Peterson A. Comparison theorems and asymptotic behavior of solutions of discrete fractional equations. *ELECTRON J QUAL THEO* 2015; 2015: 89.
- [23] Jia BG, Erbe L, Peterson A. Comparison theorems and asymptotic behavior of solutions of Caputo fractional equations. *International Journal of Difference Equations* 2016; 11: 163-178.
- [24] Jia BG, Erbe L, Peterson A. Monotonicity and convexity for nabla fractional q -differences. *DYNAM SYST APPL* 2016; 25: 47-60.
- [25] Jia BG, Erbe L, Peterson A. Asymptotic behavior of solutions of fractional nabla q -difference equations. *Georgian Math J* (in press). doi: 10.1515/gmj-2017-0030.
- [26] Jiang J, Liu L. Existence of solutions for a sequential fractional differential system with coupled boundary conditions. *BOUND VALUE PROBL* 2016; 2016: 159.
- [27] Jiang J, Liu W, Wang H. Positive solutions to singular Dirichlet-type boundary value problems of nonlinear fractional differential equations. *ADV DIFFER EQU-NY* 2018; 2018: 169.
- [28] Jiang J, Liu L, Wu Y. Multiple positive solutions of singular fractional differential system involving Stieltjes integral conditions. *ELECTRON J QUAL THEO* 2012; 2012: 43.
- [29] Jiang J, Liu L, Wu Y. Positive solutions to singular fractional differential system with coupled boundary conditions. *COMMUN NONLINEAR SCI* 2013; 18: 3061-3074.
- [30] Li M, Wang J. Exploring delayed Mittag-Leffler type matrix functions to study finite time stability of fractional delay differential equations. *APPL MATH COMPUT* 2018; 324: 254-265.
- [31] Li Y, Chen YQ, Podlubny I. Stability of fractional-order nonlinear dynamic systems: Lyapunov direct method and generalized Mittag-Leffler stability. *COMPUT MATH APPL* 2010; 59: 1810-1821.
- [32] Lizama C. l_p -maximal regularity for fractional difference equations on UMD spaces. *MATH NACHR* 2016; 288: 2079-2092.
- [33] Lizama C. The Poisson distribution, abstract fractional difference equations, and stability. *P AM MATH SOC* 2017; 145: 3809-3827.
- [34] Matignon D. Stability results for fractional differential equations with applications to control processing. *MA Comput Sci Eng* 1996; 2: 963-968.
- [35] Mozyrska D, Wyrwas M. Stability by linear approximation and the relation between the stability of difference and differential fractional systems. *MATH METHOD APPL SCI* 2016; 40: 4080-4091.
- [36] Rahmat MRS. On some (q, h) -analogues of integral inequalities on discrete time scales. *COMPUT MATH APPL* 2011; 62: 1790-1797.
- [37] Rahmat MRS. The (q, h) -Laplace transform on discrete time scales. *COMPUT MATH APPL* 2011; 62: 272-281.
- [38] Rahmat MRS, Noorani MSM. Caputo type fractional difference operator and its application on discrete time scales. *ADV DIFFER EQU-NY* 2015; 2015: 160.

- [39] Roy AZD, Xavier M. Fractional Calculus: Theory. New York, NY, USA: Nova Science, 2014.
- [40] Shen T, Xin J, Huang J. Time-space fractional stochastic Ginzburg-Landau equation driven by Gaussian white noise. STOCH ANAL APPL 2018; 36: 103-113.
- [41] Wang Y, Jiang J. Existence and nonexistence of positive solutions for the fractional coupled system involving generalized p -Laplacian. ADV DIFFER EQU-NY 2017; 2017: 337.
- [42] Wang Y, Liu L. Necessary and sufficient condition for the existence of positive solution to singular fractional differential equations. ADV DIFFER EQU-NY 2015; 2015: 207.
- [43] Wang Y, Liu L, Wu Y. Positive solutions for a nonlocal fractional differential equation. NONLINEAR ANAL-THEOR 2011; 74: 3599-3605.
- [44] Wang Y, Liu L, Wu Y. Existence and uniqueness of a positive solution to singular fractional differential equations. BOUND VALUE PROBL 2012; 2012: 81.
- [45] Wong R, Beals R. Special Functions: A Graduate Text. Cambridge, UK: Cambridge University Press, 2010.
- [46] Wu GC, Baleanu D, Luo WH. Lyapunov functions for Riemann-Liouville-like fractional difference equations. APPL MATH COMPUT 2017; 314: 228-236.
- [47] Xu R, Zhang Y. Generalized Gronwall fractional summation inequalities and their applications. J INEQUAL APPL 2015; 2015: 242.