

1-1-2018

## On the summability methods of logarithmic type and the Berezin symbol

ULAŞ YAMANCI

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

YAMANCI, ULAŞ (2018) "On the summability methods of logarithmic type and the Berezin symbol," *Turkish Journal of Mathematics*: Vol. 42: No. 5, Article 25. <https://doi.org/10.3906/mat-1803-37>  
Available at: <https://dctubitak.researchcommons.org/math/vol42/iss5/25>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

## On the summability methods of logarithmic type and the Berezin symbol

Ulaş YAMANCI\* 

Department of Statistics, Faculty of Arts and Sciences, Süleyman Demirel University, Isparta, Turkey

Received: 29.03.2018

Accepted/Published Online: 05.07.2018

Final Version: 27.09.2018

**Abstract:** We prove by means of the Berezin symbols some theorems for the  $(L)$ -summability method for sequences and series. Also, we prove a new Tauberian type theorem for  $(L)$ -summability.

**Key words:**  $(L)$ -summability, Berezin symbol,  $(e)$ -convergence, compact operator, Tauberian type theorem, Dirichlet space, diagonal operator

### 1. Introduction

In this article, by applying a new functional analytic approach based on the so-called the Berezin symbol technique, we prove the following results (see [3, 4]). Also, we give a new Tauberian type theorem for  $(L)$ -summable sequences of complex numbers.

Recall that a sequence  $(a_n)_{n \geq 0}$  of complex numbers  $a_n$  is said to be summable to a finite number  $\zeta$  by the logarithmic method  $(L)$  (or  $(L)$ -summable to  $\zeta$ ) if

$$\sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}$$

converges in the open interval  $(0, 1)$  and

$$\lim_{x \rightarrow 1^-} -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1} = \zeta.$$

The series  $\sum_{n=0}^{\infty} a_n$  is  $(L)$ -summable to  $\zeta$  if the sequence of partial sums  $s := (s_n)_{n \geq 0}$  (where  $s_n = \sum_{k=0}^n a_k$ ) is  $(L)$ -summable to  $\zeta$ .

**Theorem 1** *If  $(a_k)_{k \geq 0}$  converges to  $\zeta$ , then  $(a_k)_{k \geq 0}$   $(L)$ -converges to  $\zeta$ .*

**Theorem 2** *If the series  $\sum_{k=0}^{\infty} a_k$  converges to  $\zeta$ , then  $\sum_{k=0}^{\infty} a_k$  is  $(L)$ -summable to  $\zeta$ .*

\*Correspondence: [ulasyamanci@sdu.edu.tr](mailto:ulasyamanci@sdu.edu.tr)

2010 AMS Mathematics Subject Classification: 40D09

Before beginning the presentation, we recall some basic definitions and notations.

Recall that in [6], Karaev introduced the notions of an  $(e)$ -convergent sequence and  $(e)$ -convergent series for the complex numbers as follows.

Let  $\mathcal{H} = \mathcal{H}(\Omega)$  be a reproducing kernel Hilbert space on some suitable set  $\Omega$  with reproducing kernel

$$k_{\mathcal{H},\lambda}(z) := \sum_{n=0}^{\infty} \overline{e_n(\lambda)} e_n(z), \tag{1}$$

where  $\{e_n(z)\}_{n \geq 0}$  is an orthonormal basis of  $\mathcal{H}$ . Let  $(a_n)_{n \geq 0}$  be any sequence of complex numbers.

(1) We say that the sequence  $(a_n)_{n \geq 0}$  is  $(e)$ -convergent to  $l$  if  $\sum_{n=0}^{\infty} a_n |e_n(\lambda)|^2$  is convergent for all  $\lambda \in \Omega$  and

$$\lim_{\lambda \rightarrow \zeta} \frac{1}{\sum_{n=0}^{\infty} |e_n(\lambda)|^2} \sum_{n=0}^{\infty} a_n |e_n(\lambda)|^2 = l$$

for every  $\zeta \in \partial\Omega$ .

(2) We say that the series  $\sum_{n=0}^{\infty} a_n$  is  $(e)$ -summable to  $l$  if  $\sum_{n=0}^{\infty} a_n |e_n(\lambda)|^2$  converges for all  $\lambda \in \Omega$  and

$$\lim_{\lambda \rightarrow \zeta} \sum_{n=0}^{\infty} a_n |e_n(\lambda)|^2 = l$$

for every  $\zeta \in \partial\Omega$ .

It was shown that the definition of  $(e)$ -convergence of sequence and series coincides with the classical Abel convergence and the Borel convergence of sequence and series for the Hardy space  $\mathcal{H}(\Omega) = H^2(\mathbb{D})$  and the Fock space  $\mathcal{H}(\Omega) = \mathcal{F}(\mathbb{C})$  (see more details in [2, 5], and also [7] for related problems), respectively. One of our aims in the present article is to show in detail that  $(e)$ -summability implies the classical  $(L)$ -summability for  $\mathcal{H}(\Omega) = \mathcal{D}(\mathbb{D})$ , which is the Dirichlet space of analytic functions on  $\mathbb{D}$ , and thus to show once again the universality of the  $(e)$ -summability notion.

The associated diagonal operator  $D_a$  on  $\mathcal{H}$  for any bounded sequence  $(a_n)_{n \geq 0}$  of complex numbers is defined by the formula  $D_a e_n(z) := a_n e_n(z)$ ,  $n = 0, 1, 2, \dots$ , with respect to the orthonormal basis  $(e_n(z))_{n \geq 0}$  of  $\mathcal{H}$ . An elementary calculus shows by virtue of formula (1) that

$$\tilde{D}_a(\lambda) = \frac{1}{\sum_{n=0}^{\infty} |e_n(\lambda)|^2} \sum_{n=0}^{\infty} a_n |e_n(\lambda)|^2, \quad \lambda \in \Omega. \tag{2}$$

Following Nordgren and Rosenthal [9], we say that RKHS  $\mathcal{H}(\Omega)$  is standard if the underlying set  $\Omega$  is a subset of a topological space and the boundary  $\partial\Omega$  is nonempty and has the property that  $(k_{\mathcal{H},\lambda_n})_n$  converges weakly to 0 whenever  $(\lambda_n)_n$  is a sequence in  $\Omega$  that converges to a point in  $\partial\Omega$ . The prototypical standard RKHSs are, for example, the Hardy–Hilbert space  $H^2(\mathbb{D})$ , the Bergman–Hilbert space  $L_a^2(\mathbb{D})$ , the Fock–Hilbert space  $\mathcal{F}(\mathbb{C})$ , and the Dirichlet–Hilbert space  $\mathcal{D}(\mathbb{D})$ .

Recall that [8] the Dirichlet space  $\mathcal{D}$  is the Hilbert space of analytic functions  $f = \sum_{n=0}^{\infty} a_n z^n$  on the unit disk  $\mathbb{D}$  with  $\int_{\mathbb{D}} |f'(z)|^2 dA/\pi = \sum_{n=0}^{\infty} (n+1) |a_n|^2 < \infty$ , where  $dA$  denotes the usual Lebesgue measure on  $\mathbb{D}$ .

For any bounded linear operator  $A$  on  $\mathcal{D}$ , the Berezin symbol of  $A$  is the function  $\tilde{A}$  defined by (see [1, 9])

$$\tilde{A}(\lambda) := \left\langle A\hat{k}_\lambda, \hat{k}_\lambda \right\rangle_{\mathcal{D}} \quad (\lambda \in \Omega),$$

where  $\hat{k}_\lambda = \frac{k_\lambda}{\|k_\lambda\|}$  is the normalized reproducing kernel of the space  $\mathcal{D}$ . Since the sequence  $\{z^n/\sqrt{n+1} : n \geq 0\}$  is an orthonormal basis of the Dirichlet space, the reproducing kernel of  $\mathcal{D}$  is given by formula (1).

$$k_\lambda(z) = \sum_{n=0}^{\infty} \frac{(\bar{\lambda}z)^n}{n+1} = \frac{1}{\bar{\lambda}z} \log \frac{1}{1-\bar{\lambda}z}.$$

### 2. The results

First, we characterize the  $(L)$ -summability method in terms of the Berezin symbol.

**Theorem 3** *Let  $(a_n)_{n \geq 0}$  be a bounded sequence of complex numbers, and let  $D_a$  be the diagonal operator on the Dirichlet space  $\mathcal{D}$  with diagonal elements  $a_n$ ,  $n \geq 0$ , with respect to the orthonormal basis  $\{z^n/\sqrt{n+1}\}_{n \geq 0}$  of  $\mathcal{D}$ . Then the sequence  $(a_n)_{n \geq 0}$  is  $(L)$ -summable to  $\zeta$  if and only if*

$$\lim_{x \rightarrow 1^-} \tilde{D}_a(\sqrt{x}) = \zeta.$$

**Proof** Since  $(a_n)_{n \geq 0}$  is the bounded sequence,  $D_a$  is a bounded operator on  $\mathcal{D}$ . If  $\hat{k}_\lambda$  is the normalized reproducing kernel of  $\mathcal{D}$ , then we obtain by using formula (2) for all  $\lambda \in \mathbb{D}$  that

$$\begin{aligned} \tilde{D}_a(\lambda) &= \frac{1}{\sum_{n=0}^{\infty} \frac{(|\lambda|^2)^n}{n+1}} \sum_{n=0}^{\infty} a_n \frac{(|\lambda|^2)^n}{n+1} = \frac{1}{\frac{1}{|\lambda|^2} \log \frac{1}{1-|\lambda|^2}} \sum_{n=0}^{\infty} a_n \frac{(|\lambda|^2)^n}{n+1} \\ &= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \sum_{n=0}^{\infty} a_n \frac{(|\lambda|^2)^n}{n+1} = -\frac{1}{\log(1-|\lambda|^2)} \sum_{n=0}^{\infty} a_n \frac{(|\lambda|^2)^{n+1}}{n+1}, \end{aligned}$$

and therefore  $\tilde{D}_a$  is a radial function on  $\mathbb{D}$ ; that is,  $\tilde{D}_a(\lambda) = \tilde{D}_a(|\lambda|)$ .

Let  $|\lambda|^2 = x$ . Then

$$\tilde{D}_a(\sqrt{x}) = -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}. \tag{3}$$

We therefore get the desired assertions from (3), which proves the theorem. □

Now we are ready to prove the results.

**Proof [Proof of Theorem 1]** Let us define the diagonal operator  $D_a$  on the Dirichlet space  $\mathcal{D}$  as follows:

$$D_a \frac{z^n}{\sqrt{n+1}} = a_n \frac{z^n}{\sqrt{n+1}}, \quad n = 0, 1, 2, \dots$$

Since  $(a_k)$  is the bounded sequence,  $D_a$  is a bounded operator on  $\mathcal{D}$ . Then we get (see (3)):

$$\tilde{D}_a(\sqrt{x}) = -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1}, \quad 0 < x < 1. \tag{4}$$

Thus, we have from (4):

$$\begin{aligned} -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1} &= -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} (a_n - \zeta) \frac{x^{n+1}}{n+1} \\ &\quad + \zeta \frac{-1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{x^{n+1}}{n+1} \\ &= \tilde{D}_{a_k - \zeta}(\sqrt{x}) + \zeta. \end{aligned}$$

Since  $a_k - \zeta \rightarrow 0$  as  $n \rightarrow \infty$  by the condition of the theorem, we have that  $D_{a_k - \zeta}$  is a compact operator on  $\mathcal{D}$ . Hence, its Berezin symbol vanishes on the boundary, i.e.

$$\lim_{x \rightarrow 1^-} \tilde{D}_{a_k - \zeta}(\sqrt{x}) = 0.$$

Then we conclude from the last equality

$$\lim_{x \rightarrow 1^-} -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{a_n}{n+1} x^{n+1} = \zeta,$$

which finishes the proof. □

**Proof of Theorem 2** By using the argument to prove Theorem 1, it can easily be modified to prove the equality

$$\tilde{D}_s(\sqrt{x}) = \lim_{x \rightarrow 1^-} -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{s_n}{n+1} x^{n+1}, \tag{5}$$

where  $D_s$  denotes the diagonal operator on  $\mathcal{D}$  with diagonal elements  $s_n$ ,  $n \geq 0$ . Formula (5) means that the series  $\sum_{n=0}^{\infty} \frac{s_n}{n+1} x^{n+1}$  for all  $0 < x < 1$  is convergent. On the other hand,

$$D_s = LI + D_{s-L},$$

where the diagonal operator  $D_{s-L}$  is compact, since by the hypothesis of the theorem,  $s_k - \zeta \rightarrow 0$  as  $n \rightarrow \infty$ , and hence from (5) we get

$$\begin{aligned} \lim_{x \rightarrow 1^-} -\frac{1}{\log(1-x)} \sum_{n=0}^{\infty} \frac{s_n}{n+1} x^{n+1} &= \lim_{x \rightarrow 1^-} \tilde{D}_s(\sqrt{x}) \\ &= \lim_{x \rightarrow 1^-} \left( L + \tilde{D}_{s-L}(\sqrt{x}) \right) \\ &= \zeta + \lim_{x \rightarrow 1^-} \tilde{D}_{s-\zeta}(\sqrt{x}) = \zeta, \end{aligned}$$

which means that the series  $\sum_{k=0}^{\infty} a_k$  is  $(L)$ -summable to  $\zeta$ . The theorem is proved. □

Let  $\ell_1^2$  denote the unit sphere of the sequences space  $\ell^2$  :

$$\ell_1^2 := \left\{ (x_m)_{m \geq 0} \in \ell^2 : \|(x_m)\|_{\ell^2} = 1 \right\}.$$

Now we will prove a Tauberian theorem for  $(L)$ -summable sequences of complex numbers by applying a result due to Nordgren and Rosenthal [9, Corollary 2.8], which means that an operator  $A$  on a standard RKHS  $\mathcal{H}(\Omega)$  is compact if and only if all the Berezin symbols of unitary orbits  $U^{-1}AU$ , where  $U$  is unitary on  $\mathcal{H}(\Omega)$ , of the operator  $A$  vanish on the boundary.

**Theorem 4** *Let  $(a_n)_{n \geq 0}$  be a bounded sequence of complex numbers such that  $(a_n)_{n \geq 0}$   $(L)$ -converges to  $\zeta$ . Suppose that*

$$\sum_{m=0}^{\infty} a_m \left| \sum_{n=0}^{\infty} \frac{x_m^{(n)} \lambda^n}{\sqrt{n+1}} \right|^2 = o \left( -\frac{\log(1-|\lambda|^2)}{|\lambda|^2} \right) \tag{6}$$

for every double sequence  $(x_m^{(n)})_{m,n=0}^{\infty}$  with  $(x_m^{(n)})_{m \geq 0} \in \ell_1^2$  ( $\forall n \geq 0$ ) and  $(x_m^{(n)})_{n \geq 0} \in \ell_1^2$  ( $\forall m \geq 0$ ) whenever  $\lambda$  tends to infinity. Then  $a_n \rightarrow 0$  as  $n \rightarrow \infty$ .

**Proof** Since  $(L)$ - $a_n \rightarrow \zeta$  if and only if  $(L)$ - $(a_n - \zeta) \rightarrow 0$ , we assume without loss of generality that  $\zeta = 0$ . We will use the same method as in [7] for the proof of the theorem. Let  $U : \mathcal{D} \rightarrow \mathcal{D}$  be an arbitrary unitary operator of the Dirichlet space  $\mathcal{D}$ . Then

$$U \left( \frac{z^n}{\sqrt{n+1}} \right) = \sum_{m=0}^{\infty} b_m^{(n)} \frac{z^m}{\sqrt{m+1}}$$

with  $(b_m^{(n)})_{m \geq 0} \in \ell_1^2$  for every  $n \geq 0$ . It is easy to see then that  $(b_m^{(n)})_{n \geq 0} \in \ell_1^2$  for every  $m \geq 0$ .

Then we obtain the following:

$$\begin{aligned} U^{-1} \widetilde{D_a} U(\lambda) &= \left\langle U^{-1} D_a U \widehat{k}_\lambda, \widehat{k}_\lambda \right\rangle \\ &= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \left\langle D_a U \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \frac{z^n}{\sqrt{n+1}}, U \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \frac{z^n}{\sqrt{n+1}} \right\rangle \\ &= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \left\langle D_a \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} U \left( \frac{z^n}{\sqrt{n+1}} \right), \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} U \left( \frac{z^n}{\sqrt{n+1}} \right) \right\rangle \end{aligned}$$

$$\begin{aligned}
&= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \left\langle \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} D_a \sum_{m \geq 0} b_m^{(n)} \frac{z^m}{\sqrt{m+1}}, \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \sum_{m \geq 0} b_m^{(n)} \frac{z^m}{\sqrt{m+1}} \right\rangle \\
&= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \left\langle \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \sum_{m \geq 0} b_m^{(n)} a_m \frac{z^m}{\sqrt{m+1}}, \sum_{n \geq 0} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \sum_{m \geq 0} b_m^{(n)} \frac{z^m}{\sqrt{m+1}} \right\rangle \\
&= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \left\langle \sum_{m \geq 0} a_m \left( \sum_{n \geq 0} b_m^{(n)} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \right) \frac{z^m}{\sqrt{m+1}}, \sum_{m \geq 0} \left( \sum_{n \geq 0} b_m^{(n)} \frac{\bar{\lambda}^n}{\sqrt{n+1}} \right) \frac{z^m}{\sqrt{m+1}} \right\rangle \\
&= -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \sum_{m \geq 0} a_m \left| \sum_{n \geq 0} \frac{\bar{b}_m^{(n)}}{\sqrt{n+1}} \lambda^n \right|^2,
\end{aligned}$$

and therefore

$$U^{-1} \widetilde{D_a} U(\lambda) = -\frac{|\lambda|^2}{\log(1-|\lambda|^2)} \sum_{m \geq 0} a_m \left| \sum_{n \geq 0} \frac{\bar{b}_m^{(n)}}{\sqrt{n+1}} \lambda^n \right|^2, \quad \lambda \in \mathbb{D}. \quad (7)$$

By considering condition (6), we have from the last formula (7) that  $U^{-1} \widetilde{D_a} U$  vanishes on the boundary for every unitary operator  $U \in \mathcal{B}(\mathcal{D})$ . Then, by the above mentioned result of Nordgren and Rosenthal [9, Corollary 2.8], we conclude that  $D_a$  is a compact operator on the Dirichlet Hilbert space  $\mathcal{D}$  and as a result  $\lim_{n \rightarrow \infty} a_n = 0$ , which proves the theorem.  $\square$

### Acknowledgments

This paper was written while the author was supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK-2219). The author would like to thank the anonymous referee for helping to fix some inaccuracies.

### References

- [1] Berezin FA. Covariant and contravariant symbols of operators. *Izv Akad Nauk SSSR Ser Mat* 1972; 36: 1134-1167.
- [2] Garayev MT, Gürdal M, Yamancı U. Berezin symbols and Borel summability. *Quaest Math* 2017; 40: 403-411.
- [3] Ishiguro K. On the summability methods of logarithmic type. *P Jpn Acad* 1962; 38: 703-705.
- [4] Ishiguro K. Tauberian theorems concerning the summability methods of logarithmic type. *P Jpn Acad* 1963; 39: 156-159.
- [5] Karaev MT. Functional analysis proofs of Abel's theorems. *P Am Math Soc* 2004; 132: 2327-2329.
- [6] Karaev MT.  $(e)$ -convergence and related problem. *C R Math Acad Sci Paris* 2010; 348: 1059-1062.
- [7] Karaev MT. Tauberian-type theorem for  $(e)$ -convergent sequences. *C R Math Acad Sci Paris* 2013; 351: 177-179.
- [8] Marshall D., Sundberg C. Interpolating sequences for the multipliers of the Dirichlet space. Preprint, <http://www.math.washington.edu/~marshall/preprints/preprints.html>, 1994.
- [9] Nordgren E., Rosenthal P. Boundary values of Berezin symbols. *Oper Theory Adv Appl* 1994; 73: 362-368.