

1-1-2018

On the numerical range of square matrices with coefficients in a degree 2 Galois field extension

EDOARDO BALLICO

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

BALLICO, EDOARDO (2018) "On the numerical range of square matrices with coefficients in a degree 2 Galois field extension," *Turkish Journal of Mathematics*: Vol. 42: No. 4, Article 12. <https://doi.org/10.3906/mat-1711-103>

Available at: <https://journals.tubitak.gov.tr/math/vol42/iss4/12>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On the numerical range of square matrices with coefficients in a degree 2 Galois field extension

Edoardo BALLICO* 

Department of Mathematics, University of Trento, Povo, Trentino, Italy

Received: 28.11.2017

Accepted/Published Online: 28.03.2018

Final Version: 24.07.2018

Abstract: Let L be a degree 2 Galois extension of the field K and M an $n \times n$ matrix with coefficients in L . Let $\langle , \rangle : L^n \times L^n \rightarrow L$ be the sesquilinear form associated to the involution $L \rightarrow L$ fixing K . We use \langle , \rangle to define the numerical range $\text{Num}(M)$ of M (a subset of L), extending the classical case $K = \mathbb{R}$, $L = \mathbb{C}$, and the case of a finite field introduced by Coons, Jenkins, Knowles, Luke, and Rault. There are big differences with respect to both cases for number fields and for all fields in which the image of the norm map $L \rightarrow K$ is not closed by addition, e.g., $c \in L$ may be an eigenvalue of M , but $c \notin \text{Num}(M)$. We compute $\text{Num}(M)$ in some cases, mostly with $n = 2$.

Key words: Numerical range, sesquilinear form, formally real field, number field

1. Introduction

For any integer $n > 0$ and any field L let $M_{n,n}(L)$ be the L -vector space of all $n \times n$ matrices with coefficients in L . Let K be a field and L a degree 2 Galois extension of K . Call σ the generator of the Galois group of the extension $K \hookrightarrow L$. Thus, $\sigma : L \rightarrow L$ is a field isomorphism, $\sigma^2 : L \rightarrow L$ is the identity map, and $K = \{t \in L \mid \sigma(t) = t\}$. For any $u = (u_1, \dots, u_n) \in L^n$, $v = (v_1, \dots, v_n) \in L^n$ set $\langle u, v \rangle := \sum_{i=1}^n \sigma(u_i)v_i$. The map $\langle , \rangle : L^n \times L^n \rightarrow L$ is sesquilinear, i.e. for all $u, v, w \in L^n$ and all $c \in L$ we have $\langle u+v, w \rangle = \langle u, w \rangle + \langle v, w \rangle$, $\langle u, v+w \rangle = \langle u, v \rangle + \langle u, w \rangle$, $\langle cu, w \rangle = \sigma(c)\langle u, w \rangle$, and $\langle u, cw \rangle = c\langle u, w \rangle$. Set $C_n(1) := \{u \in L^n \mid \langle u, u \rangle = 1\}$. For any $M \in M_{n,n}(L)$ set $\text{Num}(M) := \{\langle u, Mu \rangle \mid u \in C_n(1)\}$. Since $C_n(1) \neq \emptyset$, we have $\text{Num}(M) \neq \emptyset$. As in the classical case when $K = \mathbb{R}$, $L = \mathbb{C}$, and σ is the complex conjugation the subset $\text{Num}(M)$ of L is called the *numerical range* of M [4–6]. When K is a finite field the numerical range was introduced in [1, 3]. In particular [3] built a bridge between the classical case and the finite field case and at certain points we will duly quote the parts of [3] that we adapt to our set-up.

Assume for the moment $L = K(i)$ with $K \subset \mathbb{R}$ and σ the complex conjugation. In this case, calling $\text{Num}(M)_{\mathbb{C}} \subset \mathbb{C}$ the usual numerical range of M , we have $\text{Num}(M) \subseteq \text{Num}(M)_{\mathbb{C}}$ and hence $\text{Num}(M)$ is a bounded subset of \mathbb{C} . However, even in this case there are many differences, in particular as for number fields not every element of K is a square. The main differences come from the structures of the sets Δ and Δ_n defined below.

Let $\Delta \subseteq K$ be the image of the norm map $\text{Norm}_{L/K} : L \rightarrow K$, i.e. set $\Delta := \{a\sigma(a) \mid a \in L\} \subseteq K$.

*Correspondence: ballico@science.unitn.it

2010 AMS Mathematics Subject Classification: 15A33, 15A60, 12D99, 12F99
The author was partially supported by MIUR and GNSAGA of INdAM Italy.

If $a \in K$, then $\sigma(a) = a$ and hence $\text{Norm}_{L/K}(a) = a^2$. Thus, Δ contains all squares of elements of K . In particular, $0 \in \Delta$ and $1 \in \Delta$. Since the norm map $\text{Norm}_{L/K}$ is multiplicative, Δ is closed under multiplication. If $c \in \hat{\Delta} := \Delta \setminus \{0\}$, say $c = \sigma(a)a$ for some $a \in L \setminus \{0\}$, then $1/c = \sigma(a^{-1})a^{-1}$ and hence $\hat{\Delta}$ is a multiplicative group. For any integer $n > 0$ let Δ_n be the set of all sums of n elements of Δ . If $K = \mathbb{R}$, then $\Delta = \Delta_n = \mathbb{R}_{\geq 0}$ for all $n \geq 1$. If $K = \mathbb{F}_q$ is a finite field, then $\Delta = \mathbb{F}_q$, because in this case the norm map is surjective ([1, Remark 3]); hence, $K = \Delta = \Delta_n$ if K is a finite field. If $K = \mathbb{Q}$ and $L = \mathbb{Q}(i)$, then $\Delta \subsetneq \Delta_2$ (Example 2).

For any $\delta \in \Delta_n$ set $C_n(\delta) := \{u \in L^n \mid \langle u, u \rangle = \delta\}$. We have $L^n = \sqcup_{\delta \in \Delta_n} C_n(\delta)$ and $C_n(\delta) \neq \emptyset$ for all $\delta \in \Delta_n$.

For any $M = (m_{ij}) \in M_{n,n}(L)$ let M^\dagger be the matrix $M^\dagger = (n_{ij})$ with $n_{ij} = \sigma(m_{ij})$ for all i, j . We have $(M^\dagger)^\dagger = M$ and $\langle u, Mv \rangle = \langle M^\dagger u, v \rangle$ for all $u, v \in L^n$. We say that M is unitary if $M^\dagger M = \mathbb{I}_{n,n}$ (where $\mathbb{I}_{n,n}$ is the identity $n \times n$ -matrix), i.e. if $M^\dagger = M^{-1}$. For any $U, M \in M_{n,n}(L)$ with U unitary, we have $\text{Num}(U^\dagger M U) = \text{Num}(M)$. In the case $n = 1$, say $M = (m_{11})$, we have $\text{Num}(M) = \{m_{11}\}$. We have $\text{Num}(c\mathbb{I}_{n,n}) = \{c\}$ for every $c \in L$. For any $\mu \in L$ and $c \in \Delta$, the *circle with center μ and squared-radius c* is the set of all $z \in L$ such that $\sigma(z - \mu)(z - \mu) = c$. This set is never empty, since it contains the points $\mu + b$, where $b \in L$ is such that $\sigma(b)b = c$ (two points, b and $-b$, if $\text{char}(K) \neq 2$ and $b \neq 0$). If $c = 0$, then the circle is just $\{\mu\}$, the center. If $c \in \hat{\Delta}$, then $b \neq 0$ and hence (assuming $\text{char}(K) \neq 2$) this circle has at least two points, $\mu + b$ and $\mu - b$. Hence, if $c \neq 0$, this circle is a smooth conic and (if K is infinite) it contains infinitely many points (Lemma 1 and, if $\text{char}(K) = 2$, Example 4). See Section 2 for more and in particular for its description if $K = \mathbb{Q}$ and so L is a quadratic number field.

For any integer $n > 0$ let $\hat{\Delta}_n$ denote the sum of n elements of $\hat{\Delta}$. Note that $0 \in \hat{\Delta}_2$ if and only if there is $a \in \hat{\Delta}$ with $-a \in \hat{\Delta}$. In the case $n = 1$ each matrix is a diagonal matrix and each numerical range is a singleton. The case $n > 1$ is more complicated and interesting. We prove the following results.

Proposition 1 *Assume $\text{char}(K) = 0$. If $M \in M_{n,n}(L)$ and $\text{Num}(M) = \{c\}$ for some $c \in L$, then $M = c\mathbb{I}_{n \times n}$.*

In the classical case any eigenvalue of $M \in M_{n,n}(\mathbb{C})$ is in its numerical range. When either $0 \in \hat{\Delta}_2$ or $\Delta_2 \neq \Delta$, then this is not always the case, as shown by Theorems 1 and 2.

Theorem 1 *Assume $0 \in \hat{\Delta}_2$ and take $c \in L$ and $\mu \in L^*$. Then there is $M \in M_{2,2}(L)$ with c an eigenvalue of M , $c \notin \text{Num}(M)$, and $\text{Num}(M) = c + \mu\hat{\Delta}$.*

See Proposition 6 for a description of the matrices M giving Theorem 1. We have $0 \in \hat{\Delta}_2$ for some real quadratic number fields (Lemma 5).

If M has an eigenvalue a with eigenvector u with $\langle u, u \rangle \in \hat{\Delta}$, then $a \in \text{Num}(M)$ (Remark 6).

Part (a) of the following result is an adaptation of [3, Theorem 1.2 (c)].

Theorem 2 *Assume $n = 2$ and that M has a unique eigenvalue, c . Assume that c has an eigenvector v with $\delta := \langle v, v \rangle \neq 0$ and that $M \neq c\mathbb{I}_{2,2}$.*

(a) $c \in \text{Num}(M)$ if and only if $\delta \in \Delta$.

(b) Assume $\delta \in \Delta$. There is $\mu \in L^*$ such that $(\text{Num}(M) - c)/\mu$ is the union of $\{0\}$ and all all circles $C(k(1 - k), 0)$ with $k \in \hat{\Delta} \cap (1 - \hat{\Delta})$.

For any $\delta \in \Delta_2 \setminus \{0\}$ and $v \in L^2$ with $\langle v, v \rangle = \delta$ the set of all M as in Theorem 2 is exactly the matrices M such that $\text{Ker}(M - c\mathbb{I}_{2 \times 2}) = \text{Im}(M) = Lv$.

In Section 5 we consider the case $M \in M_{n,n}(K)$. Set

$$C_n(1, K) := \{(x_1, \dots, x_n) \in K^n \mid x_1^2 + \dots + x_n^2 = 1\}.$$

Note that $C_n(1, K) := C_n(1) \cap K^n$. Note that $C_n(1, K) \neq \emptyset$ (e.g., take $x_i = 1$ and $x_j = 0$ for all $j \neq i$). The K -numerical range $\text{Num}(M)_K$ of M is the set of all $\langle u, Mu \rangle$ with $u \in C_n(1, K)$. We have $\text{Num}(M)_K \subseteq K$. The case $\text{char}(K) = 2$ is quite different from (and easier than) the case $\text{char}(K) \neq 2$.

Proposition 2 *Assume $\text{char}(K) \neq 2$. Take $M \in M_{n,n}(K)$, $n > 1$. We have $\text{Num}(M)_K = \{c\}$ if and only if the matrix $M - c\mathbb{I}_{n \times n}$ is antisymmetric.*

Proposition 3 *Assume $\text{char}(K) = 2$ and take $M = (m_{ij}) \in M_{n,n}(K)$.*

(a) *We have $\text{Num}(M)_K = \{c\}$ for some $c \in K$ if and only if $m_{ii} = c$ for all i and $m_{ij} = m_{ji}$ for all $i \neq j$.*

(b) *If $\#\text{Num}(M)_K \neq 1$ and K is infinite, then $\text{Num}(M)_K$ and K have the same cardinality.*

2. Circles

Let \bar{L} be an algebraic closure of L . In this section we assume that K is infinite and that $\text{char}(K) \neq 2$ (see Example 4 for the case $\text{char}(K) = 2$). With these assumptions there is $\alpha \in K$, which is not a square and with $L = K(\sqrt{\alpha})$. Fix $\beta \in L$ such that $\beta^2 = \alpha$. In \bar{L} the equation $t^2 = \alpha$ has β and $-\beta$ as its only solutions. L is a 2-dimensional K vector space over K with 1 and β as its basis. Hence, for any $z \in L$ there are uniquely determined $x, y \in K$ such that $z = x + y\beta$. Since $\sigma(\beta) = -\beta$ and $\sigma(t) = t$ for every $t \in K$, we have $\sigma(z) = x - y\beta$ and hence $\sigma(z)z = x^2 - y^2\alpha$. Take $k, \mu \in L$. The map $z \mapsto z - \mu$ induces a bijection between the set $\{z \in L \mid \sigma(z - \mu)(z - \mu) = k\}$ and the set $G(k, 0) := \{z \in L \mid \sigma(z)z = k\}$. Hence, it is sufficient to study the circles with center $0 \in L$. By the definition of Δ , if $k \notin \Delta$, then $G(k, 0) = \emptyset$, while if $k \in \Delta$ we have $G(k, 0) \neq \emptyset$. We have $G(0, 0) = \{0\}$, because $\sigma(z)z = 0$ if and only if $z = 0$. Write $z = x + y\beta$ and hence $\sigma(z) = x - y\beta$ and $\sigma(z)z = x^2 - \alpha y^2$. Thus, $G(k, 0) = \{(x, y) \in K^2 \mid x^2 - \alpha y^2 = k\}$. Now assume $k \in \hat{\Delta} = \Delta \setminus \{0\}$. Write $k = \sigma(c)c$ for some $c \in L^*$. Note that $\sigma(z)z = c$ if and only either $z = c$ or $z = -c$. Since $\text{char}(K) \neq 2$, the set $G(k, 0)$ contains at least two points, $-c$ and c .

Lemma 1 *If $k \in \hat{\Delta}$ the circle $G(k, 0) \subset L = K^2$ is a smooth affine conic over K . If K is infinite, then $G(k, 0)$ and K have the same cardinality.*

Proof Write $k = \sigma(c)c$ for some $c \in L^*$. We saw that $G(k, 0)$ contains the points c and $-c$ and in particular $G(k, 0) \neq \emptyset$. See x, y, z as homogeneous variables of $\mathbb{P}^2(K)$, with the line $\ell_\infty = \{z = 0\}$ as the set $\mathbb{P}^2(K) \setminus K^2$. Let $D(k, 0) \subset \mathbb{P}^2(K)$ be the conic with $g(x, y, z) := x^2 - \alpha y^2 - kz^2$ as its equation. The linear forms $2x$, $-2\alpha y$, and $-2kz$ are the partial derivatives of $g(x, y, z)$. Set $D(k, 0)_{\bar{K}} := \{(x : y : z) \in \mathbb{P}^2(\bar{L}) \mid g(x, y, z) = 0\}$. Since $\alpha \neq 0$, $k \neq 0$ and $\text{char}(K) \neq 2$, the partial derivatives of $g(x, y, z)$ have no common zero in $\mathbb{P}^2(\bar{L})$. Thus, $g(x, y, z)$ is irreducible and $D(k, 0)_{\bar{L}}$ is a smooth conic. Hence, $D(k, 0)$ is a smooth conic defined over K . Since

$D(k, 0)$ has a K -point, c , $D(k, 0)$ is isomorphic to \mathbb{P}_K^1 (use the linear projection from c) and in particular (for infinite K), K and $D(k, 0)$ have the same cardinality. The set $D(k, 0) \cap \ell_\infty$ has at most two points, because $g(x, y, z)$ is irreducible and so ℓ_∞ is not a component of $D(k, 0)$. Thus, (since K is infinite) $G(k, 0)$ and K have the same cardinality. \square

Remark 1 Take ℓ_∞ , $D(k, 0)$, and $g(x, y, z) := x^2 - \alpha y^2 - kz^2$ as in the proof of Lemma 1. We saw that $G(k, 0) = D(k, 0) \setminus \ell_\infty \cap D(k, 0)$. Here we check that $\ell_\infty \cap D(k, 0) = \emptyset$, i.e. $G(k, 0) = D(k, 0)$. We have $\ell_\infty \cap D(k, 0) = \{(x : y : 0) \in \mathbb{P}^2(K) \mid x^2 - \alpha y^2 = 0\}$. Since α is not a square in K , if $(x, y) \in K^2$ and $x^2 = \alpha y^2$, then $x = y = 0$.

Example 1 Take $K = \mathbb{Q}$. Hence, L is a quadratic number field. There is a unique square-free integer $d \notin \{0, 1\}$ such that $L = \mathbb{Q}(\sqrt{d})$ ([7, Ch 13, §1]). Take $k \in \hat{\Delta}$. If $d > 0$, then $G(k, 0)$ is a hyperbola with infinitely many points and it is unbounded. If $d < 0$, then $G(k, 0)$ is an ellipsis and in particular it is bounded; hence, each Δ_n is bounded.

3. Lemmas and examples

For any field F set $F^* := F \setminus \{0\}$. Let $e_1 = (1, 0, \dots, 0), \dots, e_n = (0, \dots, 0, 1)$ be the standard basis of L^n . For any $M \in M_{n,n}(L)$ let $\text{Num}_0(M) \subseteq L$ be the union of all $\langle u, Mu \rangle$ with $\langle u, u \rangle = 0$.

Remark 2 Take $M = (m_{ij}) \in M_{n,n}(L)$. Since $m_{ii} = \langle e_i, Me_i \rangle$, all diagonal elements of M are contained in $\text{Num}(M)$.

Remark 3 Fix $\delta \in \Delta_n$ and $a \in \Delta \setminus \{0\}$. Take $b \in L$ such that $a = b\sigma(b)$. For any $u \in L^n$ we have $\langle bu, bu \rangle = a\langle u, u \rangle$ and hence $C_n(a\delta) = bC_n(\delta)$.

Remark 4 Since $\sigma(x) = x$ for all $x \in K$, Δ contains all squares in K .

Remark 5 For any $M \in M_{n,n}(L)$ and any $c, d \in L$ we have $\text{Num}(cM + d\mathbb{I}_{n \times n}) = d + c\text{Num}(M)$.

Lemma 2 Fix $c \in \Delta_n \setminus \{0\}$. Then $1/c \in \Delta_n$.

Proof If $c = \sigma(a_1)a_1 + \dots + \sigma(a_n)a_n$ with $a_i \in K$, then $1/c = \sigma(a_1/c)a_1/c + \dots + \sigma(a_n/c)a_n/c$. \square

Lemma 3 For any $M \in M_{n,n}(L)$ we have $\text{Num}(M^\dagger) = \sigma(\text{Num}(M))$.

Proof For any $u \in C_n(1)$ we have $\langle u, Mu \rangle = \langle M^\dagger u, u \rangle = \sigma(\langle u, M^\dagger u \rangle)$. \square

Lemma 4 Fix $u \in L^n$ and assume $\delta := \langle u, u \rangle \neq 0$. There is $t \in L^*$ such that $\langle tu, tu \rangle = 1$ if and only if $\delta \in \Delta$.

Proof First assume the existence of $t \in L^*$ such that $\langle tu, tu \rangle = 1$. We have $\langle tu, tu \rangle = \sigma(t)t\delta$. Since $t \neq 0$, $\sigma(t)t \in \hat{\Delta}$. Remarks 4 and 2 give $\delta \in \Delta$. Now assume $\delta \in \Delta$. Since $\delta \neq 0$, we have $1/\delta \in \hat{\Delta}$ (Remark 2). Write $1/\delta = \sigma(t)t$ for some $t \in L^*$. We have $\langle tu, tu \rangle = 1$. \square

Remark 6 Take $M \in M_{n,n}(L)$ with an eigenvector v (say $Mv = cv$) such that $\langle v, v \rangle \in \hat{\Delta}$. Lemma 4 gives $c \in \text{Num}(M)$:

Lemma 5 Assume $\text{char}(K) \neq 2$ and take $L = K(\sqrt{\alpha})$ with α not a square in K , but α the sum of two squares in K . Then $0 \in \hat{\Delta}_2$.

Proof Note that $0 \in \hat{\Delta}_2$ if and only if there is $a \in \hat{\Delta}$ with $-a \in \hat{\Delta}$. Write $\alpha = u^2 + v^2$ with $u, v \in K^*$. Take $a := u^2 = -(v^2 - \alpha)$. □

Lemma 6 Fix integers $n > m > n/2 > 1$ and assume $\Delta_n = \Delta$. Let $M \subset L^n$ be an m -dimensional L -linear subspace. Then there are $f_1, \dots, f_{3m-2n} \in M$ such that $\langle f_i, f_i \rangle = 1$ for all i and $\langle f_i, f_j \rangle = 0$ for all $i \neq j$.

Proof Take any basis u_1, \dots, u_m of M and complete it to a basis u_1, \dots, u_n of L^n . Since the sesquilinear form $\langle \cdot, \cdot \rangle$ is nondegenerate, the matrix $E = (a_{ij})$ with $a_{ij} = \langle u_i, u_j \rangle$ has rank n . Hence, among the first m rows of E , at least $2m - n$ are linearly independent. Hence, the $m \times m$ matrix (a_{ij}) , $i, j = 1, \dots, m$, has rank at least $3m - 2n$. Permuting u_1, \dots, u_m we may assume that the matrix (a_{ij}) , $i, j = 1, \dots, 3m - 2n$, is nonsingular. Let $W \subset M$ be the linear span of u_1, \dots, u_m . Since the matrix (a_{ij}) , $i, j = 1, \dots, 3m - 2n$, is nonsingular, the restriction $\langle \cdot, \cdot \rangle_W$ of $\langle \cdot, \cdot \rangle$ to W is nondegenerate. Hence, there is $g_1 \in W$ with $\langle g_1, g_1 \rangle \neq 0$. Since $\Delta_n = \Delta$, there is $t \in L$ such that $\langle tg_1, tg_1 \rangle = 1$ (Lemma 4). Set $f_1 := tg_1$. If $3m - 2n > 1$ set $W_1 := \{w \in W \mid \langle f_1, w \rangle = 0\}$. W_1 is a codimension 1 linear subspace of W and the restriction of $\langle \cdot, \cdot \rangle$ to W_1 is nondegenerate. Therefore, there is $g_2 \in W_1$ with $\langle g_2, g_2 \rangle \neq 0$. Take $z \in L$ such that $\langle zg_2, zg_2 \rangle = 1$ (Lemma 4) and set $f_2 := zg_2$. If $3m - 2n > 2$ set $W_2 := \{w \in W_1 \mid \langle f_2, w \rangle = 0\}$ and continue in the same way. □

The definitions of numerical range and of unitary direct sum immediately give the following lemma.

Lemma 7 Fix integers $n > x > 0$, $A \in M_{x,x}(L)$, and $B \in M_{n-x,n-x}(L)$. Set $M := A \oplus B \in M_{n,n}(L)$ (unitary direct sum). The set $\text{Num}(M)$ is the union of all points $a + b$ of L with $a = \langle u, Bu \rangle$, $b = \langle v, Av \rangle$, $u \in L^x$, $v \in L^{n-x}$, and $\langle u, u \rangle + \langle v, v \rangle = 1$.

When $\Delta = \Delta_n$ we may improve Lemma 7 in the following way.

Proposition 4 Fix integers $n > x > 0$, $A \in M_{x,x}(L)$, and $B \in M_{n-x,n-x}(L)$. Set $M := A \oplus B \in M_{n,n}(L)$ (unitary direct sum). Assume $\Delta = \Delta_x = \Delta_{n-x}$. Then $\text{Num}(M)$ is the union of $\{\text{Num}_0(A) + \text{Num}(B)\} \cup \{\text{Num}(A) + \text{Num}_0(B)\}$ and all $tc + (1 - t)d$ with $t \in \hat{\Delta} \cap (1 - \hat{\Delta})$, $c \in \text{Num}(A)$, and $d \in \text{Num}(B)$.

Proof Fix $u \in L^x$, $v \in L^{n-x}$ with $\langle u, u \rangle + \langle v, v \rangle = 1$. Set $t = \langle u, u \rangle$. Hence, $\langle v, v \rangle = 1 - t$. If $t = 0$ (resp. $t = 1$), then $1 - t = 1$ (resp. $1 - t = 0$) and hence $\{\text{Num}(A)_0 + \text{Num}(B)\} \cup \{\text{Num}(A) + \text{Num}(B)_0\} \subseteq \text{Num}(M)$. Now assume $t \in \Delta \setminus \{0, 1\}$. Since $\Delta = \Delta_x = \Delta_{n-x}$, we have $t \in \hat{\Delta} \cap (1 - \hat{\Delta})$. Since $\{t, 1 - t\} \subset \hat{\Delta}$, there are $c, d \in L^*$ with $\sigma(c)c = 1/t$ and $\sigma(d)d = 1/(1 - t)$. Set $\alpha := \langle u, Au \rangle$ and $\beta := \langle v, Bv \rangle$. We have $\langle cu, cu \rangle = \langle dv, dv \rangle = 1$ and hence $\langle cu, cAu \rangle \in \text{Num}(A)$ and $\langle dv, dBv \rangle \in \text{Num}(B)$. Thus, $\alpha/t \in \text{Num}(A)$ and $\beta/(1 - t) \in \text{Num}(B)$. We get $\langle u + v, M(u + v) \rangle = tx + (1 - t)y$ with $x \in \text{Num}(A)$ and $y \in \text{Num}(B)$. The same proof done backwards gives the other inclusion. □

Proposition 4 is analogous to [3, Proposition 3.1]. Fix $c, d \in L$. The set of all $tc + (1 - t)d$ with $t \in (\hat{\Delta} \cap (1 - \hat{\Delta}))$ is called in [3] the open segment with c and d as its boundary points and we denote it

with $((c; d))$. When (as in the case of $\text{char}(L) = 0$) the set $(\hat{\Delta} \cap (1 - \hat{\Delta}))$ is nonempty (Lemma 8) we have $((c; c)) = \{c\}$ for all $c \in L$.

Lemma 8 *Assume $\text{char}(K) = 0$. Then $\hat{\Delta} \cap (1 - \hat{\Delta})$ is infinite.*

Proof For any $\delta \in \Delta$ there are x, y in K such that $x^2 - \alpha y^2 = \delta$. Note that $1 - \delta \in \Delta$ if and only if there are $w, z \in K$ such that $1 - \delta = w^2 - \alpha z^2$. Take coordinates (x, y, w, z) on K^4 . Set $T := \{(x, y, w, z) \in K^4 \mid x^2 + w^2 - \alpha(y^2 + z^2) = 1\}$. We take homogeneous coordinates x, y, w, z, t in $\mathbb{P}^4(K)$ with $H_\infty = \{t = 0\}$ and $K^4 = \mathbb{P}^4(K) \setminus H_\infty$. Let $E \subset \mathbb{P}^4(K)$ the projective quadric with equation $\{x^2 + w^2 - \alpha(y^2 + z^2) - t^2 = 0\}$. We have $E \setminus E \cap H_\infty = T$. Since $\text{char}(K) \neq 2$ and $\alpha \neq 0$, taking the partial derivatives of the polynomial $x^2 + w^2 - \alpha(y^2 + z^2) - t^2$ we get that the point $O := (1 : 0 : 0 : 0 : 1)$ is a smooth point of T . Let $M \subset \mathbb{P}^4$ be the hyperplane with equation $x - t = 0$. Note that M is the tangent space to E at O . Hence, $E \cap M$ is a quadric cone of M , which is the union of all lines of \mathbb{P}^4 contained in E and passing through O . Let $H \subset \mathbb{P}^4(K)$ be any hyperplane defined over K and with $O \notin H$. The latter condition implies $H \neq M$ and hence $N := H \cap M$ is a codimension two linear subspace of \mathbb{P}^4 . Let $\ell : \mathbb{P}^4 \setminus \{O\} \rightarrow H$ denote the linear projection from O . The morphism ℓ is defined over K , because O and H are defined over K . Hence, for each $P \in H(K)$ the line $L(O, P)$ spanned by O and P is defined over K . Since $O \in T$, the intersection $T \cap L(O, P)$ is either O with multiplicity 2 or the entire line $L(O, P)$ or the union of O and another point $O_P \in E$ defined over K . The first two cases imply $L(O, P) \subset M$. Since $O \in M$ and $O \notin H$, we have $L(O, P) \subset M$ if and only if $P \in N$. Hence, whenever we take $P \in H \setminus N$ the point $O_P \in E \setminus \{O\}$ is defined over K . Since $H \setminus N$ is 3-dimensional affine space over K , we get that E is infinite. $E \setminus T = E \cap H_\infty$. We have $O \notin H_\infty$ and hence $O \notin H_\infty \cap E$. Thus, $\ell(H_\infty \cap E)$ is a quadric hypersurface of M . If $P \in M \setminus (\ell(H_\infty \cap E))$, then $O_P \in T$. $\ell(H_\infty \cap E) \cup N$ is the union of a quadric and a hyperplane of M . Since K is infinite, the Grassmannian of all lines of $M(\bar{L})$ defined over K is Zariski dense in the Grassmannian of all \bar{L} -lines of $M(\bar{L})$. Since K is infinite, restricting to lines defined over K and contained neither in $\ell(H_\infty \cap E)$ nor in N we get that $M \setminus (N \cup \ell(E \cap H_\infty))$ is infinite. Hence, E is infinite.

(a) Assume that L has a field embedding $j : L \hookrightarrow \mathbb{C}$. We omit j and hence see L as a subfield of \mathbb{C} .

First assume that K is dense in \mathbb{C} with respect to the euclidean topology. Hence, K^4 (resp. $\mathbb{P}^4(K)$) is dense in \mathbb{C}^n (resp. $\mathbb{P}^4(\mathbb{C})$) for the euclidean topology. The topological space $N(\mathbb{C})$ is the closure of N in $N(\mathbb{C})$ with respect to the euclidean topology. Since $E \cap H_\infty$ has corank 1 with vertex $O \in \mathbb{P}^4(K)$, the closure of $E \cap H_\infty$ in the euclidean topology contains a neighborhood of O in $(E \cap H_\infty)(\mathbb{C})$. Since $(E \cap H_\infty)(\mathbb{C})$ is a cone with vertex O , it is the closure of $E \cap H_\infty$ for the euclidean topology and $(\ell(H_\infty \cap E))(\mathbb{C})$ is the closure of $\ell(H_\infty \cap E)$ for the euclidean topology. $\ell(H_\infty \cap E) \cup N$ is the union of a quadric and a hyperplane of M . We get that $E(\mathbb{C})$ is the closure of E with respect to the euclidean topology. $\hat{\Delta} \cap (1 - \hat{\Delta})$ is infinite if and only if $\Delta \cap (1 - \Delta)$ is infinite. Assume that $\Delta \cap (1 - \Delta)$ is finite, say $\Delta \cap (1 - \Delta) = \{a_1, \dots, a_s\}$ with $a_i \in K$. Set $G_i := C(a_i, 0)$ and $F_i := C(1 - a_i, 0)$. We get $E = \cup_{i=1}^s G_i \times F_i$. Hence, $\cup_{i=1}^s G_i(\mathbb{C}) \times F_i(\mathbb{C})$ is dense in $E(\mathbb{C})$ for the euclidean topology. Since $E(\mathbb{C})$ has complex dimension 3, while each $G_i(\mathbb{C}) \times F_i(\mathbb{C})$ has complex dimension 2, we get a contradiction.

Now assume that K is not dense in \mathbb{C} for the euclidean topology. Since \mathbb{Q} is dense in \mathbb{R} for the euclidean topology, the closure of K for the euclidean topology contains \mathbb{R} . Since this closure is a field, \mathbb{R} is the closure of K for the euclidean topology. We use $E(\mathbb{R})$ instead of $E(\mathbb{C})$. Since O is a smooth point of E and $O \in E(\mathbb{R})$,

$E(\mathbb{R})$ is a nonempty topological manifold of dimension 3. Hence, $E(\mathbb{R})$ cannot be the union of finitely many topological 2-manifolds $G_i(\mathbb{R}) \times F_i(\mathbb{R})$.

(b) By a theorem of Steinitz two algebraically closed fields with characteristic zero are isomorphic if and only if they have transcendental basis over \mathbb{Q} with the same cardinality [8, Theorem VIII.1.1], [10, page 125]. There are real closed fields with a transcendental basis over \mathbb{Q} with arbitrary cardinality (use that every ordered field has a real closure [2, Theorem 1.3.2] and give an ordering of $\mathbb{Q}(t_\alpha)_{\alpha \in \Gamma}$ with Γ a well-ordered set and t_α bigger than any rational function in the variable t_γ , $\gamma < \alpha$) and for any real closed field \mathbb{R} the field $\mathbb{R}(i)$ is algebraically closed [2, Theorem 1.2.2]. Hence, there is an embedding $j : L \hookrightarrow \mathbb{R}(i)$ for some real closed field \mathbb{R} . The euclidean topology on \mathbb{R}^n is the topology for which open balls form a basis of open subsets [2, Definition 2.19]. The field $\mathbb{C} := \mathbb{R}(i)$ inherits the euclidean topology. The sets \mathbb{R}^n , \mathbb{C}^n , $T(\mathbb{R})$, $T(\mathbb{C})$, $\mathbb{P}^r(\mathbb{R})$, $\mathbb{P}^r(\mathbb{C})$, $E(\mathbb{R})$, and $E(\mathbb{C})$ have the euclidean topology. Repeat the proof in step (a) with \mathbb{R} and \mathbb{C} instead of \mathbb{R} and \mathbb{C} . \square

Remark 7 Assume $\text{char}(K) = 0$. Lemma 8 says that $C_2(1)$ is infinite. Hence, $C_n(1)$ is infinite for all $n \geq 2$.

We recall that a field F is said to be *formally real* if -1 is not a sum of squares of elements of F . If F is formally real, then $\text{char}(F) = 0$.

Proposition 5 Assume that K is formally real but that L is not formally real. Then $0 \notin \hat{\Delta}_n$ for any $n > 1$.

Proof Write $L = K(\sqrt{\alpha})$ for some $\alpha \in K$. Since K is formally real but L is not formally real, there is an ordering \leq on K with $\alpha < 0$ [2, Theorem 1.1.8 and Lemma 1.1.7]. Take $z = x + y\alpha \in L$ with $x, y \in K$ and $(x, y) \neq (0, 0)$. Since $\sigma(z)z = x^2 - \alpha y^2 > 0$, $a > 0$ for every $a \in \hat{\Delta}$. Thus, $b > 0$ for every $b \in \hat{\Delta}_n$. \square

Example 2 Here we give a simple example with $\Delta_2 \neq \Delta$ and $0 \notin \hat{\Delta}$. Take $K = \mathbb{Q}$ and $L := \mathbb{Q}(i)$. For any $z = x + iy \in L$ we have $z\sigma(z) = x^2 + y^2$. Hence, Δ is the subset of $\mathbb{Q}_{\geq 0}$ formed by the sums of two squares. Every positive integer is the sum of 4 squares by a theorem of Lagrange and hence $\Delta_2 = \mathbb{Q}_{\geq 0}$. There are positive rational numbers that are not the sums of 2 or 3 squares, e.g., 7 is not the sum of 3 squares of rational numbers, because for any $n \in \{1, 2, 3\}$ a positive integer is the sum of n squares of rational numbers if and only if it is the sum of n squares of integers [9, Ch. 7]. Proposition 5 gives $0 \notin \hat{\Delta}$.

Example 3 Assume $\text{char}(K) = 0$, i.e. assume $K \supseteq \mathbb{Q}$. We have $\Delta_n \supseteq \mathbb{Q}_{\geq 0}$ for every $n \geq 4$, because every nonnegative integer is the sum of 4 integers by a theorem of Lagrange and $\Delta_n \setminus \{0\}$ is a multiplicative group.

Lemma 9 Assume $\text{char}(K) = 0$. Then there are infinitely many $m \in \Delta$ such that $1 - m$ is a square in K and $m = \sigma(z)z$ with $z \in L \setminus (K \cup K\sqrt{\alpha})$.

Proof Take $z = x + y\sqrt{\alpha} \in L^2$ with $x, y \in K$. Consider the equation in K^3 :

$$x^2 + w^2 = \alpha y^2. \tag{1}$$

As the proof of Lemma 8 we get that (1) has infinitely many solutions $(x, y, w) \in K^3$ with $y \neq 0$ and $x \neq 0$ and that the set of these solutions has infinite image under the projection $K^3 \rightarrow K$ onto the third coordinate. \square

Example 4 Assume $\text{char}(K) = 2$. We also assume that K is infinite. There is $\epsilon \in K$ such that $L = K(\beta)$, where β is any root of the polynomial $t^2 + t + \epsilon$. Note that $1 + \beta$ is the other root of the same polynomial and hence $\sigma(\beta) = 1 + \beta$ (note that $\sigma^2(\beta) = 2 + \beta = \beta$). We use $1, \beta$ as a basis of L as a 2-dimensional K -vector space. Take $z = x + y\beta \in L$. We have $\sigma(z) = x + y + y\beta$ and hence (since $xy + yx = 0$ and $\beta^2 + \beta = \epsilon$) $\sigma(z)z = x(x + y) + y^2\epsilon$. For any $c \in \Delta$ the affine conic $\{x^2 + xy + y^2\epsilon + c = 0\} \subset K^2$ is the circle with center 0 and squared-radius c . If $c = 0$, then this circle is the singleton $\{0\}$. Now assume $c \in \hat{\Delta}$. Take homogeneous coordinates x, y, z in $\mathbb{P}^2(\bar{L})$. Fix any $c \in K^*$. Set $g(x, y, z) := x^2 + xy + y^2\epsilon + cz^2$. The projective conic $T := \{g(x, y, z) = 0\} \subset \mathbb{P}^2(\bar{L})$ is smooth, because $\frac{\partial}{\partial x}g = y, \frac{\partial}{\partial y}g = x, \frac{\partial}{\partial z}g = 0$ and hence the common zero-set of the partial derivatives of $g(x, y, z)$ is the point $(0 : 0 : 1) \notin T$. Hence, if $c \in \hat{\Delta}$ any circle $\{\sigma(z - \mu)(z - \mu) = c\}$ is an affine smooth conic with at least one point P (because $c \in \Delta$). Taking the linear projection from P we see that this circle is infinite and with the cardinality of K .

4. The proofs and related results

Proof of Proposition 1. Taking $M - c\mathbb{I}_{n \times n}$ instead of M and applying Remark 5 we reduce to the case $c = 0$. Assume $\text{Num}(M) = \{0\}$ and $M \neq 0\mathbb{I}_{n \times n}$. In particular we have $n > 1$. Write $M = (m_{ij})$. Since every diagonal element of M is contained in $\text{Num}(M)$ by Remark 5, we have $m_{ii} = 0$ for all i . Hence, there is $m_{ij} \neq 0$ with $i \neq j$. Taking M^\dagger instead of M if necessary and applying Lemma 3 we reduce to the case $i < j$. A permutation of the orthonormal basis e_1, \dots, e_n is unitary and hence it preserves $\text{Num}(M)$. Permuting this basis we reduce to the case $i = 1$ and $j = 2$. Taking $(1/m_{12})M$ instead of M and applying Remark 5 we reduce to the case $m_{12} = 1$. Set $b := m_{21}$. Take $u = (x, y) \in L^2$ such that $\sigma(x)x + \sigma(y)y = 1$ and set $u := (a_1, \dots, a_n)$ with $a_1 = x, a_2 = y$ and $a_i = 0$ for all $i > 2$. We have $\langle u, Mu \rangle = b\sigma(x)y + \sigma(y)x$. Since $b \neq 0$, it is sufficient to prove the existence of $x, y \in L^*$ such that $f(x, y) := \sigma(x)x + \sigma(y)y = 1$ and $g(x, y) := b\sigma(x)y + \sigma(y)x \neq 0$. We first take $x, y \in K$ and so $\sigma(x) = x$ and $\sigma(y) = y$. Hence, $g(x, y) = 2(b + 1)xy$. Since $\text{char}(K) \neq 2$, the affine conic $D := \{x^2 + y^2 = 1\} \subset K^2$ is smooth. Since K is infinite and $(1, 0) \in D$, D has infinitely many K -points (use the linear projection from $(1, 0)$). Hence, we may find $(x, y) \in D$ with $g(x, y) \neq 0$, unless $b = -1$. Now assume $b = -1$. Write $y = tx$. We have $g(x, y) = 0$ if and only if either $xy = 0$ or $t \in K$. By Lemma 9 there is $y \in L \setminus K$, so that $1 - \sigma(y)y$ is a square in K^* and so there is $x \in K^*, y \in L \setminus K$ with $1 = \sigma(y)y + x^2$. □

Proposition 6 Take $M \in M_{2,2}(L)$ with a unique eigenvalue $c \in L$ with an eigenvector $v \neq 0$ with $\langle v, v \rangle = 0$. Then $0 \in \hat{\Delta}_2$ and either $M = c\mathbb{I}_{2,2}$ or there is $\mu \in L^*$ such that $\text{Num}(M) = c + \mu\hat{\Delta}$ and in the latter case $c \notin \text{Num}(M)$.

Proof Write $v := (a_1, a_2)$. By assumption $(a_1, a_2) \neq (0, 0)$ and $\sigma(a_1)a_1 + \sigma(a_2)a_2 = 0$. Hence, $0 \in \hat{\Delta}_2$. Since $v \neq 0$ and $\langle v, v \rangle = 0$, e_2 and v are not proportional and so they form a basis of L^2 . Since $\langle \cdot, \cdot \rangle$ is nondegenerate and $\langle v, v \rangle = 0$, we have $\langle v, e_2 \rangle \neq 0$. Taking a multiple of v if necessary we reduce to the case $\langle v, e_2 \rangle = 1$. Thus, $\langle e_2, v \rangle = 1$. Assume $M \neq c\mathbb{I}_{2,2}$. Taking $M - c\mathbb{I}_{2,2}$ instead of M and applying Remark 5 we reduce to the case $c = 0$. Write $M = (b_{ij})$, $i, j = 1, 2$, with respect to the basis v, e_2 . We have $b_{11} = b_{22} = b_{12} = 0$ and $b_{12} \neq 0$. Set $\mu := b_{12}$. Take $u = xv + ye_2$ such that $\langle u, u \rangle = 1$, i.e. $x\sigma(y) + \sigma(x)y + \sigma(y)y = 1$ (and in

particular $y \neq 0$). We have $\langle u, Mu \rangle = \langle xv + ye_2, \mu yv \rangle = \mu \sigma(y)y$. Varying $y \in L^*$ as $\sigma(y)y$ we get all elements of $\hat{\Delta}$. Hence, to conclude the proof it is sufficient to prove that for every $y \in L^*$ there is $x \in L$ such that $x\sigma(y)\sigma(x)y + \sigma(y)y = 1$. First assume $\text{char}(K) \neq 2$. Fix $e \in L \setminus K$ with $e^2 \notin K$ and write $x = u + ev$ and $y = c + ed$ with $u, v, c, d \in K$. We have $\sigma(x) = u - ev$ and $\sigma(y) = c - ed$. We find $2uc = \eta$ for some $\eta \in K$ and we always have a solution u , because we may take y giving $\sigma(y)y$ and $c \neq 0$. Now assume $\text{char}(K) = 2$. There is $e \in L \setminus K$ such that $e^2 + e \in L$ and $\sigma(e) = e + 1$. Write $x = u + ev$, $y = c + ed$ with $u, v, c, d \in K$ and $(c, d) \neq (0, 0)$. Now we find an equation $ud + vc = \eta$ for some $\eta \in K$, which may always be satisfied, because $(c, d) \neq (0, 0)$. \square

Remark 8 Assume $0 \in \hat{\Delta}_2$, say $0 = \sigma(x)x + \sigma(y)y$ with $(x, y) \in L^2 \setminus \{(0, 0)\}$, and take $c \in L$. Set $v := xe_1 + ye_2 \in L^2 \setminus \{(0, 0)\}$. Take any linear map $f : L^2 \rightarrow L^2$ with $\text{Ker}(f - c\text{Id}_{L^2}) = \text{Im}(f) = Lv$ and let M be the matrix associated to f . Then $M \neq c\mathbb{I}_{2 \times 2}$ and M satisfies the assumptions of Proposition 6.

Proof of Theorem 1. By assumption there are $a_1, a_2 \in L$ with $(a_1, a_2) \neq (0, 0)$ and $\sigma(a_1)a_1 + \sigma(a_2)a_2 = 0$. Set $v := a_1e_1 + a_2e_2$. We have $v \neq 0$ and $\langle v, v \rangle = 0$. Hence, e_2 and v are not proportional and so they form a basis of L^2 . Since $\langle \cdot, \cdot \rangle$ is nondegenerate and $\langle v, v \rangle = 0$, we have $\langle v, e_2 \rangle \neq 0$. Taking a multiple of v if necessary we reduce to the case $\langle v, e_2 \rangle = 1$. Thus, $\langle e_2, v \rangle = 1$. Take $M \in M_{n,n}(L)$ defined by $Mv = cv$ and $Me_2 = \mu v + ce_2$. Apply Proposition 6. \square

Proof [Proof of Theorem 2:] Taking $M - c\mathbb{I}_{2,2}$ instead of M and applying Remark 5 we reduce to the case $c = 0$.

Set $\delta := \langle v, v \rangle$. Write $v = a_1e_1 + a_2e_2$ and set $w := -\sigma(a_2)e_1 + \sigma(a_1)e_2$. We have $\langle v, w \rangle = -\sigma(a_1)\sigma(a_2) + \sigma(a_2)\sigma(a_1) = 0$ and hence $\langle w, v \rangle = 0$. Since $\delta = \langle v, v \rangle = \sigma(a_1)a_1 + \sigma(a_2)a_2$ and $\sigma(\sigma(a_i)) = a_i$, we have $\langle w, w \rangle = \delta$. Since $\delta \neq 0$ and $\langle v, w \rangle = 0$, v, w are a basis of L^2 . Write $M = (b_{ij})$, $i, j = 1, 2$, with respect to the basis v, w . By assumption we have $b_{11} = b_{21} = 0$. Since M has a unique eigenvalue, we have $b_{22} = 0$. Assume $M \neq 0\mathbb{I}_{2,2}$ and hence $b_{12} \neq 0$. Set $\mu := b_{12}$.

(i) If $\delta \in \Delta$, then $\delta \in \text{Num}(M)$ by Remark 6. Now assume $0 \in \text{Num}(M)$, i.e. assume the existence of $u = xv + yw$ such that $\langle u, u \rangle = 1$ (i.e. such that $\sigma(x)x + \sigma(y)y = 1/\delta$) and $\langle u, Mu \rangle = 0$ (i.e. $0 = \langle xv + yw, \mu yv \rangle = \mu \delta \sigma(x)y$). Hence, either $y = 0$ or $x = 0$. Hence, u is either a multiple of v or a multiple of w . Since $\langle v, v \rangle = \langle w, w \rangle = \delta$, Lemma 4 gives $\delta \in \Delta$.

(i) Now assume $\delta \in \hat{\Delta}$. Since $\hat{\Delta}$ is a multiplicative group, there is $t \in L$ such $\sigma(t)t = 1/\delta$. Set $v_1 := tv$ and $v_2 := tw$. We have $\langle v_i, v_i \rangle = 1$ and $\langle v_i, v_j \rangle = 0$ if $i \neq j$. Let (m_{ij}) be the matrix associated to M in the basis v_1, v_2 . We have $m_{11} = m_{12} = 0$. Since M has a unique eigenvalue, we have $m_{22} = 0$. Assume $M \neq 0\mathbb{I}_{2 \times 2}$, i.e. assume $m_{12} \neq 0$. Taking $(1/m_{12})M$ instead of M and applying Remark 5 we reduce to the case $m_{12} = 1$. Take $u = xw_1 + yw_2$ with $\langle u, u \rangle = 1$, i.e. with $\sigma(x)x + \sigma(y)y = 1$. Set $\gamma := \sigma(x)x$ and hence $\sigma(y)y = 1 - \gamma$. Since $0 \in \text{Num}(M)$, to check all other elements of $\text{Num}(M)$ we may assume $xy \neq 0$, i.e. $\gamma \notin \{0, 1\}$. Note that γ is an arbitrary element of $\hat{\Delta} \cap (1 - \hat{\Delta})$. We fix γ , but we only take x, y with $\sigma(x)x = \gamma$ and $\sigma(y)y = 1 - \gamma$. We have $\langle u, Mu \rangle = \langle xw_1 + yw_2, yw_1 \rangle = \sigma(x)y$. Note that $\sigma(x)y \cdot \sigma(\sigma(x)y) = \gamma(1 - \gamma)$. Fix $w \in L$ such that $\sigma(w)w = \gamma(1 - \gamma)$. Take any x with $\sigma(x)x = \gamma$. Note that $x \neq 0$, because $\gamma \neq 0$. Take $y := w/\sigma(x)$. To conclude the proof of part (b) it is sufficient to prove that $\sigma(y)y = 1 - \gamma$. We have $\sigma(y)y = \sigma(w)w/x\sigma(x) = \gamma(1 - \gamma)/\gamma$. \square

Lemma 10 and Proposition 7 are, respectively, the analogues of [3, Lemma 3.6 and Theorem 1.2 (d)]. In their statements the last \bigcup is a union of circles with center 0, in which if we take $d \in \hat{\Delta} \cap (1 - \hat{\Delta})$ as a parameter space the circles coming from d and $1 - d$ are the same and we do not claim that \bigcup is a disjoint union (see [3, Example 3.7]).

Lemma 10 Fix $b \in L^*$ and let $M = (m_{ij})$ be the 2×2 matrix with $a_{11} = 1$, $a_{21} = a_{22} = 0$, and $a_{12} = b$. Then

$$\text{Num}(M) = \{0, 1\} \cup \bigcup_{d(1-d), d \in \hat{\Delta} \cap (1-\hat{\Delta})} C(\sigma(b)bd(1-d), d).$$

Proof The vector e_1 gives $1 \in \text{Num}(M)$. The vector e_2 gives $0 \in \text{Num}(M)$. Take $u = xe_1 + ye_2 \in L^2$ such that $\langle u, u \rangle = 1$, i.e. such that $\sigma(x)x + \sigma(y)y = 1$. Set $d := \sigma(x)x \in \Delta$. We have $\sigma(y)y = 1 - d$ and hence $d \in (1 - \Delta)$. We have $\langle u, Mu \rangle = \langle xe_1 + ye_2, (x + by)e_1 \rangle = d + b\sigma(x)y$. Set $m := b\sigma(x)y$. We have $\sigma(m)m = \sigma(b)b\sigma(y)y\sigma(x)x = \sigma(b)bd(1-d)$. Thus, $m \in C(bd(1-d), 0)$ and hence $\langle u, Mu \rangle \in d + C(\sigma(b)bd(1-d), 0)$. Since $0, 1 \in \text{Num}(M)$, from now on we may assume $d \notin \{0, 1\}$ and prove the other inclusion. Take any $m' \in C(\sigma(b)bd(1-d), 0)$, i.e. with $\sigma(m')m' = \sigma(b)bd(1-d)$. Take $x_1 \in C(d, 0)$. Since $d \neq 0$, we have $x_1 \neq 0$. Set $y_1 := m'/(b\sigma(b)x_1)$ and $u' := x_1e_1 + y_1e_2$. We have $y_1 \in C((1-d), 0)$, $\langle u', Mu' \rangle = d + m'$ and $\sigma(x_1)x_1 + \sigma(y_1)y_1 = \sigma(b)bd(1-d)$. \square

Proposition 7 Assume $n = 2$ and that M has two eigenvalues $c_1, c_2 \in L$, $c_1 \neq c_2$, with eigenvectors v_i for c_i with $\langle v_1, v_1 \rangle \in \hat{\Delta}$.

(i) If $\langle v_1, v_2 \rangle = 0$, then M is unitarily equivalent to $c_1\mathbb{I}_{1 \times 1} \oplus c_2\mathbb{I}_{1 \times 1}$.

(ii) Assume $\langle v_1, v_2 \rangle \neq 0$. Then there is $\mu \in L^*$ such that $\text{Num}(M)$ is the union of $\{c_1, c_2\}$ and a union of circles $C(\sigma(\mu)\mu d(1-d), d)$ with $d \in \hat{\Delta} \cap (1 - \hat{\Delta})$.

Proof Set $\delta := \langle v_1, v_1 \rangle$. Since $1/\delta \in \hat{\Delta}$ (Remark 2), there is $t \in L^*$ such that $\sigma(t)t = 1/\delta$. Set $w_1 := tv_1$. We have $\langle w_1, w_1 \rangle = 1$ and $Mw_1 = c_1w_1$. Write $w_1 = a_1e_1 + a_2e_2$ for some $a_1, a_2 \in L$. Set $w_2 := -\sigma(a_2)e_1 + \sigma(a_1)e_2$. Note that $\langle w_2, w_1 \rangle = 0$ and $\langle w_2, w_2 \rangle = 1$. Taking $M - c_2\mathbb{I}_{2,2}$ instead of M and applying Remark 5 we reduce to the case $c_2 = 0$ and hence $c := c_1 - c_2 \neq 0$. Taking $(1/c)M$ instead of M and applying Remark 5 we reduce to the case $c = c_1 - c_2 = 1$. Write $M = (m_{ij})$, $i = 1, 2$, in the orthonormal basis w_1 and w_2 . We have $m_{11} = 1$ and $m_{21} = m_{22} = 0$. If $m_{12} = 0$, then M is unitary equivalent to the matrix $c_1\mathbb{I}_{1 \times 1} \oplus c_2\mathbb{I}_{1 \times 1}$. We have $m_{12} = 0$ if and only if w_2 is proportional to v_2 , i.e. (being v_2 linearly independent from $tv_1 = w_1$) if and only if $\langle v_1, v_2 \rangle = 0$. Apply Lemma 8 with $\mu := m_{12}/(c_1 - c_2)$. \square

Proposition 8 Take $M \in M_{2,2}(L)$ with eigenvalues $c_1, c_2 \in L$, $c_1 \neq c_2$, and take $v_i \in L^2$ such that $Mv_i = c_iv_i$ and $v_i \neq 0$. Assume $\langle v_i, v_i \rangle = 0$ for all i . Set $\mathcal{D} := \{t \in L \mid t + \sigma(t) = 1\}$. Then $\text{Num}(M) = \mathcal{D}$

Proof Taking $(1(c_2 - c_1)(M - c_1\mathbb{I}_{2 \times 2}))$ and applying Remark 5 we reduce to the case $c_1 = 0$ and $c_2 = 1$. Since $\langle \cdot, \cdot \rangle$ is nondegenerate and $\langle v_i, v_i \rangle = 0$ for all i , we have $\langle v_1, v_2 \rangle \neq 0$. Taking $(1/\langle v_1, v_2 \rangle)v_2$ instead of v_2 we reduce to the case $\langle v_1, v_2 \rangle = 1$. Thus, $\langle v_2, v_1 \rangle = 1$. Take $u = xv_1 + yv_2$ with $\langle u, u \rangle = 1$, i.e. with $\sigma(x)y + \sigma(y)x = 1$. Note that $x \neq 0$ and $y \neq 0$. We have $\langle u, Mu \rangle = \langle xv_1 + yv_2, yv_1 \rangle = \sigma(x)y$. For any $b \in L^*$ let $N(b)$ be the set of all $\sigma(x)b$ with $\sigma(x)b + \sigma(b)x = 1$. Fix $y \in L^*$. Set $t := x/y$.

Note that $t \neq 0$. Since $\sigma(x)y + \sigma(y)x = 1$, we have $\sigma(t)y\sigma(y) + ty\sigma(y) = 1$ and $\langle u, Mu \rangle = \sigma(t)y\sigma(y)$. Take $t_0, y_0, y \in L^*$ such that $t_0\sigma(t_0)y_0\sigma(y_0) + t_0y_0\sigma(y_0) = 1$ and set $t_1 := t_0y_0\sigma(y_0)/(y\sigma(y))$. We have $\sigma(t_1)y\sigma(y) + t_1y\sigma(y) = t_0\sigma(t_0)\sigma(t_0) + t_0y_0\sigma(y_0)$. Hence, $N(y) \supseteq N(y_0)$. By symmetry we get $N(y) = N(y_0)$. Taking $y_0 = 1$ we get $\text{Num}(M) = N(1)$. Note that $t + \sigma(t) = 1$ if and only if $\sigma(t) + t = 1$. Hence, $\sigma(N(1)) = N(1)$. Thus, $\text{Num}(M) = \mathcal{D}$. \square

Proposition 9 Fix $n > 2$ and assume $\Delta_n = \Delta$. Take $M \in M_{n,n}(L)$ with an eigenvalue $c \in L$ with an eigenspace of dimension $n - 1$. Then one of the following cases occurs:

1. M is unitarily equivalent to $c\mathbb{I}_{n-1,n-1} \oplus d\mathbb{I}_{1 \times 1}$ (unitary direct sum) for some $d \neq c$;
2. M is unitarily equivalent to $c\mathbb{I}_{n-2,n-2} \oplus M'$ (unitary direct sum) with M' as in case (ii) of Proposition 7;
3. M is unitarily equivalent to $c\mathbb{I}_{n-2,n-2} \oplus M'$ (unitary direct sum) with M' as in Theorem 2;
4. M is unitarily equivalent to $c\mathbb{I}_{n-2,n-2} \oplus M'$ (unitary direct sum) with M' as in Proposition 8;
5. M is unitarily equivalent to $c\mathbb{I}_{n-2,n-2} \oplus M'$ (unitary direct sum) with M' as in Proposition 6.

Proof Let $W \subset L^n$ be the c -eigenspace space of M . Taking $M - c\mathbb{I}_{n,n}$ instead of M and applying Remark 5 we reduce to the case $c = 0$. Hence, $Mw = 0$ for all $w \in W$. By Lemma 6 there are $f_1, \dots, f_{n-2} \in W$ such that $\langle f_i, f_i \rangle = 1$ for all i and $\langle f_i, f_j \rangle = 0$ for all $i \neq j$. Let V be the linear span of f_1, \dots, f_{n-2} . Set $V^\perp := \{x \in L^n \mid \langle w, x \rangle = 0 \text{ for all } w \in W\}$. By the choice of f_1, \dots, f_{n-2} we have $V \cap V^\perp = \{0\}$. Since $\langle \cdot, \cdot \rangle$ is nondegenerate, we have $\dim(V) + \dim(V^\perp) = n$ and so $L^n = V \oplus V^\perp$ (unitary direct sum). Fix $w, v \in V$. We have $\langle v, M^\dagger w \rangle = \langle Mv, w \rangle = \langle 0, w \rangle = 0$. Since the restriction of $\langle \cdot, \cdot \rangle$ to V is nondegenerate, we get $M^\dagger w = 0$. Hence, $M^\dagger w = 0$ for all $w \in W$. Fix $m \in V^\perp$ and $v \in W$. We have $\langle v, Mm \rangle = \langle M^\dagger v, m \rangle = \langle 0, m \rangle = 0$. Since this is true for all $v \in W$, we get $Mm \in V^\perp$. Hence, $MV^\perp \subseteq V^\perp$. Set $B := M|_{V^\perp}$, seen as a map $V^\perp \rightarrow V^\perp$. All the eigenvalues of M are in L and we call d the other eigenvalue. The matrix B has eigenvalues 0 and d with 0 the eigenspace that contains $u \in W \cap V^\perp, u \neq 0$.

(a) First assume $d \neq 0$ and hence there is $v \in V^\perp$ with $Mv = dv$ and $v \neq 0$. We have $\langle z, z \rangle \in \Delta_n = \Delta$ for all $z \in V^\perp$. Hence, if either $\langle u, u \rangle \neq 0$ or $\langle v, v \rangle \neq 0$, then we apply Proposition 9 and get that we are either in case (1) or case (2). If $\langle u, u \rangle = \langle v, v \rangle = 0$, then we apply Proposition 8.

(b) Now assume $d = 0$. By assumption Lu is the only one-dimensional subspace of V^\perp sent into itself by B . If $\langle u, u \rangle = 0$, then we apply Proposition 6. If $\langle u, u \rangle \neq 0$, then we apply Theorem 2. \square

5. Matrices with coefficients in K

Take $M = (m_{ij}) \in M_{n,n}(K)$. The set $C_n(1, K)$ is the set of all solutions $(x_1, \dots, x_n) \in K^n$ of the equation

$$x_1^2 + \dots + x_n^2 = 1. \tag{2}$$

Thus, $\text{Num}(M)_K$ is the set of all

$$\sum_{i,j} m_{ij} x_i x_j \tag{3}$$

with x_1, \dots, x_n satisfying (2).

Lemma 11 Take $M = (m_{ij}), N = (n_{ij}) \in M_{n,n}(K)$.

1. If $m_{ii} = n_{ii}$ for all i and $m_{ij} + m_{ji} = n_{ij} + n_{ji}$ for all $i \neq j$, then $\text{Num}(M)_K = \text{Num}(N)_K$.
2. We have $\text{Num}(B)_K = \text{Num}(M)_K$ for the matrix $B := (b_{ij})$ with $b_{ii} = m_{ii}$ for all i , $b_{ij} = 0$ for all $i < j$, and $b_{ij} = m_{ij} + m_{ji}$ for all $i > j$.
3. If $\text{char}(K) \neq 2$, the matrix $A := (a_{ij})$ with $a_{ij} = (m_{ij} + m_{ji})/2$ for all i, j is symmetric and $\text{Num}(A)_K = \text{Num}(M)_K$.

Proof Equation (3) is the same for M (i.e. with m_{ij} as coefficients) and for N (i.e. with n_{ij} as coefficients). The last two assertions of Lemma 11 follow from the first one. \square

Remark 9 For all $c, d \in K$ and all $M \in M_{n,n}(K)$ we have $\text{Num}(cM + d\mathbb{I}_{n,n})_K = c\text{Num}(M)_K + d$.

Remark 10 The vectors e_1, \dots, e_n prove that for any $M \in M_{n,n}(K)$ the diagonal elements of M are contained in $\text{Num}(M)_K$.

Proof [Proof of Proposition 2:] Write $M = (m_{ij})$. Taking $M - m_{11}\mathbb{I}_{n,n}$ instead of M we reduce to the case $m_{11} = 0$ by Remark 9. If M is antisymmetric and $m_{ii} = 0$ for all i , then $\text{Num}(M)_K = \text{Num}(0\mathbb{I}_{n \times n})_K = \{0\}$ by Lemma 11.

Now assume $\sharp(\text{Num}(M)_K) = 1$. Since the diagonal elements of M are contained in $\text{Num}(M)_K$ by Remark 10, we have $m_{ii} = 0$ for all i . Assume $m_{ij} \neq 0$ for some $i \neq j$. The first part of the proof of Proposition 1 with e_i, e_j instead of e_1, e_2 (i.e. the part with $x, y \in K$) gives $m_{ji} = -m_{ij}$. \square

Remark 11 Assume $\text{char}(K) = 2$. Then $x_1^2 + \dots + x_n^2 = (x_1 + \dots + x_n)^2$. Hence, the elements of (3) coming from the solutions of (2) are the ones coming from the solutions of

$$x_1 + \dots + x_n = 1. \tag{4}$$

Substituting $x_n = 1 + x_1 + \dots + x_{n-1}$ in (3) we get that $\text{Num}(M)_K$ is the image of a map $f_M : K^{n-1} \rightarrow K$ with f_M a polynomial in x_1, \dots, x_{n-1} with $\deg(f_M) \leq 2$. If $\deg(f_M) = 1$, then f_M is surjective, i.e. $\text{Num}(M)_K = K$. If $\deg(f_M) = 0$, then f_M is constant and hence $\sharp(\text{Num}(M)_K) = 1$. Let g_M be the homogeneous degree 2 part of f_M and let $A = (a_{ij}), i, j = 1, \dots, n-1$, be the matrix associated to g_M with $a_{ij} = 0$ if $i < j$. We have $a_{ii} = m_{ii} + m_{nn}$ and $a_{ij} = m_{ij} + m_{ji}$ for all $i \neq j$ with $i, j < n$. Since $\text{char}(K) = 2$, we have $a_{ii} = 0$ if and only if $m_{ii} = m_{nn}$ and $a_{ij} = 0$ (with $i \neq j$) if and only if $m_{ij} = m_{ji}$. Thus, $g_M = 0$ if and only if all diagonal elements of M are the same and the top $(n-1) \times (n-1)$ principal submatrix of M is symmetric.

(a) Assume $g_M \neq 0$ and that K is infinite.

Claim 1: If $a_{ij} \neq 0$ for some $i \neq j$, then $\text{Num}(M)_K = K$.

Proof of Claim 1: Up to a permutation of e_1, \dots, e_{n-1} we may assume $a_{12} \neq 0$. We have $f_M(x_1, x_2, 0, \dots, 0) = (m_{11} + m_{nn})x_1^2 + a_{12}x_1x_2 + (m_{22} + m_{nn})x_2^2 + \beta x_1 + \gamma x_2 + \delta$ for some $\beta, \gamma, \delta \in K$. Hence, it is sufficient to prove that the image of the map $\psi : K^2 \rightarrow K$ induced by the polynomial $f_M(x_1, x_2, 0, \dots, 0)$ has the cardinality of K . We will prove that ψ is surjective. Take $b \in K$ such that $a_{11}b \neq -\gamma$. The polynomial

$f_M(b, x_2, 0, \dots, 0)$ is a nonconstant degree 1 polynomial and hence it induces a surjection $K \rightarrow K$. Thus, ψ is surjective. Now assume $a_{ii} \neq 0$ for some i , i.e. $m_{ii} \neq m_{nn}$ for some $i < n$.

Claim 2: Assume $a_{ij} = 0$ for all $i \neq j$, but $a_{ii} \neq 0$ for some $i < n-1$. If K is infinite, then $\text{Num}(M)_K$ has the cardinality of K .

Proof of Claim 2: We have $g_M(x_1, \dots, x_{n-1}) = \sum_{i=1}^{n-1} a_{ii}x_i^2 \neq 0$ and

$$f_M(x_1, \dots, x_{n-1}) = g_M(x_1, \dots, x_{n-1}) + \ell(x_1, \dots, x_{n-1}) + \gamma$$

for some $\gamma \in K$ and a linear form $\ell \in K[x_1, \dots, x_{n-1}]$. Up to a permutation of the indices we may assume that $a_{11} \neq 0$. Fix any $(b_2, \dots, b_{n-1}) \in K^{n-2}$ and call $\phi : K \rightarrow K$ the map induced by $f_M(x_1, b_2, \dots, b_{n-1})$. Since $f_M(x_1, b_2, \dots, b_{n-1})$ is a degree 2 nonconstant polynomial, each fiber of ϕ has at most cardinality 2. Hence, $\phi(K)$ and K have the same cardinality.

(b) Assume $g_M \equiv 0$. In particular, $m_{ii} = m_{nn}$ for all $i < n$. We have $f_M(0, \dots, 0) = a_{nn}$ and $f_M(x_1, \dots, x_m) = b_1x_1 + \dots + b_{n-1}x_{n-1} + a_{nn}$ with $b_i = m_{ni} + m_{in}$. Hence, f_M is surjective if and only if $m_{ni} \neq m_{in}$ for some $i < n$, while f_M is constant if $m_{ni} = m_{in}$ for all $i < n$.

Proof of Proposition 3. The proposition was proved in Remark 11, with as a bonus the discussion of some cases with $\text{Num}(M)_K = K$. □

References

- [1] Ballico E. On the numerical range of matrices over a finite field. *Linear Algebra Appl* 2016; 512: 162-171.
- [2] Bochnak J, Coste M, Roy MF. *Real Algebraic Geometry*. Berlin, Germany: Springer, 1998.
- [3] Coons JI, Jenkins J, Knowles D, Luke RA, Rault PX. Numerical ranges over finite fields. *Linear Algebra Appl* 2016; 501: 37-47.
- [4] Gustafson KE, Rao DKM. *Numerical Range: The Field of Values of Linear Operators and Matrices*. New York, NY, USA: Springer, 1997.
- [5] Horn RA, Johnson CR. *Matrix Analysis*. New York, NY, USA: Cambridge University Press, 1985.
- [6] Horn RA, Johnson CR. *Topics in Matrix Analysis*. Cambridge, UK: Cambridge University Press, 1991.
- [7] Ireland K, Rosen M. *A Classical Introduction to Modern Number Theory*. New York, NY, USA: Springer, 1982
- [8] Lang S. *Algebra*. Revised Third Edition. Berlin, Germany: Springer, 2002.
- [9] LeVeque W. *Topics in Number Theory, Vol. 1*. Reading, MA, USA: Addison-Wesley, 1956.
- [10] Steinitz E. *Algebraische Theorie der Körper*. Berlin, Germany: De Gruyter, 1930 (in German).