

1-1-1998

Latest Cretaceous Magmatism in the Central Anatolian Crystalline Complex: Review of Field, Petrographic and Geochemical Features

NURDAN S. AYDIN

M. CEMAL GÖNCÜOĞLU

AYHAN ERLER

Follow this and additional works at: <https://journals.tubitak.gov.tr/earth>



Part of the [Earth Sciences Commons](#)

Recommended Citation

AYDIN, NURDAN S.; GÖNCÜOĞLU, M. CEMAL; and ERLER, AYHAN (1998) "Latest Cretaceous Magmatism in the Central Anatolian Crystalline Complex: Review of Field, Petrographic and Geochemical Features," *Turkish Journal of Earth Sciences*: Vol. 7: No. 3, Article 11. Available at: <https://journals.tubitak.gov.tr/earth/vol7/iss3/11>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Earth Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Latest Cretaceous Magmatism in the Central Anatolian Crystalline Complex: Review of Field, Petrographic and Geochemical Features

Nurdan S. AYDIN*

Mersin Üniversitesi, Jeoloji Mühendisliği Bölümü, 33160 Mersin-TURKEY

M. Cemal GÖNCÜOĞLU, Ayhan ERLER

Orta Doğu Teknik Üniversitesi, Jeoloji Mühendisliği Bölümü, 06531 Ankara-TURKEY

Received: 21.05.1998

Abstract: Magmatic, metamorphic and ophiolitic rock assemblages in Central Anatolia are collectively named as the Central Anatolian Crystalline Complex (CACC). Magmatic rocks form significant portion of the CACC and display a range of fabrics, mineralogies and compositions. However, composite granitoid intrusions, such as the Yozgat, Ağaören, Ekecikdağ and Cefalıkdağ, dominate the geology of the complex. Overall, there are two main types of Central Anatolian Granitoids (CAG): a) the C-type (crustal) leucogranites and b) the H-type (hybrid) hornblende ± K-feldspar megacrysts ± mafic microgranular enclave bearing granites. The emplacement of granitoid magmatism was followed by the syenitoid magmatism where the quartz syenitoids (e.g., the İdişdağı Intrusion) predate the feldspathoid bearing syenitoids (e.g., the Atdere Intrusion). The type of magmatism in the CACC varies from peraluminous, metaluminous to alkaline/peralkaline through time.

The C-type granitoids represent the early granitoid phase of the CACC and are considered to be products of the syn-collisional magmatism. The H-type granitoids and the syenitoid intrusions represent the advanced and final stages of the post-collisional magmatism, respectively. The C-type granitoids are likely derived from upper crustal sedimentary protolith by partial melting induced by crustal thickening with or without intrusion of mantle-derived mafic melts. Most of the field and petrographic features of the H-type granitoids require mantle-derived mafic magma contributions in the genesis of these rocks which can be explained in terms of mafic magma underplating of lower crust as a result of lithospheric delamination. Transition from silica-saturated to silica-undersaturated magmatism can be attributed to variation of availability of water in the melting zone. In general, the nature of magmatism in the CACC and related magmatic processes and the relative contributions of source material varied through time.

Orta Anadolu Kristalen Karma 1 ında En Geç Kretase Magmatizması: Arazi, Petrografi ve Jeokimya Bulgularının Gözden Geçirilmesi

Özet: Orta Anadolu'daki magmatik, metamorfik ve ofiyolitik kayalık toplulukları hep birlikte Orta Anadolu Kristalen Karmaşığı (OAKK) olarak adlanırlar. Magmatik kayalar, OAKK'nın önemli bir bölümünü oluştururlar ve farklı dokular, mineralojik içerikler ve bileşimler sergilerler. Ancak, Yozgat, Ağaören, Ekecikdağ ve Cefalıkdağ gibi kompozit granitoid intrüzyonları karmaşığın jeolosinde egemen durumdadır. Genel olarak Orta Anadolu Granitoidleri (OAG) iki ana gruba ayrılır: a) C-tipi (kabuksal) lökograditler ve b) H-tipi (hibrid) hornblend±K-feldspat megakristleri±mafik mikrogranüler anklavlar içeren granitler. Granitoid magmatizmasının yerleşmesini izleyen siyenitoid magmatizma evresinde ilk fazı kuvars siyenitoidler (örneğin, İdişdağı intrüzyonu) ikinci fazı ise feldspatoidli siyenitoidler (örneğin, Atdere intrüzyonu) oluşturur. Zaman boyunca OAKK'ndaki magmatizma türleri peralüminaldan metaalüminalya ve giderek alkalın/peralkaline doğru değişir.

C-tipi granitoidler OAKK'ndaki erken granitoid evresini temsil ederler ve çarpışma-sırası magmatizmasının ürünleri olarak kabul edilirler. H-tipi granitoidler ve siyenitoid intrüzyonları çarpışması-sonrası magmatizmasının, sırasıyla, ilerlemiş ve son evrelerini temsil ederler. C-tipi granitoidler, olasılıkla üst kabuğa ait kayalarda kabuksal kalınlaşmanın yol açtığı kısmi ergimeyle oluşmuşlardır. Bu evrede manto-kökenli mafik ergiyiklerin katkısı belirgin değildir. H-tipi granitoidlerin arazi ve petrografik özelliklerinin çoğu bu kayaların evriminde, litosferik sıyrılma sonucu olarak alt kabuğun mafik magma ile altlanması biçiminde açıklanabilen, manto-kökenli mafik magma katkıları gerektirir. Silikaca-doygun magmatizmadan silikaca-doymamış magmatizmaya geçişin nedeni ergime zonundaki su miktarındaki değişimler olabilir. Genel olarak, OAKK'ndaki magmatizmanın türü, ilişkili magmatik süreçler ve kaynak malzemenin göreceli katkıları zaman boyunca değişmiştir.

* Present Address: Department of Earth Sciences, University of Hong Kong, Pokfulam Road-HONG KONG.

Introduction

Magmatic, metamorphic and ophiolitic rock assemblages in Central Anatolia are collectively named as the Central Anatolian Crystalline Complex (CACC; Göncüoğlu et al., 1991). The complex lies in a triangular area bounded by the Tuzgölü fault to the west, the Ecemiş fault to the east and the İzmir-Ankara-Erzincan suture to the north (Fig. 1). The CACC has been recently re-mapped by the METU-CACC Study Group (Göncüoğlu et al., 1994 and the references therein). Magmatic rocks dominate the geology of the complex and intrude the ophiolitic and metamorphic rocks of the complex. The complex is overlain by Uppermost Maastrichtian, Lower Paleocene and Eocene volcanic, clastic and carbonate rocks, Oligocene-Miocene evaporite and clastic rocks and Miocene-Pliocene continental clastic rocks (Göncüoğlu et al., 1991; 1992; 1993).

Granitoids and syenitoids form a significant portion of the complex. Although they have been subjected to numerous geological, petrographical and geochemical studies, their tectonic setting and nature of the relevant magmatism are still unresolved. This paper summarizes

main field, petrographic and geochemical features of the selected granitoid and syenitoid intrusions of the CACC mainly based on the studies of the METU-CACC Study Group. The main goal of this compilation is to construct a common framework upon which a petrogenetic model for the diverse magmatism of the CACC can be based. In this study, the Yozgat, Ağaçören, Ekecikdağ and Cefalıklıdağ intrusions are reviewed to characterize the granitoid magmatism of the CACC, whereas the İdişdağı and Atdere intrusions are discussed to represent the syenitoid magmatism of the complex. Note that this study is presented (Aydın et al., 1997) during the TÜBİTAK-BAYG/NATO-D program on Alkaline Magmatism in Central Anatolia in September 1997 in Sivas.

Geologic and Petrographic Framework

The granitoid rocks dominate the geology of the complex and reveal a wide range of fabrics, mineralogies and compositions. They crop out within three distinct belts (Fig. 1). These are: a) a large number of individual




Figure 1. Location map and simplified geological setting of the Central Anatolian Crystalline Complex.

small plutons exposed along the western margin extending from NE-SW to NW-SE ; b) a relatively narrow and smaller set of disconnected plutons exposed along the eastern margin; and c) a large intrusion exposed along the northern margin (Erlar et al., 1991). The granitoid rocks can be grouped as a) two-mica leucogranites; b) biotite/hornblende granites; c) K-feldspar megacryst-bearing granites; d) granodiorites; e) tonalites; and f) aplitic K-feldspar granites (Erlar and Göncüoğlu, 1996). Most of the granitoid intrusion of the CACC, e.g. the Yozgat, Ağaören, Ekecikdağ and Cefalıkdağ intrusions, comprises several subunits (Erlar et al., 1991; Erlar and Bayhan, 1995; Erlar and Göncüoğlu, 1996; Kadioğlu, 1996; Kadioğlu and Güleç, 1996; Tatar and Boztuğ, 1997; Aydın, 1997; Aydın et al., 1997; 1998). However, overall they contain two main granitoid types: a) locally garnet bearing leucogranites and b) hornblende ± biotite ± clinopyroxene bearing granites.

Leucogranites forming only a minor part of the complex represent the early magmatic phase of the complex (Akıman et al., 1993; Erlar and Göncüoğlu 1996; Boztuğ, 1997). Their size and form vary significantly from meter scale patches to several km scale geographically distinct subunits within Ağaören, Ekecikdağ and Yozgat intrusions. Main minerals are K-feldspar, quartz and plagioclase, the accessory minerals are apatite, zircon, garnet, muscovite and opaque phases. Presence of garnet ± muscovite ± tourmaline, clearly suggest their peraluminous nature. The leucogranites are typically free of K-feldspar megacrysts and mafic microgranular enclaves, and have been classified as crustal-type (C-type) (Aydın, 1997; Aydın et al., 1997; 1998) according to the classification system proposed by Barbarin (1990).

Hornblende ± biotite ± clinopyroxene bearing granites display compositional variation from quartz monzonites to monzogranites. Main minerals are perthitic K-feldspar, plagioclase, quartz and hornblende, the accessory minerals are apatite, zircon, sphene and opaque minerals. In general, lack of riebeckite, garnet, muscovite, sillimanite or cordierite imply that these granitoid bodies are neither peralkaline nor peraluminous. They typically contain variable size K-feldspar megacrysts (2-15 cm in length) and mafic microgranular enclaves. The examined K-feldspar megacrysts from the granitoids of the CACC are euhedral and ubiquitously display simple twinning and poikilitic texture suggesting their magmatic nature. Mafic microgranular enclaves of the CACC (on the basis of the data particularly from the Ağaören and Yozgat intrusions) are characterized by the following physical features. The presence of a) primary igneous features,

i.e. lack of deformation and recrystallization; b) acicular apatite grains; and c) blade-shape biotite ± quartz ocelli rimmed by mafic minerals (Kadioğlu and Güleç, 1996; Tatar and Boztuğ, 1997). These features clearly suggest that commingling of contrasting magmas play a significant role in the formation of the hornblende ± biotite ± clinopyroxene bearing rocks of the CACC (Kadioğlu and Güleç, 1996; Tatar and Boztuğ, 1997; Aydın et al., 1997). Thus, this type granitoid rocks of the CACC have been classified as hybrid-type (H-type) to emphasize their nature (Aydın, 1997; Aydın et al., 1997; 1998) according to the classification system proposed by Barbarin (1990).

Following the granitoid intrusions, an extensive syenitoid magmatism was developed. The syenitoid magmatism has been subjected to many studies (e.g., Özkan, 1987; Erlar et al., 1991; Özkan and Erkan, 1994; Bayhan and Tolluoğlu, 1987; Boztuğ, 1997; Otlar and Boztuğ, 1997; Göncüoğlu et al., 1997). The syenitoid magmatism includes both silica-oversaturated /saturated syenitoids and monzonites and silica-undersaturated syenitic rocks. İdişdağı Intrusion represents the silica-saturated/oversaturated syenitoids, whereas the Atdere Intrusion characterize the silica-undersaturated syenitoids.

The following features of the İdişdağı Intrusion are summarized from Göncüoğlu et al. (1997). The intrusion is composed of quartz and K-feldspar bearing syenites. They are medium- to fine- grained and locally display porphyritic texture and composed of K-feldspar, quartz, plagioclase and amphibole with minor amounts of biotite, muscovite and clinopyroxene. Sphene and opaque minerals are ubiquitous accessory phases.

The Atdere Intrusion (Kayseri) was investigated by Özkan (1987) and Özkan and Erkan (1994). These authors divided the Atdere Intrusion into four different subunits on the basis of their modal mineralogy. These are nepheline syenite, cancrinite-nepheline syenite, sodalite-nepheline syenite, and melanite-nepheline syenite. Although there is no direct dating on feldspathoid bearing syenitoid rocks, it is generally accepted that these rocks are younger than the granitoids and quartz bearing syenitoids (Ayan, 1963; Seymen, 1982; Bayhan and Tolluoğlu, 1987; Erlar et al, 1991; Boztuğ, 1997; Otlar and Boztuğ, 1997). The Atdere Syenitoids display holocrystalline-hypidiomorphic granular texture. Alkali-feldspar and feldspathoid minerals including nepheline, sodalite, cancrinite and melanite are the main minerals. Main accessory minerals are biotite, aegirine-augite, zircon and sphene.

Radiometric data on the age of the Central Anatolian Granitoids is very limited. In general, Rb/Sr whole-rock isochron data from Niğde-Üçkapılı (95 ± 11 Ma, Göncüoğlu, 1986) and Ağaçören granitoids (110 ± 14 Ma, Güleç, 1994) indicate a late early Cretaceous to early late Cretaceous intrusion age. Syenitic rocks from the Bayındır area, that intrude the granitoids yield Rb/Sr whole rock isochron ages of 85.1 ± 3.6 Ma (Kuruç, 1990). Rb/Sr whole rock-mineral isochron ages from granitoids (Cefalık Dağ: 71 Ma, Ataman, 1972; Üçkapılı: 78 Ma, Göncüoğlu, 1986) as well as K/Ar biotite and muscovite ages (Üçkapılı, 78 to 75 Ma, Göncüoğlu, 1986) on the other hand are interpreted as cooling ages. According to this data, the radiometric ages for the granitoid intrusions range between 110 to 85 Ma. This data is in limited accordance with the recent findings, at least with those from the central (Terlemez) and northern (Çiçekdağ) parts of the CACC, where fossil-bearing epiophiolitic sediments of early Middle Turonian to latest Coniacian are intruded by monzonitic rocks. K/Ar mineral data from the Terlemez quartz-monzonite (81 to 67 Ma, Yalınız et al., 1997) as well as the regional transgression of latest Maastrichtian-earliest Paleocene marine-lacustrine sediments on the CACC (Göncüoğlu, 1986; Göncüoğlu et al, 1991) puts further restraints on the age of the granitic intrusions.

Geochemical Framework

Geochemical data presented in this paper are compiled from published as well as unpublished data sets. Data for the Yozgat, Ağaçören, Ekecikdağ, Cefalıkdağ, İdişdağı and Atdere intrusions are from Aydın et al. (1998), Kadioğlu (1996), Türeli (1991), Geven (1992), Türeli et al. (1993), Göncüoğlu et al., (1997) and Özkan and Erkan (1994), respectively.

The main whole rock geochemical features of the granitoids of the CACC are as follows. The granitoids are subalkaline (Fig. 2) and exhibit well-developed calc-alkaline trend on the AFM diagram of Irvine and Baragar (1971) (Fig. 3). The H-type granitoids of the CACC are metaluminous, whereas C-type granitoids are slightly to strongly peraluminous (Fig. 4). Ocean ridge granite (ORG) normalized spider diagram is utilized to present the trace element chemistry of the granitoids of the CACC (Fig. 5). In general, samples from the complex reveal large ion lithophile elements (LILE- e.g., K_2O , Rb) enrichment relative to the high field strength elements (HFSE- e.g., Zr, Y). However, the C-type granitoids and aplitic dykes show strong negative Ba-anomalies together with moderate Zr anomalies, whereas the H-type

granitoids display slightly to moderately developed negative Ba- and Zr anomalies.

The İdişdağı syenitoids are mostly subalkaline to slightly alkaline, whereas the feldspathoid bearing syenitoids (the Atdere Intrusion) are strongly alkaline (Fig. 2). Both of the syenitoid intrusions are typically rich in alkaline content and concentrated on the alkaline corner of the AFM diagram (Fig. 3) of the Irvine and Baragar (1971). The İdişdağı syenitoids are metaluminous, whereas the Atdere syenitoids are slightly peralkaline (Fig. 4). On the ORG normalized spider diagram, the İdişdağı Intrusion display weakly developed negative Ba- and Nb-anomalies (Fig. 5). Note that there is no enough data from the Atdere Intrusion to compare the trace-element geochemistry of the intrusion with those of the İdişdağı Intrusion.

Trace element classification diagrams of Pearce et al. (1984) are utilized to present the tectonic settings of the granitoids and syenitoids of the CACC (Fig. 6). Most of the investigated samples plot close to triple junction of WPG (Within Plate Granite), Syn-COLG (Syn-Collisional Granite) and VAG (Volcanic Arc Granite). In detail, the C-type granitoid rocks plot on WPG and Syn-COLG fields, whereas the H-type granitoids fall into WPG and Syn-COLG fields. Samples from the İdişdağı Intrusion plot mainly on the VAG, whereas samples from the Atdere Intrusion fall within the Syn-COLG and WPG fields.

Summary and Conclusions

Granitoid and syenitoid rocks dominate the geology of the CACC and a wide range of composition. However, there are two main granitoid types in the complex: a) the C-type leucogranites and b) the H-type, hornblende \pm K-feldspar megacrysts \pm mafic microgranular enclave bearing granites. The emplacement of granitoid magmatism was followed by the syenitoid magmatism where the quartz syenitoids (e.g., the İdişdağı Intrusion) predate the feldspathoid bearing syenitoids (e.g., the Atdere Intrusion). The type of magmatism varies from peraluminous, metaluminous to alkaline/ peralkaline through time.

Petrographic, field and geochemical features of the leucogranites of the CACC including: a) their peraluminous nature; b) the presence of garnet; c) the lack of ferromagnesian minerals, K-feldspar megacrysts and mafic microgranular enclaves, are typical of crustal derived granitoids (Barbarin, 1990). Although the C-type granitoids of the complex plot within both Syn-COLG and WPG, they typically have low Zr, Ce, Sm, and Ba contents




Figure 2. Plots of the total alkali versus SiO₂ diagram where alkaline and subalkaline fields are from Irvine and Baragar (1971).

which is inconsistent with WPG. Thus, the C-type granitoids can be considered to be products of syn-collisional magmatism which can be attributed to crustal thickening with or without any direct contribution from the mantle-derived mafic magma. They may have been derived from upper crustal sedimentary rocks.

The H-type granitoids of the CACC are metaluminous, locally includes mafic microgranular enclaves and/or K-feldspar megacrysts and contain hornblende ± clinopyroxene ± biotite as the dominant ferromagnesian minerals. These features require a mantle derived mafic magma contribution in the genesis of these rocks which can be explained in terms of mafic magma underplating




Figure 3. Plots of AFM diagram where tholeiitic and calc-alkaline fields are from Irvine and Baragar (1971). Symbols as in Fig. 2.

of lower crust as a result of lithospheric delamination following the crustal thickening. With this in mind, the H-type granitoids can be considered products of post-collisional magmatism of the CACC.

Syenitoid rocks are slightly to strongly alkaline. Although, they display similar trace element geochemical features with those of the WPG, they are considered post-collisional alkaline magmatism in the CACC on the




Figure 4. Shand Index diagram after Maniar and Piccoli (1989). Symbols as in Fig. 2.

basis of their mineralogical and geochemical features (Göncüoğlu et al., 1997; Boztuğ, 1997). Note that type of syenitoid magmatism vary from silica-(over)saturated to silica-undersaturated intrusions. This transition might reflect source characteristics or can simply be explained in terms of variation in availability of water in the melting zone (cf., Bonnin, 1988; Boztuğ, 1997).

The main implications of these observations are as follows. Through time, a) type of magmatism in the CACC varies from peraluminous, metaluminous to alkaline; b) the nature of the magmatic processes and relative contributions of source materials vary; c) relative contribution of mantle-derived mafic magma in the genesis of the magmatic rocks of the complex increases;




Figure 5. Ocean ridge granite (ORG) normalized spider diagrams. Normalizing values are from Pearce et al. (1984). Symbols as in Fig. 2. Note that samples from the Cefalikdağ intrusion represent only the H-type granitoids (due to simply availability of samples).




Figure 6. Tectonic discrimination diagrams of Rb versus Y+N and Nb versus Y (after Pearce et al., 1984). Symbols as in Fig. 2.

d) tectonic setting vary from syn-collisional to post-collisional; e) products of post-collisional magmatism vary from the H-type granitoids to the syenitoids ; f) syenitoid magmatism vary from silica-oversaturated/saturated to silica-undersaturated.

However, the presented common framework for the magmatism in the CACC which is summarized in Table 1, need to be substantiated by relevant geochronologic and isotopic data.




Table 1. Summary of diverse magmatism in the Central Anatolian Crystalline Complex (CACC).

References

- Akıman, O., Erler, A., Göncüoğlu, M.C., Güleç, N., Geven, A., Türeli, T.K., Kadioğlu, Y.K., 1993. Geochemical characteristics of the granitoids along the western margin of the Central Anatolian Crystalline Complex and their tectonic implications. *Geol. Jour.*, 28: 371-382.
- Ataman, G., 1972. Ankara'nın güneydoğusundaki granitik granodiyoritik kütlelerden Cefalık Dağı'nın radyometrik yaşı hakkında ön çalışma. *Hacettepe Fen ve Mühendislik Bilimleri Dergisi*, 2, 44 - 49.
- Ayan, M., 1963. Contribution a l'étude petrographique et geologique de la region situee au Nord-Est de Kaman, *Mineral Research Exploration Publ.*, No.115, 332 p.
- Aydın, N.S., 1997. Petrography, trace-element geochemistry and petrology of the Yozgat Batholith. *Çukurova Univ. 20 th Anniv. Geol. Edu. Symp.*, Adana, Abstracts, p.244 .
- Aydın, N.S., Göncüoğlu, M.C., and Erler., E., 1997. Granitoid magmatism in Central Anatolia. *Proceedings TÜBİTAK-BAYG/NATO-D Program on Alkaline Magmatism in Central Anatolia* (ed): Boztuğ, D., Yılmaz-Şahin, S., Otlu, N., and Tatar, S., Sivas, Turkey. p.202.
- Aydın, N.S., Göncüoğlu, M.C., Erler, A. and Malpas, J., 1998. Diverse magmatism in the Central Anatolian Crystalline Complex, Turkey. *Int. Volcanological Congress, Abstracts*.
- Barbarin, B. 1990. Granitoids: main petrogenetic classifications in relation to origin and tectonic setting. *Geol. Jour.*, 25: 227-238.
- Bayhan, H. and Tolluoğlu, A.Ü., 1987. Çayağzı siyenitoidinin (Kırşehir Kuzeybatısı) mineralojik-petrografik ve jeokimyasal özellikleri. *Hacettepe Üniversitesi Yerbilimleri Dergisi*, 14, 109-120 (in Turkish).
- Bonnin, B., 1988. From orogenic to anorogenic environments: evidence from associated magmatic episodes. *Schweiz. Mineral. Petrogr. Mitt.*, 68, 301-311.
- Boztuğ, D., 1997. Post-collisional Central Anatolian alkaline plutonism, Turkey. *Proceedings TÜBİTAK-BAYG/NATO-D Program on Alkaline Magmatism in Central Anatolia* (ed): Boztuğ, D., Yılmaz-Şahin, S., Otlu, N., and Tatar, S., Sivas, Turkey. 105-146.
- Erler, A. and Bayhan, H., 1995. General evaluation and problems of the Central Anatolian granitoids. *Yerbilimleri*, 17: 49-67 (in Turkish).
- Erler, A. and Göncüoğlu M.C., 1996. Geologic and tectonic setting of the Yozgat Batholith, Northern Central Anatolian Crystalline Complex, Turkey. *Inter. Geol. Review*, 38: 714-726.
- Erler, A., Akıman, O., Unan, C., Dalkılıç, F., Dalkılıç, B., Geven, A., and Önen, P., 1991. Petrology and geochemistry of the magmatic rocks of the Kırşehir Massif at Kaman (Kırşehir) and Yozgat. *Doğa, Türk. Jour. Eng. and Env. Sci.*, 15: 76-100 (in Turkish).
- Geven, A., 1992. Mineralogy, petrography and geochemistry of the Cefalıkdağ plutonic rocks (Kaman region, Central Anatolia). Ph.D. Thesis, METU-Ankara, 165 p. (Unpublished).
- Göncüoğlu, M. C., 1986. Geochronological data from the southern part (Niğde Area) of the Central Anatolian Massif. *Mineral Research Exploration Bull.*, 105 / 106, 83 - 96.
- Göncüoğlu, M.C., and Türeli, T.K., 1994. Alpine collision type granitoids in the western Central Anatolian Crystalline Complex. *Jour. Kocaeli Univ.* 1: 39-46.

- Göncüoğlu, M.C., Toprak, V., Kuşcu, I., Erler, A. and Olgun E., 1991. Geology of the western part of the Central Anatolian Massif, Part 1: Southern Section: Unpubl. Report No.2909, Turkish Petroleum Company Report (in Turkish).
- Göncüoğlu, M.C., Erler, A., Dirik, K., Olgun, E., and Rojay, B., 1992. Geology of the western part of the Central Anatolian Massif, Part: 2: Central Section: Unpubl. Report No. 3155. Turkish Petroleum Company Report (in Turkish)
- Göncüoğlu, M.C., Erler, A., Toprak, V., Olgun, E., Yalınız, K., Kuşcu, I., Köksal, S., and Dirik, K., 1993. Geology of the central part of the Central Anatolian Massif, Part 3: Geologic evolution of the Central Kızılırmak Tertiary Basin: Unpubl. Report No.3313, Turkish Petroleum Company Report (in Turkish).
- Göncüoğlu, M.C., Dirik, K., Erler, A., and Yalınız, K., 1994. Geology of the Eastern part of the Central Anatolian Massif, Part 4: The relationship between the Central Anatolian Massif and Sivas Basin. METU-TPAO Report, 135 p. (in Turkish).
- Göncüoğlu, M.C., Köksal, S., and Floyd, P.A., 1997. Post-collisional A-type magmatism in the Central Anatolian Crystalline Complex: Petrology of the İdişdağı intrusives (Avanos, Turkey). Turkish Journal of Earth Sciences, 6, 65-76.
- Güleç, N., 1994. Rb-Sr Isotope Data From the Ağaçören Granitoid (East of Tuz Gölü): Geochronological and Genetical implications, Turkish Journal of Earth Sciences, 3, 39-43.
- Irvine, T.N. and Baragar, W.R.A., 1971. A guide to the chemical classification of the common volcanic rocks. Can. Jour. Earth Sci., 8: 523-548.
- Kadioğlu, Y.K., 1996. Genesis of Ağaçören Intrusive Suite and its enclaves (Central Anatolia): Constraints from geological, petrographic, geophysical and geochemical data. Ph.D. Thesis. METU-Ankara, 242 p. (unpublished).
- Kadioğlu, Y.K., and Güleç, N., 1996. Mafic microgranular enclaves and interaction between felsic and mafic magmas in the Ağaçören Intrusive Suite: Evidence from petrographic features and mineral chemistry. Inter. Geol. Review, 38: 854-867.
- Kuruç, A., 1990. Rb/Sr geochemistry of syenitoids from Kaman-Kırşehir region. Hacettepe University, M.S. Thesis, 97 p., (unpublished, in Turkish).
- Maniar, P.D. and Piccoli, P.M., 1989. Tectonic discrimination of granitoids. Geol. Soc. America Bull., 101: 635-643.
- Otlu, N., and Boztuğ, D., 1997. The coexistence of the silica oversaturated (Alkos) and undersaturated alkaline (Alkus) rocks in the Kortundağ and Baranadağ plutons from the central Anatolian alkaline plutonism, E Kaman/NW Kırşehir, Turkey. Proceedings TÜBİTAK-BAYG/NATO-D Program on Alkaline Magmatism in Central Anatolia (ed): Boztuğ, D., Yılmaz-Şahin, S., Otlu, N., and Tatar, S., Sivas, Turkey. 85-104.
- Özkan, H.M., 1987. Petrographical and geochemical investigation of Hayriye (Kayseri) nepheline-syenite intrusion. M.S. Thesis, Ankara University-Ankara, 33 p. (unpublished, in Turkish)
- Özkan, H.M. and Erkan, Y., 1994. A petrological study on a foid syenite intrusion in Central Anatolia (Kayseri, Turkey). Turkish Journal of Earth Sciences, 3, 45-55.
- Pearce, J.A. Harris, N.B.W. and Tindle, A.G. 1984. Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. Journal of Petrology, 25: 956-983.
- Seymen, I., 1982. Geology of the Kırşehir Massif around Kaman. Unpubl. Hab. Thesis, İst. Tech. Univ., Faculty of Mines, 164 p. (in Turkish).
- Tatar, S., and Boztuğ, D., 1997. Fractional crystallization and magma mingling/mixing processes in the monzonitic association in the SW part of the composite Yozgat Batholith (Şefaatli-Yerköy, SW Yozgat). Proceedings TÜBİTAK-BAYG/NATO-D Program on Alkaline Magmatism in Central Anatolia (ed): Boztuğ, D., Yılmaz-Şahin, S., Otlu, N., and Tatar, S., Sivas, Turkey. 146-164.
- Türel, T.K., 1991. Geology, petrography and geochemistry of Ekecikdağ plutonic rocks (Aksaray Region-Central Anatolia). Ph. D. Thesis, METU-Ankara, 194 p. (unpublished).
- Türel, T.K., Göncüoğlu, M.C. and Akıman, O., 1993. Ekecikdağ Granitoidinin petrolojisi ve kökeni. Mineral Research Exploration Bull., 115, 15-28.
- Yalınız, K., Parlak, O., Özkan-Altın, S., and Göncüoğlu, M.C., 1997. Formation and emplacement ages of supra-subduction-type ophiolites in Central Anatolia; Sarıkaraman Ophiolites, Central Turkey. Çukurova Univ. 20 th Anniv. Geol. Edu. Symp., Adana, Abstracts, 61-62.