

1-1-2018

## The Brezis-Lieb lemma in convergence vector lattices

MOHAMMAD MARABEH

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

MARABEH, MOHAMMAD (2018) "The Brezis-Lieb lemma in convergence vector lattices," *Turkish Journal of Mathematics*: Vol. 42: No. 3, Article 54. <https://doi.org/10.3906/mat-1708-7>

Available at: <https://dctubitak.researchcommons.org/math/vol42/iss3/54>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

## The Brezis–Lieb lemma in convergence vector lattices

Mohammad MARABEH\*

Department of Applied Mathematics, College of Sciences and Arts, Palestine Technical University-Kadoorie,  
Tulkarem, Palestine

Received: 03.08.2017

Accepted/Published Online: 05.01.2018

Final Version: 08.05.2018

**Abstract:** Recently measure-free versions of the Brezis–Lieb lemma were proved for unbounded order convergence in vector lattices. In this article, we extend these versions to convergence vector lattices.

**Key words:** Vector lattice, unbounded order convergence, almost order bounded set, Brezis–Lieb lemma, convergence vector space, convergence vector lattice

### 1. Introduction

The Brezis–Lieb lemma [4, Theorem 2] has numerous applications mainly in calculus of variations (see for example [5, 9]). We begin with its statement. Let  $j : \mathbb{C} \rightarrow \mathbb{C}$  be a continuous function with  $j(0) = 0$ . In addition, let  $j$  satisfy the following hypothesis: for every sufficiently small  $\varepsilon > 0$ , there exist two continuous, nonnegative functions  $\varphi_\varepsilon$  and  $\psi_\varepsilon$  such that

$$|j(a+b) - j(a)| \leq \varepsilon \varphi_\varepsilon(a) + \psi_\varepsilon(b) \quad (1)$$

for all  $a, b \in \mathbb{C}$ . The following result has been stated and proved by Brezis and Lieb in [4].

**Theorem 1.1** (Brezis–Lieb lemma, [4, Theorem 2]). *Let  $(\Omega, \Sigma, \mu)$  be a measure space. Let the mapping  $j$  satisfy the above hypothesis, and let  $f_n = f + g_n$  be a sequence of measurable functions from  $\Omega$  to  $\mathbb{C}$  such that:*

- (i)  $g_n \xrightarrow{\text{a.e.}} 0$ ;
- (ii)  $j \circ f \in L_1$ ;
- (iii)  $\int \varphi_\varepsilon \circ g_n d\mu \leq C < \infty$  for some  $C$  independent of  $\varepsilon$  and  $n$ ;
- (iv)  $\int \psi_\varepsilon \circ f d\mu < \infty$  for all  $\varepsilon > 0$ .

Then, as  $n \rightarrow \infty$ ,

$$\int |j(f + g_n) - j(g_n) - j(f)| d\mu \rightarrow 0. \quad (2)$$

\*Correspondence: m.maraabeh@gmail.com

2010 AMS Mathematics Subject Classification: 54A20, 46A40, 46B42

Recall that a subset  $A$  in a normed lattice  $(X, \|\cdot\|)$  is said to be *almost order bounded* if for any  $\varepsilon > 0$  there is  $u_\varepsilon \in X_+$  such that  $\sup_{a \in A} \| |a| - u_\varepsilon \wedge |a| \| < \varepsilon$ . Motivated by the proof of Theorem 1.1 the following results were proven in [7].

**Proposition 1.2** [7, Proposition 1.2] (*Brezis–Lieb lemma for mappings on  $L_0$* ). Let  $(\Omega, \Sigma, \mu)$  be a measure space,  $f_n = f + g_n$  be a sequence in  $L_0(\mu)$  such that  $g_n \xrightarrow{\text{a.e.}} 0$ , and  $J : L_0(\mu) \rightarrow L_0(\mu)$  be a mapping satisfying  $J(0) = 0$  that preserves almost everywhere convergence and such that the sequence  $(J(f_n) - J(g_n))_{n \in \mathbb{N}}$  is almost order bounded. Then, as  $n \rightarrow \infty$ ,

$$\int |J(f + g_n) - (J(g_n) + J(f))| d\mu \rightarrow 0. \quad (3)$$

**Proposition 1.3** [7, Proposition 1.3] (*Brezis–Lieb lemma for uniform integrable sequence  $(J(f_n) - J(g_n))_{n \in \mathbb{N}}$* ). Let  $(\Omega, \Sigma, \mu)$  be a finite measure space,  $f_n = f + g_n$  be a sequence in  $L_0(\mu)$  such that  $g_n \xrightarrow{\text{a.e.}} 0$ , and  $J : L_0(\mu) \rightarrow L_0(\mu)$  be a mapping satisfying  $J(0) = 0$  that preserves almost everywhere convergence and such that the sequence  $(J(f_n) - J(g_n))_{n \in \mathbb{N}}$  is uniformly integrable. Then

$$\lim_{n \rightarrow \infty} \int |J(f + g_n) - (J(g_n) + J(f))| d\mu = 0. \quad (4)$$

Recall that a sequence  $(x_n)$  in a vector lattice  $X$  is *order convergent* (or *o-convergent*, for short) to  $x \in X$  if there is a sequence  $(z_n)$  in  $X$  satisfying  $z_n \downarrow 0$  and  $|x_n - x| \leq z_n$  for all  $n \in \mathbb{N}$  (we write  $x_n \xrightarrow{o} x$ ); see e.g. [10, Theorem 16.1]. In a vector lattice  $X$ , a sequence  $(x_n)$  is *unbounded order convergent* (or *uo-convergent*, for short) to  $x \in X$  if  $|x_n - x| \wedge y \xrightarrow{o} 0$  for all  $y \in X_+$  (we write  $x_n \xrightarrow{uo} x$ ). It is well known that if  $(\Omega, \Sigma, \mu)$  is a measure space, then in  $L_p$  spaces ( $1 \leq p \leq \infty$ ), *uo-convergence* of sequences is the same as the almost everywhere convergence; see e.g. [8].

**Definition 1.1** [7, page 23] A mapping  $f : X \rightarrow Y$  between vector lattices is said to be  *$\sigma$ -unbounded order continuous* (in short,  *$\sigma$ uo-continuous*) if  $x_n \xrightarrow{uo} x$  in  $X$  implies  $f(x_n) \xrightarrow{uo} f(x)$  in  $Y$ .

In [7] we gave two variants of the Brezis–Lieb lemma in vector lattice setting by replacing *a.e.*-convergence by *uo-convergence*, integral functionals by strictly positive functionals, and the continuity of the scalar function  $j$  (in Theorem 1.1) by the  $\sigma$ -unbounded order continuity of the mapping  $J : X \rightarrow Y$  between vector lattices  $X$  and  $Y$ . As standard references for basic notions on vector lattices we adopt the books [1, 10, 14] and on unbounded order convergence the paper [8]. In this article, all vector lattices are assumed to be Archimedean.

Let  $Y$  be a vector lattice and  $l$  be a strictly positive linear functional on  $Y$ . Define the following norm on  $Y$ :

$$\|y\|_l := l(|y|). \quad (5)$$

Then the  $\|\cdot\|_l$ -completion  $(Y_l, \|\cdot\|_l)$  of  $(Y, \|\cdot\|_l)$  is an *AL-space*, and so it is an order continuous Banach lattice. The following result is a measure-free version of Proposition 1.2.

**Proposition 1.4** [7, Proposition 2.2] (A Brezis–Lieb lemma for strictly positive linear functionals). Let  $X$  be a vector lattice and  $Y_l$  be the AL-space constructed above. Let  $J : X \rightarrow Y_l$  be  $\sigma uo$ -continuous with  $J(0) = 0$ , and  $(x_n)$  be a sequence in  $X$  such that:

1.  $x_n \xrightarrow{uo} x$  in  $X$ ;
2. the sequence  $(J(x_n) - J(x_n - x))_{n \in \mathbb{N}}$  is almost order bounded in  $Y_l$ .

Then

$$\lim_{n \rightarrow \infty} \|J(x_n) - J(x_n - x) - J(x)\|_l = 0. \quad (6)$$

Similar to Proposition 1.4 one can easily show the following result.

**Proposition 1.5** Let  $X$  and  $Y$  be vector lattices and  $l$  be a strictly positive linear functional on  $Y$ . Let  $J : X \rightarrow Y$  be  $\sigma uo$ -continuous with  $J(0) = 0$ , and  $(x_n)$  be a sequence in  $X$  such that:

1.  $x_n \xrightarrow{uo} x$  in  $X$ ;
2. the sequence  $(J(x_n) - J(x_n - x))_{n \in \mathbb{N}}$  is almost order bounded in  $Y$ .

Then

$$\lim_{n \rightarrow \infty} \|J(x_n) - J(x_n - x) - J(x)\|_l = 0. \quad (7)$$

The next result is another measure-free version of Proposition 1.2.

**Proposition 1.6** [7, Proposition 2.3] (A Brezis–Lieb lemma for  $\sigma uo$ -continuous linear functionals). Let  $X, Y$  be vector lattices and  $l$  be a  $\sigma uo$ -continuous functional on  $Y$ . Assume further  $J : X \rightarrow Y$  is a  $\sigma uo$ -continuous mapping with  $J(0) = 0$  and  $(x_n)$  is a sequence in  $X$  such that  $x_n \xrightarrow{uo} x$ . Then

$$\lim_{n \rightarrow \infty} l(J(x_n) - J(x_n - x) - J(x)) = 0. \quad (8)$$

## 2. Two variants of the Brezis–Lieb lemma in convergence vector lattices

Recently, order convergence in vector lattices has been studied from the viewpoint of convergence structures; see e.g. [2, 12, 13], and in this section we recall basic notions related to convergence structures and use them to investigate variants of the Brezis–Lieb lemma. For unexplained notions and results we refer the reader to the monograph [3].

A filter  $\mathcal{F}$  on a set  $X$  is a nonempty collection of subsets of  $X$  that does not contain the empty set and is closed under the formation of finite intersections and supersets. A subset  $\mathcal{B}$  of a filter  $\mathcal{F}$  is called a *basis* of  $\mathcal{F}$  and  $\mathcal{F}$  the filter *generated* by  $\mathcal{B}$  if each set in  $\mathcal{F}$  contains a set of  $\mathcal{B}$ . If  $\mathcal{B}$  is a collection of nonempty subsets of  $X$  that is directed downward with respect to inclusion, then  $[\mathcal{B}] = \{F \subseteq X : B \subseteq F \text{ for some } B \in \mathcal{B}\}$  is the filter generated by  $\mathcal{B}$ . If  $\mathcal{B} = \{x\}$  then we write  $[x]$  for  $[\{x\}]$ . If  $f : X \rightarrow Y$  is a mapping then  $\{f(F) : F \in \mathcal{F}\}$  is the basis of a filter called the *image filter* of  $\mathcal{F}$  under  $f$  and denoted by  $f(\mathcal{F})$ .

**Definition 2.1** Let  $X$  be a set. A mapping  $\lambda$  from  $X$  into the power set of the set of all filters on  $X$  is called a *convergence structure* on  $X$  and  $(X, \lambda)$  a *convergence space* if the following hold for all  $x \in X$ :

1.  $[x] \in \lambda(x)$ ;
2. for all filters  $\mathcal{F}, \mathcal{G} \in \lambda(x)$  the intersection  $\mathcal{F} \cap \mathcal{G} \in \lambda(x)$ ;
3. if  $\mathcal{F} \in \lambda(x)$ , then  $\mathcal{G} \in \lambda(x)$  for all filters  $\mathcal{G}$  on  $X$  for which  $\mathcal{G} \supseteq \mathcal{F}$ .

Let  $(X, \lambda)$  be a convergence space. A filter  $\mathcal{F}$  on  $X$  converges to  $x$  in the space  $X$  if  $\mathcal{F} \in \lambda(x)$ . We write  $\mathcal{F} \rightarrow x$  in  $(X, \lambda)$  or  $\mathcal{F} \rightarrow x$  in  $X$ . The element  $x$  is called the *limit* of  $\mathcal{F}$ .

Every topological space is a convergence space; see e.g. [3, Examples 1.1.2(i)], but the converse is not necessarily true; see e.g. [3, Examples 1.1.2(iii)]. Throughout this article,  $\mathbb{R}$  refers to the convergence space induced by its standard topology. That is, a filter  $\mathcal{F} \rightarrow x$  in  $\mathbb{R}$  iff  $\mathcal{F} \supseteq \mathcal{U}_x$ , where  $\mathcal{U}_x$  is the neighbourhood filter of  $x$  (the collection of all topological neighbourhoods of  $x$ ).

Let  $X$  and  $Y$  be convergence spaces. A mapping  $f : X \rightarrow Y$  is called *continuous at a point*  $x \in X$  if  $f(\mathcal{F}) \rightarrow f(x)$  in  $Y$  whenever  $\mathcal{F} \rightarrow x$  in  $X$ . The mapping  $f$  is called *continuous* if it is continuous at every point of  $X$ .

If  $\mathcal{F}$  is a filter on  $X_1$  and  $\mathcal{G}$  is a filter on  $X_2$  then the *product filter*  $\mathcal{F} \times \mathcal{G}$  is the filter on  $X_1 \times X_2$  generated by  $\{F \times G : F \in \mathcal{F}, G \in \mathcal{G}\}$ . Let  $X_1$  and  $X_2$  be convergence spaces; then a filter  $\mathcal{F}$  on  $X_1 \times X_2$  converges to  $(x_1, x_2)$  iff  $p_i(\mathcal{F})$  converges to  $p_i((x_1, x_2))$ , for each  $i \in \{1, 2\}$ , where  $p_i$  is the projection of  $X_1 \times X_2$  onto  $X_i$ .

Given a convergence space  $(X, \lambda)$ , for a sequence  $(x_n)$  in  $X$  the *Fréchet filter* of  $(x_n)$  denoted by  $\langle (x_n) \rangle$  is the filter generated by  $\{A_k\}_{k \in \mathbb{N}}$ , where  $A_k = \{x_n : n \geq k\}$ . A sequence  $(x_n)$  converges to a point  $x \in X$  if the Fréchet filter  $\langle (x_n) \rangle$  converges to  $x$  in  $X$ . In this case we write  $x_n \xrightarrow{\lambda} x$ .

Let  $X$  be a real vector space. A convergence structure  $\lambda$  on  $X$  is called a *vector space convergence structure* and  $(X, \lambda)$  a *convergence vector space* if addition and scalar multiplication are continuous.

**Definition 2.2** Let  $X$  be a vector lattice. A vector space convergence structure  $\lambda$  on  $X$  is called a *vector lattice convergence structure* and  $(X, \lambda)$  a *convergence vector lattice* whenever lattice operations are continuous.

It is obvious by definition that every convergence vector lattice is a convergence vector space. The converse is not necessarily true. Let  $X = (L_1[0, 1], \tau_w)$ , where  $\tau_w$  denotes the weak topology on  $L_1[0, 1]$ . Then  $X$  is a linear topological vector space and so it is a convergence vector space, but  $X$  is not a convergence vector lattice.

**Definition 2.3** Let  $(X, \lambda)$  be a convergence vector lattice. A sequence  $(x_n)$  in  $X$  is unbounded convergent to  $x \in X$  if  $|x_n - x| \wedge y \xrightarrow{\lambda} 0$  for all  $y \in X_+$ . We write  $x_n \xrightarrow{u\lambda} x$  and say  $x_n$   *$u\lambda$ -converges* to  $x$ .

Given a locally solid vector lattice  $(X, \tau)$  and a sequence  $(x_n)$  in  $X$ , then  $X$  is a convergence vector lattice and  $x_n$   *$u\lambda$ -converges* to  $x \in X$  iff  $x_n$   *$u\tau$ -converges* to  $x$ ; see [6]. The following notion is motivated by Definition 1.1.

**Definition 2.4** Let  $X$  and  $Y$  be two convergence vector lattices. A mapping  $f : X \rightarrow Y$  is called  *$\sigma u$ -continuous* if  $f(x_n) \xrightarrow{u\lambda} f(x)$  in  $Y$  whenever  $x_n \xrightarrow{u\lambda} x$  in  $X$ .

The next result should be compared with Proposition 3.7 in [8].

**Lemma 2.1** Let  $(X, \lambda)$  be a convergence vector lattice and  $\|\cdot\| : X \rightarrow \mathbb{R}$  be a continuous lattice norm. If a sequence  $(x_n)$  is almost order bounded in  $(X, \|\cdot\|)$  and  $x_n \xrightarrow{u\lambda} x$ , then  $x_n \xrightarrow{\|\cdot\|} x$ .

**Proof** It is easy to see that the sequence  $(|x_n - x|)_{n \in \mathbb{N}}$  is almost order bounded. Since  $x_n \xrightarrow{u\lambda} x$  then  $|x_n - x| \wedge y \xrightarrow{\lambda} 0$  for all  $y \in X_+$ . The continuity of the lattice norm assures that  $|x_n - x| \wedge y \xrightarrow{\|\cdot\|} 0$  for all  $y \in X_+$ . The remaining part of the proof is the same as in [8, Proposition 3.7].  $\square$

The following two results generalize Propositions 1.5 and 1.6, respectively.

**Theorem 2.2** Let  $X$  and  $Y$  be convergence vector lattices and  $l : Y \rightarrow \mathbb{R}$  be a continuous strictly positive linear functional. Suppose  $J : X \rightarrow Y$  is  $\sigma u$ -continuous mapping with  $J(0) = 0$  and  $(x_n)$  is a sequence in  $X$  such that:

1.  $x_n \xrightarrow{u\lambda} x$  in  $X$ ;
2. the sequence  $(J(x_n) - J(x_n - x))_{n \in \mathbb{N}}$  is almost order bounded in  $(Y, \|\cdot\|_l)$ .

Then

$$\lim_{n \rightarrow \infty} \|J(x_n) - J(x_n - x) - J(x)\|_l = 0. \quad (9)$$

**Proof** Since  $x_n \xrightarrow{u\lambda} x$  and  $J$  is  $\sigma u$ -continuous, then  $J(x_n) \xrightarrow{u\lambda} J(x)$  and  $J(x_n - x) \xrightarrow{u\lambda} J(0) = 0$ . Thus,  $J(x_n) - J(x_n - x) \xrightarrow{u\lambda} J(x)$ . Since  $l : Y \rightarrow \mathbb{R}$  is a continuous linear functional then the norm  $\|\cdot\|_l$  given by (5) is continuous on  $Y$ . Thus it follows from Lemma 2.1 that  $\|J(x_n) - J(x_n - x) - J(x)\|_l \rightarrow 0$  as  $n \rightarrow \infty$ .  $\square$

**Theorem 2.3** Let  $X$  and  $Y$  be convergence vector lattices. Supposing  $l : Y \rightarrow \mathbb{R}$  is a  $\sigma u$ -continuous linear functional,  $J : X \rightarrow Y$  is  $\sigma u$ -continuous mapping with  $J(0) = 0$  and  $x_n \xrightarrow{u\lambda} x$  in  $X$ . Then

$$\lim_{n \rightarrow \infty} l(J(x_n) - J(x_n - x) - J(x)) = 0. \quad (10)$$

**Proof** Since  $x_n \xrightarrow{u\lambda} x$  and  $J$  is  $\sigma u$ -continuous with  $J(0) = 0$ , then  $J(x_n) - J(x_n - x) - J(x) \xrightarrow{u\lambda} 0$ . The conclusion now follows from the  $\sigma u$ -continuity of the linear functional  $l$ .  $\square$

Finally, we show that Propositions 1.5 and 1.6 follow from Theorems 2.2 and 2.3.

Let  $X$  be a vector lattice and  $x \in X$ . The following relation

$$\mathcal{F} \in \lambda_o(x) \Leftrightarrow \{[u_n, v_n] : n \in \mathbb{N}\} \subseteq \mathcal{F}, \quad (11)$$

with  $(u_n), (v_n) \subseteq X$  increasing and decreasing to  $x$ , respectively, defines a vector lattice convergence structure on  $X$  and so  $(X, \lambda_o)$  is a convergence vector lattice; see [12, Corollary 12, Theorem 14, and Proposition 15]. Moreover, a sequence  $(x_n)$  in  $X$  converges to  $x \in X$  iff  $(x_n)$  order converges to  $x$ ; see e.g. [12, Corollary 12], [2, Theorem 16(iii)], or [11, Definition II.1.7].

Let  $X$  and  $Y$  be vector lattices equipped with the vector lattice convergence structure  $\lambda_o$  given in (11). Then  $x_n \xrightarrow{u\lambda} x$  in  $(X, \lambda_o)$  iff  $x_n \xrightarrow{uo} x$  in  $X$  and a mapping  $f : X \rightarrow Y$  is  $\sigma u$ -continuous iff it is  $\sigma uo$ -continuous (in the sense of Definition 1.1).

Therefore, if we equip the vector lattices  $X$  and  $Y$  in Theorems 2.2 and 2.3 with the vector lattice convergence structure  $\lambda_o$ , then we obtain Propositions 1.5 and 1.6, respectively.

**References**

- [1] Aliprantis CD, Burkinshaw O. Positive Operators. Dordrecht, the Netherlands: Springer, 2006.
- [2] Anguelov R, van der Walt JH. Order convergence structure on  $C(X)$ . Quaest Math 2005; 28: 425-457.
- [3] Beattie R, Butzmann HP. Convergence Structures and Applications to Functional Analysis. Dordrecht, the Netherlands: Springer-Science+Business Media, B.V., 2002.
- [4] Brezis H, Lieb E. A relation between pointwise convergence of functions and convergence of functionals. P Am Math Soc 1983; 88: 486-490.
- [5] Brezis H, Nirenberg L. Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. Commun Pur Appl Math 1983; 36: 437-477.
- [6] Dabboorasad YA, Emelyanov EY, Marabeh MAA.  $u\tau$ -Convergence in locally solid vector lattices. preprint 2017; arXiv:1706.02006v3 [math.FA].
- [7] Emelyanov EY, Marabeh MAA. Two measure-free versions of the Brezis-Lieb Lemma. Vladikavkaz Mat Zh 2016; 18: 21-25.
- [8] Gao N, Xanthos F. Unbounded order convergence and application to martingales without probability. J Math Anal Appl 2014; 415: 931-947.
- [9] Lieb E. Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities. Ann Math 1983; 118: 349-374.
- [10] Luxemburg WAJ, Zaanen AC. Riesz Spaces, Volume I. Amsterdam, the Netherlands: North-Holland Publishing Company, 1971.
- [11] Schaefer HH. Banach Lattices and Positive Operators. Berlin, Germany: Springer-Verlag, 1974.
- [12] van der Walt JH. The order convergence structure. Indagat Math New Ser 2011; 21: 138-155.
- [13] van der Walt JH. Applications of convergence spaces to vector lattice theory. Topol P 2014; 41: 311-331.
- [14] Zaanen AC. Riesz Spaces II. Amsterdam, the Netherlands: North-Holland Publishing Company, 1983.