

1-1-2015

The assembly of virus-like particles of porcine transmissible gastroenteritisvirus in vitro by baculovirus expression system

SONG ZHENHUI

HAIBO FENG

ZHI ZHU

XIANJIN DAI

YUE ZHOU

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

Recommended Citation

ZHENHUI, SONG; FENG, HAIBO; ZHU, ZHI; DAI, XIANJIN; ZHOU, YUE; LI, YUNTIAN; and CAO, XINZHI (2015)

"The assembly of virus-like particles of porcine transmissible gastroenteritisvirus in vitro by baculovirus expression system," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 39: No. 3, Article 8.

<https://doi.org/10.3906/vet-1411-42>

Available at: <https://journals.tubitak.gov.tr/veterinary/vol39/iss3/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The assembly of virus-like particles of porcine transmissible gastroenteritisvirus in vitro by baculovirus expression system

Authors

SONG ZHENHUI, HAIBO FENG, ZHI ZHU, XIANJIN DAI, YUE ZHOU, YUNTIAN LI, and XINZHI CAO

The assembly of virus-like particles of porcine transmissible gastroenteritis virus in vitro by baculovirus expression system

Song ZHENHUI^{1*}, Haibo FENG¹, Zhi ZHU², Xianjin DAI¹, Yue ZHOU¹, Yuntian LI¹, Xinzhi CAO¹

¹Department of Veterinary Medicine, Southwest University, Rongchang Campus, Chongqing, P.R. China

²Department of Animal Science, Southwest University, Rongchang Campus, Chongqing, P.R. China

Received: 20.11.2014

Accepted/Published Online: 16.02.2015

Printed: 10.06.2015

Abstract: Virus-like particles (VLPs) produced by recombinant expression of the major viral structural proteins could be an attractive method for transmissible gastroenteritis virus (TGEV) assembly and release. In this study, the *M* gene, *sM* gene, and *N* gene of TGEV were inserted into donor plasmid pFastBac Dual. Those recombinant donor plasmids were transformed into *E.coli* DH10Bac respectively. Recombinant baculoviruses were obtained by transfection of the Sf9 cells with Bacmid-M, Bacmid-sM, and Bacmid-N. The results of indirect immunofluorescence assay showed that recombinant M protein, sM protein, and N protein were expressed successfully. In vitro experiments of VLP assembly of TGEV were performed by infection of insect cells with various combinations of recombinant baculoviruses identified. It is shown that expression of M protein alone, coexpression of M plus sM protein, and coexpression of M plus N protein in Sf9 insect cells infected by recombinant baculovirus could result in the occurrence of a virion morphologically similar to TGEV. However, the sM and N protein could not generate a structure similar to the virion of TGEV when expressed alone. This study provides a theoretical foundation for further study on the role of TGEV structural proteins in the assembly process of virions.

Key words: Transmissible gastroenteritis virus, virus-like particles, baculovirus expression system

1. Introduction

Transmissible gastroenteritis virus (TGEV) is the etiological agent of transmissible gastroenteritis, which is a condition associated with high morbidity in animals of all ages and high mortality in sucking piglets (1). TGEV is a member of the Coronaviridae family, possesses a large 28.5-kb single-stranded sense RNA genome, and comprises four structural proteins encoded by the spike (*S*), membrane (*M*), envelope (*sM*), and nucleoprotein (*N*) genes. The *S* protein forms the peplomers on the virion envelope and features a major antigenic site (2,3). The *M* protein is embedded in the lipid envelope, taking part in virus-like particle (VLP) assembly, and the *N* protein is associated with the genomic RNA to form the nucleocapsid, inducing cell immunity of infected animals (4,5). The small *sM* protein is localized in the perinuclear region of infected cells and plays an important but not fully understood role in viral morphogenesis and budding. Previous studies on coronaviruses, including TGEV, have demonstrated that the *sM* protein is essential for coronavirus particle assembly (6). Coronaviruses are assembled intracellularly at membranes of the intermediate compartment, between the endoplasmic reticulum (ER) and the Golgi complex (7–11). The highly basic *N* interacts

with genomic RNA to form helical nucleocapsids (12), and *M* interacts with nucleocapsids on cell membranes at the ER or Golgi complex (13,14). *S* and *sM* are also translated on membrane bound polysomes, inserted into the ER, and transported to the Golgi complex. This complex is where *sM* and *M* proteins interact and trigger virion budding (15). *S* is incorporated into virions via interactions with *M*. Virions accumulated in vesicles are fused with the plasma membrane and released into the extracellular space eventually (16).

Work on TGEV has established that *M* protein may serve to initiate the viral particle assembly process through interactions with genomic RNA and nucleoprotein in pre-Golgi compartments. Baudoux et al. found that coexpression of the *M* and *sM* proteins led to the formation of VLPs in cells, and pseudoparticles resembling authentic virions were released in the culture medium (17). Here we demonstrate that expression of *M* protein alone in insect Sf9 cells infected by recombinant baculovirus could form a virion morphologically similar to TGEV. This finding is not in accordance with previous results. Our studies also found that TGEV VLPs could be recovered in insect cells coexpressing a combination of *M* plus *sM* or *M* plus *N* proteins.

* Correspondence: szh7678@126.com

2. Materials and methods

2.1. Cell and culture conditions

Sf9 cells were maintained at 27 °C in Grace's insect medium supplemented with 10% fetal bovine serum and 10 µg/mL gentamicin.

2.2. Plasmid constructions

The plasmid genes encoding the membrane protein (*M*), the small envelope protein (*sM*), and the nucleocapsid protein (*N*) of TGEV, cloned into a PMD18-T vector (TaKaRa, Japanese), were constructed in our laboratory. The full-length *M*, *sM*, and *N* genes were amplified by polymerase chain reaction (PCR) using primers as follows: *M* gene forward primer 5'-GGGGAT-CCCCACCATGAAGATTTTGTTAAT-3'; *M* gene reverse primer 5'-GGGGAATTCTTATAC-CATATGTAATAATTTTCTTG-3'; *sM* gene forward primer 5'-GGGCCCCGGGATGAC-GTTTCCTAGGGCATTG-3'; *sM* gene reverse primer 5'-GGGGGTACCTCAAGCAAGGAGT-GCTCCATC-3'; *N* gene forward primer 5'-AACCCCGGGATGGCCAACC-3'; *N* gene reverse primer 5'-AGCTCGAGCATCTCGTTAG -3'. The *M* gene was subcloned into the pFastBac Dual plasmid vector (Invitrogen, USA) between the *Bam*H I and *Eco*R I restriction sites, the *sM* gene was subcloned into the pFastBac Dual plasmid vector between the *Sma* I and *Kpn* I restriction sites, and the *N* gene was subcloned into the pFastBac Dual plasmid vector between the *Sma* I and *Xho* I restriction sites. The correct orientation of the insertions was examined by PCR and restriction enzyme analysis. Plasmid pFastBac Dual-M, pFastBac Dual-sM, and pFastBac Dual-N were transformed into *E. coli* DH10Bac respectively. Recombinant baculovirus rBac-M, rBac-sM, and rBac-N were obtained by transfection of the Sf9 cells with the constructs. The recombinant baculoviruses individually expressed the *M*, *sM*, and *N* proteins.

2.3. Immunofluorescence analysis

The Sf9 cells were grown in chamber slides and infected with the recombinant baculovirus at a multiplicity of infection of 5 for 48 h, the supernatant was removed, the cell sheet was fixed in cold acetone for 20 min at -20 °C and stained by an immunofluorescence assay (IFA) technique, and then immunofluorescence analysis was performed as described previously on a Zeiss LSH 510 microscope.

2.4. Transient transfections and production of TGEV VLPs

A total of 8×10^5 cells were incubated in 35-mm plates and transfected with plasmid constructs using Cellfection II Reagent (Invitrogen) according to the manufacturer's instructions. Briefly, 8 µL of Cellfection II Reagent was mixed with Grace's medium followed by incubation for 30 min at room temperature, and then 1 µg of plasmid was

added. The Cellfection II-plasmid mixture was incubated for 30 min at room temperature and then 210 µL of Cellfection II-plasmid mixture was added dropwise onto the cells. After 5 h of incubation at 27 °C, the transfection mixture was discarded and replaced with 2 mL of complete growth medium (Grace's insect medium, supplemented with 10% FBS). Cells were incubated at 27 °C for 72 h.

In vitro experiments of VLP assembly of TGEV were performed by infecting insect cells with various combinations of identified recombinant baculoviruses (rBac-M, rBac-sM, rBac-N, rBac-M+rBac-sM, rBac-M+rBac-N). The VLPs were observed in insect cells using a transmission electron microscope.

2.5. Purification of TGEV VLPs

At 72 h after transfection, cell lysates were collected and cleared by centrifugation at a low speed ($1000 \times g$ for 10 min) to pull down cell debris. After passage through a filter with 0.45 µm pore size, cleared cell medium was loaded on top of 20% sucrose cushions and ultracentrifuged for 3 h at 30,000 rpm using a SW41 rotor (Beckman Coulter, Inc., USA). VLP-containing pellets were resuspended in TNE buffer (50 mM Tris-HCl, 100 mM NaCl, 0.5 mM EDTA, pH 7.4).

2.6. Electron microscopy

Negative staining was performed by standard techniques described previously. Briefly, cell lysates were adsorbed to UV-activated copper grids for 2 min at room temperature. Grids were washed with DB twice and stained with 2% uranyl acetate for 30 s. Samples were visualized in a JEOL 1200 EX II transmission electron microscope.

3. Results

3.1. Amplification and construction of the recombinant bacmid plasmid

The target fragments of the *M*, *sM*, and *N* genes, with an expected size of 800 bp, 260 bp, and 1200 bp, were amplified from recombinant plasmid templates of PMD18-M, PMD18-sM, and PMD18-N by PCR (Figures 1A–1C). The *M*, *sM*, and *N* PCR products were cut with *Bam*H I and *Eco*R I, *Sma* I and *Kpn* I, and *Sma* I and *Xho* I, respectively, and subcloned into the pFastBac Dual vector. The cloning was confirmed by enzyme restriction (Figures 1D–1F). Plasmid pFastBac Dual-M, pFastBac Dual-sM, and pFastBac Dual-N were transformed into *E. coli* DH10Bac, followed by extraction of genome DNA, and expectant fragments were obtained by PCR identification using M13 primers (Figures 1G–1I).

3.2. Identification of *M*, *sM*, and *N* protein expressed in Sf9 cells

The venoms of rBac-M, rBac-sM, and rBac-N were inoculated respectively into Sf9 cells and then detected by IFA. Sf9 cells infected with recombinant baculovirus




Figure 1. Construction and identification of recombinant bacmid by restriction digestion and electrophoresis. (A) Electrophoresis of *M* gene. M, DL-2000 marker; 1, *M* gene. (B) Electrophoresis of *sM* gene. M, DL-2000 marker; 1, 2, *sM* gene. (C) Electrophoresis of *N* gene. M, DL-2000 marker; 1, *N* gene. (D) Identification of recombinant plasmid pFastBac Dual-M by restriction digestion. M1, DL-2000 marker; M2, λ -EcoI-14 marker; 1, pFastBac Dual-M digested by *Bam*H I and *Eco*R I. (E) Identification of recombinant plasmid pFastBac Dual-sM by restriction digestion. M1, λ -EcoI-14 marker; M2, DL-2000 marker; 1, pFastBac Dual-sM digested by *Sma* I and *Kpn* I. (F) Identification of recombinant plasmid pFastBac Dual-N by restriction digestion. M1, λ -EcoI-14 marker; M2, DL-2000 marker; 1, pFastBac Dual-N digested by *Sma* I and *Xho* I. (G) Identification of recombinant Bacmid-M. M1, λ -EcoI-14 marker; 1, PCR product of recombinant bacmid Bacmid-M. (H) Identification of recombinant Bacmid-sM. M2, λ -EcoI-14 marker; 2, PCR product of recombinant bacmid Bacmid-sM. (I) Identification of recombinant Bacmid-N. M3, λ -EcoI-14 marker; 3, PCR product of recombinant bacmid Bacmid-N.

showed green fluorescence, but the normal Sf9 cells did not. The results suggested that recombinant proteins were expressed by the baculovirus expression system (Figure 2).

3.3. Assembly of VLPs

To determine if these recombinant proteins could interact with each other and assemble into particulate

structures, Sf9 cells were infected with recombinant baculoviruses expressing M, sM, or N alone or coinfecting with the combination of M and sM or M and N. Electron microscopic analysis of the negatively stained infected cell sections exhibited no particulate structures when the sM and N proteins were expressed alone; there was only




Figure 2. IFA of Sf9 cells infected by recombinant baculovirus rBac-M, rBac-sM, and rBac-N (200×). (A) Normal Sf9 insect cells. (B) Sf9 insect cells infected by rBac-M. (C) Sf9 insect cells infected by rBac-sM. (D) Sf9 insect cells infected by rBac-N.

baculovirus virion in infected Sf9 cells. However, cells infected with recombinant viruses expressing M protein alone and coexpressing M and sM proteins could lead to the occurrence of virions morphologically similar to TGEV. The diameter of VLPs was variable from 61 nm to 101 nm, and envelope-like structures appeared in some virions. Coexpression of M and N proteins in insect Sf9 cells infected by recombinant baculovirus could also assemble into VLPs. The results demonstrated that the M protein was sufficient for the assembly of particles (Figure 3).

4. Discussion

The integral membrane protein, the envelope protein (sM), and the nucleocapsid protein (N) are major structural proteins of TGEV. These structural proteins play a vital role in the processes of viral replication, pathogenesis, and assembly. The M protein is mainly embedded in the lipid envelope and its function is similar to that of nonglycosylated membrane protein in rhabdovirus, orthomyxovirus, or paramyxovirus. M protein, containing a high mannose asparagine-linked side chain, is responsible for the connection of the nucleocapsid and envelope, and it plays an important role in the assembly of viral particles by affecting the sites of assembly and budding. The sM protein may regulate the assembly and release of TGEV and is also needed in effective replication of the virus. The nucleocapsid protein is bound up with viral RNA forming the ribonucleoprotein complex in the assembly process of the virus particle, and then it interacts with membrane protein to migrate to the mature region of the viral packaging-ER-Golgi intermediate compartment (18). Currently, morphogenesis of coronavirus containing mouse hepatitis virus (MHV), infectious bronchitis virus (IBV), and severe acute respiratory (SARS) have been reported (19–22). Research results have shown that M protein and sM protein have more effects on the process of viral morphogenesis, and the coronavirus-like particles can be formed by coexpressing M protein and sM protein in mammalian cells (23). The M protein plays a key role in the morphological processes of coronaviruses, and the point mutation in the N-terminus and C-terminus of this protein will significantly affect the assembly of virus particles. In particular, deletion of the last C-terminal

amino acid residue could not make M protein participate in the assembly of VLPs, but the stability of virus particles may remain constant (24). In studies on the assembly of IBV and MHV particles, it was found that VLPs were released into the supernatant when IBV sM protein was expressed in BHK-21 alone (15), and the results suggested that even without M protein, the sM protein could promote the formation of VLPs independently. M protein (IBV and MHV) expressed alone cannot generate VLPs, but an envelope-like structure can be observed in infected cells (25). Tseng et al. showed that M protein has the capability of assembling and releasing SARS in 293T cells; spherical particles (100 nm in diameter) were observed in transfected supernatants and M alone formed particles that had densities of 1.13 g/mL, suggesting that M alone is sufficient for particle formation (26). Consistent with our results, the TGEV M is capable of assembling virus particles in the absence of other viral components.

The M protein is an important component of the core structures in TGEV, and the C-terminal amino acid sequence is also essential for the stability of the viral core structures. Baudoux et al. showed that VLPs are released into the culture medium of cells coexpressing M and sM proteins *in vitro*, and the VLPs formed by M and sM proteins could induce IFN- α to protect from the infection of TGEV (17). In this study, three structural proteins of TGEV were expressed in Sf9 insect cells by the baculovirus expression system. *In vitro* experiments for VLP assembly of TGEV indicated that expression of M protein alone and coexpression of M plus sM protein and M plus N protein in Sf9 insect cells infected by recombinant baculovirus could result in the occurrence of virions morphologically similar to TGEV. Unlike other coronaviruses (27,28), the M protein of TGEV could be assembled into VLPs just by being expressed *in vitro* alone. This finding is not in accordance with previous results (17,29,30), and the reason is probably that the M protein of TGEV is different from that of other coronaviruses (such as IBV and MHV) in the assembly of virion particles. Electron microscope analysis, as shown in Figure 3, demonstrated M protein in the form of membrane-enveloped vesicles with densities lower than those of VLPs formed by M plus sM or M plus N. This suggests that TGEV M is not a major determinant of virus




Figure 3. Assembly of virus-like particles by expression of structural proteins of TGEV. (A) Negative-staining electron microscopy of virus-like particles formed by expression of recombinant *M* gene. (B) Negative-staining electron microscopy of virus-like particles formed by coexpression of recombinant *M/sM* genes. (C) Negative-staining electron microscopy of virus-like particles formed by coexpression of recombinant *M/N* genes. (D) Negative-staining electron microscopy of virus-like particles formed by expression of recombinant *sM* gene. (E) Negative-staining electron microscopy of virus-like recombinant *N* gene. (F) Negative-staining electron microscopy of virus-like normal Sf9 insect cells.

particle density, such as those reported in self-assembly of the SARS M protein (26). Obvious VLPs were not observed in Sf9 cells when sM or N protein was expressed alone by using the baculovirus expression system. Consistent with this result, experiments on the assembly of coronavirus particles have shown that the N protein is a nonessential component in the formation of VLPs (19). Mammals are the natural hosts of TGEV, but there are some subtle differences between baculovirus-insect cell expression systems and mammalian cell expression systems in the process of translation, processing, and transport for protein expression. Therefore, further study is needed to

confirm the function and mechanism of TGEV structural protein expressed in mammalian cells for the assembly of VLPs.

Acknowledgments

Financial support for this work was provided by grants from the Chongqing Basic Research Program (cstc2014jcyjA80015), the Fundamental Research Funds for the Central Universities (XDJK2014B039), the Fundamental Research Funds for the Xinjiang Uighur Autonomous Region Universities (XJENU2012118), and the Urumqi Research Program (Y121210005).

References

1. Goodwin RF, Jennings AR. A highly infectious gastroenteritis of pig. *Vet Rec* 1985; 70: 271–272.
2. Correa I, Jimenez G, Sune C, Bullido MJ, Enjuanes L. Antigenic structure of the E2 glycoprotein from transmissible gastroenteritis coronavirus. *Virus Res* 1998; 10: 77–93.
3. Delmas B, Rasschaert D, Godet M, Gelfi J, Laude H. Four major antigenic site of the coronavirus transmissible gastroenteritis virus are located on the amino-terminal half of spike glycoprotein S. *J Gen Virol* 1990; 71: 1313–1323.
4. Regenmortel MH, Fauquel CM, Bioshop DH, Carstens EB, Estes MK, Lemon SM, Maniloff J, Mayo MA, McGeoch DJ, Pringle CR et al. *Virus Taxonomy*, 2nd ed. New York, NY, USA: Academic Press; 2000.
5. Siddell SG. *The Coronaviridae*. 1st ed. New York, NY, USA: Plenum Press; 1995.
6. Godet M, Haridon RL, Vautherot JF, Laude H. TGEV coronavirus ORF4 encodes a membrane protein that is incorporated in virions. *Virology* 1992; 188: 666–675.
7. Hurst KR, Kuo L, Koetzner CA, Ye R, Hsue B, Masters PS. A major determinant membrane protein interaction localizes to the carboxy-terminal domain of the mouse coronavirus nucleocapsid protein. *J Virol* 2005; 79: 13285–13297.
8. Sturman LS, Holmes KV, Behnke J. Isolation of coronavirus envelope glycoprotein and interaction with the viral nucleocapsid. *J Virol* 1980; 33: 449–462.
9. Risco C, Anton IM, Enjuanes L, Carrascosa JL. The transmissible gastroenteritis coronavirus contains a spherical core shell consisting of M and N proteins. *J Virol* 1996; 70: 4773–4777.
10. Narayanan K, Maeda A, Maeda J, Makino S. Characterization of the coronavirus M protein and nucleocapsid interaction in infected cell. *J Virol* 2000; 74: 8127–8134.
11. Kuo L, Masters PS. Genetic evidence for a structural interaction between the carboxy termini of the membrane and nucleocapsid. *J Virol* 2002; 76: 4987–4999.
12. de Haan CA, Rottier PJ. Molecular interactions in the assembly of coronaviruses. *Adv Virus Res* 2005; 64: 165–230.
13. Machamer CE, Mentone SA, Rose JK, Farquhar MG. The E1 glycoprotein of an avian coronavirus is targeted to the cis Golgi complex. *P Natl Acad Sci USA* 1990; 87: 6944–6948.
14. Klumperman J, Locker JK, Meijer A, Horzinek MC, Geuze HJ, Rottier PJ. Coronavirus M proteins accumulate in the Golgi complex beyond the site of virion budding. *J Virol* 1994; 68: 6523–6534.
15. Corse E, Machamer CE. Infectious bronchitis virus E protein is targeted to the Golgi complex and directs release of virus-like particles. *J Virol* 2000; 74: 4319–4326.
16. Masters PS. The molecular biology of coronaviruses. *Adv Virus Res* 2006; 66: 193–292.
17. Baudoux P, Carrat C, Besnardeau L, Charley B, Laude H. Coronavirus pseudo-particles formed with recombinant M and E proteins induce alpha interferon synthesis by leukocytes. *J Virol* 1998; 72: 8636–8643.
18. Cologna R, Hogue BG. Coronavirus nucleocapsid protein RNA interactions. *Adv Exp Med Biol* 1998; 440: 355–359.
19. Vennema H, Godeke GJ, Rossen JW, Voorhout WF, Horzinek MC, Opstelten DJ, Rottier PJ. Nucleocapsid-independent assembly of coronavirus-like particles by co-expression of virus envelope protein genes. *EMBO J* 1996; 15: 2020–2028.
20. Bost AG, Prentice E, Denison MR. Mouse hepatitis virus replicase protein complexes are translocated to sites of M protein accumulation in the ERGIC at late times of infection. *Virology* 2001; 285: 21–29.
21. Escors D, Camafeita E, Ortego J, Laude H, Enjuanes L. Organization of two transmissible gastroenteritis coronavirus membrane protein topologies within the virion and core. *J Virol* 2001; 75: 12228–12240.
22. Triplet B, Howard MW, Jobling M, Holmes KV, Hodges RS. Structural characterization of the SARS-coronavirus spike S fusion protein core. *Biol Chem* 2004; 279: 20836–20849.
23. De HC, Vennema H, Rottier PJ. Coronavirus envelope assembly is sensitive to changes in the terminal regions of the viral M protein. *Adv Exp Med Biol* 1998; 440: 365–375.
24. de Haan CA, Vennema H, Rottier PJ. Coronavirus particles assembly: primary structure requirements of the membrane protein. *J Virol* 1998; 72: 6838–6850.
25. Maeda J, Maeda A, Makino S. Release of coronavirus E protein in membrane vesicles from virus-infected cells and E protein-expressing cells. *Virology* 1999; 263: 265–272.
26. Tseng YT, Wang SM, Huang KJ, Lee AI, Chiang CC, Wang CT. Self-assembly of severe acute respiratory syndrome coronavirus membrane protein. *J Biol Chem* 2010; 285: 12862–12872.
27. Bos EC, Luytjes W, van der Meulen HV, Koerten HK, Spaan WJ. The production of recombinant infectious DI-particles of a murine coronavirus in the absence of helper virus. *Virology* 1996; 218: 52–60.
28. Boscarino JA, Logan HL, Lacny JL, Gallagher TM. Envelope protein palmitoylations are crucial for murine coronavirus assembly. *J Virol* 2008; 82: 2989–2999.
29. Francoise F, Stegen CF, Masters PS, Samsonoff WA. Analysis of constructed E gene mutants of mouse hepatitis virus confirms a pivotal role for E protein in coronavirus assembly. *J Virol* 1998; 72: 7885–7894.
30. Liu G, Lv L, Yin L, Li X, Luo D, Liu K, Xue C, Cao Y. Assembly and immunogenicity of coronavirus-like particles carrying infectious bronchitis virus M and S proteins. *Vaccine* 2013; 31: 5524–5530.