

1-1-2018

Nonstandard hulls of lattice-normed ordered vector spaces

ABDULLAH AYDIN

SVETLANA GOROKHOVA

HASAN GÜL

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

AYDIN, ABDULLAH; GOROKHOVA, SVETLANA; and GÜL, HASAN (2018) "Nonstandard hulls of lattice-normed ordered vector spaces," *Turkish Journal of Mathematics*: Vol. 42: No. 1, Article 14. <https://doi.org/10.3906/mat-1612-59>

Available at: <https://dctubitak.researchcommons.org/math/vol42/iss1/14>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

Nonstandard hulls of lattice-normed ordered vector spaces

Abdullah AYDIN^{1,2}, Svetlana GOROKHOVA^{3,*}, Hasan GÜL¹

¹Department of Mathematics, Middle East Technical University, Ankara, Turkey

²Department of Mathematics, Muş Alparslan University, Muş, Turkey

³Laboratory of Functional Analysis, Sobolev Institute of Mathematics, Novosibirsk, Russia

Received: 16.12.2016

Accepted/Published Online: 29.03.2017

Final Version: 22.01.2018

Abstract: Nonstandard hulls of a vector lattice were introduced and studied in many papers. Recently, these notions were extended to ordered vector spaces. In the present paper, following the construction of associated Banach–Kantorovich space due to Emelyanov, we describe and investigate the nonstandard hull of a lattice-normed space, which is the foregoing generalization of Luxemburg’s nonstandard hull of a normed space.

Key words: Vector lattice, ordered vector space, lattice-normed space, decomposable lattice norm, associated Banach–Kantorovich space, lattice-normed ordered vector space, nonstandard hull

1. Introduction

Nonstandard analysis provides a natural approach to various branches of functional analysis (see, for example, [1,9,13–15,21–24,26]). Luxemburg’s construction of the nonstandard hull of a normed space (cf. [1,23,29]) is one of the most important and elegant illustrations of the said approach. Recall that, given an internal normed space $(X, \|\cdot\|)$, an element $x \in X$ is called *infinitesimal* if $\|x\| \approx 0$ and *finite* if $\|x\| \leq n$ for some $n \in \mathbb{N}$. Denote the set of infinitesimal elements and the set of finite elements of X by $\mu(X)$ and $\text{fin}(X)$, respectively. Since $\mu(X)$ is a vector subspace of a vector space $\text{fin}(X)$, we may define \hat{X} to be a quotient $\text{fin}(X)/\mu(X)$. Note that \hat{X} is a real vector space and also a Banach space (cf. [23, p.33]) under the norm defined by

$$\| [x] \| = \text{st}\|x\| = \inf_{\mathbb{R}} \{ a \in \mathbb{R} : \|x\| \leq a \} \quad (x \in \text{fin}(X)). \quad (1)$$

In the case when $X = {}^*Y$ for some standard normed space $Y = (Y, \|\cdot\|)$, the normed space $\hat{Y} := {}^*\hat{Y} = ({}^*\hat{Y}, \|\cdot\|)$ is called the *nonstandard hull* of Y . In the present paper, we develop the notion of the nonstandard hull of a normed space further by generalizing it to the case of a lattice-normed space (abbreviated by LNS).

In the 1990s, Luxemburg’s construction was extended to vector lattices (see [7–9]). Note that a vector lattice E can be seen as the corresponding LNS $(E, |\cdot|, E)$. Lattice-normed vector lattices (abbreviated by LNVLS) have attracted attention in [4,5,10,12,14,26,27]. The general theory of lattice-normed ordered vector spaces (abbreviated by LNOVSs) is still under investigation. The present paper contributes to the study of this theory by using nonstandard analysis, namely by using nonstandard hulls of LNOVSs normed by Dedekind complete vector lattices.

*Correspondence: lanagor71@gmail.com

2010 AMS Mathematics Subject Classification: 46A40, 46B40, 46S20

The scheme of nonstandard analysis used below has been introduced by Luxemburg and Stroyan [30]. In our paper, we deal only with nonstandard enlargements satisfying the general saturation principle (such nonstandard enlargements are called *polysaturated* [1, p.47]). Since the basic methods of nonstandard analysis are well developed and presented in many textbooks (see, for example, [1,21,23,24,27,29,32]), we refer the reader for corresponding notions and terminology to these standard sources. We also refer to [2,3,14,26,31,33,34,36] for theory of ordered vector spaces (abbreviated by OVSs) and [10,12,14,16,18] for nonstandard hulls of vector lattices and OVSs.

The structure of the paper is as follows. In Section 2, we include elementary theory of LNOVSs in parallel with theory of LNVs (see, for example, [4,5,26]). In Section 3, we introduce and investigate the nonstandard hull of an LNS normed by a Dedekind complete vector lattice. This notion is closely related to the construction of an associated Banach–Kantorovich space [10,12,14]. In Section 4, we investigate nonstandard hulls of LNOVSs. The main result here is Theorem 5, that is the nonstandard hull of a p -semimonotone LNOVS is also p -semimonotone with the same constant of semimonotonicity.

2. Preliminaries

In the present paper, all standard OVSs are assumed to be real, Archimedean, and equipped with the generating positive cone [3]. We define and study certain necessary notions such as p -normality and op -continuity in LNOVSs, p -Levi spaces, etc. (see also [4–6] for their lattice versions).

The following notions in lattice-normed vector spaces (abbreviated by LNSs) are motivated by their analogies in normed spaces.

Definition 1 (see also [4]) *Given an LNS (X, p, E) and $A, B \subseteq X$.*

- (a) *A is said to be p -dense in B if, for any $b \in B$ and for any $0 \neq u \in p(X)$, there is $a \in A$ such that $p(a - b) \leq u$.*
- (b) *A is said to be p -closed if, for any net a_α in A such that $p(a_\alpha - x) \rightarrow 0$ in X (abbreviated by $z_\alpha \xrightarrow{p} x$), it holds that $x \in A$.*
- (c) *B is said to be the p -closure of A if B is the intersection of all p -closed subsets of X containing A .*

In what follows, $X = (X, p, E)$ is an LNOVS. The next property is an analogy of the well-known property of normed OVSs. It is a direct extension of [4, Prop.1] and it has a similar proof, which is omitted.

Proposition 1 *Let the positive cone X^+ in an LNOVS $X = (X, p, E)$ be p -closed. Then any monotone p -convergent net in X is o -convergent to its p -limit.*

We continue with further basic notions in LNOVSs, which are motivated by their analogies for vector lattices and for LNVs (see, for example, [4–6,19,20,25–27]).

Definition 2 (a) *A subset $A \subseteq X$ is p -bounded if there exists $e \in E$ such that $p(a) \leq e$ for all $a \in A$.*

(b) *X is p -semimonotone if there is $M \in \mathbb{R}$ such that $0 \leq y \leq x \in X$ implies $p(y) \leq Mp(x)$.*

(c) *X is p -normal if $x_\alpha \leq y_\alpha \leq z_\alpha$ in X , $x_\alpha \xrightarrow{p} u$, and $z_\alpha \xrightarrow{p} u$ imply $y_\alpha \xrightarrow{p} u$.*

(d) *X is a p -Levi-space if every p -bounded increasing net in X^+ is p -convergent.*

(e) *X is op -continuous if $x_\alpha \xrightarrow{o} 0$ implies that $x_\alpha \xrightarrow{p} 0$.*

- (f) X is σ - op -continuous if $x_n \xrightarrow{o} 0$ implies that $x_n \xrightarrow{p} 0$.
- (g) A net $(x_\alpha)_{\alpha \in A}$ in X is said to be p -Cauchy if $(x_\alpha - x_{\alpha'})_{(\alpha, \alpha') \in A \times A} \xrightarrow{p} 0$.
- (h) X is p -complete if every p -Cauchy net in X is p -convergent.

Lemma 1 Let X be a p -semimonotone LNOVS with the semimonotonicity constant M . Then form $a \leq x \leq b$ in X follows that $p(x) \leq 2(M + 1)(p(a) \vee p(b))$.

Proof Since $a \leq x \leq b$, then $0 \leq x - a \leq b - a$ and

$$p(x) - p(a) \leq p(x - a) \leq Mp(b - a) \leq M(p(b) + p(a)).$$

Hence

$$p(x) \leq M(p(b) + p(a)) + p(a) \leq (M + 1)(p(b) + p(a)) \leq 2(M + 1)(p(a) \vee p(b)).$$

□

Lemma 2 Let X be a p -semimonotone LNOVS and $\pm x_\alpha \leq y_\alpha \xrightarrow{p} 0$. Then $x_\alpha \xrightarrow{p} 0$.

Proof By Lemma 1, $-y_\alpha \leq x_\alpha \leq y_\alpha$ implies $p(x_\alpha) \leq 2(M + 1)p(y_\alpha)$. Since $y_\alpha \xrightarrow{p} 0$, then $2(M + 1)p(y_\alpha) \xrightarrow{o} 0$ and hence $p(x_\alpha) \xrightarrow{o} 0$. Thus, $x_\alpha \xrightarrow{p} 0$. □

Definition 2(c) is motivated by the property (cf. [3, Thm.2.23]) of normal normed OVSs. Note that, without lost of generality, one may suppose that, in Definition 2(c), $u = 0$ and $x_\alpha \equiv 0$. Therefore, Lemma 2 ensures that any p -semimonotone LNOVS is p -normal (in particular, any LNVL is p -normal). Thus, the p -normality coincides with the usual normality in a normed OVS $(X, p, E) = (X, \|\cdot\|)$. In this case X is p -normal iff it is p -semimonotone (cf. [35, Thm.IV.2.1]).

It was established in [4, Lm.2] that an LNVL (X, p, E) is op -continuous iff $X \ni w_\beta \downarrow 0 \Rightarrow w_\beta \xrightarrow{p} 0$.

In order to extend this result, we need the following lemma.

Lemma 3 Let an LNOVS $X = (X, p, E)$ be p -semimonotone and w_β be a net in X . If $w_\beta \downarrow 0$ implies $w_\beta \xrightarrow{p} 0$, then X is op -continuous.

Proof Let $x_\alpha \xrightarrow{o} 0$. Then there are two nets $y_\beta \downarrow 0$ and $z_\gamma \downarrow 0$ in X such that, for every β and γ , there exists $\alpha_{\beta, \gamma}$ with

$$-y_\beta \leq x_\alpha \leq z_\gamma \quad (\forall \alpha \geq \alpha_{\beta, \gamma}).$$

By Lemma 1,

$$p(x_\alpha) \leq 2(M + 1)(p(y_\beta) \vee p(z_\gamma)) \quad (\forall \alpha \geq \alpha_{\beta, \gamma}). \tag{2}$$

By the assumption, $p(y_\beta) \xrightarrow{o} 0$ and $p(z_\gamma) \xrightarrow{o} 0$. Then $p(y_\beta) \vee p(z_\gamma) \xrightarrow{o} 0$. It follows from (2) that $p(x_\alpha) \xrightarrow{o} 0$. Therefore, X is op -continuous. □

Hence we have the following result.

Theorem 1 A p -semimonotone LNOVS $X = (X, p, E)$ is op -continuous iff, for any net $x_\alpha \in X$, the condition $x_\alpha \downarrow 0$ implies $x_\alpha \xrightarrow{p} 0$.

Clearly, the *op*-continuity in LNOVSs is equivalent to the order continuity in the sense of [26, 2.1.4, p.48]. For *p*-complete LNOVSs, we have more conditions for *op*-continuity (see also [4, Thm.1] for the LNVL case).

Theorem 2 *Let an LNOVS $X = (X, p, E)$ be *p*-complete and *p*-semimonotone. The following conditions are equivalent:*

- (i) *X is op-continuous;*
- (ii) *if $0 \leq x_\alpha \uparrow \leq x$ holds in X , then x_α is a *p*-Cauchy net;*
- (iii) *$x_\alpha \downarrow 0$ in X implies $x_\alpha \xrightarrow{p} 0$.*

The proof is similar to the proof of [4, Thm.1] and we omit it.

The following two results generalize [4, Cor.1], [4, Cor.2], and [4, Prop.2] respectively.

Theorem 3 *Let an LNOVS (X, p, E) be *op*-continuous, *p*-complete, and *p*-semimonotone. Then X is Dedekind complete.*

Proof Assume $0 \leq x_\alpha \uparrow \leq u$; then, by Theorem 2(ii), x_α is a *p*-Cauchy net and, since X is *p*-complete, there exists x such that $x_\alpha \xrightarrow{p} x$. It follows from Proposition 1 that $x_\alpha \uparrow x$, and so X is Dedekind complete. \square

Theorem 4 *Any *p*-semimonotone *p*-Levi LNOVS (X, p, E) with *p*-closed X^+ is *op*-continuous.*

The proof is similar to the proof of [4, Cor.2] and therefore it is omitted.

Proposition 2 *Any *p*-semimonotone *p*-Levi LNOVS (X, p, E) with *p*-closed X^+ is Dedekind complete.*

Proof Let $0 \leq x_\alpha \uparrow \leq z \in X$. Then $p(x_\alpha) \leq Mp(z)$. Hence the net x_α is *p*-bounded and therefore $x_\alpha \xrightarrow{p} x$ for some $x \in X$. By Proposition 1, $x_\alpha \uparrow x$. \square

3. Nonstandard hulls of LNSs and of dominated operators acting between them

Order- and *regular-nonstandard hulls* of LNSs were introduced in [10] as certain generalizations of Luxemburg’s nonstandard hull of a normed space [29]. Here we employ a different approach for extending Luxemburg’s construction to LNSs. In the rest of the paper, we suppose all LNSs to be normed by Dedekind complete vector lattices. To be certain, we fix a Dedekind complete vector lattice E for the norming lattice for all LNSs in what follows. While considering an internal LNS $(\mathcal{X}, p, \mathcal{E})$, we always assume that its norming lattice is standard, i.e. $\mathcal{E} = {}^*E$.

3.1. Some external vector spaces associated with OVSs and LNSs

We begin with several basic constructions from [10, 16, 18]. Let Y be an OVS. Consider the following external real vector subspaces of *Y [18].

$$\text{fin}({}^*Y) := \{\kappa \in {}^*Y : (\exists y \in Y) -y \leq \kappa \leq y\},$$

$$\eta({}^*Y) := \{\kappa \in {}^*Y : \inf_Y \{y \in Y : -y \leq \kappa \leq y\} = 0\},$$

$$\text{o-pns}({}^*Y) := \{\kappa \in {}^*Y : \inf_Y \{y' - y : Y \ni y \leq \kappa \leq y' \in Y\} = 0\},$$

and $\bar{Y} := \text{fin}({}^*Y)/\eta({}^*Y)$.

Let $\mathcal{X} = (\mathcal{X}, p, {}^*E)$ be an internal LNS. In accordance with [10, 12], consider the following external subspaces of \mathcal{X} :

$$\begin{aligned} \mathbf{fn}(\mathcal{X}) &= \{x \in \mathcal{X} : p(x) \in \mathbf{fn}({}^*E)\}, \\ \mathbf{n}(\mathcal{X}) &= \{x \in \mathcal{X} : p(x) \in \eta({}^*E)\}, \end{aligned}$$

In the case of a standard LNS $X = (X, p, E)$,

$$\mathbf{o} - \mathbf{pns}({}^*X) = \{\kappa \in {}^*X : \inf_E \{p(\kappa - x) : x \in X\} = 0\}.$$

Remark that, similarly to the case in which $X = (X, p, E)$ is a normed space (cf. [1, Prop.2.2.2.]), it can be easily shown that X is p -complete iff $\mathbf{o} - \mathbf{pns}({}^*X) = X + \mathbf{n}({}^*X)$.

3.2. Nonstandard hull of an LNS

For an internal LNS $\mathcal{X} = (\mathcal{X}, p, {}^*E)$, consider the quotient $\overline{\mathcal{X}} := \mathbf{fn}(\mathcal{X})/\mathbf{n}(\mathcal{X})$ and define the mapping $\overline{p} : \overline{\mathcal{X}} \rightarrow E$ by the following rule motivated by formula (1) (see also [12, Thm.2.3.5.] and [10, 3.1]):

$$\overline{p}([x]) := \inf_E \{e \in E : e \geq p(x)\} \quad (x \in \mathbf{fn}(\mathcal{X})). \tag{1^*}$$

It is easy to see that this mapping is a well-defined E -valued norm on $\overline{\mathcal{X}}$.

Definition 3 *Given an LNS (X, p, E) . The LNS $(\overline{{}^*X}, \overline{p}, E)$ is called the nonstandard hull of (X, p, E) .*

According to [10, Thm.3.5], the nonstandard hull of (X, p, E) is a Banach–Kantorovich space, when p is decomposable. The main reason for using the term "nonstandard hull" lies in [12, Thm.2.4.1.] (see also [10, Thm.4.3]), saying that, in the case of decomposable LNS (X, p, E) , the p -completion of (X, p, E) can be obtained by natural embedding of (X, p, E) into $(\overline{{}^*X}, \overline{p}, E)$, and then by just taking its p -closure there.

3.3. Nonstandard hulls of dominated operators between decomposable LNSs

Given two LNSs (X, p, E) and (Y, m, E) . Let $T : X \rightarrow Y$ be a dominated operator (cf. [26, 4.4.1.]). Under the assumption of decomposable X , T has an exact dominant $|T|$ (cf. [26, 4.4.2.]) and, in that case, the space $M(X, Y)$ can be considered a decomposable LNS $(M(X, Y), |\cdot|, L_b(E))$.

Denote by $\mathcal{M}_n({}^*X, {}^*Y)$ the set of all internal linear operators from *X into *Y that admit standard order-continuous dominants, that is: $T \in \mathcal{M}_n({}^*X, {}^*Y)$ iff there is an operator $S \in L(E, F)$ satisfying ${}^*m(T\kappa) < {}^*S({}^*p(\kappa))$ for all $\kappa \in {}^*X$. The following lemma was proved in [12, Lm.2.4.2.].

Lemma 4 *For every operator $T \in \mathcal{M}_n({}^*X, {}^*Y)$, $T(\mathbf{fn}({}^*X)) \subseteq \mathbf{fn}({}^*Y)$ and $T(\mathbf{n}({}^*X)) \subseteq \mathbf{n}({}^*Y)$.*

Lemma 4 ensures that, for any operator $T \in \mathcal{M}_n({}^*X, {}^*Y)$, the rule

$$\overline{T}([\kappa]) := [T\kappa] \quad (\kappa \in \mathbf{fn}({}^*X))$$

defines a mapping $\overline{T} : \overline{X} \rightarrow \overline{Y}$. By [12, Thm.2.4.3.], \overline{T} is a linear dominated operator from $(\overline{{}^*X}, \overline{p}, E)$ into $(\overline{{}^*Y}, \overline{m}, E)$ with the least dominant $|\overline{T}|$ satisfying

$$|\overline{T}| \leq \inf \{S \in L_n(E) : {}^*S \geq |T|\}, \tag{3}$$

where $|T|$ is the least internal dominant of T . The operator \overline{T} is said to be the *nonstandard hull* of T . Since $T \in M_n(X, Y)$ iff $*T \in \mathcal{M}_n(*X, *Y)$, inequality (3) implies that $|\overline{T}| = |T|$ for any $T \in M_n(X, Y)$ (see also [12, Thm.2.4.4.]).

4. Nonstandard hulls of LNOVSs

4.1. Nonstandard hull of an LNOVS

Let $\mathcal{Y} = (\mathcal{Y}, p, *E)$ be an internal p -semimonotone LNOVS with a finite constant $\mathcal{M} \in \text{fin}(\mathbb{R})$ of the semimonotonicity. The key step is the following technical lemma.

Lemma 5 $\mathbf{n}(\mathcal{Y})$ is an order ideal in $\mathbf{fin}(\mathcal{Y})$.

Proof Since $\mathbf{n}(\mathcal{Y})$ is a real vector subspace of $\mathbf{fin}(\mathcal{Y})$, it is enough to show that $\mathbf{n}(\mathcal{Y})$ is order convex. Let $\xi \leq \kappa \leq \zeta$ with $\kappa \in \mathcal{Y}$, and $\xi, \zeta \in \mathbf{n}(\mathcal{Y})$. By Lemma 1,

$$p(\kappa) \leq 2(\mathcal{M} + 1)(p(\xi) \vee p(\zeta)) \leq 2(\text{st}(\mathcal{M}) + 2)(p(\xi) \vee p(\zeta)) \in \eta(*E),$$

and hence $\kappa \in \mathbf{n}(\mathcal{Y})$. □

Theorem 5 Let $(\mathcal{Y}, p, *E)$ be a p -semimonotone LNOVS with a finite constant \mathcal{M} of semimonotonicity. Then $\mathbf{fin}(\mathcal{Y})/\mathbf{n}(\mathcal{Y})$ is an OVS. Moreover, the LNOVS $(\overline{\mathcal{Y}}, \overline{p}, E)$ is p -semimonotone with a constant $M = \text{st}(\mathcal{M})$ of semimonotonicity.

Proof $\overline{\mathcal{Y}} = \mathbf{fin}(\mathcal{Y})/\mathbf{n}(\mathcal{Y})$ is an OVS, by Lemma 4. Now let $0 \leq [\kappa] \leq [\xi] \in \overline{\mathcal{Y}}$. By the definition of ordering in the quotient space $\mathbf{fin}(\mathcal{Y})/\mathbf{n}(\mathcal{Y})$ (cf. [17, p.3]), we may assume that $0 \leq \kappa \leq \xi$. Hence $\frac{1}{\mathcal{M}}p(\kappa) \leq p(\xi)$ and then, for any $n \in \mathbb{N}$,

$$\begin{aligned} \overline{p}([\xi]) &= \inf_E \{e \in E : e \geq p(\xi)\} \geq \inf_E \{e \in E : e \geq \mathcal{M}^{-1}p(\kappa)\} \geq \\ &\inf_E \{e \in E : e \geq (\frac{1}{\mathcal{M}} - \frac{1}{2n})p(\kappa)\} \geq \inf_E \{e \in E : e \geq (\frac{1}{M} - \frac{1}{n})p(\kappa)\} = \\ &(\frac{1}{M} - \frac{1}{n}) \inf_E \{e \in E : e \geq p(\kappa)\} = (\frac{1}{M} - \frac{1}{n})\overline{p}([\kappa]). \end{aligned}$$

Since the inequality is true for all $n \in \mathbb{N}$, we obtain $\overline{p}([\xi]) \geq (\frac{1}{M}\overline{p}([\kappa]))$ or $\overline{p}([\kappa]) \leq M\overline{p}([\xi])$, as desired. □

Corollary 1 Let (Y, p, E) be a p -semimonotone LNOVS. Then $(*\overline{Y}, \overline{p}, E)$ is also a p -semimonotone LNOVS with the same constant of semimonotonicity.

Proof Let M be a semimonotonicity constant of (Y, p, E) . By the transfer principle, $\mathcal{M} = M$ is a semimonotonicity constant of $(*\overline{Y}, \overline{p}, E)$. Now apply Theorem 5. □

Corollary 2 Let $(Y, \|\cdot\|)$ be a normal OVS. Then its nonstandard hull $*\overline{Y}$ is a normal Banach space.

Proof Note that any OVS is normal iff it is semimonotone (cf. [35, Thm.IV.2.1.]) and apply Theorem 5. □

4.2. Internal LNVLS

Here we consider some properties of LNS $(\bar{\mathcal{Y}}, \bar{p}, E)$ in the case where $(\mathcal{Y}, p, {}^*E)$ is an internal LNVL.

Theorem 6 *Let $(\mathcal{Y}, p, {}^*E)$ be an internal LNVL. Then $(\bar{\mathcal{Y}}, \bar{p}, E)$ is also an LNVL.*

Proof Note that the quotient $\mathbf{fn}(\mathcal{Y})/\mathbf{n}(\mathcal{Y})$ of a vector lattice $\mathbf{fn}(\mathcal{Y})$ by an order ideal $\mathbf{n}(\mathcal{Y})$ is a vector lattice. Since $(\mathcal{Y}, p, {}^*E)$ has a semimonotonicity constant $\mathcal{M} = 1$, then, by Theorem 5, the LNOVS $(\bar{\mathcal{Y}}, \bar{p}, E)$ has a semimonotonicity constant $M = 1$, which means that $\bar{p}([\kappa]) \leq \bar{p}([\xi])$ for all κ, ξ with $|\kappa| \leq |\xi|$. Therefore, \bar{p} is an E -valued lattice norm on $\bar{\mathcal{Y}}$ and $(\bar{\mathcal{Y}}, \bar{p}, E)$ is an LNVL. \square

The following proposition generalizes [23, Prop.4.7] for LNVL.

Proposition 3 *Let $(\mathcal{Y}, p, {}^*E)$ be an internal LNVL. Then the LNVL $(\bar{\mathcal{Y}}, \bar{p}, E)$ is σ -op-continuous, and every monotone p -bounded sequence in $\bar{\mathcal{Y}}$ is order-bounded.*

Proof First we show σ -op-continuity. Clearly, it is enough to show that $\bar{\mathcal{Y}} \ni x_n \downarrow 0$ implies $\bar{p}(x_n) \downarrow 0$. Suppose in contrast that $\bar{p}(x_n) \not\downarrow 0$ for all n and some $0 \neq u \in E_+$. The monotonicity of the lattice norm p ensures the existence of a sequence $\kappa_n \downarrow$ in \mathcal{Y}_+ with $[\kappa_n] = x_n$ and $2p(\kappa_n) \geq u$ for all $n \in \mathbb{N}$. Consider the sequence of nonempty internal sets

$$A_n = \{\chi \in \mathcal{Y} : 2p(\chi) \geq u \ \& \ 0 \leq \chi \leq \kappa_n\} \quad (n \in \mathbb{N}).$$

By saturation principle there exists $\chi \in \bigcap_{n=1}^{\infty} A_n$. Then $0 < [\chi] \leq [\kappa_n] = x_n$ violating $x_n \downarrow 0$. Therefore, $\bar{p}(x_n) \downarrow 0$.

For the second part, let $\bar{\mathcal{Y}} \ni x_n \downarrow$ and $\bar{p}(x_n) \leq u \in E$ for all $n \in \mathbb{N}$. The monotonicity p gives a sequence $\kappa_n \downarrow$ in \mathcal{Y} with $[\kappa_n] = x_n$ and $p(\kappa_n) \leq 2u$ for all $n \in \mathbb{N}$. By the saturation principle there is $\chi \in \mathcal{Y}$ with $p(\chi) \leq 2u$ and $\chi \leq \kappa_n \leq \kappa_1$ for all $n \in \mathbb{N}$. Hence $x_n = [\kappa_n] \in [[\chi], [\kappa_1]]$ for all $n \in \mathbb{N}$, which is required. \square

4.3. Nonstandard criterion for op-continuity

The following theorem generalizes [12, Thm.4.5.3].

Theorem 7 *An LNVL (X, p, E) is op-continuous iff $\eta({}^*X) \subseteq \mathbf{n}({}^*X)$.*

Proof Suppose that (X, p, E) is op-continuous and fix $\kappa \in \eta({}^*X)$. Then there exists a net $x_\alpha \in {}^*X$ such that $x_\alpha \downarrow 0$ and $0 \leq \kappa \leq x_\alpha$. Clearly, $x_\alpha \overset{o}{\rightarrow} 0$, and so we have $x_\alpha \overset{p}{\rightarrow} 0$ since p is op-continuous. Since $0 \leq p(\kappa) \leq p(x_\alpha)$, it follows that $p(\kappa) \in \eta({}^*E)$ or $\kappa \in \mathbf{n}({}^*X)$. Hence $\eta({}^*X) \subseteq \mathbf{n}({}^*X)$.

Now suppose $\eta({}^*X) \subseteq \mathbf{n}({}^*X)$ and $X \ni x_\alpha \overset{o}{\rightarrow} 0$. Then there are two nets $y_\beta \downarrow 0$ and $z_\gamma \downarrow 0$ in X such that, for every β and γ , there exists $\alpha_{\beta,\gamma}$ with

$$-y_\beta \leq x_\alpha \leq z_\gamma \quad (\alpha \geq \alpha_{\beta,\gamma}).$$

Thus, $x_\alpha \in \eta({}^*X)$ for all infinitely large α . Hence, by the hypothesis, $x_\alpha \in \mathbf{n}({}^*X)$ for all infinitely large α . Therefore, $p(x_\alpha) \rightarrow 0$ and (X, p, E) is op-continuous. \square

We finish with a discussion of the p -Levi property. Let (X, p, E) be p -Levi and $(x_\alpha)_{\alpha \in A}$ be a monotone p -bounded net in X . Then $x_\alpha \xrightarrow{p} x$ for some $x \in X$. By the transfer principle, $x_\alpha \in \mathbf{fin}(*X)$ for all $\alpha \in *A$. Given an infinitely large ν . Since $x_\alpha \xrightarrow{p} x$, then $x_\nu \in x + \mathbf{n}(*X) \subseteq \mathbf{o} - \mathbf{pns}(*X)$. We do not know under which conditions on (X, p, E) the converse is also true.

References

- [1] Albeverio S, Hoegh-Krohn R, Fenstad JE, Lindstrom T. Nonstandard Methods in Stochastic Analysis and Mathematical Physics. Orlando, FL, USA: Academic Press, 1986.
- [2] Aliprantis CD, Burkinshaw O. Positive Operators. Orlando, FL, USA: Academic Press, 1985.
- [3] Aliprantis CD, Tourky R. Cones and duality. Providence, RI, USA: American Mathematical Society, 2007.
- [4] Aydin A, Emelyanov EY, Erkursun-Ozcan N, Marabeh MAA. Unbounded p -convergence in lattice-normed vector lattices. arXiv:1609.05301v3. 2016.
- [5] Aydin A, Emelyanov EY, Erkursun-Ozcan N, Marabeh MAA. Compact-like operators in lattice-normed spaces. Indag Math 2017; <https://doi.org/10.1016/j.indag.2017.11.002>
- [6] Dabboorasad YA, Emelyanov EY, Marabeh MAA. um-Topology in multi-normed vector lattices, Positivity 2017; <https://doi.org/10.1007/s11117-017-0533-6>
- [7] Emelyanov EY. Nonstandard hulls of vector lattices. Sibirsk Mat Zh 1994; 35: 83-95.
- [8] Emelyanov EY. The order and regular hulls of Riesz spaces. Sibirsk Mat Zh 1994; 35: 1243-1252.
- [9] Emelyanov EY. Order hulls of vector lattices. Dokl Math 1995; 340: 303-304.
- [10] Emelyanov EY. Banach-Kantorovich spaces associated with order-hulls of decomposable lattice-normed spaces. Sibirsk Mat Zh 1995; 36: 72-85.
- [11] Emelyanov EY. An infinitesimal approach to the representation of vector lattices by spaces of continuous functions on a compactum. Dokl Math 1995; 344: 9-11.
- [12] Emelyanov EY. Infinitesimal analysis and vector lattices. Siberian Adv Math 1996; 6: 19-70.
- [13] Emelyanov EY. Invariant homomorphisms of nonstandard extensions of Boolean algebras and vector lattices. Sibirsk Mat Zh 1997; 38: 286-296.
- [14] Emelyanov EY. Infinitesimals in vector lattices. In: Kutateladze SS, editor. Nonstandard analysis and vector lattices 2000. Dordrecht, the Netherlands: Kluwer Academic Publishers, 2000, pp. 161-230.
- [15] Emelyanov EY. Some conditions for a C_0 -semigroup to be asymptotically finite-dimensional. Sibirsk Mat Zh 2003; 44: 1015-1020.
- [16] Emelyanov EY. Infinitesimals in ordered vector spaces. Vladikavkaz Mat Zh 2013; 15: 18-22.
- [17] Emelyanov EY. Archimedean Cones in Vector Spaces. J Convex Anal 2017; 24: 169-183.
- [18] Emelyanov EY, Gul H. Nonstandard hulls of ordered vector spaces. Positivity 2016; 20: 413-433.
- [19] Emelyanov EY, Marabeh MAA. Two measure-free versions of the Brezis-Lieb lemma. Vladikavkaz Mat Zh 2016; 18: 21-25.
- [20] Gao N, Troitsky VG, Xanthos F. Uo-convergence and its applications to Cesaro means in Banach lattices. Israel J Math 2017; 220: 649-689.
- [21] Gordon EI, Kusraev AG, Kutateladze SS. Infinitesimal analysis. Dordrecht, the Netherlands: Kluwer Academic Publishers, 2002.
- [22] Gorokhova SG, Emelyanov EY. On the concept of stability of order convergence in vector lattices. Sibirsk Mat Zh 1994; 35: 1026-1031.

- [23] Henson CW, Moore LCJr. Nonstandard analysis and the theory of Banach spaces. In: Hurd AE, editor. *Lecture Notes in Mathematics* 983. Berlin-New York, Germany-USA: 1983; pp. 27-112.
- [24] Hurd AE, Loeb PA. *An Introduction to Nonstandard Real Analysis*. Orlando, FL, USA: Academic Press, 1985.
- [25] Kandić M, Marabeh MAA, Troitsky VG. Unbounded norm topology in Banach lattices. *J Math Anal Appl* 2017; 451: 259-279.
- [26] Kusraev AG. *Dominated Operators*. Dordrecht, the Netherlands: Kluwer Academic Publishers, 2000.
- [27] Kusraev AG, Kutateladze SS. *Boolean Valued Analysis*. Dordrecht, the Netherlands: Kluwer Academic Publishers, 1999.
- [28] Loeb PA, Wolff MPH. *Nonstandard Analysis for the Working Mathematician*. Dordrecht, the Netherlands: Kluwer Academic Publishers, 2000.
- [29] Luxemburg WAJ. A general theory of monads. In: *Applications of Model Theory to Algebra, Analysis, and Probability*. Internat. Symposium. Pasadena, Calif., 1967; NJ, USA: Holt, Rinehart and Winston, 1969, pp. 18-86.
- [30] Luxemburg WAJ, Stroyan KD. *Introduction to the Theory of Infinitesimals*. New York, NY, USA: Academic Press, 1976.
- [31] Luxemburg WAJ, Zaanen AC. *Riesz Spaces, I*. Amsterdam, the Netherlands: North-Holland Publishing Co., 1971.
- [32] Robinson A. *Non-standard Analysis*. Amsterdam, the Netherlands: North-Holland Publishing Co., 1966.
- [33] Schaefer HH, Wolff MPH. *Topological Vector Spaces, second edition*. New York, NY, USA: Springer-Verlag, 1999.
- [34] Vulikh BZ. *Introduction to the Theory of Partially Ordered Spaces*. Groningen, the Netherlands: Wolters-Noordhoff Scientific Publications, 1967.
- [35] Vulikh BZ. *Introduction to the Theory of Cones in Normed Spaces*. Kalinin, Russia: Kalinin State University Press, 1977.
- [36] Zaanen AC. *Riesz Spaces, II*. Amsterdam, the Netherlands: North-Holland Publishing Co., 1983.