

1-1-2015

Relationships between sarcomere length and basic composition of infraspinatus and longissimus dorsi muscle

DOMINIKA GUZEK

DOMINIKA GLABSKA

KRZYSZTOF GLABSKI

GRZEGORZ POGORZELSKI

JERZY BARSZCZEWSKI

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/veterinary>



Part of the [Animal Sciences Commons](#), and the [Veterinary Medicine Commons](#)

Recommended Citation

GUZEK, DOMINIKA; GLABSKA, DOMINIKA; GLABSKI, KRZYSZTOF; POGORZELSKI, GRZEGORZ; BARSZCZEWSKI, JERZY; and WIERZBICKA, AGNIESZKA (2015) "Relationships between sarcomere length and basic composition of infraspinatus and longissimus dorsi muscle," *Turkish Journal of Veterinary & Animal Sciences*: Vol. 39: No. 1, Article 15. <https://doi.org/10.3906/vet-1405-69>
Available at: <https://journals.tubitak.gov.tr/veterinary/vol39/iss1/15>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Veterinary & Animal Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Relationships between sarcomere length and basic composition of infraspinatus and longissimus dorsi muscle

Authors

DOMINIKA GUZEK, DOMINIKA GLABSKA, KRZYSZTOF GLABSKI, GRZEGORZ POGORZELSKI, JERZY BARSZCZEWSKI, and AGNIESZKA WIERZBICKA

Relationships between sarcomere length and basic composition of *infraspinatus* and *longissimus dorsi* muscle

Dominika GUZEK^{1*}, Dominika GŁĄBSKA², Krzysztof GŁĄBSKI³, Grzegorz POGORZELSKI¹,
Jerzy BARSZCZEWSKI⁴, Agnieszka WIERZBICKA¹

¹Division of Engineering in Nutrition, Faculty of Human Nutrition and Consumer Sciences, Warsaw University of Life Sciences, Warsaw, Poland

²Department of Dietetics, Faculty of Human Nutrition and Consumer Sciences, Warsaw University of Life Sciences, Warsaw, Poland

³Department of Microbial Biochemistry, Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland

⁴Grasslands Division, Institute of Technology and Life Sciences in Falenty, Raszyn, Poland

Received: 22.05.2014 • Accepted: 10.08.2014 • Published Online: 12.01.2015 • Printed: 09.02.2015

Abstract: The purpose of this study was to determine the relationship between sarcomere length and basic composition (water, protein, ash, collagen, and fat content including marbling) of beef muscles with fast-twitch glycolytic (*longissimus dorsi muscle*) and slow-twitch oxidative fibers (*infraspinatus muscle*) in Limousin bulls. Samples (n = 15) were obtained from federally inspected slaughter facilities, and animals used in the experiment were treated according to standard ethical norms. The shortest sarcomeres were observed for *longissimus dorsi* from striploin, and the longest for *infraspinatus* muscle. Additionally, the influence of both cut and animal was observed. In both cases of muscles, sarcomere length was positively correlated with marbling level and negatively correlated with protein content. Moreover, sarcomere length was very strongly correlated with marbling level in the case of *longissimus dorsi et lumborum*. Differences in sarcomere length were associated with cut and animal, and this correlation may partly explain differences in composition, especially in protein content.

Key words: Beef, computer image analysis, marbling, protein, sarcomere

1. Introduction

Meat quality varies to a great extent within and between animals as well as in individual muscles. This results from the fact that meat features depend on breed, genotype, sex, age, nutrition, and slaughter procedures. Individual muscles from various anatomical locations are characterized by different metabolic fiber types. *Longissimus dorsi* muscle (*LDM*) is characterized by an abundance of fast-twitch glycolytic fibers (type IIB), while *infraspinatus* muscle (*IM*) consists mostly of slow-twitch oxidative fibers (type I) (1). Several muscles are also characterized by diverse sarcomere length (2), chemical composition (3), and intramuscular fat quantity (marbling) (4). These factors influence beef tenderness, whereas sarcomere length plays a crucial role in the mechanical structure of meat (5).

It has been indicated that sarcomere length determines 16% of the variation in tenderness of *semitendinosus* muscle and 44%–55% of the variation in tenderness of *LDM* (6) when measured with the method of Warner–Brazler share force (7). At the same time, the evaluation of *LDM* tenderness with sensory methods suggests that

sarcomere length explains between 14% and 38% (7) of variation in tenderness.

According to Stolowski et al. (2), sarcomere length is not affected by breed. Instead, it may depend to a large extent on the specific muscle, as sarcomere length is associated with the type of fibers dominant in the muscle. It has been indicated that type IIB fibers are characterized by a shorter sarcomere length than type I fibers (8). Li et al. (4) reported that sarcomere length increases with advancing marbling score. At the same time, it should be emphasized that until now little attention has been paid to muscles other than *LDM* and to the physiological function and metabolic types of fibers.

The purpose of this study was to determine the relationship between sarcomere length and basic composition of beef meat in muscles that are characterized by different physiological functions and metabolic types of fibers: *LDM* (fast-twitch glycolytic fibers – type IIB) and *IM* (slow-twitch oxidative fibers – type I) for U conformation class and 2–3 fat class animals.

* Correspondence: dominika_guzek@sggw.pl

2. Materials and methods

2.1. Preparation of samples

The analyzed samples were obtained from the federally inspected slaughter facilities, and animals used in the experiment were treated according to standard ethical norms. This study presents preliminary results, which we plan to subsequently verify in a larger group of animals.

The samples for this study originated from 5 Limousin bulls with an average hot carcass weight of 432 ± 31 kg (388–466 kg), U conformation class, and 2–3 fat class. The animals were raised according to typical practices. They were dam-reared to the age of natural weaning (8–9 months) and subsequently raised on silage and corn. All animals were slaughtered between 25 and 27 months of age and carcasses were stored at 4 °C for 5 days postmortem. Cuts were obtained from each carcass: *IM* from the blade (slow-twitch oxidative fibers — type I) and *LDM* (fast-twitch glycolytic fibers — type IIB) as the longest muscle in beef cattle. *LDM* was divided into 2 cuts: anterior from the cube roll (*longissimus dorsi et thoracis* muscle (*LDETM*) collected from Th7–Th9 vertebrae region), and posterior from the striploin (*longissimus dorsi et lumborum* muscle (*LDELM*) collected from Th12–L4 vertebrae region). Cuts were obtained from a commercial abattoir with a limited amount of intramuscular fat tissue and connective tissue. Collected meat was stored in vacuum at 0 °C for 5 days, 2 days in carcass form and 3 days in steak form. After aging, steaks were stored in vacuum at –18 °C. For the texture analysis, the representative beef steaks were thawed in a refrigerator (Küppersbusch Hausgeräte GmbH, Germany) until an internal temperature of 2 °C was reached.

2.2. Sarcomere length

Sarcomere length values were measured according to the method of Cross et al. (9) with some modifications (10). For each cut, 5 g of representative meat samples were collected and microscopic preparations were obtained from the sucrose solution. Samples were homogenized in 30 mL of cold 0.25 M sucrose at a low speed of 5000 rpm for 60 s with a PRO 200 mechanical homogenizer (PRO Scientific Inc., USA). The evaluation of the sarcomere length was conducted using the Carl Zeiss Axio Imager M2 microscope (Carl Zeiss, Germany) with the EC Plan-Neofluar 100×/1.30 Oil Ph 3 M27 objective and AxioCamMR5 camera. Microscopic preparations were observed in the differential interference contrast (DIC) (Figure 1). Sarcomere length was measured using AxioVision Rel.4.8.2 software (Carl Zeiss, Germany). Three myofibrils were selected for every sample, and the length of 25 sarcomeres was measured for each myofibril.

2.3. Near-infrared spectroscopy analyses

Basic composition was determined with the near-infrared spectroscopy (NIR) method. In order to obtain homogeneous mixtures for each cut, 150 g of representative




Figure 1. An example of microscopic preparation of myofibril with visible sarcomeres observed in the differential interference contrast.

meat samples were collected and homogenized with a blender for 30–60 s. Samples were placed on a petri dish (90 mm in diameter) in the NIR device (NIR Flex Solids N-500, spectral range 800–2500 nm). Spectral analysis software (NIR Ware 1.1, NIR Cal 5.1) was used to measure water, fat, protein, ash, and collagen content (%).

2.4. Computer image analysis

Marbling (%) of meat samples was determined by computer image analysis (CIA) according to the widely applied methodology (11). After blooming for 30 min, the representative beef steaks were placed on a matte green background to ensure easier segmentation. Pictures of beef steaks were taken using a CD QImaging MicroPublisher 5.0 RTV camera under fluorescent light in standard conditions (color temperature 5400 K). Each image was captured and saved in .tif format, and the area of intramuscular fat tissue was calculated using Image-Pro Plus 7 (Figure 2).




Figure 2. An example of intramuscular fat tissue (marbling) area calculation.

2.5. Statistical analysis

The W Shapiro–Wilk test was performed to verify the normality of distribution. Differences between traits were assessed using Student t-test or the Mann–Whitney U test. In order to characterize the relationships, 2-way analysis of variance (ANOVA) and post hoc tests (Scheffe test) were conducted. Analysis of correlation was carried out using Pearson’s correlation or Spearman’s rank correlation. Statistical analysis was conducted using Statistica software version 8.0 (StatSoft, USA), and a $P \leq 0.05$ level of significance was applied.

3. Results

Mean sarcomere length for the samples was obtained from 3 beef cuts (*IM*, *LDETM*, and *LDELM*) of 5 bulls (Table 1) ranging from 1.82 μm (observed for *LDELM*, the 2nd animal) to 2.87 μm (*LDETM*, the 3rd animal).

Beef characteristics from the 3 analyzed cuts, *IM*, *LDETM*, and *LDELM* (cumulative analysis for all the animals), are presented in Table 2.

Analysis of correlation between sarcomere length and beef characteristics (cumulative analysis for all the cuts and animals) and comparison of beef characteristics between samples characterized by various sarcomere lengths is presented in Table 3.

4. Discussion

4.1. Sarcomere length for analyzed samples

The observed results are similar to results obtained in previous studies, such as that of Rhee et al. (12), where sarcomere length was 1.80 μm for *LDM* and 2.25 μm for *IM*.

ANOVA revealed that differences in sarcomere length were associated with both analyzed factors, cut (P

Table 1. Mean \pm SD and median (min–max) sarcomere length for beef samples from 3 analyzed cuts and 5 analyzed animals (μm).

Animal	<i>Infraspinatus</i>		<i>Longissimus dorsi et thoracis</i>		<i>Longissimus dorsi et lumborum</i>	
	Mean \pm SD	Median (min–max)	Mean \pm SD	Median (min–max)	Mean \pm SD	Median (min–max)
1	2.68 \pm 0.10 A/a	2.68 (2.49–2.87)	2.40 \pm 0.16 A/b	2.39 (2.06–2.68)	2.22 \pm 0.11 A/c	2.25 (1.96–2.40)
2	2.65 \pm 0.10 AB/a	2.65 (2.46–2.86)	2.72 \pm 0.20 B/a	2.73 (2.21–3.05)	1.82 \pm 0.11 B/b	1.79 (1.66–2.01)
3	2.50 \pm 0.09 B/a	2.50 (2.33–2.65)	2.87 \pm 0.15 B/b	2.88 (2.46–3.11)	2.48 \pm 0.13 C/a	2.48 (2.22–2.75)
4	2.66 \pm 0.14 AB/a	2.66 (2.45–2.89)	1.95 \pm 0.09 C/b	1.93 (1.80–2.16)	2.08 \pm 0.11 A/b	2.07 (1.81–2.31)
5	2.85 \pm 0.13 C/a	2.86 (2.54–3.05)	2.33 \pm 0.15 A/b	2.34 (2.03–2.61)	2.47 \pm 0.11 C/b	2.46 (2.29–2.70)

For all groups: normal distribution (verification on the basis of W Shapiro–Wilk test, $P \leq 0.05$).

Mean values marked with capital letters (A, B, C) in columns and different lower case letters (a, b, c) in rows differ on the basis of post hoc Scheffe test criteria for $P \leq 0.05$.

Table 2. Mean \pm SD and median (min–max) values for beef characteristics from 3 analyzed cuts (cumulative analysis for all animals): content of water, fat, protein, ash, and collagen in samples and marbling in surface.

Basic composition [%]	<i>Infraspinatus</i>		<i>Longissimus dorsi et thoracis</i>		<i>Longissimus dorsi et lumborum</i>	
	Mean \pm SD	Median (min–max)	Mean \pm SD	Median (min–max)	Mean \pm SD	Median (min–max)
Water	74.4 \pm 2.3A	75.4 (70.6–76.3)	75.8 \pm 0.4A	75.7 (75.5–76.4)	75.0 \pm 1.7A	75.3 (72.2–76.3)
Fat	3.4 \pm 2.6A	2.8 (1.4–7.9)	1.0 \pm 0.5B	1.1 (0.4–1.6)	1.4 \pm 1.5AB	0.7A (0.5–4.0) ^a
Protein	19.8 \pm 0.9A	19.8 (18.6–20.9)	21.7 \pm 0.6B	21.9 (21.1–22.4)	22.8 \pm 0.5C	22.7 (22.1–23.6)
Ash	1.1 \pm 0.2A	1.2 (0.9–1.2)	1.1 \pm 0.0A	1.1 (1.1–1.2)	1.1 \pm 0.1A	1.1 (1.0–1.3)
Collagen	2.0 \pm 0.7A	1.7A (1.6–3.3) ^a	1.3 \pm 0.2B	1.2 (1.1–1.6)	1.3 \pm 0.3AB	1.3 (0.9–1.7)
Marbling	3.5 \pm 1.6A	3.2 (1.6–5.4)	2.6 \pm 0.7A	2.9 (1.4–3.3)	1.2 \pm 0.3B	1.1 (0.8–1.6)

^aDistribution different than normal (verification on the basis of W Shapiro–Wilk test, $P \leq 0.05$).

Mean values marked with capital letters (A, B, C) in rows differ on the basis of Student t-test (normal distribution) or Mann–Whitney U test (distribution different than normal) criteria for $P \leq 0.05$.

Table 3. Analysis of correlation between sarcomere length and characteristics of beef (cumulative analysis for all cuts and animals) accompanied by comparison of beef characteristics between samples characterized by varying sarcomere lengths.

Basic composition	Correlation between sarcomere length and characteristics of beef		Comparison between samples characterized by varying sarcomere lengths				
	P-value	R coefficient	Mean sarcomere length lower than 2.5 µm		Mean sarcomere length higher than 2.5 µm		P-value
			Mean ± SD	Median (min–max)	Mean ± SD	Median (min–max)	
Water	0.771 ^a	-0.0821	75.3 ± 1.3	75.6 (72.2–76.3) ^a	74.9 ± 2.1	75.5 (70.6–76.4) ^a	1.0000
Fat	0.101 ^a	0.4393	1.3 ± 1.1	0.9 (0.5–4.0) ^a	2.7 ± 2.5	1.9 (0.4–7.9) ^a	0.1480
Protein	0.039 [*]	-0.5365	22.3 ± 0.8	22.5 (21.1–23.6)	20.4 ± 1.3	20.3 (18.6–22.0)	0.0046 ^{**}
Ash	0.345	0.2621	1.1 ± 0.1	1.1 (1.0–1.3)	1.1 ± 0.1	1.1 (0.8–1.2)	0.7881
Collagen	0.167 ^a	0.3763	1.3 ± 0.3	1.3 (0.9–1.7)	1.7 ± 0.7	1.6 (1.1–3.3) ^a	0.1832
Marbling	0.011 [*]	0.6353	1.7 ± 0.9	1.4 (0.8–3.3)	3.3 ± 1.4	2.9 (1.6–5.4)	0.0226 ^{**}

^aDistribution different than normal (verification on the basis of W Shapiro–Wilk test, P ≤ 0.05).

^{*}: P ≤ 0.05 for Pearson’s correlation (normal distribution) or Spearman’s rank correlation (distribution different than normal).

^{**}: P ≤ 0.05 for Student t-test (normal distribution) or Mann–Whitney U test (distribution different than normal).

<0.01) and animal (P < 0.01). A combined effect of these factors, cut × animal, was also observed (P < 0.01). In paired comparisons, an analysis of sarcomere length was conducted with the same cut and animal. Sarcomeres from *LDETM* were longest in the third animal. In the case of the 4 other animals, the longest sarcomeres were observed for *IM*, which was consistent with results of other studies (13). At the same time, the shortest sarcomeres were observed in *LDELM*. *LDETM* and *LDELM* are parts of the same muscle, yet they have different characteristics, observed by other researchers in cases of purge loss, cooking loss, and sensory features (14). Due to the observed differences between *LDETM* and *LDELM*, it was deemed appropriate in this study to divide the muscle into cuts and analyze the differences between the single parts. Given that the above data do indicate such differences, it can be concluded that both muscle and cut are important factors influencing sarcomere length. The results indicate differences between *LDETM* and *LDELM* for 3 of 5 analyzed animals. However, some general conclusions may be presented for cuts. The shortest sarcomeres are observed for *LDELM* and the longest for *IM*. These results are in general agreement with previous studies (12).

In comparisons made between animals within the same cut, differences among the paired comparisons were also observed. The longest sarcomeres were observed for animals 5 (*IM* muscles compared), 3, and 2 (*LDETM*), and 3 and 5 (*LDELM*), respectively, while the shortest were observed for animals 4, 2, and 3 (*IM*) and 4 (*LDETM*) and 2 (*LDELM*), respectively. Similarly, Koohmaraie (15) observed the greatest variation in sarcomere length between animals, not regarding other factors.

It should be emphasized that the longest sarcomeres (*LDETM*) and the shortest sarcomeres (*IM*) were observed in animals 2 and 3. Simultaneously, in the case of *LDELM*, the longest sarcomeres were identified in animals 3 and 5, and the shortest in animal 2. The mean value for striploin (*LDELM*) in the case of animal 2 was the lowest among all presented sarcomere length values, and the minimum and maximum values of sarcomere length for animal 2 were the lowest of all. This variation in sarcomere length had been reported by other authors in studies on different muscles such as *gluteus medius* and *semitendinosus* (12). However, in the case of *semitendinosus* muscle Weaver et al. found no variation in sarcomere length (16). Thus, it is essential to broaden the knowledge and understanding of all factors influencing sarcomere length.

Given that the studied animals originated from a single herd and were characterized by the same breed, sex, age, and comparable body weight, it can be suggested that features affecting sarcomere length are associated with their detailed genetics. Sarcomere length also depends on ultimate pH (17), shortening process during rigor mortis development (18), and stretch or contraction (16). However, these features were not taken into account as variables in the present study.

4.2. Basic composition of analyzed samples

In the case of certain analyzed features, such as water and ash content, no differences were observed between cuts. This was in agreement with previous reports by other researchers (19). It may be concluded that water and ash content are relatively constant within the entire carcass.

In the case of fat and collagen content, differences were observed between *IM* and *LDETM*. Both fat and collagen contents were higher in the case of *IM* (3.4% and 1.7%, respectively) than in *LDETM* (1.0% and 1.3%, respectively). In the case of *LDELM*, fat and collagen content did not differ from the content observed for *IM* and *LDETM*. However, relationships were not the same for all features. Differences were inverse for marbling levels. Marbling for *LDELM* (1.2%) was lower than for *IM* and *LDETM* (2.6%–3.5% for the analyzed cuts). It may be stated that differences between cuts are not identical for all features. Instead, they may depend on the construction and function of the muscle. It is generally observed that differences in muscle characteristics also occur among muscles of various types, even if animals are raised under similar production conditions (20).

The most significant differences between muscles were observed in the case of protein content. *IM* was characterized by the lowest protein content (19.8%), *LDELM* was characterized by the highest protein content (22.8%), and the content value was intermediate for *LDETM* (21.7%). It can be concluded that protein content is the most variable feature for the analyzed cuts, in all likelihood depending on the muscle fibers.

4.3. Relationship between sarcomere length and basic composition for analyzed samples

Correlation between sarcomere length and beef characteristics is observed only in the case of protein content (analyzed with NIR) and marbling (analyzed with CIA). The comparison of beef characteristics between samples characterized by varying sarcomere lengths confirmed the obtained results. It can be stated that in the case of beef samples characterized by longer sarcomeres, lower protein content ($P = 0.039$, $R = -0.5365$ for correlation; $P = 0.005$ for comparison) and higher marbling level ($P = 0.011$, $R = 0.6353$ for correlation; $P = 0.023$ for comparison) are observed. In several studies it was proven that increase in sarcomere length is positively correlated with marbling score (4). Correlations were

moderate for protein content and marbling level ($R < 0.7$), whereas sarcomere length for *LDELM* was very strongly correlated with marbling level ($P = 0.006$; $R = 0.9692$, Pearson's correlation). It may be suggested that other unknown factors influence correlation, and the prediction of sarcomere length solely on the basis of these analyzed factors might be impossible.

In conclusion, differences in beef sarcomere length are associated with the cut and animal. The combined effect of both factors, cut \times animal, was also observed. In most animals, the longest sarcomeres were observed for the *infraspinatus* muscle. In the case of *longissimus dorsi*, differences in sarcomere length between its parts (*longissimus dorsi et thoracis* and *longissimus dorsi et lumborum*) were also observed. Sarcomere length is positively correlated with content of intramuscular fat in the surface of steak and negatively correlated with content of protein. Several other beef features (content of fat, protein, collagen, and marbling) also depend on the cut, which may be partially influenced by differences in sarcomere length. Additional knowledge about sarcomere length may explain the variability of beef characteristics. Therefore, further analysis of beef sarcomere length in the case of various cuts and animals is crucial for expanding the understanding of the associations presented.

Acknowledgments

This research was supported by the program 'Improving Polish sources of plant protein, their production, trading, and application in pasturages – Production of high-quality pasturages from the permanent pastures' (2011–2015) [Ulepszenie krajowych źródeł białka roślinnego, ich produkcji, systemu obrotu i wykorzystania w paszach – Produkcja wysokiej jakości pasz z trwałych użytków zielonych (2011–2015)]. The authors thank research assistants Małgorzata Konarska, Karolina Kozan, Jacek Pietras, and Anna Sakowska for their technical support during this study.

References

1. Kirchofer KS, Calkins CR, Gwartney BL. Fiber-type composition of muscles of the beef chuck and round. *J Anim Sci* 2002; 80: 2872–2878.
2. Stolarski GD, Baird BE, Miller RK, Savell JW, Sams AR, Taylor JE, Sanders JO, Smith SB. Factors influencing the variation in tenderness of seven major beef muscles from three Angus and Brahman breed crosses. *Meat Sci* 2003; 73: 475–483.
3. Hwang YH, Kim GD, Jeong JY, Hur SJ, Joo ST. The relationship between muscle fiber characteristics and meat quality traits of highly marbled Hanwoo (Korean native cattle) steers. *Meat Sci* 2010; 86: 456–461.
4. Li C, Zhou G, Xu X, Zhang J, Xu S, Ji Y. Effects of marbling on meat quality characteristics and intramuscular connective tissue of beef *longissimus* muscle. *Asian Austral J Anim* 2006; 19: 1799–1808.
5. Wick MP, Marriott NG. The Relationship of the Sarcomeric Architecture to Meat Tenderness. Special Circular 172-99. Columbus, OH, USA: Ohio State University; 1999.
6. Wheeler TL, Shackelford SD, Koohmaraie M. Variation in proteolysis, sarcomere length, collagen content, and tenderness among major pork muscles. *J Anim Sci* 2000; 78: 958–965.

7. Maher SC, Mullen AM, Moloney AP, Reville W, Buckley DJ, Kerry JP, Troy DJ. Ultrastructural variation in beef *M. longissimus dorsi* as an explanation of the variation in beef tenderness. *J Food Sci* 2005; 70: 579–584.
8. Christensen M, Kok C, Ertbjerg P. Mechanical properties of type I and type IIB single porcine muscle fibres. *Meat Sci* 2006; 73: 422–425.
9. Cross HR, West RL, Dutson TR. Comparison of methods for measuring sarcomere length in beef *semitendinosus* muscle. *Meat Sci* 1981; 5: 261–266.
10. Guzek D, Głabska D, Głabski K, Plewa P, Plewa R, Wierzbicka A. Comparison of sarcomere length for two types of meat from animal family *Suidae* – analysis of measurements carried out by microscopic technique. *Adv Sci Tech Res J* 2012; 6: 13–17.
11. Yang XJ, Albrecht E, Ender K, Zhao RQ, Wegner J. Computer image analysis of intramuscular adipocytes and marbling in the *longissimus* muscle of cattle. *J Anim Sci* 2006; 84: 3251–3258.
12. Rhee MS, Wheeler TL, Shackelford SD, Koohmaraie M. Variation in palatability and biochemical traits within and among eleven beef muscles. *J Anim Sci* 2004; 82: 534–550.
13. Purchas RW, Zou M. Composition and quality differences between the *longissimus* and *infraspinatus* muscles for several groups of pasture-finished cattle. *Meat Sci* 2008; 80: 470–479.
14. Hergenreder JE, Hosch JJ, Varnold KA, Haack AL, Senaratne LS, Pokharel S, Beauchamp C, Lobaugh B, Calkins CR. The effects of freezing and thawing rates on tenderness, sensory quality, and retail display of beef subprimals. *J Anim Sci* 2013; 91: 483–490.
15. Koohmaraie M. Biochemical factors regulating the toughening and tenderization processes of meat. *Meat Sci* 1996; 43: 193–201.
16. Weaver AD, Bowker BC, Gerrard DE. Sarcomere length influences postmortem proteolysis of excised bovine *semitendinosus* muscle. *J Anim Sci* 2008; 86: 1925–1932.
17. Costa RS, Henry FC, Ferreira KS, do Valle FRAF, Quirino CR. Characterization of *rigor mortis* of *longissimus dorsi* and *triceps brachii* muscles of male cattle carcasses. *Afr J Biotechnol* 2012; 11: 8127–8132.
18. Koohmaraie M, Doumit ME, Wheeler TL. Meat toughening does not occur when rigor shortening is prevented. *J Anim Sci* 1996; 74: 2935–2942.
19. Patten LE, Hodgen JM, Stelzleni AM, Calkins CR, Johnson DD, Gwartney BL. Chemical properties of cow and beef muscles: benchmarking the differences and similarities. *J Anim Sci* 2008; 86: 1904–1916.
20. Jurie C, Martin JF, Listrat A, Jailler R, Culioli J, Picard B. Effects of age and breed of beef bulls on growth parameters, carcass and muscle characteristics. *Anim Sci* 2005; 80: 257–263.