

1-1-2011

## In vitro antagonistic activity of fungi isolated from sclerotia on potato tubers against *Rhizoctonia solani*

ERKOL DEMİRCİ

ELİF DANE

CAFER EKEN

Follow this and additional works at: <https://journals.tubitak.gov.tr/biology>



Part of the [Biology Commons](#)

---

### Recommended Citation

DEMİRCİ, ERKOL; DANE, ELİF; and EKEN, CAFER (2011) "In vitro antagonistic activity of fungi isolated from sclerotia on potato tubers against *Rhizoctonia solani*," *Turkish Journal of Biology*. Vol. 35: No. 4, Article 9. <https://doi.org/10.3906/biy-1004-98>

Available at: <https://journals.tubitak.gov.tr/biology/vol35/iss4/9>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Biology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

## In vitro antagonistic activity of fungi isolated from sclerotia on potato tubers against *Rhizoctonia solani*

Erkol DEMİRÇİ<sup>1</sup>, Elif DANE<sup>2</sup>, Cafer EKEN<sup>1,3</sup>

<sup>1</sup>Department of Plant Protection, Faculty of Agriculture, Atatürk University, 25240 Erzurum - TURKEY

<sup>2</sup>Provincial Directorate of Agriculture, Section of Plant Protection, Manisa - TURKEY

<sup>3</sup>Ardahan University, Ardahan - TURKEY

Received: 08.04.2010

**Abstract:** Forty-five fungal isolates were obtained from sclerotia of *Rhizoctonia solani* on potato tubers in Erzurum, Turkey. The interaction between fungal isolates and *R. solani* was studied in dual culture technique. Some fungal isolates affected *R. solani* by antibiosis and/or parasitism. Results of the antagonism tests showed that *Acremonium* sp., *Gliocladium viride*, *Paecilomyces marquandii*, *Paecilomyces sulphurellus*, *Penicillium camemberti*, *Penicillium expansum*, *Penicillium frequentans* (ME-50), *Penicillium nigricans*, *Penicillium olsonii*, *Penicillium phialosporum*, *Sporothrix* sp. (MCY-4), *Sporothrix schenckii*, and *Verticillium dahliae* isolates produced an inhibition zone in front of the *R. solani* colony to a varying degree. *Trichoderma harzianum* isolates were able to overgrow the mycelium of *R. solani*. Physical colony contact was observed between the remaining 21 fungal isolates and *R. solani*. Furthermore, coiling of hyphae of *Acremonium* sp., *Acremonium strictum*, *Gliocladium catenulatum*, *G. viride*, and *T. harzianum* around those of *R. solani* was commonly observed.

**Key words:** Biocontrol, potato, antibiosis, parasitism, *Rhizoctonia solani*

### Patates yumrularındaki sklerotiumlardan izole edilen fungusların *Rhizoctonia solani*'ye in vitro antagonistik etkileri

**Özet:** Patates yumruları üzerinde bulunan *Rhizoctonia solani*'nin sklerotiumlarından 45 fungal izolat elde edilmiştir. Fungal izolatlar ve *R. solani* arasındaki etkileşim ikili kültür yöntemi ile çalışılmıştır. Bazı fungal izolatlar *R. solani*'ye antibiosis ve/veya parazitizm yolu ile etki yapmıştır. Antagonizm test sonuçlarına göre, *Acremonium* sp., *Gliocladium viride*, *Paecilomyces marquandii*, *Paecilomyces sulphurellus*, *Penicillium camemberti*, *Penicillium expansum*, *Penicillium frequentans* (ME-50), *Penicillium nigricans*, *Penicillium olsonii*, *Penicillium phialosporum*, *Sporothrix* sp. (MCY-4), *Sporothrix schenckii* ve *Verticillium dahliae* izolatları ile *R. solani* kolonileri arasında değişen derecelerde inhibisyon zonu oluşmuştur. *Trichoderma harzianum* izolatları ise *R. solani* miselyumunun üzerinde gelişmiştir. Geri kalan 21 izolat ile *R. solani* kolonileri arasında fiziksel koloni teması gerçekleşmiştir. Ayrıca, *Acremonium* sp., *Acremonium strictum*, *Gliocladium catenulatum*, *G. viride* ve *T. harzianum* izolatlarına ait hiflerin *R. solani* hiflerini sarması yaygın olarak gözlenmiştir.

**Anahtar sözcükler:** Biyokontrol, patates, antibiosis, parazitizm, *Rhizoctonia solani*

## Introduction

*Rhizoctonia solani* Kühn [teleomorph: *Thanatephorus cucumeris* (Frank) Donk.] is an important pathogen responsible for serious damage in many crops including potato (*Solanum tuberosum* L.) (1). *R. solani* causes stem canker and black scurf of potato and occurs in all potato growing areas of the world (2). The pathogen is also found to be widely distributed on the potato plants and tubers in Erzurum province (3-5). Rhizoctonia disease of potato is mainly caused by *R. solani* anastomosis group (AG)-3, but isolates that belong to other AGs, such as AG-2 type 1, AG-2 type 2, AG-4, and AG-5, also infect potato stems and/or tubers (4).

Both soil-borne and tuber-borne inoculum of *R. solani* is important in disease development on potato (6,7). Present chemical and cultural control methods have reduced the soil-borne and tuber-borne inoculum, although research has been directed toward the use of antagonists for biocontrol of *R. solani* on potato. A biocontrol agent may act against pathogens by using one or more of the following mechanisms: competition, antibiosis, and parasitism as well as activating host defense mechanisms (8). In fact, several fungi have been reported to be effective biocontrol agents of *R. solani* on potato. Among these are species of *Gliocladium* (9-11), *Trichoderma* (9-16), and *Verticillium* (17,18). *Chaetomium olivaceum*, *Cylindrocarpon destructans*, *Epicoccum nigrum*, *Fusarium culmorum*, *Fusarium moniliforme*, *Gliocladium viride* (syn. *Gliocladium deliquescens*), *Gliocladium roseum*, *Penicillium cyclopium*, *Penicillium nigricans*, *Trichoderma harzianum*, and *Trichothecium roseum* were frequently isolated from sclerotia of *R. solani* (12). In another study, fungal isolates from the sclerotia of *R. solani* were identified as *Alternaria*, *Aspergillus*, *Cladosporium*, *Coniothyrium*, *Curvularia*, *Gliocladium*, *Fusarium*, *Metarhizium*, *Penicillium*, *Phoma*, *Phytophthora*, and *Trichoderma* genera (19). Additionally, *Verticillium biguttatum* was reported from sclerotia of *R. solani* on potato tubers in Turkey (20).

The objective of the present study was to isolate and identify fungi from sclerotia of *R. solani* on potato tubers, and to determine the efficacy of their potential as biocontrol agents on interaction with *R. solani* in pure culture.

## Materials and methods

### Isolation and identification of fungal isolates

Potato tubers of cv. Marfona affected with black scurf were obtained from 2 randomly selected potato storage facilities in Erzurum, Turkey, and 100 sclerotia from each sample were divided into 5 sub-samples having 20 sclerotia. Isolation of fungi was studied using a method modified from Chand and Logan (12). Each sub-sample was placed on autoclaved moist coarse sand in petri dishes. After incubation at 20 °C for 4 weeks, the sclerotia of each sub-sample were blended together in 200 mL of sterile distilled water together with agar (2%). Serial dilutions ( $10^{-2}$ ,  $10^{-3}$ ,  $10^{-4}$ , and  $10^{-5}$ ) were made and 1 mL aliquots of each dilution were pipetted into 10 petri dishes. Each of them was mixed with 15-20 mL of cooled (40-45 °C) potato dextrose agar (PDA) by gentle swirling. The plates were incubated at 25 °C in the dark for 7 days or longer and then emerging fungi were repeatedly sub-cultured to obtain pure culture. The fungi isolates were identified using cultural and morphological features (21,22). Single-spore isolates were transferred to PDA medium in tubes for preservation at 4 °C.

### Dual culture tests

Fungal isolates from sclerotia on potato tubers were tested in a dual culture assay against *R. solani* representing AG-3 (isolate R-99), which was recovered from potato tubers with black scurf. A mycelial disc (4 mm diam.) of each fungal isolate was transferred on fresh PDA, 1 cm from the wall of each petri dish, immediately or after 2-4 days, depending upon the growth rate of the fungus, a 4 mm disc of *R. solani* removed from the colony margin of actively growing cultures on PDA was placed 6 cm away from the disc of fungal isolate in the same petri dish. All petri dishes were incubated in the dark at 25 °C for 7 days or longer according to the growth rate of fungi. The percentage inhibition of radial growth [ $100 \times (r_1 - r_2)/r_1$ ] of *R. solani* and the width of the zone of inhibition between both colonies were recorded as described by Royse and Ries (23). Mode of inhibition was assessed on a scale from 1 to 4, in which 1 = mycelial growth of *R. solani* ceased due to overgrowth of interacting fungus, 2 = partial inhibition of *R. solani* and interacting fungus but both grow slowly over each other, 3 = mutual inhibition at a distance <

2 mm, 4 = growth of *R. solani* inhibited at a distance > 2 mm (12). The mean of the 4 measurements was recorded for each fungus.

### Hyphal interaction between fungal isolates and *R. solani*

Hyphal interactions were observed in 5-day-old *R. solani* cultures on 2% water agar (WA) in 9 cm petri dishes, which were inoculated with a 4 mm disc removed from actively growing cultures on PDA of each fungal isolate. Plates were incubated at 25 °C in a sterile humidity chamber (100% r.h.). After 7-14 days, 3 rectangular blocks (about 4 cm × 2 cm) from each plate were cut, mounted on glass slides, and examined for hyphal interaction between fungal isolates and *R. solani* by a phase contrast microscope at 400× magnification.

### Results and discussion

Totally, 45 fungal isolates (Table) were obtained from tuber-borne sclerotia of *R. solani* in Erzurum province. They were identified by cultural and morphological characteristics and tested for their potential as biocontrol agents.

The isolates were identified as *Acremonium* sp. (1 isolate), *Acremonium strictum* (1 isolate), *Alternaria alternata* (1 isolate), *Chaetomium* sp. (1 isolate), *Cladosporium cladosporioides* (1 isolate), *Fusarium equiseti* (3 isolates), *Fusarium oxysporum* (1 isolate), *Fusarium solani* (2 isolates), *Fusarium verticillioides* (1 isolate), *Gliocladium catenulatum* (1 isolate), *Gliocladium viride* (2 isolates), *Paecilomyces* sp. (2 isolates), *Paecilomyces marquandii* (2 isolates), *Paecilomyces sulphurellus* (1 isolate), *Penicillium camemberti* (1 isolate), *Penicillium expansum* (7 isolates), *Penicillium frequentans* (2 isolates), *P. nigricans* (1 isolate), *Penicillium olsonii* (1 isolate), *Penicillium phialosporum* (2 isolates), *Plectosporium tabacinum* (5 isolates), *Sporothrix* sp. (2 isolates), *Sporothrix schenckii* (1 isolate), *T. harzianum* (2 isolates), and *Verticillium dahliae* (1 isolate).

All isolates were tested against *R. solani* AG-3 isolate in dual culture. Colony and hyphal interactions of these fungal isolates with *R. solani* on PDA are listed in the Table. Results of the antagonism tests showed that *T. harzianum* isolates were able to overgrow (Mode 1) the mycelium of *R. solani*.

Physical colony contact (Mode 2) was observed between the 21 isolates of fungi and *R. solani*. No physical contact was observed between 22 isolates of fungi and *R. solani*; moreover, *Acremonium* sp., *G. viride*, *P. marquandii*, *P. sulphurellus*, *P. camemberti*, *P. expansum*, *P. frequentans* (ME-50), *P. nigricans*, *P. olsonii*, *P. phialosporum*, *Sporothrix* sp. (MCY-4), *S. schenckii*, and *V. dahliae* isolates produced an inhibition zone (Mode 3 and 4) in front of the *R. solani* colony (Table). An inhibition zone was observed, which indicates the presence of fungistatic metabolites secreted by these fungi. Different isolates of the same fungus species showed the same mode of inhibition to the pathogen, except for *P. expansum*, *P. frequentans*, and *P. phialosporum* isolates, which showed different inhibition modes, with interaction modes 3 and 4, 2 and 3, and 3 and 4, respectively.

A biocontrol agent may act against pathogens by using one or more of the following mechanisms: competition, antibiosis, and parasitism as well as activating host defense mechanisms (8). Antagonistic activity by *Penicillium* species against *R. solani* has been observed, and it has been reported in relation to the production of toxic metabolites (12,24-27). According to reports in the literature, all *Penicillium* species in the present study except *P. expansum* and *P. nigricans* are new records as possible antagonists of *R. solani*.

Coiling of hyphae of *Acremonium* sp., *A. strictum*, *G. catenulatum*, *G. viride*, and *T. harzianum* around those of *R. solani* was commonly observed (Table). Of these results, *Acremonium* sp. and *G. viride* also affected *R. solani* by antibiosis and parasitism. *R. solani* can be parasitized by parasites such as *Gliocladium* spp. (11,12,18), *Trichoderma* spp. (11,12,15), and *Verticillium* spp. (11,17,18,20). *A. strictum* is notable for parasitizing *Helminthosporium solani* (28) and *Botrytis cinerea* (29).

Several microorganisms including fungi have been shown to be effective antagonists of *R. solani*. The genera *Gliocladium* and *Trichoderma* contain many mycoparasitic species that are considered good biocontrol agents against soil-borne pathogens (9). Within the genus *Trichoderma*, species such as *T. hamatum*, *T. harzianum*, *T. reesei*, *T. virens*, and *T. viride* have demonstrated excellent antagonistic activity against *R. solani* on potato pot and/

Table. Colony and hyphal interactions of fungal isolates with *Rhizoctonia solani* in vitro.

Fungal species	Isolate number	Mode of inhibition*	Width of the zone (mm)	Inhibition of <i>R. solani</i> (%)	Hyphal interaction <sup>‡</sup>
<i>Trichoderma harzianum</i>	MEY-1	1	-	69	+
<i>Trichoderma harzianum</i>	MCY-1	1	-	72	+
<i>Acremonium strictum</i>	MEY-10	2	-	20	+
<i>Alternaria alternata</i>	MCY-2	2	-	29	-
<i>Chaetomium</i> sp.	ME-8	2	-	30	-
<i>Cladosporium cladosporioides</i>	AE-4	2	-	13	-
<i>Fusarium equiseti</i>	MEY-4	2	-	47	-
<i>Fusarium equiseti</i>	AE-9	2	-	48	-
<i>Fusarium equiseti</i>	MC-9	2	-	37	-
<i>Fusarium oxysporum</i>	MEY-6	2	-	35	-
<i>Fusarium solani</i>	AE-8	2	-	42	-
<i>Fusarium solani</i>	AE-7	2	-	40	-
<i>Fusarium verticillioides</i>	MEY-3	2	-	52	-
<i>Gliocladium catenulatum</i>	MEY-2	2	-	37	+
<i>Paecilomyces</i> sp.	MC-1	2	-	22	-
<i>Paecilomyces</i> sp.	AE-3	2	-	50	-
<i>Penicillium frequentans</i>	ME-57	2	-	26	-
<i>Plectosporium tabacinum</i>	MC-10	2	-	33	-
<i>Plectosporium tabacinum</i>	MC-4	2	-	34	-
<i>Plectosporium tabacinum</i>	MC-11	2	-	38	-
<i>Plectosporium tabacinum</i>	MEY-5	2	-	35	-
<i>Plectosporium tabacinum</i>	AE-6	2	-	27	-
<i>Sporothrix</i> sp.	MC-5	2	-	34	-
<i>Paecilomyces marquandii</i>	MEY-8	3	1.0	39	-
<i>Paecilomyces marquandii</i>	MEY-9	3	1.0	37	-
<i>Penicillium expansum</i>	MC-61	3	0.8	17	-
<i>Penicillium frequentans</i>	ME-50	3	0.9	24	-
<i>Penicillium nigricans</i>	AE-62	3	0.5	21	-
<i>Penicillium phialosporum</i>	ME-51	3	1.1	14	-
<i>Acremonium</i> sp.	AE-5	4	3.9	47	+
<i>Gliocladium viride</i>	ME-7	4	2.3	35	+
<i>Gliocladium viride</i>	ME-10	4	2.1	31	+
<i>Paecilomyces sulphurellus</i>	MC-2	4	2.9	21	-
<i>Penicillium camemberti</i>	AE-63	4	6.1	31	-
<i>Penicillium expansum</i>	ME-56	4	5.5	35	-
<i>Penicillium expansum</i>	MC-60	4	5.3	29	-
<i>Penicillium expansum</i>	ME-52	4	3.8	33	-
<i>Penicillium expansum</i>	AE-64	4	7.3	28	-
<i>Penicillium expansum</i>	ME-53	4	12.0	40	-
<i>Penicillium expansum</i>	ME-59	4	7.8	19	-
<i>Penicillium olsonii</i>	ME-58	4	7.8	35	-
<i>Penicillium phialosporum</i>	ME-55	4	8.3	29	-
<i>Sporothrix</i> sp.	MCY-4	4	3.5	32	-
<i>Sporothrix schenckii</i>	MCY-3	4	4.0	25	-
<i>Verticillium dahliae</i>	ME-2	4	2.5	33	-

\* 1 = Mycelial growth of *R. solani* ceased due to overgrowth of interacting fungus, 2 = Partial inhibition of *R. solani* and interacting fungus but both grow slowly over each other, 3 = Mutual inhibition at a distance < 2 mm, 4 = Growth of *R. solani* inhibited at a distance > 2 mm (12).

<sup>‡</sup>+: Coiling of *R. solani* hyphae, -: No coiling of *R. solani* hyphae.

or field tests (10,12-16). *T. harzianum*, the most commonly studied species, was tested with varying degrees of success against *R. solani* on potato. *G. virens* successfully controlled *R. solani* on potato in greenhouse and field tests (10). Currently, commercial biocontrol products including several *Trichoderma* and *Gliocladium* species are used for the control of soil-borne pathogens (e.g. *R. solani*) in the United States (30,31) and some other countries (31).

In another study (20), *V. biguttatum* was isolated from sclerotia of *R. solani* on potato tubers in Erzurum, and it also significantly reduced the disease severity of *R. solani* on potato sprouts in pot experiments. As a matter of fact, the mycoparasite *V. biguttatum* is an effective biocontrol agent against *Rhizoctonia* disease of potato in field experiments (17). It was shown to quickly establish itself on the host fungus by colonizing the sclerotia, and killing the moniliod cells (32). This is the first observation except *V. biguttatum* on potential biocontrol agents from sclerotia of *R. solani* on potato tubers in Erzurum province.

Some of the fungal species encountered in this study (e.g. *G. catenulatum*, *G. viride*, *P. expansum*, *P. nigricans*, and *T. harzianum*) have previously been studied as antagonists of *R. solani*. Other species, such as *Acremonium* sp., *A. strictum*, *Paecilomyces*

*marquandii*, *P. sulphurellus*, *Penicillium camemberti*, *P. frequentans* (ME-50), *P. olsonii*, *P. phialosporum*, *Sporothrix* sp. (MCY-4), and *S. schenckii* appear to be candidates for in vivo investigations to check their suitability as biocontrol agents. Other fungi (e.g. *A. alternata*, *Chaetomium* sp., *C. cladosporioides*, *F. equiseti*, *F. oxysporum*, *F. solani*, *F. verticillioides*, *Paecilomyces* sp., *P. frequentans* (ME-57), *P. tabacinum*, and *Sporothrix* sp. (MC-5)) are not antagonists of *R. solani*, and some of them and *V. dahliae* are pathogens on potato.

### Acknowledgements

The authors would like to thank the Atatürk University Scientific Research Projects Unit for its financial support. We are also grateful to Professor İsmet Hasenekoğlu (Kazım Karabekir Education Faculty, Atatürk University, 25240 Erzurum, Turkey) for identification of some of the fungi species.

### Corresponding author:

Erkol DEMİRCİ

Department of Plant Protection,

Faculty of Agriculture, Atatürk University,

25240 Erzurum - TURKEY

E-mail: edemirci@atauni.edu.tr

### References

- Ogoshi A. Introduction - the genus *Rhizoctonia*. In: Sneh B, Jabaji-Hare S, Neate S, Dijst G. eds. *Rhizoctonia* Species: Taxonomy, Molecular Biology, Ecology, Pathology and Disease Control. Kluwer Academic Publishers, Dordrecht: 1996, pp. 1-9.
- Frank JA. *Rhizoctonia* canker (Black scurf). In: Hooker WC. ed. *Compendium of Potato Diseases*. The American Phytopathological Society; 1986, pp. 52-54.
- Demirci E, Döken MT. Studies on the determination and identification of fungal agents of the diseases isolated from the roots, stems and leaves of potatoes in Erzurum Region. *J Turkish Phytopath* 18: 51-60, 1989.
- Demirci E, Döken MT. Anastomosis groups and pathogenicity of *Rhizoctonia solani* Kühn isolates from potatoes in Erzurum-TÜRKİYE. *J Turkish Phytopath* 22: 95-102, 1993.
- Eken C, Demirci E, Şahin F. Pathogenicity of the fungi determined on tubers from potato storages in Erzurum, Türkiye. *J Turkish Phytopath* 29: 61-69, 2000.
- Frank JA, Leach SS. Comparison of tuberborne and soilborne inoculum in the *Rhizoctonia* disease of potato. *Phytopathology* 70: 51-53, 1980.
- Demirci E, Eken C. The effect of tuberborne and soilborne inoculum of *Rhizoctonia solani* Kühn on disease severity on potato. 7th Turkish Phytopathological Congress (26-29 September 1995, Adana-Turkey): 39-43 1995 (Turkish, with English abstr.).
- Papavizas GC, Lumsden RD. Biological-control of soilborne fungal propagules. *Annu Rev Phytopathol* 18: 389-413, 1980.
- Papavizas GC. *Trichoderma* and *Gliocladium*: biology, ecology, and potential for biocontrol. *Annu Rev Phytopathol* 23: 23-54, 1985.

10. Beagle-Ristaino JE, Papavizas GC. Biological controls of *Rhizoctonia* stem canker and black scurf of potato. *Phytopathology* 75: 560-564, 1985.
11. Van den Boogert PHJF. Mycoparasitism and biocontrol In: Sneh B, Jabaji-Hare S, Neate S, Dijst G. eds. *Rhizoctonia* Species: Taxonomy, Molecular Biology, Ecology, Pathology and Disease Control. Kluwer Academic Publishers, Dordrecht: 1996, pp. 485-493.
12. Chand T, Logan C. Antagonists and parasites of *Rhizoctonia solani* and their efficacy in reducing stem cancer of potato under controlled conditions. *Trans British Mycol Soc* 83: 107-112, 1984.
13. Tsrar (Lahkim) L, Barak R, Sneh B. Biological control of black scurf on potato under organic management. *Crop Prot* 20: 145-150, 2001.
14. Brewer MT, Larkin RP. Efficacy of several potential biocontrol organisms against *Rhizoctonia solani* on potato. *Crop Prot* 24: 939-950, 2005.
15. Grosch R, Scherwinski K, Lottmann J et al. Fungal antagonists of the plant pathogen *Rhizoctonia solani*: selection, control efficacy and influence on the indigenous microbial community. *Mycol Res* 110: 1464-1474, 2006.
16. Wilson PS, Ketola EO, Ahvenniemi PM et al. Dynamics of soilborne *Rhizoctonia solani* in the presence of *Trichoderma harzianum*: effects on stem canker, black scurf and progeny tubers of potato. *Plant Pathol* 57: 152-161, 2008.
17. Jager G, Velvis H. Biological control of *Rhizoctonia solani* on potatoes by antagonists. 4. Inoculation of seed tubers with *Verticillium biguttatum* and other antagonists in field experiments. *Netherlands J Plant Pathol* 91: 49-63, 1985.
18. Turhan G. Further hyperparasites of *Rhizoctonia solani* Kühn as promising candidates for biological control. *J Plant Dis Protect* 97: 208-215, 1990.
19. Feng SJ, Yang M, Zhou DB et al. Fungal and bacterial species isolated from sclerotia of *Rhizoctonia solani* and their effects on the growth of *R. solani*. *Acta Phytopathol Sin* 38: 557-560, 2008.
20. Demirci E, Eken C, Dane E. Biological control of *Rhizoctonia solani* on potato by *Verticillium biguttatum*. *Afr J Biotechnol* 8: 2503-2507, 2009.
21. Gerlach W, Nirenberg H. The Genus *Fusarium*, a Pictorial Atlas. Biologische Bundesanstalt Land- und Forstwirtschaft Inst. Mikrobiologie, Berlin-Dahlem; 1982.
22. Hasenekoğlu İ. Toprak Mikrofungusları. Atatürk Üniversitesi Yayınları, No, 689, Kazım Karabekir Eğitim Fakültesi Yayınları No, 11, Cilt, 1-7, Erzurum; 1991.
23. Royse DJ, Ries SM. The influence of fungi isolated from peach twigs on the pathogenicity of *Cytospora cincta*. *Phytopathology* 68: 603-607, 1978.
24. Anderson HA, Bracewell JM, Fraser AR et al. 5-Hydroxymaltol and mycophenolic acid, secondary metabolites from *Penicillium echinulatum*. *Trans British Mycol Soc* 91: 649-651, 1988.
25. Kharbanda PD, Dahiya JS. A metabolite of *Penicillium verrucosum* inhibitory to growth of *Leptosphaeria maculans* and *Rhizoctonia solani*. *Can J Plant Pathol* 12: 335, 1990.
26. De Stefano S, Nicoletti R, Milone A et al. 3-O-Methylfunicone, a fungitoxic metabolite produced by the fungus *Penicillium pinophilum*. *Phytochemistry* 52: 1399-1401, 1999.
27. Nicoletti R, De Stefano M, De Stefano S et al. Antagonism against *Rhizoctonia solani* and fungitoxic metabolite production by some *Penicillium* isolates. *Mycopathologia* 158: 465-474, 2004.
28. Rivera-Varas VV, Freeman TA, Gudmestad NC et al. Mycoparasitism of *Helminthosporium solani* by *Acremonium strictum*. *Phytopathology* 97: 1331-1337 2007.
29. Choi GJ, Kim JC, Jang KS et al. Mycoparasitism of *Acremonium strictum* BCP on *Botrytis cinerea*, the gray mold pathogen. *J Microbiol Biotechnol* 18: 167-170, 2008.
30. McSpadden Gardener BB, Fravel DR. Biological control of plant pathogens: Research, commercialization, and application in the USA. *Plant Health Progress*. Online, publication doi:10.1094/PHP-2002-0510-01-RV, 2002.
31. Eken C, Demirci E. Fungusların biyolojik mücadelede kullanımı. Atatürk Üniversitesi Ziraat Fakültesi Dergisi 28: 138-152, 1997.
32. Velvis H, Jager G. Biological control of *Rhizoctonia solani* on potatoes by antagonists. 1. Preliminary experiments with *Verticillium biguttatum*, a sclerotium-inhabiting fungus. *Neth J Plant Pathol* 89: 113-123, 1983.