

1-1-2017

On generalized Kropina change of s th root Finsler metrics with special curvature properties

BANKTESHWAR TIWARI

GHANASHYAM KR. PRAJAPATI

Follow this and additional works at: <https://journals.tubitak.gov.tr/math>



Part of the [Mathematics Commons](#)

Recommended Citation

TIWARI, BANKTESHWAR and PRAJAPATI, GHANASHYAM KR. (2017) "On generalized Kropina change of s th root Finsler metrics with special curvature properties," *Turkish Journal of Mathematics*: Vol. 41: No. 6, Article 19. <https://doi.org/10.3906/mat-1512-14>
Available at: <https://journals.tubitak.gov.tr/math/vol41/iss6/19>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

On generalized Kropina change of m th root Finsler metrics with special curvature properties

Bankteshwar TIWARI, Ghanashyam Kr. PRAJAPATI*

Department of Science and Technology-Centre for Interdisciplinary Mathematical Sciences (DST-CIMS),
Institute of Science, Banaras Hindu University, Varanasi, India

Received: 04.12.2015

Accepted/Published Online: 07.02.2017

Final Version: 23.11.2017

Abstract: In the present paper, we consider generalized Kropina change of m th root Finsler metrics and prove that every generalized Kropina change of m th root Finsler metrics with isotropic Berwald curvature, isotropic mean Berwald curvature, relatively isotropic Landsberg curvature, and relatively isotropic mean Landsberg curvature reduces to the Berwald metric, weakly Berwald metric, Landsberg metric, and weakly Landsberg metric, respectively. We also show that every generalized Kropina change of m th root Finsler metrics with almost vanishing \mathbf{H} -curvature has vanishing \mathbf{H} -curvature.

Key words: Finsler space, Kropina metrics, m th root metrics, Berwald curvature, Landsberg curvature, \mathbf{H} -curvature

1. Introduction

The theory of m th root Finsler metrics was developed by Shimada [10] in 1979 as a generalization of the Riemannian metric, applied to ecology by Antonelli [2] and studied by several authors [3–6, 9, 18]. Tiwari and Kumar [16] studied the Randers change of a Finsler space with m th root metric. Tayebi et al. [11–13] studied the m th root Finsler metric with several non-Riemannian quantities of Berwald curvature, Landsberg curvature, \mathbf{H} -curvature, etc., and they established a necessary and sufficient condition to be projectively flat and locally dually flat for Kropina change of m th root metrics [15]. Xu and Li [17] studied a class of Finsler metrics called special generalized fourth root metrics and established a necessary and sufficient condition for this Finsler metrics to be projectively flat with constant flag curvature $\mathbf{K} = 1$. Recently Tayebi and Shahbazi Nia [14] considered a new class of Finsler metrics, namely Kropina change of generalized m th root metrics, and classified such metrics, which are projectively flat and dually flat.

Let $(M, F) = F^n$ be an n -dimensional Finsler manifold. For a 1-form $\beta(x, y) = b_i(x)y^i$ on M , define a Finsler change as follows:

$$F(x, y) \rightarrow \bar{F}(x, y) = f(F, \beta),$$

where $f(F, \beta)$ is a positively homogeneous function of degree one in F and β . A Finsler change is called a Kropina change if $f(F, \beta) = \frac{F^2}{\beta}$ and a generalized Kropina change if $f(F, \beta) = \frac{F^{k+1}}{\beta^k}$, where k is any positive integer. The purpose of this paper is to investigate the generalized Kropina change of m th root metrics, defined

*Correspondence: gspbhu@gmail.com

2010 AMS Mathematics Subject Classification: 53B40, 53C60

by

$$\overline{F} = \frac{F^{k+1}}{\beta^k}, \tag{1.1}$$

where $F = \sqrt[m]{A}$ is an m th root metric on the manifold M and 1-form $\beta = b_i(x)y^i$ are positive on the whole manifold M .

A geodesic curve $c = c(t)$ of a Finsler metric $F = F(x, y)$ on a smooth manifold M is given by $\ddot{c}^i(t) + 2G^i(c(t), \dot{c}(t)) = 0$, where the local functions $G^i = G^i(x, y)$ are called the spray coefficients given by

$$G^i = \frac{1}{4}g^{il}\{[F^2]_{x^k y^l y^k} - [F^2]_{x^l}\}. \tag{1.2}$$

Let F^n be a Finsler space. For a tangent vector $y \in T_x M$, in the local coordinate system, define $B_y : T_x M \otimes T_x M \otimes T_x M \rightarrow T_x M$ and $E_y : T_x M \otimes T_x M \rightarrow \mathbb{R}$ by $B_y(u, v, w) = B_{jkl}^i(x, y)u^j v^k w^l \frac{\partial}{\partial x^i}$ and $E_y(u, v) = E_{jk}(x, y)u^j v^k$, respectively, where $B_{jkl}^i = [G^i]_{y^j y^k y^l}$, $E_{jk} = \frac{1}{2}B_{mjk}^m = \frac{1}{2} \frac{\partial^2}{\partial y^j \partial y^k} [\frac{\partial G^m}{\partial y^m}]$, $u = u^i \frac{\partial}{\partial x^i}$, $v = v^i \frac{\partial}{\partial x^i}$, and $w = w^i \frac{\partial}{\partial x^i}$. Then $\mathbf{B} = \{B_y | y \in TM_0\}$ and $\mathbf{E} = \{E_y | y \in TM_0\}$ are called the Berwald curvature and mean Berwald curvature, respectively. A Finsler metric is called a Berwald metric and a mean Berwald metric if $\mathbf{B} = 0$ and $\mathbf{E} = 0$, respectively.

For a tangent vector $y \in T_x M$, in the local coordinate system, define $L_y(u, v, w) = L_{ijk}(x, y)u^i v^j w^k$, where $L_{ijk} = -\frac{1}{2}F F_{y^s} [G^s]_{y^i y^j y^k}$, $u = u^i \frac{\partial}{\partial x^i}$, $v = v^i \frac{\partial}{\partial x^i}$, and $w = w^i \frac{\partial}{\partial x^i}$. Then $\mathbf{L} = \{L_y | y \in TM_0\}$ is called the Landsberg curvature. A Finsler metric is called a Landsberg metric if $\mathbf{L} = 0$.

In this paper, we consider the generalized Kropina change of m th root Finsler metrics and, more precisely, we establish the following theorems:

Theorem 1.1 *Let $\overline{F} = \frac{F^{k+1}}{\beta^k}$ be a generalized Kropina change of m th root Finsler metrics on the manifold, where $m \geq 2$. Suppose that \overline{F} has isotropic Berwald curvature (isotropic mean Berwald curvature). Then \overline{F} reduces to a Berwald metric (weakly Berwald metric).*

Theorem 1.2 *Let $\overline{F} = \frac{F^{k+1}}{\beta^k}$ be a generalized Kropina change of m th root Finsler metrics on the manifold, where $m \geq 2$. Suppose that \overline{F} is of relatively isotropic Landsberg curvature. Then \overline{F} reduces to a Landsberg metric.*

Theorem 1.3 *Let $\overline{F} = \frac{F^{k+1}}{\beta^k}$ be a generalized Kropina change of m th root Finsler metrics on the manifold, where $m \geq 2$. Suppose that \overline{F} is of relatively isotropic mean Landsberg curvature. Then \overline{F} reduces to a weakly Landsberg metric.*

In [1], Akbar-Zadeh introduced the non-Riemannian quantity \mathbf{H} , which is obtained from the mean Berwald curvature by covariant horizontal differentiation along geodesics. More precisely, the non-Riemannian quantity $\mathbf{H} = H_{ij} dx^i \otimes dx^j$ is defined by $H_{ij} := E_{ij|s} y^s$. It is proved that for a Finsler manifold of scalar flag curvature \mathbf{K} with dimension $n \geq 3$, $\mathbf{K} = \text{constant}$ if and only if $\mathbf{H} = 0$. The notion of almost vanishing \mathbf{H} -curvature was first introduced by Shen et al. [8]. They investigated that the non-Riemannian quantity \mathbf{H} is closely related to the flag curvature.

In the present paper, we prove the following theorem:

Theorem 1.4 Let $\bar{F} = \frac{F^{k+1}}{\beta^k}$ be a generalized Kropina change of m th root Finsler metrics on the manifold, where $m \geq 2$. Suppose that \bar{F} has almost vanishing \mathbf{H} -curvature. Then $\mathbf{H} = 0$.

Throughout the paper we call the Finsler metric \bar{F} a generalized Kropina change of m th root metric and $\bar{F}^n = (M, \bar{F})$ a generalized Kropina transformed Finsler space. We restrict ourselves for $m \geq 2$ throughout the paper and the quantities corresponding to the generalized Kropina transformed Finsler space \bar{F}^n will be denoted by putting a bar on the top of that quantity.

2. Preliminaries

Let M be an n -dimensional C^∞ -manifold; $T_x M$ denotes the tangent space of M at x . The tangent bundle TM is the union of tangent spaces, $TM := \bigcup_{x \in M} T_x M$. We denote the elements of TM by (x, y) , where $x = (x^i)$ is a point of M and $y \in T_x M$. We denote $TM_0 = TM \setminus \{0\}$.

Definition: A Finsler metric on M is a function $F : TM \rightarrow [0, \infty)$ with the following properties:

- (i) F is C^∞ on TM_0 ,
- (ii) F is positively 1-homogeneous on the fibers of tangent bundle TM , and
- (iii) the Hessian of F^2 with element $g_{ij} = \frac{1}{2} \frac{\partial^2 F^2}{\partial y^i \partial y^j}$ is positive definite on TM_0 .

The pair $F^n = (M, F)$ is called a Finsler space of dimension n . F is called the fundamental function and g_{ij} is called the fundamental tensor of the Finsler space F^n .

The normalized supporting element l_i and angular metric tensor h_{ij} of F are defined respectively as:

$$l_i = \frac{\partial F}{\partial y^i}, \quad h_{ij} = F \frac{\partial^2 F}{\partial y^i \partial y^j}. \tag{2.1}$$

Let F be a Finsler metric defined by $F = \sqrt[m]{A}$, where A is given by $A = a_{i_1 i_2 \dots i_m}(x) y^{i_1} y^{i_2} \dots y^{i_m}$ with $a_{i_1 \dots i_m}$ symmetric in all its indices [10]. Then F is called an m th root Finsler metric. Clearly, A is homogeneous of degree m in y . Let

$$A_i = \frac{\partial A}{\partial y^i}, A_{ij} = \frac{\partial^2 A}{\partial y^i \partial y^j}, A_{x^i} = \frac{\partial A}{\partial x^i}, A_0 = A_{x^i} y^i. \tag{2.2}$$

Then the following holds:

$$g_{ij} = \frac{A^{\frac{2}{m}-2}}{m^2} [mA A_{ij} + (2 - m)A_i A_j],$$

$$y^i A_i = mA, y^i A_{ij} = (m - 1)A_j, y_i = \frac{1}{m} A^{\frac{2}{m}-1} A_i,$$

$$A^{ij} A_{jk} = \delta_k^i, A^{ij} A_i = \frac{1}{m-1} y^j, A_i A_j A^{ij} = \frac{m}{m-1} A.$$

3. Fundamental metric tensors and geodesic sprays of generalized Kropina changed m th root metrics

The differentiation of equation (1.1) with respect to y^i yields the normalized supporting element \bar{l}_i given by

$$\bar{l}_i = \bar{F} \left[\frac{k+1}{mA} A_i - \frac{k}{\beta} b_i \right] \tag{3.1}$$

and the angular metric tensor

$$\bar{h}_{ij} = (k + 1)\bar{F}^2 \left[\frac{1}{mA}A_{ij} + \frac{(k + 1 - m)}{m^2A^2}A_iA_j - \frac{k}{mA\beta}(A_ib_j + A_jb_i) + \frac{k}{\beta^2}b_ib_j \right]. \tag{3.2}$$

The fundamental metric tensor \bar{g}_{ij} of Finsler space \bar{F}^n is given by

$$\bar{g}_{ij} = \bar{h}_{ij} + \bar{l}_i\bar{l}_j.$$

Therefore, using equations (3.1) and (3.2), we obtain metric tensor \bar{g}_{ij} as

$$\begin{aligned} \bar{g}_{ij} = \bar{F}^2 & \left[\frac{k + 1}{mA}A_{ij} + \frac{(k + 1)(2k + 2 - m)}{m^2A^2}A_iA_j - \frac{2k(k + 1)}{mA\beta}(A_ib_j + A_jb_i) \right. \\ & \left. + \frac{k(2k + 1)}{\beta^2}b_ib_j \right]. \end{aligned} \tag{3.3}$$

The above equation can be written as

$$\begin{aligned} \bar{g}_{ij} = \bar{F}^2 & \left[\frac{k + 1}{mA}A_{ij} + \frac{(k + 1)(2k + 2 - (2k + 1)m)}{(2k + 1)m^2A^2}A_iA_j \right. \\ & \left. + \left(\frac{2(k + 1)\sqrt{k}}{\sqrt{2k + 1}mA}A_i - \frac{\sqrt{k(2k + 1)}}{\beta}b_i \right) \left(\frac{2(k + 1)\sqrt{k}}{\sqrt{2k + 1}mA}A_j - \frac{\sqrt{k(2k + 1)}}{\beta}b_j \right) \right]. \end{aligned} \tag{3.4}$$

Let

$$X_{ij} = \frac{k + 1}{mA}A_{ij} + \frac{(k + 1)(2k + 2 - (2k + 1)m)}{(2k + 1)m^2A^2}A_iA_j. \tag{3.5}$$

We know that [7] $A_{ij} = B_{ij} + \delta C_iC_j$, and then

$$A^{ij} = B^{ij} - \frac{\delta C^iC^j}{1 + \delta C^2}, \text{ where } C^2 = B^{ij}C_iC_j \text{ and } C^i = B^{ij}C_j.$$

Using the above results, we obtain

$$X^{ij} = \frac{mA}{k + 1}A^{ij} - \frac{(2k + 2 - (2k + 1)m)}{(k + 1)(m - 1)}y^iy^j. \tag{3.6}$$

Thus, in view of equations (3.4) and (3.5), \bar{g}_{ij} can be written as $\bar{g}_{ij} = \bar{F}^2 [X_{ij} + Y_iY_j]$, where $Y_i =$

$\left(\frac{2(k+1)\sqrt{k}}{\sqrt{2k+1}mA}A_i - \frac{\sqrt{k(2k+1)}}{\beta}b_i \right)$. By direct computation, we have the contravariant metric tensor

$$\bar{g}^{ij} = \frac{1}{\bar{F}^2} [a_0A^{ij} + a_1y^iy^j + a_2B^iB^j + a_3(y^iB^j + y^jB^i)], \tag{3.7}$$

where

$$\begin{aligned}
 B^i &= A^{ij}b_j, B^2 = B^i b_i, a_0 = \frac{mA}{k+1}, a_1 = - \left[\frac{(2k+2) - (2k+1)m}{(k+1)(m-1)} \right. \\
 &\quad \left. + \frac{k(2k+1)m^2\beta^2}{(k+1)(m-1)[\beta^2\{k(2m-3) + (m-1-2k^2)\} + km(m-1)(2k+1)AB^2]} \right], \\
 a_2 &= - \left[\frac{k(2k+1)m^2(m-1)^2A^2}{(k+1)(m-1)[\beta^2\{k(2m-3) + (m-1-2k^2)\} + km(m-1)(2k+1)AB^2]} \right], \\
 a_3 &= - \left[\frac{k(2k+1)m^2(m-1)^2A\beta}{(k+1)(m-1)[\beta^2\{k(2m-3) + (m-1-2k^2)\} + km(m-1)(2k+1)AB^2]} \right]. \tag{3.8}
 \end{aligned}$$

Thus, we obtain the following proposition:

Proposition 3.1 The covariant metric tensor \bar{g}_{ij} and the contravariant metric tensor \bar{g}^{ij} of generalized Kropina transformed Finsler space \bar{F}^n are given by the following:

$$\bar{g}_{ij} = \bar{F}^2 \left[\frac{k+1}{mA} A_{ij} + \frac{(k+1)(2k+2-m)}{m^2A^2} A_i A_j - \frac{2k(k+1)}{mA\beta} (A_i b_j + A_j b_i) + \frac{k(2k+1)}{\beta^2} b_i b_j \right]$$

and

$$\bar{g}^{ij} = \frac{1}{\bar{F}^2} [a_0 A^{ij} + a_1 y^i y^j + a_2 B^i B^j + a_3 (y^i B^j + y^j B^i)],$$

where a_0, a_1, a_2, a_3, B^i , and B^2 are given by equation (3.8).

In view of equation (1.2), to calculate spray coefficients \bar{G}^i of $\bar{F} = \frac{F^{k+1}}{\beta^k}$, we need the following:

$$\left[\bar{F}^2 \right]_{x^t} = \bar{F}^2 \left[\frac{2(k+1)A_{x^t}}{mA} - \frac{2k\beta_{x^t}}{\beta} \right] \tag{3.9}$$

and

$$\begin{aligned}
 \left[\bar{F}^2 \right]_{x^t y^l} y^t &= \bar{F}^2 \left[\frac{2(k+1)(A_l)_0}{mA} + \frac{(k+1)(4(k+1) - 2m)}{m^2A^2} A_l A_0 - \frac{2k}{\beta} (b_l)_0 \right. \\
 &\quad \left. + \frac{2k(2k+1)}{\beta^2} b_l \beta_0 - \frac{4k(k+1)}{mA\beta} (A_l \beta_0 + A_0 b_l) \right]. \tag{3.10}
 \end{aligned}$$

In view of equations (1.2), (3.9), and (3.10) and Proposition (3.1), we have

$$\begin{aligned}
 \bar{G}^i &= \frac{1}{4} [a_0 A^{il} + a_1 y^i y^l + a_2 B^i B^l + a_3 (y^i B^l + y^l B^i)] \\
 &\quad \times \left[\frac{2(k+1)(A_l)_0}{mA} + \frac{(k+1)(4(k+1) - 2m)}{m^2A^2} A_l A_0 - \frac{2k}{\beta} (b_l)_0 + \frac{2k(2k+1)}{\beta^2} b_l \beta_0 \right. \\
 &\quad \left. - \frac{4k(k+1)}{mA\beta} (A_l \beta_0 + A_0 b_l) \right]. \tag{3.11}
 \end{aligned}$$

Proposition 3.2 Let $\bar{F} = \frac{F^{k+1}}{\beta^k}$ be a generalized Kropina change of m th root Finsler metric F on a manifold, with $m \geq 2$. Then the spray coefficients \bar{G}^i of \bar{F}^n are given by equation (3.11).

Remark 3.1 It is remarkable to note that the metric tensors \bar{g}_{ij} and \bar{g}^{ij} of \bar{F}^n are not necessarily rational functions in y but spray coefficients \bar{G}^i of \bar{F}^n are rational functions in y .

4. Berwald curvature

A Finsler metric F is said to be an isotropic Berwald metric if its Berwald curvature is in the following form:

$$B_{jkl}^i = c(F_{y^j y^k} \delta_l^i + F_{y^k y^l} \delta_j^i + F_{y^l y^j} \delta_k^i + F_{y^j y^k y^l} y^i), \tag{4.1}$$

and an isotropic mean Berwald metric if its mean curvature is in the form $E_{ij} = \frac{n+1}{2F} c h_{ij}$, where $c = c(x)$ is a scalar function on M and h_{ij} is the angular metric tensor. A Finsler metric F is said to be weakly Berwald metric if its mean curvature $E_{ij} = 0$.

Proof of Theorem 1.1 Let $\bar{F} = \frac{F^{k+1}}{\beta^k}$ be a generalized Kropina change of the m th root Finsler metric. Suppose that \bar{F} has isotropic Berwald curvature given by equation (4.1). By Remark (3.1), the left-hand side of equation (4.1) is a rational function in y , while the right-hand side is an irrational function. This implies that $c = 0$ and hence \bar{F} reduces to a Berwald metric.

Moreover, $\bar{F} = \frac{F^{k+1}}{\beta^k}$ is of isotropic mean Berwald curvature, that is,

$$\bar{E}_{ij} = \frac{n+1}{2\bar{F}} c \bar{h}_{ij}, \tag{4.2}$$

where $c = c(x)$ is a scalar function on M and \bar{h}_{ij} is the angular metric.

By putting the angular metric \bar{h}_{ij} given by equation (3.2) in (4.2), we have

$$\bar{E}_{ij} = \frac{(n+1)(k+1)c}{2} \bar{F} \left[\frac{1}{mA} A_{ij} + \frac{(k+1-m)}{m^2 A^2} A_i A_j - \frac{k}{mA\beta} (A_i b_j + A_j b_i) + \frac{k}{\beta^2} b_i b_j \right]. \tag{4.3}$$

The left side of equation (4.3) is a rational function in y while its right side is an irrational function in y . Thus, equation (4.3) implies either $c = 0$ or

$$\left[\frac{1}{mA} A_{ij} + \frac{(k+1-m)}{m^2 A^2} A_i A_j - \frac{k}{mA\beta} (A_i b_j + A_j b_i) + \frac{k}{\beta^2} b_i b_j \right] = 0. \tag{4.4}$$

If equation (4.4) holds, then $\bar{h}_{ij} = 0$, which is not possible. Hence, $c = 0$ and $\bar{E}_{ij} = 0$. This completes the proof of Theorem 1.1.

5. Landsberg curvature

For a nonzero vector $y \in T_p M$, the Cartan torsion $C_y = C_{ijl} dx^i \otimes dx^j \otimes dx^l : T_p M \otimes T_p M \otimes T_p M \rightarrow \mathbb{R}$ is defined by $C_{ijl} = \frac{1}{2} \frac{\partial g_{ij}}{\partial y^l}$.

Proof of Theorem 1.2 Let $\bar{F} = F^{k+1}/\beta^k$ be a relatively isotropic Landsberg metric, that is,

$$\bar{L}_{ijl} = c \bar{F} \bar{C}_{ijl}, \tag{5.1}$$

where $c = c(x)$ is a scalar function on M . The Cartan torsion \bar{C}_{ijl} of \bar{F} is given by the following:

$$\begin{aligned} \bar{C}_{ijl} = & \frac{(k+1)\bar{F}^2}{2} \left[\frac{1}{mA} A_{ijl} + \frac{2(2k+2-m)(k+1-m)}{m^3 A^3} A_i A_j A_l \right. \\ & + \frac{(2k+2-m)}{m^2 A^2} \sum_{(i,j,l)} A_{ij} A_l - \frac{2k}{mA\beta} \sum_{(i,j,l)} A_{ij} b_l - \frac{2k(2k+2-m)}{m^2 A^2 \beta} \sum_{(i,j,l)} A_i A_j b_l \\ & \left. + \frac{2k(2k+1)}{mA\beta^2} \sum_{(i,j,l)} A_i b_j b_l - \frac{2k(2k+1)}{\beta^3} b_i b_j b_l \right], \end{aligned} \tag{5.2}$$

where $\sum_{(i,j,l)}$ denotes the cyclic interchange of suffices i, j, l and summation, for instance $\sum_{(i,j,l)} A_{ij} A_l = A_{ij} A_l + A_{jl} A_i + A_{li} A_j$. Also, we know that

$$\bar{L}_{ijl} = -\frac{1}{2} \bar{F} \bar{F}_{y^s} \bar{G}_{y^i y^j y^l}. \tag{5.3}$$

In our case,

$$\bar{L}_{ijl} = -\frac{\bar{F}^2}{2} \left[\frac{(k+1)}{mA} A_s - \frac{k}{\beta} b_s \right] \bar{G}_{y^i y^j y^l}. \tag{5.4}$$

Using equations (5.1), (5.2), and (5.4), we obtain

$$\begin{aligned} \left(\frac{2A_s}{mA} - \frac{b_s}{\beta} \right) G_{y^i y^j y^l}^s = & -c\bar{F} \left[\frac{1}{mA} A_{ijl} + \frac{2(2k+2-m)(k+1-m)}{m^3 A^3} A_i A_j A_l \right. \\ & + \frac{(2k+2-m)}{m^2 A^2} \sum_{(i,j,l)} A_{ij} A_l - \frac{2k}{mA\beta} \sum_{(i,j,l)} A_{ij} b_l - \frac{2k(2k+2-m)}{m^2 A^2 \beta} \sum_{(i,j,l)} A_i A_j b_l \\ & \left. + \frac{2k(2k+1)}{mA\beta^2} \sum_{(i,j,l)} A_i b_j b_l - \frac{2k(2k+1)}{\beta^3} b_i b_j b_l \right]. \end{aligned} \tag{5.5}$$

The left side of equation (5.5) is a rational function in y while its right side is an irrational function in y . Therefore, either $c = 0$ or A satisfies the following PDEs:

$$\begin{aligned} & \left[\frac{1}{mA} A_{ijl} + \frac{2(2k+2-m)(k+1-m)}{m^3 A^3} A_i A_j A_l + \frac{(2k+2-m)}{m^2 A^2} \sum_{(i,j,l)} A_{ij} A_l \right. \\ & - \frac{2k}{mA\beta} \sum_{(i,j,l)} A_{ij} b_l - \frac{2k(2k+2-m)}{m^2 A^2 \beta} \sum_{(i,j,l)} A_i A_j b_l + \frac{2k(2k+1)}{mA\beta^2} \sum_{(i,j,l)} A_i b_j b_l \\ & \left. - \frac{2k(2k+1)}{\beta^3} b_i b_j b_l \right] = 0. \end{aligned}$$

If the above equation holds, then $\bar{C}_{ijl}=0$. Hence, \bar{F} is a Riemannian metric, which contradicts our assumption. Therefore, $c = 0$. This completes the proof of Theorem (1.2).

The mean Cartan torsion $I_y = I_i(x, y)dx^i : T_pM \rightarrow \mathbb{R}$ is defined by

$$I_i = g^{jl}C_{ijl}. \tag{5.6}$$

Thus, the mean Cartan torsion of \bar{F} is given by

$$\begin{aligned} \bar{I}_i &= \frac{1}{2} [a_0A^{jl} + a_1y^jy^l + a_2B^jB^l + a_3(y^jB^l + y^lB^j)] \times \left[\frac{1}{mA}A_{ijl} \right. \\ &+ \frac{2(2k+2-m)(k+1-m)}{m^3A^3}A_iA_jA_l + \frac{(2k+2-m)}{m^2A^2} \sum_{(i,j,l)} A_{ij}A_l - \frac{2k}{mA\beta} \sum_{(i,j,l)} A_{ij}b_l \\ &\left. - \frac{2k(2k+2-m)}{m^2A^2\beta} \sum_{(i,j,l)} A_iA_jb_l + \frac{2k(2k+1)}{mA\beta^2} \sum_{(i,j,l)} A_ib_jb_l - \frac{2k(2k+1)}{\beta^3}b_ib_jb_l \right]. \end{aligned} \tag{5.7}$$

Lemma 5.1 *Let \bar{F}^n be a generalized Kropina change of an m th root Finsler manifold. Then mean Cartan torsions \bar{I}_i of \bar{F}^n are rational functions in y .*

The horizontal covariant derivative of \mathbf{I} along a vector $u \in T_xM$ gives rise to the mean Landsberg curvature $J_y(u) := J_i(y)u^i$, where $J_i = I_{i|s}y^s$. A Finsler metric with $\mathbf{J} = 0$ is called a weakly Landsberg metric.

The mean Landsberg curvature of \bar{F} is given by

$$\bar{J}_i = \bar{g}^{jl}\bar{L}_{ijl}. \tag{5.8}$$

In view of equations (3.7), (5.4), and (5.8), we have

$$\bar{J}_i = -\frac{1}{2} [a_0A^{jl} + a_1y^jy^l + a_2B^jB^l + a_3(y^jB^l + y^lB^j)] \left[\frac{(k+1)}{mA}A_s - \frac{k}{\beta}b_s \right] \bar{G}_{y^iy^jy^l}^s. \tag{5.9}$$

Lemma 5.2 *Let \bar{F}^n be a generalized Kropina change of an m th root Finsler manifold. Then mean Landsberg curvatures \bar{J}_i of \bar{F}^n are rational functions in y .*

In view of equations (5.1), (5.6), and (5.8), we have

$$\bar{J}_i = c\bar{F}\bar{I}_i. \tag{5.10}$$

Proof of Theorem 1.3 Using Lemma 5.1 and Lemma 5.2, the left side of equation (5.10) is a rational function in y while its right side is an irrational function in y . Thus, if equation (5.10) holds, then either $c = 0$ or $\bar{I}_i = 0$. If $\bar{I}_i = 0$, \bar{F} is a Riemannian metric, which contradicts our assumption. Therefore, $c = 0$. This completes the proof of Theorem 1.3.

6. H-curvature

A Finsler metric on an n -dimensional manifold M is said to be of almost vanishing **H**-curvature if $H_{ij} = \frac{n+1}{2F}\theta h_{ij}$, for some 1-form θ on M , where h_{ij} is the angular metric.

Proof of Theorem 1.4 Let $\bar{F} = \frac{F^{k+1}}{\beta^k}$ be of almost vanishing $\bar{\mathbf{H}}$ -curvature, that is,

$$\bar{H}_{ij} = \frac{n+1}{2\bar{F}}\theta\bar{h}_{ij}, \quad (6.1)$$

for some 1-form θ on M , where \bar{h}_{ij} is the angular metric.

Plugging the angular metric \bar{h}_{ij} given by equation (3.2) into (6.1), we have

$$\bar{H}_{ij} = \frac{(n+1)(k+1)}{2}\theta\bar{F} \left[\frac{1}{mA}A_{ij} + \frac{(k+1-m)}{m^2A^2}A_iA_j - \frac{k}{mA\beta}(A_ib_j + A_jb_i) + \frac{k}{\beta^2}b_ib_j \right]. \quad (6.2)$$

The left side of equation (6.2) is a rational function in y while its right side is an irrational function in y . Thus, equation (6.2) implies either $\theta = 0$ or

$$\left[\frac{1}{mA}A_{ij} + \frac{(k+1-m)}{m^2A^2}A_iA_j - \frac{k}{mA\beta}(A_ib_j + A_jb_i) + \frac{k}{\beta^2}b_ib_j \right] = 0. \quad (6.3)$$

If equation (6.3) holds, then $\bar{h}_{ij} = 0$, which is not possible. Hence, $\theta = 0$ and $\bar{H}_{ij} = 0$. This completes the proof of Theorem (1.4).

References

- [1] Akbar-Zadeh H. Sur les espaces de Finsler a courbures sectionnelles constantes. Bull Acad Roy Belg Cl Sci 1988; 84: 281-322 (in French).
- [2] Antonelli PL, Ingarden RS, Matsumoto M. The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology. Dordrecht, the Netherlands: Kluwer Academic Publishers, 1993.
- [3] Asanov GS. Finslerian Extension of General Relativity. Dordrecht, the Netherlands: Reidel, 1984.
- [4] Balan V. Notable submanifolds in Berwald-Moór spaces. In: BSG Proceedings 17. Bucharest, Romania: Geometry Balkan Press, 2010, pp. 21-30.
- [5] Balan V, Brinzei N. Einstein equations for (h, v) -Berwald-Moór relativistic models. Balkan J Geom Appl 2006; 11: 20-26.
- [6] Li B, Shen Z. Projectively flat fourth root Finsler metrics. Can Math Bulletin 2012; 55: 138-145.
- [7] Shen Z, Chern SS. Riemann-Finsler Geometry. Nankai Tracts in Mathematics. Singapore: World Scientific, 2004.
- [8] Shen Z, Najafi B, Tayebi A. On Finsler metrics of scalar curvature with some non-Riemannian curvature properties. Geom Dedicata 2008; 131: 87-97.
- [9] Shibata C. On invariant tensors of β -changes of Finsler metrics. J Math Kyoto Univ 1984; 24: 163-188.
- [10] Shimada H. On Finsler spaces with the metric $L = \sqrt[n]{a_{i_1 \dots i_m} y^{i_1} \dots y^{i_m}}$. Tensor N S 1979; 33: 365-372.
- [11] Tayebi A, Najafi B. On m -th root Finsler metrics. J Geom Phys 2011; 61: 1479-1484.
- [12] Tayebi A, Najafi B. On m -th root metrics with special curvature properties. C R Acad Sci Paris Ser I 2011; 349: 691-693.

- [13] Tayebi A, Nankali A, Peyghan E. Some properties of m -th root Finsler metrics. J Contemp Math Anal 2014; 49: 157-166.
- [14] Tayebi A, Shahbazi Nia M. A new class of projectively flat Finsler metrics with constant flag curvature $\mathbf{K} = 1$. Diff Geom Appl 2015; 41: 123-133.
- [15] Tayebi A, Tabatabaeifar T, Peyghan E. On Kropina-change of m -th root Finsler metrics. Ukrainian J Math 2011; 66: 140-144.
- [16] Tiwari B, Kumar M. On Randers change of a Finsler space with m -th root metric. Int J Geom Methods M 2014; 11: 1-13.
- [17] Xu B, Li B. On a class of projectively flat Finsler metrics with flag curvature $\mathbf{K} = 1$. Diff Geom Appl 2013; 31: 524-532.
- [18] Yu Y, You Y. On Einstein m -th root metrics. Diff Geom Appl 2010; 28: 290-294.