

1-1-2017

A note on locally graded minimal non-metahamiltonian groups

SEVGİ ATLIHAN

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

ATLIHAN, SEVGİ (2017) "A note on locally graded minimal non-metahamiltonian groups," *Turkish Journal of Mathematics*: Vol. 41: No. 5, Article 5. <https://doi.org/10.3906/mat-1503-31>

Available at: <https://dctubitak.researchcommons.org/math/vol41/iss5/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

A note on locally graded minimal non-metahamiltonian groups

Sevgi ATLIHAN*

Department of Mathematics, Faculty of Education, Gazi University, Teknikokullar, Ankara, Turkey

Received: 09.03.2015

Accepted/Published Online: 25.10.2016

Final Version: 28.09.2017

Abstract: We prove that a nonperfect locally graded minimal non-metahamiltonian group G is a soluble group with derived length of at most 4. On the other hand, if G is perfect, then $G/\Phi(G)$ is isomorphic to A_5 , where $\Phi(G)$ is the Frattini subgroup of G and A_5 is the alternating group. Moreover, we show that under some conditions, if G is a p -group, then G is metabelian, where p is a prime integer.

Key words: Locally graded minimal non-metahamiltonian, soluble, metabelian, d -maximal

1. Introduction

A group G is called *metahamiltonian* if every nonabelian subgroup of G is normal. Metahamiltonian groups were introduced and investigated in a series of papers by Romalis and Sesekin (see [17–19]); they proved in particular that the commutator subgroup of any group with such property is finite with prime-power order. In [6, Theorem 3.4], De Mari and De Giovanni proved that a locally graded metahamiltonian group is soluble with derived length of at most 3. In [14, Theorem 1], Mahnev proved that for prime integer p , the commutator subgroup of a finite metahamiltonian p -group is abelian. For more details and the current knowledge related to these kinds of groups, we recommend the book by Kuzennyi and Semko [12].

A group G is called *minimal non-metahamiltonian* if every proper subgroup of G is a metahamiltonian but G itself is not. In [5, Lemma 4.2], De Falco et al. proved that a locally graded group with such property is finite. In [5], the authors also gave the alternating group A_5 as an example that shows that there exist finite insoluble minimal non-metahamiltonian groups. Further, A_5 is perfect simple with every proper subgroup metabelian. Corollary 2 shows that a simple locally graded minimal non-metahamiltonian group is isomorphic to A_5 .

Therefore, every minimal non-metahamiltonian group may not be soluble. It is natural to ask: “Is there a soluble minimal non-metahamiltonian group?” The answer to this question is yes. It is well known that the general linear group $GL(2, 3)$ is a soluble group with derived length 4. On the other hand, this group is also a minimal non-metahamiltonian group: indeed, its subgroups of order sixteen are nonabelian and not normal. Also, one can see that every proper subgroup of $GL(2, 3)$ is metahamiltonian. Theorem 1(ii) shows that a nonperfect locally graded minimal non-metahamiltonian group is soluble with derived length of at most 4.

The second crucial question is “Does there exist an insoluble nonsimple minimal non-metahamiltonian group?” The answer to this question is yes. If we consider the unique nonsplitting central extension of a group

*Correspondence: asevgi@gazi.edu.tr

2010 AMS Mathematics Subject Classification: 20M19.

of order 2, which is the Schur multiplier of A_5 , by the group A_5 itself, this group is perfect, but not simple. It is easy to show that it is a minimal non-metahamiltonian group with its only proper nontrivial normal subgroup being its center, and also, the Frattini factor of it is isomorphic to A_5 . Theorem 1(i) shows that the Frattini factor of a perfect nonsimple locally graded minimal non-metahamiltonian group is isomorphic to A_5 .

The generalized dihedral group for the direct product of Z_4 and Z_4 is known as a metabelian 2-group, but it is also a minimal non-metahamiltonian group: indeed, it has nonnormal nonabelian subgroups of order eight. One can also see that every proper subgroup is metahamiltonian.

Another question is "Does there exist a non-metabelian locally graded minimal non-metahamiltonian 2-group?" The technique used by O'Brein in [15] shows that there is no non-metabelian minimal non-metahamiltonian group of order dividing 2^9 , which means that there is no such a group of order dividing 2^9 . Furthermore, we could not find any example that shows that there exists a metabelian (or not) minimal non-metahamiltonian group of order at least 2^{10} in the literature. As a consequence, we have no answer to this question, neither yes nor no. In this study, for any prime integer p , we show that a locally graded minimal non-metahamiltonian p -group for which every proper subgroup is metacyclic is metabelian (see Theorem 3). We also prove for any prime integer p that a d -maximal locally graded minimal non-metahamiltonian p -group is metabelian (see Theorem 4). A group G is said to be d -maximal if $d(H) < d(G)$ for any proper subgroup H of G , where $d(G)$ denotes the cardinality of a minimal generating set of G .

Therefore, we are going to deal with the conditions that make the answer to this question be no.

Recall that a group G is called *locally graded* if every finitely generated nontrivial subgroup of G has a proper subgroup of finite index. Locally graded groups form a wide class of generalized soluble groups, containing in particular all locally (soluble-by-finite) groups.

Most of our notation is standard and can be found in [16].

2. The main conclusions

Now we are ready to give and prove Theorem 1, Theorem 2, and Theorem 3.

Theorem 1 *Let G be a locally graded minimal non-metahamiltonian group.*

- (i) *If G is perfect, then the Frattini factor group of G , i.e. $G/\Phi(G)$, is isomorphic to A_5 .*
- (ii) *If G is not perfect, then G is a soluble group with derived length of at most 4.*

Proof

- (i) Let G be perfect, i.e. $G' = G$.

Then $\bar{G} = G/\Phi(G)$ is a simple nonabelian group by [6, Theorem 3.4] and [9, Lemma 3.2], which is of course minimal non-metahamiltonian. Since \bar{G} is a finite nonabelian simple group in which every proper subgroup is soluble with derived length of at most 3 by [6, Theorem 3.4], then to complete the proof, it is enough to show that $PSL(3, 3)$, $Sz(q)$, where $q = 2^p$ for any odd prime integer p , $PSL(2, p)$, where $p > 3$ is any prime integer such that $p^2 + 1 \equiv 0 \pmod{5}$, $PSL(2, 2^p)$, and $PSL(2, 3^p)$, where p is any odd integer, are not minimal non-metahamiltonian groups by [22, Corollary 1] and [8, Proposition 2.2].

Since $PSL(3, 3)$ contains a subgroup of derived length 5 (see [8, pp. 4-5]), then $PSL(3, 3)$ is not a minimal non-metahamiltonian group.

By [20, Theorem 9], $Sz(q)$ contains a non-metabelian Frobenius group F of order $q^2(q-1)$ and $Sz(q)$ has only one abelian subgroup of order dividing $q^2(q-1)$ that is a cyclic group of order $q-1$. It follows that the commutator subgroup of F has not prime-power order. Therefore, $Sz(q)$ is not a minimal non-metahamiltonian group by [6, Theorem 3.4].

$PSL(2, p)$, where $p > 3$ is any prime integer $p^2 + 1 \equiv 0 \pmod{5}$ and $p^2 - 1 \equiv 0 \pmod{16}$, has a proper subgroup isomorphic to S_4 by [21, Theorem 6.25 and Theorem 6.26]. Therefore, $PSL(2, p)$ is not a minimal non-metahamiltonian group by S_4 with a nonnormal subgroup of order 8 to isomorphic D_8 .

$PSL(2, p)$, where $p > 3$ is any prime integer $p^2 + 1 \equiv 0 \pmod{5}$ and $p^2 - 1 \not\equiv 0 \pmod{16}$, $PSL(2, 2^p)$, and $PSL(2, 3^p)$, where p is any odd integer, contain a maximal subgroup that is dihedral and has a proper nonabelian subgroup that is not normal (see [7]), and so these groups cannot be minimal non-metahamiltonian groups.

- (ii) If G is not perfect, i.e. $G' \neq G$, then G' is metahamiltonian by the hypothesis and so G is a soluble group with derived length of at most 4 by [6, Theorem 3.4].

□

Corollary *Let G be a simple locally graded minimal non-metahamiltonian group. Then G is isomorphic to A_5 .*

Theorem 2 *Let G be a locally graded minimal non-metahamiltonian p -group, for a prime integer p . If every proper subgroup of G is metacyclic, then G is metabelian.*

Proof First, let $p = 2$. Assume that G is not metacyclic. Since G is a minimal non-metahamiltonian group by the hypothesis, the order of G must be 2^5 by [2, Theorem 66.1]. Also, every subgroup of order 8 is abelian by the proof of [2, Theorem 66.1]. However, since G is not metahamiltonian, that is a contradiction. Therefore, G is metacyclic and so it is metabelian.

Now let $p > 2$. In the case of $d(G) \geq 4$, since every proper subgroup is metacyclic by the hypothesis, then G is metabelian by [1, Theorem 4]. Let $d(G) \leq 3$. Since every proper subgroup is metacyclic by the hypothesis, then every proper subgroup can be generated by two elements by [12, Theorem 2.3.1]. If $d(G) = 2$, then G is metabelian by [3, Theorem 4]. If $d(G) = 3$, since every proper subgroup of G can be generated by two elements, G is d -maximal and therefore G is metabelian by $p > 2$ and by [13, Theorem]. This completes the proof of this theorem. □

Theorem 3 *Let G be a locally graded minimal non-metahamiltonian p -group for a prime integer p . If G is d -maximal, then G is metabelian.*

Proof Suppose that G is d -maximal.

For $p > 2$, since G is a finite p -group, then G is metabelian by [13, Theorem].

Now let be $p = 2$. Assume that G is not metabelian. Then G' cannot be generated by a commutative set. Hence, there exist elements a, b, c , and d of G such that $[[a, b], [c, d]] \neq 1$, which implies that $\langle a, b, c, d \rangle$ is a non-metabelian finite p -group, and then $G = \langle a, b, c, d \rangle$ by [14, Theorem 1]. It follows that $d(G) \leq 4$. If $d(G) \leq 3$, since G is a d -maximal 2-group, then it follows by [10, Lemma A2] and [4, Theorem 3.2] that the nilpotent class of G is at most 2, which implies a contradiction by G not being metabelian. Hence, we

obtain $d(G) = 4$. It follows that the nilpotent class of G is at most 2 by [11, Proposition 5.1], which implies a contradiction again by G not being metabelian. This completes the proof. \square

We would like to make a note here that the generalized dihedral group for the direct product of Z_4 and Z_4 shows that the converse of the statement in Theorem 3 is not true in general.

References

- [1] Alperin JL. On a special class of regular p -group. T Am Math Soc 1963; 106: 77-99.
- [2] Berkovich Y, Janko Z. Groups of Prime Power Order. Berlin, Germany: Walter de Gruyter, 2008.
- [3] Blackburn N. On prime-power groups in which the derived group has two generators. P Camb Philos Soc 1957; 53: 19-27.
- [4] Blackburn N. Generalizations of certain elementary theorems on p -groups. P Lond Math Soc 1961; 11: 1-22.
- [5] De Falco M, de Giovanni F, Musella C. Groups whose finite homomorphic images are metahamiltonian. Comm Algebra 2009; 37: 2468-2476.
- [6] De Mari F, de Giovanni F. Groups with few normalizer subgroups. Irish Math Soc Bull 2005; 56: 103-113.
- [7] Dickson LE. Linear Groups, with an Exposition of the Galois Field Theory. New York, NY, USA: Dover Pub. Co., 1958.
- [8] Ersoy K, Tortora A, Tota M. On groups with all subgroups subnormal or soluble of bounded derived length. Glasgow Math J 2014; 56: 221-227.
- [9] Guralnick RM, Tiep PH. Lifting in Frattini covers and a characterization of finite solvable groups. J Reine Angew Math 2015; 708: 49-72.
- [10] Kahn B. The total Stiefel-Whitney classes of a regular representation. J Algebra 1991; 144: 214-247.
- [11] Kahn B, Minh PA. A description of the second mod p chomology group of a p -group. J Pure Appl Algebra 1998; 132: 73-104.
- [12] Kuzennyi NF, Semko NN. Metahamiltonian Groups and Their Generalizations. Kiev, Ukraine: National Academy of Sciences, 1996 (in Russian).
- [13] Laffey TJ. The minimum number of generators of a finite p -group. B Lond Math Soc 1973; 5: 288-290.
- [14] Mahnev AA. Finite metahamiltonian groups. Ural Gos Univ Mat Zap 1976; 10: 60-75.
- [15] O'Brien EA. The p -group generation algorithm. J Symbolic Comput 1990; 9: 677-698.
- [16] Robinson DJS. Finiteness Conditions and Generalized Soluble Groups. Berlin, Germany: Springer, 1972.
- [17] Romalis GM, Sesekin NF. Metahamiltonian groups I. Ural Gos Univ Mat Zap 1966; 5: 101-106.
- [18] Romalis GM, Sesekin NF. Metahamiltonian groups II. Ural Gos Univ Mat Zap 1968; 6: 50-52.
- [19] Romalis GM, Sesekin NF. Metahamiltonian groups III. Ural Gos Univ Mat Zap 1970; 7: 195-199.
- [20] Suzuki M. On a class of doubly transitive groups. Ann Math 1962; 75: 105-145.
- [21] Suzuki M. Group Theory I. Berlin, Germany: Springer-Verlag, 1982.
- [22] Thompson JG. Nonsolvable finite groups all of whose local subgroups are solvable. B Am Math Soc 1968; 74: 383-437.