

1-1-2017

## New inequalities of Opial type for conformable fractional integrals

MEHMET ZEKİ SARIKAYA

HÜSEYİN BUDAK

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

---

### Recommended Citation

SARIKAYA, MEHMET ZEKİ and BUDAK, HÜSEYİN (2017) "New inequalities of Opial type for conformable fractional integrals," *Turkish Journal of Mathematics*: Vol. 41: No. 5, Article 8. <https://doi.org/10.3906/mat-1606-91>

Available at: <https://dctubitak.researchcommons.org/math/vol41/iss5/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

## New inequalities of Opial type for conformable fractional integrals

Mehmet Zeki SARIKAYA, Hüseyin BUDAK\*

Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey

Received: 20.06.2016

Accepted/Published Online: 07.11.2016

Final Version: 28.09.2017

**Abstract:** In this paper, some Opial-type inequalities for conformable fractional integrals are obtained using the remainder function of Taylor's theorem for conformable integrals.

**Key words:** Opial inequality, Hölder's inequality, conformable fractional integrals

### 1. Introduction

In 1960, Opial established the following interesting integral inequality [10]:

**Theorem 1** *Let  $x(t) \in C^{(1)}[0, h]$  be such that  $x(0) = x(h) = 0$ , and  $x(t) > 0$  in  $(0, h)$ . Then the following inequality holds:*

$$\int_0^h |x(t)x'(t)| dt \leq \frac{h}{4} \int_0^h (x'(t))^2 dt \quad (1.1)$$

*The constant  $h/4$  is the best possible.*

Opial's inequality and its generalizations, extensions, and discretizations play a fundamental role in establishing the existence and uniqueness of initial and boundary value problems for ordinary and partial differential equations as well as difference equations. Over the last 20 years a large number of papers have appeared in the literature that deals with the simple proofs, various generalizations, and discrete analogues of Opial's inequality and its generalizations; see [2,4,5,11–14,16,17].

The purpose of this paper is to establish some Opial-type inequalities for conformable integrals. The structure of this paper is as follows. In Section 2, we give the definitions of conformable derivatives and conformable integrals and introduce several useful notations for conformable integrals used in our main results. In Section 3, the main result is presented. Using the remainder function of Taylor's theorem for conformable integrals, we establish several Opial-type inequalities.

### 2. Definitions and properties of conformable fractional derivatives and integrals

The following definitions and theorems with respect to conformable fractional derivatives and integrals were referred to (see [1], [3], [6]–[9]).

\*Correspondence: [hsyn.budak@gmail.com](mailto:hsyn.budak@gmail.com)

2010 AMS Mathematics Subject Classification: 26D15, 26A51; Secondary 26A33, 26A42.

**Definition 1 (Conformable fractional derivative)** Given a function  $f : [0, \infty) \rightarrow \mathbb{R}$ . Then the “conformable fractional derivative” of  $f$  of order  $\alpha$  is defined by

$$D_\alpha(f)(t) = \lim_{\epsilon \rightarrow 0} \frac{f(t + \epsilon t^{1-\alpha}) - f(t)}{\epsilon} \quad (2.1)$$

for all  $t > 0$ ,  $\alpha \in (0, 1)$ . If  $f$  is  $\alpha$ -differentiable in some  $(0, a)$ ,  $\alpha > 0$ ,  $\lim_{t \rightarrow 0^+} f^{(\alpha)}(t)$  exist, then define

$$f^{(\alpha)}(0) = \lim_{t \rightarrow 0^+} f^{(\alpha)}(t). \quad (2.2)$$

We can write  $f^{(\alpha)}(t)$  for  $D_\alpha(f)(t)$  to denote the conformable fractional derivatives of  $f$  of order  $\alpha$ . In addition, if the conformable fractional derivative of  $f$  of order  $\alpha$  exists, then we simply say  $f$  is  $\alpha$ -differentiable.

**Theorem 2** Let  $\alpha \in (0, 1]$  and  $f, g$  be  $\alpha$ -differentiable at a point  $t > 0$ . Then

i.  $D_\alpha(af + bg) = aD_\alpha(f) + bD_\alpha(g)$ , for all  $a, b \in \mathbb{R}$ ,

ii.  $D_\alpha(\lambda) = 0$ , for all constant functions  $f(t) = \lambda$ ,

iii.  $D_\alpha(fg) = fD_\alpha(g) + gD_\alpha(f)$ ,

iv.  $D_\alpha\left(\frac{f}{g}\right) = \frac{fD_\alpha(g) - gD_\alpha(f)}{g^2}$ .

If  $f$  is differentiable, then

$$D_\alpha(f)(t) = t^{1-\alpha} \frac{df}{dt}(t). \quad (2.3)$$

**Definition 2 (Conformable fractional integral)** Let  $\alpha \in (0, 1]$  and  $0 \leq a < b$ . A function  $f : [a, b] \rightarrow \mathbb{R}$  is  $\alpha$ -fractional integrable on  $[a, b]$  if the integral

$$\int_a^b f(x) d_\alpha x := \int_a^b f(x) x^{\alpha-1} dx \quad (2.4)$$

exists and is finite. All  $\alpha$ -fractional integrable on  $[a, b]$  is indicated by  $L_\alpha^1([a, b])$ .

**Remark 1**

$$I_\alpha^a(f)(t) = I_1^a(t^{\alpha-1}f) = \int_a^t \frac{f(x)}{x^{1-\alpha}} dx,$$

where the integral is the usual Riemann improper integral, and  $\alpha \in (0, 1]$ .

**Theorem 3** Let  $f : (a, b) \rightarrow \mathbb{R}$  be differentiable and  $0 < \alpha \leq 1$ . Then, for all  $t > a$  we have

$$I_\alpha^a D_\alpha^a f(t) = f(t) - f(a). \quad (2.5)$$

**Theorem 4 (Integration by parts)** Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be two functions such that  $fg$  is differentiable. Then

$$\int_a^b f(x) D_\alpha^a(g)(x) d_\alpha x = fg|_a^b - \int_a^b g(x) D_\alpha^a(f)(x) d_\alpha x. \quad (2.6)$$

**Theorem 5** Assume that  $f : [a, \infty) \rightarrow \mathbb{R}$  such that  $f^{(n)}(t)$  is continuous and  $\alpha \in (n, n + 1]$ . Then, for all  $t > a$  we have

$$D_\alpha^a I_\alpha^a f(t) = f(t).$$

We can give Hölder's inequality in conformable integral as follows:

**Lemma 1** Let  $f, g \in C[a, b]$ ,  $p, q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ , then

$$\int_a^b |f(x)g(x)| d_\alpha x \leq \left( \int_a^b |f(x)|^p d_\alpha x \right)^{\frac{1}{p}} \left( \int_a^b |g(x)|^q d_\alpha x \right)^{\frac{1}{q}}.$$

**Remark 2** If we take  $p = q = 2$  in Lemma 1, then we have the Cauchy-Schwarz inequality for conformable integrals.

**Theorem 6 (Taylor's Formula)** [3] Let  $\alpha \in (0, 1]$  and  $n \in \mathbb{N}$ . Suppose  $f$  is  $n + 1$  times  $\alpha$ -fractional differentiable on  $[0, \infty)$ , and  $s, t \in [0, \infty)$ . Then we have

$$f(t) = \sum_{k=0}^n \frac{1}{k!} \left( \frac{t^\alpha - s^\alpha}{\alpha} \right)^k D_\alpha^k f(s) + \frac{1}{n!} \int_s^t \left( \frac{t^\alpha - \tau^\alpha}{\alpha} \right)^n D_\alpha^{n+1} f(\tau) d_\alpha \tau.$$

Using Taylor's theorem, we define the remainder function by

$$R_{-1, f}(\cdot, s) := f(s),$$

and for  $n > -1$ ,

$$\begin{aligned} R_{n, f}(t, s) & : = f(s) - \sum_{k=0}^n \frac{1}{k!} \left( \frac{t^\alpha - s^\alpha}{\alpha} \right)^k D_\alpha^k f(s) \\ & = \frac{1}{n!} \int_s^t \left( \frac{t^\alpha - \tau^\alpha}{\alpha} \right)^n D_\alpha^{n+1} f(\tau) d_\alpha \tau. \end{aligned} \quad (2.7)$$

Opial's inequality can be represented for conformable fractional integral forms as follows [15]:

**Theorem 7** Let  $\alpha \in (0, 1]$  and  $u$  be an  $\alpha$ -fractional differentiable function on  $(0, h)$  with  $u(0) = u(h) = 0$ . Then the following inequality for conformable fractional integrals holds:

$$\int_0^h |u(t) D_\alpha(u)(t)| d_\alpha t \leq \frac{h^\alpha}{4\alpha} \int_0^h |D_\alpha(u)(t)|^2 d_\alpha t. \quad (2.8)$$

Now we present the main results,

### 3. Opial-type inequalities for conformable fractional integrals

Let  $\alpha \in (0, 1]$ . In the following we adapt to the  $\alpha$ -fractional setting some results from [2] by applying the fractional Opial inequality.

**Theorem 8** *Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n + 1$  times  $\alpha$ -fractional differentiable function,  $p, q > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $t \geq x_0$ ,  $t, x_0 \in [a, b]$ . Then we have the following inequality:*

$$\begin{aligned} & \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\ & \leq \frac{(t^\alpha - x_0^\alpha)^{n+2/p}}{\alpha^{n+2/p} 2^{\frac{1}{q}} n! [(np+1)(np+2)]^{1/p}} \left( \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau \right)^{\frac{2}{q}}. \end{aligned} \quad (3.1)$$

**Proof** From (2.7), we have

$$R_{n,f}(x_0, t) = \frac{1}{n!} \int_{x_0}^t \left( \frac{t^\alpha - \tau^\alpha}{\alpha} \right)^n D_\alpha^{n+1} f(\tau) d_\alpha \tau, \quad x_0, t \in [a, b].$$

By using Hölder's inequality for conformable integrals, it follows that

$$\begin{aligned} |R_{n,f}(x_0, t)| & \leq \frac{1}{\alpha^n n!} \int_{x_0}^t (t^\alpha - \tau^\alpha)^n |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\ & \leq \frac{1}{\alpha^n n!} \left( \int_{x_0}^t (t^\alpha - \tau^\alpha)^{np} d_\alpha \tau \right)^{\frac{1}{p}} \left( \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau \right)^{\frac{1}{q}} \\ & = \frac{1}{\alpha^{n+1/p} n!} \frac{(t^\alpha - x_0^\alpha)^{n+1/p}}{(np+1)^{1/p}} (z(t))^{\frac{1}{q}} \end{aligned} \quad (3.2)$$

where

$$z(t) = \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau, \quad x_0 \leq t \leq b, \quad z(x_0) = 0.$$

Thus,

$$D_\alpha z(t) = |D_\alpha^{n+1} f(t)|^q$$

and

$$|D_\alpha^{n+1} f(t)| = (D_\alpha z(t))^{1/q}. \quad (3.3)$$

By (3.2) and (3.3), we get

$$|R_{n,f}(x_0, t)| |D_\alpha^{n+1} f(t)| \leq \frac{1}{\alpha^{n+1/p} n!} \frac{(t^\alpha - x_0^\alpha)^{n+1/p}}{(np+1)^{1/p}} (z(t) D_\alpha z(t))^{\frac{1}{q}}. \quad (3.4)$$

Integrating the inequality (3.4) and using Hölder’s inequality for conformable integrals, we have

$$\begin{aligned} & \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\ & \leq \frac{1}{\alpha^{n+1/p} n! (np+1)^{1/p}} \int_{x_0}^t (\tau^\alpha - x_0^\alpha)^{n+1/p} (z(\tau) D_\alpha z(\tau))^{\frac{1}{q}} d_\alpha \tau \\ & \leq \frac{1}{\alpha^{n+1/p} n! (np+1)^{1/p}} \left( \int_{x_0}^t (\tau^\alpha - x_0^\alpha)^{np+1} d_\alpha \tau \right)^{\frac{1}{p}} \left( \int_{x_0}^t z(\tau) D_\alpha z(\tau) d_\alpha \tau \right)^{\frac{1}{q}} \\ & = \frac{(t^\alpha - x_0^\alpha)^{n+2/p}}{\alpha^{n+2/p} n! [(np+1)(np+2)]^{1/p}} \frac{(z(t))^{\frac{2}{q}}}{2^{\frac{1}{q}}} \end{aligned}$$

which completes the proof. □

**Corollary 1** Under the assumption of Theorem 8 with  $p = q = 2$ , we get

$$\int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \leq \frac{(t^\alpha - x_0^\alpha)^{n+1}}{2\alpha^{n+2} n! \sqrt{(2n+1)(n+1)}} \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^2 d_\alpha \tau.$$

**Theorem 9** Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n + 1$  times  $\alpha$ -fractional differentiable function,  $p, q > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $t \leq x_0$ ,  $t, x_0 \in [a, b]$ . Then we have the following inequality:

$$\begin{aligned} & \int_t^{x_0} |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \tag{3.5} \\ & \leq \frac{(x_0^\alpha - t^\alpha)^{n+2/p}}{\alpha^{n+1+2/p} 2^{\frac{1}{q}} n! [(np+1)(np+2)]^{1/p}} \left( \int_t^{x_0} |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau \right)^{\frac{2}{q}}. \end{aligned}$$

**Proof** From (2.7), we have

$$\begin{aligned} |R_{n,f}(x_0, t)| &= \frac{1}{\alpha^n n!} \left| \int_{x_0}^t (t^\alpha - \tau^\alpha)^n D_\alpha^{n+1} f(\tau) d_\alpha \tau \right| \tag{3.6} \\ &\leq \frac{1}{\alpha^n n!} \int_t^{x_0} (\tau^\alpha - t^\alpha)^n |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\ &\leq \frac{1}{\alpha^n n!} \left( \int_t^{x_0} (\tau^\alpha - t^\alpha)^{np} d_\alpha \tau \right)^{\frac{1}{p}} \left( \int_t^{x_0} |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau \right)^{\frac{1}{q}} \\ &= \frac{1}{\alpha^{n+1/p} n!} \frac{(x_0^\alpha - t^\alpha)^{n+1/p}}{(np+1)^{1/p}} (z(t))^{\frac{1}{q}} \end{aligned}$$

where

$$z(t) = \int_t^{x_0} |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau, \quad a \leq t \leq x_0, \quad z(x_0) = 0.$$

Therefore,

$$D_\alpha z(t) = - |D_\alpha^{n+1} f(t)|^q$$

and

$$|D_\alpha^{n+1} f(t)| = (-D_\alpha z(t))^{1/q}. \quad (3.7)$$

From (3.6) and (3.7), it follows that

$$|R_{n,f}(x_0, t)| |D_\alpha^{n+1} f(t)| \leq \frac{1}{\alpha^{n+1/p} n!} \frac{(x_0^\alpha - t^\alpha)^{n+1/p}}{(np+1)^{1/p}} (-z(t) D_\alpha z(t))^{1/q}. \quad (3.8)$$

Integrating the inequality (3.8) and using Hölder's inequality for conformable integrals, we have

$$\begin{aligned} & \int_t^{x_0} |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\ & \leq \frac{1}{\alpha^{n+1/p} n! (np+1)^{1/p}} \int_t^{x_0} (x_0^\alpha - \tau^\alpha)^{n+1/p} (z(\tau) D_\alpha z(\tau))^{1/q} d_\alpha \tau \\ & \leq \frac{1}{\alpha^{n+1/p} n! (np+1)^{1/p}} \left( \int_t^{x_0} (x_0^\alpha - \tau^\alpha)^{np+1} d_\alpha \tau \right)^{\frac{1}{p}} \left( \int_t^{x_0} (-z(\tau) D_\alpha z(\tau)) d_\alpha \tau \right)^{\frac{1}{q}} \\ & = \frac{(x_0^\alpha - t^\alpha)^{n+2/p}}{\alpha^{n+2/p} n! [(np+1)(np+2)]^{1/p}} \frac{(z(t))^{2/q}}{2^{1/q}}. \end{aligned}$$

This completes the proof.  $\square$

**Corollary 2** Under the assumption of Theorem 9 with  $p = q = 2$ , we get

$$\int_t^{x_0} |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \leq \frac{(x_0^\alpha - t^\alpha)^{n+1}}{2\alpha^{n+2} n! \sqrt{(2n+1)(n+1)}} \int_t^{x_0} |D_\alpha^{n+1} f(\tau)|^2 d_\alpha \tau.$$

**Theorem 10** Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n+1$  times  $\alpha$ -fractional differentiable function,  $p, q > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $t, x_0 \in [a, b]$ . Then we have the following inequality:

$$\begin{aligned} & \left| \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \right| \\ & \leq \frac{|t^\alpha - x_0^\alpha|^{n+2/p}}{\alpha^{n+2/p} 2^{1/q} n! [(np+1)(np+2)]^{1/p}} \left| \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau \right|^{2/q}. \end{aligned} \quad (3.9)$$

**Proof** Combining Theorem 8 and Theorem 9, we can easily get the required result.  $\square$

**Corollary 3** Under the assumption of Theorem 10 with  $p = q = 2$ , we get

$$\left| \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \right| \leq \frac{|t^\alpha - x_0^\alpha|^{n+1}}{2\alpha^{n+1}n!\sqrt{(n+1)(2n+1)}} \left| \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^2 d_\alpha \tau \right|.$$

Using Theorem 10 and Corollary 3, we obtain the following important inequality.

**Corollary 4** Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n + 1$  times  $\alpha$ -fractional differentiable function,  $p, q > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $t, x_0 \in [a, b]$ . If  $D_\alpha^k f(x_0) = 0$ ,  $k = 0, 1, \dots, n$ , then we have the following Opial-type inequality:

$$\begin{aligned} & \left| \int_{x_0}^t |f(\tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \right| \\ & \leq \min \left\{ \frac{|t^\alpha - x_0^\alpha|^{n+2/p}}{\alpha^{n+2/p} 2^{\frac{1}{q}} n! [(np+1)(np+2)]^{1/p}} \left| \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^q d_\alpha \tau \right|^{\frac{2}{q}}, \right. \\ & \quad \left. \frac{|t^\alpha - x_0^\alpha|^{n+1}}{2\alpha^{n+1}n!\sqrt{(n+1)(2n+1)}} \left| \int_{x_0}^t |D_\alpha^{n+1} f(\tau)|^2 d_\alpha \tau \right| \right\}. \end{aligned}$$

**Corollary 5** If we choose  $n = 0$  Corollary 4, then we have the following inequality:

$$\begin{aligned} & \left| \int_{x_0}^t |f(\tau)| |D_\alpha f(\tau)| d_\alpha \tau \right| \\ & \leq \frac{1}{2} \min \left\{ \frac{|t^\alpha - x_0^\alpha|^{2/p}}{2\alpha^{2/p}} \left| \int_{x_0}^t |D_\alpha f(\tau)|^q d_\alpha \tau \right|^{\frac{2}{q}}, \frac{|t^\alpha - x_0^\alpha|}{2\alpha} \left| \int_{x_0}^t |D_\alpha f(\tau)|^2 d_\alpha \tau \right| \right\}. \end{aligned}$$

**Theorem 11** Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n + 1$  times  $\alpha$ -fractional differentiable function,  $p = 1$ ,  $q = \infty$  and  $t \in [x_0, b]$ . Then we have the inequality

$$\int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \leq \frac{(t^\alpha - x_0^\alpha)^{n+2}}{\alpha^{n+2}(n+2)!} \|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}^2 \quad (3.10)$$

where

$$\|D_\alpha^{n+1} f\|_{\infty} := \sup_{x \in [a, b]} |D_\alpha^{n+1} f(x)|.$$



**Proof** From (2.7), we have

$$\begin{aligned}
 |R_{n,f}(x_0, t)| &\leq \frac{1}{\alpha^n n!} \int_{x_0}^t (t^\alpha - \tau^\alpha)^n |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\
 &\leq \frac{1}{\alpha^n n!} \|D_\alpha^{n+1} f\|_{\infty, [x_0, b]} \int_{x_0}^t (t^\alpha - \tau^\alpha)^n d_\alpha \tau \\
 &= \frac{\|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}}{\alpha^{n+1} (n+1)!} (t^\alpha - x_0^\alpha)^{n+1}.
 \end{aligned} \tag{3.11}$$

Moreover, we get

$$|D_\alpha^{n+1} f(t)| \leq \|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}$$

for all  $t \in [x_0, b]$ .

Therefore it follows that

$$|R_{n,f}(x_0, t)| |D_\alpha^{n+1} f(t)| \leq \frac{\|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}^2}{\alpha^{n+1} (n+1)!} (t^\alpha - x_0^\alpha)^{n+1}. \tag{3.12}$$

Integrating the inequality (3.12), we have

$$\begin{aligned}
 \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau &\leq \frac{\|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}^2}{\alpha^{n+1} (n+1)!} \int_{x_0}^t (\tau^\alpha - x_0^\alpha)^{n+1} d_\alpha \tau \\
 &= \frac{(t^\alpha - x_0^\alpha)^{n+2}}{\alpha^{n+2} (n+2)!} \|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}^2
 \end{aligned}$$

This completes the proof of the inequality (3.10). □

**Theorem 12** Let  $p = 1$ ,  $q = \infty$  and  $t \in [a, x_0]$ . Then we have the inequality

$$\int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \leq \frac{(x_0^\alpha - t^\alpha)^{n+2}}{\alpha^{n+2} (n+2)!} \|D_\alpha^{n+1} f\|_{\infty, [a, x_0]}^2. \tag{3.13}$$

**Proof** From (2.7), we get

$$\begin{aligned}
 |R_{n,f}(x_0, t)| &= \left| \frac{1}{\alpha^n n!} \int_{x_0}^t (t^\alpha - \tau^\alpha)^n D_\alpha^{n+1} f(\tau) d_\alpha \tau \right| \\
 &\leq \frac{1}{\alpha^n n!} \int_t^{x_0} (\tau^\alpha - t^\alpha)^n |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \\
 &\leq \frac{1}{\alpha^n n!} \|D_\alpha^{n+1} f\|_{\infty, [x_0, b]} \int_t^{x_0} (\tau^\alpha - t^\alpha)^n d_\alpha \tau \\
 &= \frac{\|D_\alpha^{n+1} f\|_{\infty, [x_0, b]}}{\alpha^{n+1} (n+1)!} (x_0^\alpha - t^\alpha)^{n+1}.
 \end{aligned} \tag{3.14}$$

Furthermore, we have

$$|D_\alpha^{n+1} f(t)| \leq \|D_\alpha^{n+1} f\|_{\infty, [a, x_0]} \tag{3.15}$$

for all  $t \in [a, x_0]$ .

Thus, we obtain

$$\begin{aligned}
 \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau &\leq \frac{\|D_\alpha^{n+1} f\|_{\infty, [a, x_0]}^2}{\alpha^{n+1} (n+1)!} \int_{x_0}^t (x_0^\alpha - \tau^\alpha)^{n+1} d_\alpha \tau \\
 &= \frac{(x_0^\alpha - t^\alpha)^{n+2}}{\alpha^{n+2} (n+2)!} \|D_\alpha^{n+1} f\|_{\infty, [a, x_0]}^2
 \end{aligned}$$

which completes the proof of the inequality (3.13).  $\square$

Combining Theorem 11 and Theorem 12, we have the following result.

**Corollary 6** Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n + 1$  times  $\alpha$ -fractional differentiable function,  $p = 1$ ,  $q = \infty$  and  $t \in [a, b]$ . Then the following inequality holds:

$$\left| \int_{x_0}^t |R_{n,f}(x_0, \tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \right| \leq \frac{|t^\alpha - x_0^\alpha|^{n+2}}{\alpha^{n+2} (n+2)!} \|D_\alpha^{n+1} f\|_{\infty}^2.$$

Using the Corollary 6, we obtain the following important inequality.

**Corollary 7** Let  $\alpha \in (0, 1]$ ,  $f : [a, b] \rightarrow \mathbb{R}$  be an  $n + 1$  times  $\alpha$ -fractional differentiable function,  $p = 1$ ,  $q = \infty$  and  $t \in [a, b]$ . If  $D_\alpha^k f(x_0) = 0$ ,  $k = 0, 1, \dots, n$ , then we have the following Opial-type inequality:

$$\left| \int_{x_0}^t |f(\tau)| |D_\alpha^{n+1} f(\tau)| d_\alpha \tau \right| \leq \frac{|t^\alpha - x_0^\alpha|^{n+2}}{\alpha^{n+2} (n+2)!} \|D_\alpha^{n+1} f\|_{\infty}^2.$$

**Corollary 8** *If we choose  $n = 0$  Corollary 7, then we have the following inequality:*

$$\left| \int_{x_0}^t |f(\tau)| |D_\alpha f(\tau)| d_\alpha \tau \right| \leq \frac{|t^\alpha - x_0^\alpha|^2}{2\alpha^2} \|D_\alpha f\|_\infty^2.$$

#### 4. Conclusions

In this study, we presented some Opial-type inequalities for conformable fractional integrals via using the remainder function of Taylor's theorem for conformable integrals. A further study could assess weighted versions of these inequalities.

#### References

- [1] Abdeljawad T. On conformable fractional calculus. *J Comput Appl Math* 2015; 279: 57-66.
- [2] Anastassiou G. Opial type inequalities for vector valued functions. *Bull Greek Math Soc* 2008; 55: 1-8.
- [3] Anderson DR. Taylor's formula and integral inequalities for conformable fractional derivatives, *Contributions in Mathematics and Engineering, in Honor of Constantin Caratheodory*, Springer, to appear.
- [4] Cheung WS. Some new Opial-type inequalities. *Mathematika* 1990; 37: 136-142.
- [5] Cheung WS. Some generalized Opial-type inequalities. *J Math Anal Appl* 1991; 162: 317- 321.
- [6] Hammad MA, Khalil R. Conformable fractional heat differential equations. *Int J Differ Equ Appl* 2014; 13: 177-183.
- [7] Hammad MA, Khalil R. Abel's formula and wronskian for conformable fractional differential equations. *Int J Differ Equ Appl* 2014; 13: 177-183.
- [8] Iyiola OS, Nwaeze ER. Some new results on the new conformable fractional calculus with application using D'Alambert approach. *Progr Fract Differ Appl* 2016; 2: 115-122.
- [9] Khalil R, Al Horani M, Yousef A, Sababheh M. A new definition of fractional derivative. *J Comput Appl Math* 2014; 264: 65-70.
- [10] Opial Z. Sur une inegaliti. *Ann Polon Math* 1960; 8: 29-32.
- [11] Pachpatte BG. On Opial-type integral inequalities. *J Math Anal Appl* 1986; 120: 547-556.
- [12] Pachpatte BG. Some inequalities similar to Opial's inequality. *Demonstratio Math* 1993; 26: 643-647.
- [13] Pachpatte BG. A note on some new Opial type integral inequalities. *Octagon Math Mag* 1999; 7: 80-84.
- [14] Pachpatte BG. On some inequalities of the Weyl type. *An Stiint Univ "Al.I. Cuza" Iasi* 1994; 40: 89-95.
- [15] Sarikaya MZ, Budak H. Opial type inequalities for conformable fractional integrals. *RGMIA Research Report Collection* 2016; 19: Article 93, 11 pp.
- [16] Traple J. On a boundary value problem for systems of ordinary differential equations of second order. *Zeszyty Nauk Univ Jagiello Prace Mat* 1971; 15: 159-168.
- [17] Zhao CJ, Cheung WS. On Opial-type integral inequalities and applications. *Math Inequal Appl* 2014; 17: 223-232.