

1-1-2010

Species composition and diversity of epipelagic algae in Balık Lake (Şavşat-Artvin, Turkey)

BÜLENT ŞAHİN

BÜLENT AKAR

İMİRAN BAHÇECİ

Follow this and additional works at: <https://journals.tubitak.gov.tr/botany>



Part of the [Botany Commons](#)

Recommended Citation

ŞAHİN, BÜLENT; AKAR, BÜLENT; and BAHÇECİ, İMRAN (2010) "Species composition and diversity of epipelagic algae in Balık Lake (Şavşat-Artvin, Turkey)," *Turkish Journal of Botany*. Vol. 34: No. 5, Article 8. <https://doi.org/10.3906/bot-0912-290>

Available at: <https://journals.tubitak.gov.tr/botany/vol34/iss5/8>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Botany by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Species composition and diversity of epipellic algae in Balık Lake (Şavşat-Artvin, Turkey)

Bülent ŞAHİN*, Bülent AKAR, İmran BAHÇECİ

Karadeniz Technical University, Fatih Education Faculty, Department of Biology 61335 Söğütü, Trabzon - TURKEY

Received: 30.12.2009

Accepted: 05.05.2010

Abstract: The species composition and diversity of the epipellic algae of Balık Lake were investigated between May and October 2008. A total of 39 taxa were determined, and *Cyclotella meneghiniana*, *Craticula cuspidata*, *Closterium strigosum*, *Cocconeis placentula* var. *euglypta*, and *Amphora ovalis* were the most conspicuous algae, regarding frequency of occurrence and relative abundance in the epipellic algal flora. While the highest total cell number of the epipellic community was 26,214 cells/cm² in October, the lowest was 23,644 cells/cm² in July. The highest species diversity index (H') value was 2.26, in June.

Key words: Epipellic algae, seasonal variation, diversity, Balık Lake, Turkey

Balık Gölü (Şavşat-Artvin, Türkiye) epipelik alglerinin tür kompozisyonu ve çeşitliliği

Özet: Balık Gölü epipelik alglerinin tür kompozisyonu ve çeşitliliği 2008 yılının Mayıs ve Ekim ayları arasında incelenmiştir. Toplam 39 takson tespit edilmiştir. *Cyclotella meneghiniana*, *Craticula cuspidata*, *Closterium strigosum*, *Cocconeis placentula* var. *euglypta* ve *Amphora ovalis* epipelik alg florasının nispi bolluğu ve bulunuş sıklığı açısından göze en çok çarpan türleri olmuştur. Epipelik kommunitenin en yüksek yoğunluğu Ekim ayında 26.214 hücre/cm² iken, en düşük yoğunluk Temmuz ayında 23.644 hücre/cm²'dir. En yüksek çeşitlilik indeksi (H') Haziran ayında 2,26'dır.

Anahtar sözcükler: Epipelik algler, mevsimsel değişim, çeşitlilik, Balık Göl, Türkiye

Introduction

Epipellic algae live on the soft bottom sediments and cover a considerable area in most lakes (Björk-Ramberg, 1984). They perform a range of ecosystem functions including biostabilisation of sediments,

regulation of benthic-pelagic nutrient cycling, and primary production (Poulickova et al., 2008). Despite their significance, many aspects of the geographical distribution, biodiversity, and ecology of epipellic algae are poorly understood (Hasler et al., 2008).

* E-mail: bsahin@ktu.edu.tr

Although there are a number of high mountain lakes in the Black Sea region of Turkey, there is only little information on the ecology and composition of epipelagic algae (Şahin, 2000, 2001, 2002, 2004, 2008, 2009; Şahin & Akar, 2005; Gönüloğlu et al., 2009). Apart from the study done by Kolaylı and Şahin (2008, 2009), there are no published articles on the lakes of Artvin.

The aim of this study was to investigate the species composition and diversity of epipelagic algae of Balık Lake.

Materials and methods

Balık Lake is located at latitude 41°17'24"N and longitude 42°29'82"E in Artvin in the Eastern Black Sea region of Turkey (Figure 1). The surface area of the lake is approximately 0.006 km², the average depth is 15 m, and the elevation is 1656 m above sea level.

In order to examine the epipelagic algal flora of Balık Lake one station was chosen. The number of sampling stations depended on the size of the lake. Collections were made during the snow-free period from May to October in 2008. The samples were performed on a monthly basis from 15-30 cm depth and 50-100 cm offshore. Vascular plants and stones were absent at the station.

The epipelagic algal community was sampled by means of a glass tube 0.8 cm in diameter and 100 cm in length. The pipe was moved in a circular direction over the surface of the sediment, releasing the thumb to take up sediment (Round, 1953). Samples were transferred into plastic bottles and fixed with 4% formalin.

At least 3 water-mounted slides were examined for algae and living diatoms from the station to obtain an estimate of algal relative abundance (Round, 1953). At least 300 algal cells were counted at 400×




Figure 1. Map of the study area.

magnification. Permanent slides for the identification of diatoms were prepared from the same sample after boiling in a 1:1 mixture of concentrated H₂SO₄ and HNO₃. The acid cleaned diatoms were mounted in Naphrax medium with a high refractive index (Round, 1953). Identifications were carried out at 400× magnification.

Taxonomic identifications were carried out according to John et al. (2003), Krammer and Lange-Bertalot (1986, 1988, 1991a, 1991b), Round et al. (1990), Sims (1996), Lenzenweger (1996), and Wotowski and Hindák (2005).

During sampling, water temperature and pH were measured using a mercury thermometer and a WTW Digi 88 model pH meter, respectively. Dissolved oxygen concentration was measured according to the method of Winkler (Yaramaz, 1988). Photographs were taken with a Leica DM 2500 microscope.

To evaluate the richness and diversity of the epipellic community of Balık Lake, the Shannon-Weaver species diversity index (H) was used (Shannon and Weaver, 1949):

$$H = \sum_{i=1}^s P_i \log P_i$$

where s is total number of species in the sample, and P_i is the proportion of the number of individuals in the i -th species to the total number of individuals.

Results and discussion

Environmental conditions

The water temperature of the Balık Lake ranged from 8 to 24 °C (mean 16 °C) during the sampling period. pH values varied between 7.2 and 7.9 (mean 7.5). Accordingly, the lake water was slightly alkaline. Dissolved oxygen concentrations were measured between 8.20 and 11.40 mg L⁻¹.

Epipellic algal flora

The epipellic algal flora of Balık Lake consisted of 39 taxa. Of these, 31 belong to the Bacillariophyta, 4 to the Euglenozoa, 3 to the Cyanobacteria, and 1 to the Charophyta divisions. The identified species are listed in Table 1. Photographs of some species are shown in Figure 2.

In the present study, all taxa were found to belong to 22 families. The best represented family is *Bacillariaceae* and comprises 12.82% of all recorded taxa. Medvedeva (2001) pointed out that diverse families are of great diagnostic importance in floristic analyses.

Bacillariophyta was the dominant group in terms of species number and density, and comprised 79.48% of all recorded taxa. The same result was observed in many lakes in the East Black Sea region of Turkey (Şahin, 2000, 2001, 2002, 2004, 2008; Şahin & Akar, 2005). Moore (1974) and Round (1984) pointed out that diatoms are usually the most common elements of epipellic communities. It is well known that the members of the Bacillariophyta are sensitive to a wide range of limnological and environmental variables, and that their community structure may quickly respond to changing physical, chemical, and biological conditions in the environment (Mooser et al., 1996).

During the study period, while the highest total cell number of the epipellic community was 26,214 cells/cm² (in October), the lowest was 23,644 cells/cm² (in July) (Figure 3). Most of the species in the epipellic community exhibited variations during the study. *Craticula cuspidata*, for example, reached its peak (14,906 cells/cm²) and represented 59.18% of the total taxa in September. The same pattern was observed with *Cyclotella meneghiniana* (12,336 cells/cm²) in May and represented 48% of the total taxa. While *Closterium strigosum* reached its peak (4626 cells/cm²) in July, *Cocconeis placentula* var. *euglypta* reached its peak (2056 cells/cm²) in June, August, and October. *Amphora ovalis* reached its peak (1799 cells/cm²) in August (Figure 4).

Nather Khan (1990) explained that the species of *Craticula* were common and abundant in both organically enriched and non-enriched areas. Round (1956a) reported that species of *Amphora*, *Cocconeis*, *Cymbella*, and *Navicula* were common in calcareous and slightly alkaline waters. The present study supported this finding, since the results of chemical analyses of the Balık Lake water showed that it is slightly alkaline. Round (1984) reported that *Cyclotella* species are considered planktonic. However, *Cyclotella meneghiniana* was found in the epipellic community of Balık Lake. The same situation was

Table 1. List of the epipelagic algae in Balık Lake.

Taxa	
Division:	Bacillariophyta
Class:	Coscinodiscophyceae
Order:	Aulacoseriales
Family:	Aulacoseriaceae
	<i>Aulacoseira muzzanensis</i> (Meister) Krammer
Class:	Mediophyceae
Order :	Thalassiosirales
Family:	Stephanodiscaceae
	<i>Cyclotella meneghiniana</i> Kützing (Figure 2a, b)
Class:	Bacillariophyceae
Order:	Fragilariales
Family:	Fragilariaceae
	<i>Staurosira construens</i> Ehrenberg (Figure 2c)
	<i>Ulnaria ulna</i> (Nitzsch) P.Compère in Jahn et al.
Order :	Achnanthales
Family:	Cocconeidaceae
	<i>Cocconeis placentula</i> Ehrenberg var. <i>euglypta</i> (Ehrenberg) Grunow (Figure 2d)
Order :	Thalassiophysales
Family:	Catenulaceae
	<i>Amphora ovalis</i> (Kützing) Kützing (Figure 2e)
Order :	Cymbellales
Family:	Cymbellaceae
	<i>Cymbella cistula</i> (Hemprich & Ehrenberg) Kirchner
	<i>C. cymbiformis</i> C.Agardh (Figure 2f)
	<i>Encyonema minutum</i> (Hilse in Rabenhorst) D.G.Mann in Round et al. (Figure 2g)
Family:	Gomphonemataceae
	<i>Gomphonema acuminatum</i> Ehrenberg
	<i>G. clavatum</i> Ehrenberg
	<i>G. olivaceum</i> (Hornemann) Brébisson var. <i>olivaceoides</i> (Hustedt) lange-Bertalot
Order:	Naviculales
Family :	Diploneidaceae
	<i>Diploneis elliptica</i> (Kützing) Cleve
Family:	Naviculaceae
	<i>Navicula cryptocephala</i> Kützing
	<i>N. gysingensis</i> Foged
Family:	Neidiaceae
	<i>Neidium affine</i> (Ehrenberg) Pfitzer
Family :	Pleurosigmatophyceae
	<i>Gyrosigma acuminatum</i> (Kützing) Rabenhorst
Family :	Pinnulariaceae
	<i>Pinnularia appendiculata</i> (C.Agardh) Cleve
	<i>P. brebissonii</i> (Kützing) Rabenhorst
	<i>P. maior</i> (Kützing) Rabenhorst
	<i>P. viridis</i> (Nitzsch) Ehrenberg (Figure 2h)

Table 1. (Continued).

Taxa	
Family:	Sellaphoraceae <i>Sellaphora pupula</i> (Kützing) Mereschkovsky
Family:	Stauroneidaceae <i>Craticula cuspidata</i> (Kützing) D.G.Mann in Round et al. (Figure 2i) <i>Stauroneis anceps</i> Ehrenberg (Figure 2j)
Order:	Surirellales
Family :	Surirellaceae <i>Cymatopleura solea</i> (Brebisson) W.Smith
Order:	Rhopalodiales
Family :	Rhopalodiaceae <i>Rhopalodia gibba</i> (Ehrenberg) O.Müller
Order:	Bacillariales
Family :	Bacillariaceae <i>Hantzschia amphioxys</i> (Ehrenberg) Grunow (Figure 2k) <i>Surirella robusta</i> (Ehrenberg) van Heurck <i>S. spiralis</i> Kützing <i>S. tenera</i> W.Gregory <i>S. turgida</i> W.Smith (Figure 2l)
Division:	Charophyta
Class:	Zygnematophyceae
Order:	Zygnematales
Family:	Closteriaceae <i>Closterium strigosum</i> Brébisson (Figure 2m)
Division:	Cyanobacteria
Class:	Cyanophyceae
Order:	Nostocales
Family:	Nostocaceae <i>Anabaena</i> sp.
Order:	Oscillatoriales
Family:	Phormidiaceae <i>Phormidium limosum</i> (Dillwyn) P.C.Silva
Order:	Pseudanabaenales
Family:	Pseudanabaenaceae <i>Spirulina maior</i> (Kützing) Gomont
Division:	Euglenozoa
Class:	Euglenophyceae
Order:	Euglenales
Family:	Euglenaceae <i>Euglena gracilis</i> Klebs <i>E. hemichromata</i> Skuja <i>E. limnophila</i> Lemmermann <i>Phacus caudatus</i> Hübner




Figure 2. a and b. *Cyclotella meneghiniana*, c. *Staurosira construens*, d. *Cocconeis placentula* var. *euglypta*, e. *Amphora ovalis*, f. *Cymbella cymbiformis*, g. *Encyonema minutum*, h. *Pinnularia viridis*, i. *Craticula cuspidata*, j. *Stauroneis anceps*, k. *Hantzschia amphioxys*, l. *Surirella turgida*, m. *Closterium strigosum*.

observed in Gölbaşı Lake (Çetin et al., 2002). We can say that the presence of epipelagic algae species together with those of phytoplankton may be a result of wind affecting the surface of water in shallow lakes. *Cyclotella* spp. are indicator species for transition to eutrophy (Round, 1956b).

Euglenozoa are known to be abundant in eutrophic waters and sediments polluted with organic

matter (Round, 1984). In the Balık Lake, Euglenozoa were represented by 4 species and constituted 10.25% of the total taxa.

Round (1984) notified that Cyanobacteria grow in eutrophic waters and on organically polluted sediment in summer and autumn. In addition, it has been suggested that *Phormidium* spp., which are indicators of pollution, are more prolific in polluted




Figure 3. The seasonal variations in the total organism and species diversity of the epipelagic algal community.




Figure 4. Seasonal variations in the density of common epipelagic algae.

regions (Şen et al., 1990). Cyanobacteria were represented by 3 species and constituted 7.69% of the total taxa in our study area.

Charophyta were represented only by 1 species and make up an insignificant portion (2.56%) of the epipelagic algal flora in Balık Lake.

A number of previous reviews focused on phytoplankton diversity (Harris, 1986). In general, when using a diversity index such as Shannon-Weaver, phytoplankton diversity has a strong seasonal component (Margelef, 1958). However, little is known

concerning the diversity of epipelagic algae in lakes in Turkey (Şahin, 2004; Akar & Şahin, 2006). In Balık Lake, it was observed that the seasonal changes in diversity showed an inverse pattern with species number (Figure 3). This means that species evenness decreased with increasing size of the algal population. *Craticula cuspidata* and *Staurosira construens*, for example, increased and occupied 55.87% of the total algal flora in October, when total cell number increased. The same trend was true in May, when *Cyclotella meneghiniana* occupied 48% of the algal flora (Figure 3). The decrease in diversity index values during these months was probably caused by the high relative abundance of the dominant species. In June, when total species number was small, the dominant species (*Cyclotella meneghiniana*) occupied only 36.45% of total species number, and such a low relative abundance resulted in a high diversity index (Figure 3). According to May (1975), the Shannon-Weaver diversity index is related to both the total number of species and their relative abundances, and can be designated as a positive function of total number of species.

In conclusion, the species found in Balık Lake mainly reflect the trophic state of the lake, which is eutrophic, as can be seen from the existence of the species and their abundance.

Acknowledgement

We are grateful to Yücel BAHÇECİ for his help with obtaining samples.

References

- Akar B & Şahin B (2006). Benthic algal flora of Karanlık Lake and diversity of epipellic algae. *Fresen Environ Bull* 15: 48-54.
- Björk-Ramberg S (1984). Species composition and biomass of an epipellic algal community in a subarctic lake before and during lake fertilization. *Holarctic Ecology* 7: 195-201.
- Çetin AK, Şen B & Yıldırım V (2002). Seasonal variations of epipellic diatoms in Gölbaşı Lake with relation to physical-chemical variables. *Fresen Environ Bull* 11: 306-311.
- Gönülol A, Ersanlı E & Baytut Ö (2009). Taxonomical and numerical comparison of epipellic Algae from Balık and Uzun Lagoon, Turkey. *J Environ Biol* 30: 777-784.
- Harris GP (1986). *Phytoplankton Ecology, Structure, Function and Fluctuation*. London: Chapman and Hall Press.
- Hasler P, Stepankova J, Spackova J, Neustupa J, Kitner M, Hereka P, Vesela J, Burian J & Poulickova A (2008). Epipellic cyanobacteria and algae: a case study from Czech fishponds. *Fottea* 8: 133-146.
- John DM, Whitton BA & Brook AJ (2003). *The Freshwater Algal Flora of the British Isles*. Cambridge: Cambridge University Press.
- Kolaylı S & Şahin B (2008). Seasonal variations of benthic diatoms with relation to physical and chemical variables in Karagöl Lake (Borçka-Artvin/Turkey). *Fresen Environ Bull* 17: 956-961.
- Kolaylı S & Şahin B (2009). Benthic algae (except Bacillariophyta) and their seasonal variations in Karagöl Lake (Borçka, Artvin-Turkey). *Turk J Bot* 33: 27-32.
- Krammer K & Lange-Bertalot H (1986). *Subwasserflora von Mitteleuropas, Bacillariophyceae. Band 2/1, 1. Teil: Naviculaceae*. Stuttgart: Gustav Fischer Verlag.
- Krammer K & Lange-Bertalot H (1988). *Subwasserflora von Mitteleuropas, Bacillariophyceae. Band 2/2, 2. Teil: Bacillariaceae, Epithemiaceae, Surirellaceae*. Stuttgart: Gustav Fischer Verlag.
- Krammer K & Lange-Bertalot H (1991a). *Subwasserflora von Mitteleuropas, Bacillariophyceae. Band 2/3, 3. Teil: Centrales, Fragilariaceae, Eunotiaceae*. Stuttgart: Gustav Fischer Verlag.
- Krammer K & Lange-Bertalot H (1991b). *Subwasserflora von Mitteleuropas, Bacillariophyceae. Band 2/4, 4. Teil: Achmanthaceae, Kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema Gesamtliteraturverzeichnis*. Stuttgart: Gustav Fischer Verlag.
- Lenzenweger R (1996). *Desmidiaceenflora von Österreich. Teil 1*. Stuttgart: J Cramer.
- Margalef R (1958). Temporal succession and spatial heterogeneity in phytoplankton. In: Buzzati-Traverso A (ed.) *Perspectives in Marine Biology*, pp. 323-349, Berkeley: University of California Press.
- May RM (1975). Patterns of species abundance and diversity. In: Cody ML & Diamond JM (eds.) *Ecology and Evolution of Communities*. Cambridge, Massachusetts: The Belknap Press of Harvard University Press.
- Medvedeva LA (2001). Biodiversity of aquatic algal communities in the Sikhote-Alin biosphere reserve (Russia). *Cryptogamie Algologie* 22: 65-100.
- Moore JW (1974). Benthic algae of Southern Baffin Island, II. The epipellic communities in temporary ponds. *J Ecol* 62: 809-819.
- Mooser KA, Macdonald GM & Smol JP (1996). Applications of freshwater diatoms to geographical research. *Progress Physical Geog* 20: 21-52.
- Nather-Khan ISA (1990). Assessment of water pollution using diatoms community structure and species distribution: a case study in a tropical river. *Int Revue Ges Hydrobiol* 75: 317-338.
- Poulickova A, Hasler P, Lyskova M & Spears B (2008). The ecology of freshwater epipellic algae: an update. *Phycologia* 47: 437-450.
- Round FE (1953). An investigation of two benthic algal communities in Malharm Tarn, Yorkshire. *J Ecol* 41: 97-174.
- Round FE (1956a). A note on some communities of the littoral zone lakes. *Arch F Hydrobiol* 52: 3-28.
- Round FE (1956b). The phytoplankton of three water supply reservoirs in Central Wales. *Arch F Hydrobiol* 52: 220-232.
- Round FE (1984). *The Ecology of Algae*. Cambridge: Cambridge University Press.
- Round FE, Crawford RM & Mann DG (1990). *Diatoms: Biology and Morphology of the Genera*. Cambridge: Cambridge University Press.
- Shannon CE & Weaver W (1949). *The Mathematical Theory of Communication*. Urbana: University of Illinois Press.
- Sims PA (1996). *An Atlas of British Diatoms*. Biopress Ltd.
- Şahin B (2000). Algal flora of Lakes Aygır and Balıklı (Trabzon, Turkey). *Turk J Bot* 24: 35-45.
- Şahin B (2001). Epipellic and epilithic algae of Dağbaşı Lake (Rize-Turkey). *Turk J Bot* 25: 187-194.
- Şahin B (2002). Epipellic and epilithic algae of the Yedigöller Lakes (Erzurum-Turkey). *Turk J Biol* 26: 221-228.
- Şahin B (2004). Species composition and diversity of epipellic algae in Çatal Lake (Şebinkarahisar-Giresun, Turkey). *Turk J Biol* 28: 103-109.
- Şahin B & Akar B (2005). Epipellic and epilithic algae of Küçükgöl Lake (Gümüşhane-Turkey). *Turk J Biol* 29: 57-63.
- Şahin B (2008). Species composition and diversity of epipellic algae in Limni Lake (Gümüşhane, Turkey). *Acta Bot Hung* 50: 397-405.
- Şahin B (2009). Contribution to the desmid flora of Turkey. *Turk J Bot* 33: 457-460.
- Şen B, Çetin AK & Nacar YA (1990). Study on the growth patterns of algae in the wastewaters of a small water-collecting channel. X. *Ulusal Biyoloji Kong Erzurum*, pp. 86-94.
- Wotowski K & Hindák F (2005). *Atlas of Euglenophytes*. Bratislava: House of the Slovak Academy of Sciences VEDA/ Publishing.
- Yaramaz Ö (1988). *Su Kalitesi*. İzmir: Ege Üniversitesi Su Ürünleri Yüksek Okulu.