

1-1-2019

The role of hepcidin, GDF15, and mitoferrin-1 in iron metabolism of polycythemia vera and essential thrombocytosis patients

CANAN ALBAYRAK YAŞAR

PINAR TARKUN

ELİF BİRTAŞ ATEŞOĞLU

FATMA CEYLA ERALDEMİR

ÖZGÜR DOĞA ÖZSOY

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>

 Part of the [Medical Sciences Commons](#)

Recommended Citation

YAŞAR, CANAN ALBAYRAK; TARKUN, PINAR; ATEŞOĞLU, ELİF BİRTAŞ; ERALDEMİR, FATMA CEYLA; ÖZSOY, ÖZGÜR DOĞA; DEMİRİSOY, ESRA TERZİ; MEHTAP, ÖZGÜR; GEDÜK, AYFER; and HACİHANİFİOĞLU, ABDULLAH (2019) "The role of hepcidin, GDF15, and mitoferrin-1 in iron metabolism of polycythemia vera and essential thrombocytosis patients," *Turkish Journal of Medical Sciences*: Vol. 49: No. 1, Article 12.

<https://doi.org/10.3906/sag-1803-13>

Available at: <https://journals.tubitak.gov.tr/medical/vol49/iss1/12>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The role of hepcidin, GDF15, and mitoferrin-1 in iron metabolism of polycythemia vera and essential thrombocytosis patients

Authors

CANAN ALBAYRAK YAŐAR, PINAR TARKUN, ELİF BİRTAŐ ATEŐOĐLU, FATMA CEYLA ERALDEMİR, ÖZGÜR DOĐA ÖZSOY, ESRA TERZİ DEMİRSOY, ÖZGÜR MEHTAP, AYFER GEDÜK, and ABDULLAH HACİHANİFİOĐLU

The role of hepcidin, GDF15, and mitoferrin-1 in iron metabolism of polycythemia vera and essential thrombocytosis patients

Canan ALBAYRAK¹, Pınar TARKUN^{2*}, Elif BİRTAŞ ATEŞOĞLU², Ceyla ERALDEMİR³, Özgür Doğa ÖZSOY³,
Esra TERZİ DEMİRSOY², Özgür MEHTAP², Ayfer GEDÜK², Abdullah HACIHANEFİOĞLU²

¹Department of Internal Medicine, School of Medicine, Kocaeli University, Kocaeli, Turkey

²Department of Hematology, School of Medicine, Kocaeli University, Kocaeli, Turkey

³Department of Biochemistry, School of Medicine, Kocaeli University, Kocaeli, Turkey

Received: 02.03.2018 • Accepted/Published Online: 03.11.2018 • Final Version: 11.02.2019

Background/aim: GDF15, hepcidin and mitoferrin-1 (mfrn-1) are proteins involved in systemic iron regulation. There are no studies in the literature demonstrating the serum mfrn-1 levels in polycythemia vera (PV) and essential thrombocythemia (ET) patients. The aim of this study was to investigate GDF15, hepcidin and mfrn-1 levels in PV and ET patients.

Materials and methods: Ten PV, 17 ET patients, and 27 healthy controls (HCs) were enrolled. GDF15, hepcidin and mfrn-1 values were measured with enzyme-linked immunosorbent assay (ELISA).

Results: GDF15 levels were higher in the myeloproliferative neoplasm (MPN) group (P = 0.002). Hepcidin levels were not different between MPN patients and HCs. The mfrn-1 levels were lower in MPN patients (P = 0.039). Hepcidin, GDF15, and mfrn-1 levels were not different between PV and ET patients. mfrn-1 levels were lower in ET patients than HCs (P = 0.038).

Conclusion: Increased erythropoiesis in MPNs may lead to high GDF15 levels in these patients. However, hepcidin was not suppressed despite the increased GDF15 levels and erythropoiesis in these patients. Decrease in mfrn-1 in MPNs can be the result of its increased turnover due to increased myelopoiesis. It can be hypothesized that similar hepcidin levels in patients and controls and low mfrn-1 levels in patients may be a defense mechanism against erythroid activity and thromboembolic complications.

Key words: Polycythemia vera, essential thrombocythemia, chronic myeloproliferative neoplasm, GDF15, hepcidin, mitoferrin-1

1. Introduction

Myeloproliferative neoplasms (MPN) are clonal hematopoietic stem cell disorders (1). In polycythemia vera (PV) and essential thrombocythemia (ET), there is an increase in all hematopoietic series, but essentially in erythroid and megakaryocytoid series (2). JAK V617F mutation exists in PV, ET, and primary myelofibrosis patients (3). The JAK2 gene is located on 9p24 (4,5).

Hepcidin is a small protein primarily released from the liver. It is the main regulator of iron absorption and transportation to the tissues. Plasma iron level and erythropoietic activity regulate the hepcidin level. The increase in the iron level increases the hepcidin production. High hepcidin level reduces dietary iron absorption and the use of iron within the cell; therefore, hepcidin prevents iron accumulation in tissues. High erythropoietic activity suppresses hepcidin. In cases of inflammation and infection, as a host defense hepcidin production increases in order to prevent iron transportation to microorganisms (6).

* Correspondence: pinartarkun@hotmail.com

GDF15 (growth differentiation factor-15) is a member of the TGF- β (transforming growth factor β) family. It is released from mature erythrocytes. It is defined as a hepcidin suppression factor. In disease states where ineffective erythropoiesis is observed (e.g., thalassemia, sideroblastic anemia, congenital dyserythropoietic anemia, etc.), its level increases. In these patients, when GDF15 reaches extremely high levels, it suppresses hepcidin and, therefore, by increasing iron absorption, it contributes to iron overload in the tissues (7).

Iron is an element that is stored and utilized in cellular organelles. Mitochondria use iron in the synthesis of heme molecules (8). Previously, it was not known how the iron molecule was transported to mitochondria. In 2006, Shaw et al. identified a mutated gene in frascati zebrafish with deep anemia and also the protein that this gene synthesizes (3). The protein that is the product of this gene is in the mitochondrial agent transporter family (SLC 25) and identified as mitoferrin (9). Mitoferrin-1 (also known as

Slc 25a37) transports the iron necessary for heme synthesis in the developing erythrocytes through the mitochondrial membrane. Mitoferrin-1 in mice is synthesized in high amounts in hematopoietic organs like the fetal liver, bone marrow, and spleen. Mitoferrin-2 is analogous with mitoferrin-1. It is detected in low amounts in all tissues but it is necessary primarily in nonerythroid cells for heme synthesis in hemoproteins and the formation of Fe-sulfide compound (9,10). During erythrocyte maturation, Abcb 10 prolongs the half-life of mitoferrin-1 and therefore increases its concentration. Thus, it enables more iron molecules in the mitochondria for heme and Fe-S compound synthesis (11). Abcb 10 is inhibited by the formed heme molecule and therefore mitoferrin-1 activity reduces. (8). Recent studies have shown that mitoferrin-1 is located as an oligomeric complex together with ferrochelatase and Abcb 10 in mitochondria (11,12).

The purpose of this study is to determine the differences between the iron metabolism of PV and ET patients and healthy controls (HCs) using mitoferrin-1, GDF15 and hepcidin levels since there is an increase in hematopoiesis in MPNs.

2. Materials and methods

In our study, patients who were evaluated in the Department of Hematology and who were diagnosed with PV or ET were included. Patients were diagnosed according to the 2008 WHO criteria (2). All of the enrolled patients were newly diagnosed and patients who received any treatment that could affect iron metabolism, who were previously treated by phlebotomy or hydroxyurea, who had hematologic diseases other than PV or ET, who had active infection or inflammation, and who had decompensated liver, heart, or kidney diseases were excluded from the study. The control group of the study included 27 individuals presenting to the General Internal Medicine Department with no organic diseases. Our study was approved by the local ethics committee. Written consent was obtained from all the participants.

Real-time PCR with a QIAGEN Rotor-Gene Q device was used for the analysis of JAK2 V617F gene mutation. Mitoferrin-1 levels were studied with a Cusabio human mitoferrin-1 (SLC25A37) enzyme-linked immunosorbent assay (ELISA) kit (Cusabio Inc., Wuhan, Hubei, China). Hepcidin levels were studied with a DRG Hepcidin 25 (bioactive) ELISA kit (DRG International Inc., Marburg, Germany). GDF15 levels were studied with a RayBio human GDF15 ELISA kit (RayBiotech Inc., Norcross GA, USA). The results were recorded in ng/mL.

For statistical analyses, SPSS 20.0 for Windows was used. In comparison of normally distributed variables, the independent sample t-test with two averages or one-way ANOVA analysis was used. The Mann-Whitney U

test, Kruskal-Wallis test, and Tukey multiple comparison tests were used in comparison of nonnormally distributed variables. In the comparison of qualitative data, Pearson chi-square and Fisher exact tests were used. $P < 0.05$ was accepted as statistically significant.

3. Results

Twenty-seven patients diagnosed with MPNs [10 PV (37%) and 17 ET (63%) cases] and 27 HCs were included in the study. General characteristics of MPN patients and HCs are shown in Table 1. As seen in Table 1, serum creatinine, uric acid, LDH levels, leukocyte and thrombocyte counts of the patient group were significantly higher than those of HCs. MCV, serum transferrin saturation, iron and ferritin levels were significantly lower than in HCs.

General characteristics of patients diagnosed with PV and ET in the MPN group are shown in Table 2. In the PV group, the hematocrit levels ($P = 0.07$) were significantly higher and the thrombocyte counts were significantly lower than in ET patients ($P < 0.001$) (Table 2). Splenomegaly rate was significantly higher in PV patients ($P = 0.003$)

When serum hepcidin, GDF15, and mitoferrin-1 levels were compared between the MPN patients and HCs, serum hepcidin levels were similar in the two groups. However, the GDF15 levels of MPN patients were significantly higher than those of the HCs ($P = 0.002$). In the MPN patients, serum mitoferrin-1 levels were detected to be significantly lower than in the HCs ($P = 0.039$). An analysis was performed between MPN subgroups and HCs and a subgroup analysis was performed in MPN itself. The data of these analyses are shown in Table 3. No difference was detected between PV and ET patients and HCs in terms of serum hepcidin levels ($P > 0.05$). However, serum GDF15 levels were significantly higher than those of HCs in both PV and ET patients ($P = 0.013$, $P = 0.034$, respectively). The GDF15 levels of PV and ET patients were similar. While the serum mitoferrin-1 levels of ET patients were significantly lower than those of HCs ($P = 0.038$), no significant difference was observed in other subgroups.

When hepcidin, GDF15 and mitoferrin-1 levels were analyzed according to the presence of JAK2 mutation and splenomegaly, it was found that hepcidin levels were significantly lower in the patients with JAK2 mutation ($P = 0.007$) (Table 4). There was no difference in terms of hepcidin, GDF15, and mitoferrin levels in patients with and without splenomegaly (Table 4). As the number of patients was low, no analysis was performed according to the thrombosis/hemorrhage history and hepcidin, GDF15, and mitoferrin-1 levels (the number of patients with a history of thrombosis and hemorrhage was 3 and 1, respectively).

Table 1. Characteristics of patients with MPNs and HC group.

	MPNs, mean \pm SD Median (25th–75th percentile)	HCs, mean \pm SD Median (25th–75th percentile)	P
Age (years)	57.07 \pm 13.39	56.74 \pm 10.97	0.921
Sex (female/male)	11/16	11/16	1.00
Creatinine ² (0.6–1.3 mg/dL) ³	0.91 \pm 0.23 0.83 (0.72–0.97)	0.77 \pm 0.11 (0.76) 0.76 (0.68–0.84)	0.017*
Uric acid ² (2.5–7.7 mg/dL) ³	5.85 \pm 0.99 5.85 (4.97–6.90)	4.97 \pm 1.34 5.20 (3.70–6.00)	0.027*
LDH ¹ (125–220 U/L) ³	236.20 \pm 55.95	167.25 \pm 25.61	<0.001*
Hb ² (12.2–18.1 g/dL) ³	14.37 \pm 1.91 15.10 (12.8–15.6)	14.41 \pm 1.23 14.00 (13.5–15.6)	0.736
Hct ² (37.7–53.7%) ³	44.27 \pm 7.02 44.60 (38.8–50.6)	42.98 \pm 3.66 41.60 (40.7–44.9)	0.324
MCV ¹ (80.0–97.0 fL) ³	80.90 \pm 10.26	90.86 \pm 3.68	<0.001*
Leukocytes ¹ (4.60–10.2 $\times 10^3/\mu\text{L}$) ³	10.75 \pm 4.58	6.86 \pm 1.04	<0.001*
Platelets ¹ (142–424 $\times 10^3/\text{L}$) ³	731.55 \pm 252.12	240.92 \pm 57.50	<0.001*
CRP ² (0–0.5 mg/dL) ³	0.39 \pm 0.65 0.13 (0.07–0.41)	0.28 \pm 0.21 0.25 (0.11–0.42)	0.444
ESR ² (<20 mm/h) ³	10.18 \pm 12.7 4.50 (4.00–11.25)	8.18 \pm 7.36 6.00 (2.00–13.00)	0.441
Serum iron ¹ (F: 50–170 $\mu\text{g}/\text{dL}$) ³ (M: 65–175 $\mu\text{g}/\text{dL}$) ³	57.42 \pm 34.74	91.18 \pm 28.60	<0.001*
Ferritin ² (F: 11–306.8 ng/mL) ³ (M: 23.9–336.2 ng/mL) ³	53.45 \pm 73.24 26.80 (7.30–61.30)	79.59 \pm 51.92 63.00 (38.60–99.00)	0.003*
Total iron-binding capacity ¹ (228–428 $\mu\text{g}/\text{dL}$) ³	344.69 \pm 64.72	318.92 \pm 35.31	0.081
Transferrin saturation ² (%) (20–50) ³	18.07 \pm 12.17 17.47 (5.65–27.48)	28.84 \pm 9.07 27.38 (22.60–32.76)	0.003*

* P < 0.05, statistically significant.

¹Independent sample t-test, ²Mann–Whitney U test, ³values given in parentheses are the normal reference ranges. LDH: Lactate dehydrogenase, Hb: hemoglobin, Hct: hematocrit, MCV: mean corpuscular volume, CRP: C reactive protein, ESR: erythrocyte sedimentation rate, F: female, M: male

4. Discussion

Hepcidin is a peptide containing 25 amino acids, which was first identified in human urine and plasma (13). There are a few studies in the literature investigating the hepcidin level measurements in MPNs. In one of those studies, four myelofibrosis patients' urine hepcidin levels were measured and their hepcidin levels were detected to be

low (14). In another study by Kwaipasz et al., prohepcidin levels in PV patients were measured and low values were obtained when compared to the control group (15). Especially in patients with iron deficiency, hepcidin levels were lower. It was thought that reduction in hepcidin due to iron deficiency would increase iron absorption and therefore would stimulate erythropoiesis. This would

Table 2. Characteristics of patients with PV and ET.

	PV, mean \pm SD Median (25th–75th percentile) (n = 10 patients)	ET, mean \pm SD Median (25th–75th percentile) (n = 17 patients)	P
Age (years)	57.80 \pm 15.33	56.66 \pm 12.59	0.970
Sex (female/male)	4 / 6	7 / 10	1.000
Hb ² (12.2–18.1 g/dL) ³	15.34 \pm 1.65 15.30 (14.32–16.57)	13.81 \pm 1.86 14.30 (11.95–15.40)	0.052
Hct ² (%) (37.7–53.7) ³	48.75 \pm 5.96 50.15 (45.37–51.95)	41.64 \pm 6.35 43.60 (36.65–45.90)	0.007*
MCV ¹ (80.0–97.0 fL) ³	77.99 \pm 11.06	82.61 \pm 9.69	0.289
Leukocytes ¹ (4.60–10.2 \times 10 ³ / μ L) ³	12.18 \pm 5.39	9.91 \pm 3.97	0.199
Platelets ¹ (142–424 \times 10 ³ /L) ³	567.20 \pm 194.31	828.23 \pm 235.04	<0.001*
CRP ² (0–0.5 mg/dL) ³	0.58 \pm 1.01 0.20 (0.09–0.64)	0.29 \pm 0.35 0.11 (0.05–0.46)	0.401
Serum iron ¹ (F: 50–170 μ g/dL) ³ (M: 65–175 μ g/dL) ³	49.50 \pm 42.03	62.37 \pm 29.73	0.577
Ferritin ² (F: 11–306.8 ng/mL) ³ (M: 23.9–336.2 ng/mL) ³	47.53 \pm 79.92	56.92 \pm 71.34	0.400
Splenomegaly, n (%)	8 (80 %)	3 (17.6 %)	0.003*
JAK2 V617F mutation positivity, n (%)	10 (100 %)	14 (82.4 %)	0.274
Hemorrhage, n (%)	2 (20.0 %)	1 (5.9 %)	0.535
Thrombosis, n (%)	0 (0 %)	1 (5.9 %)	1.000

*P < 0.05, statistically significant.

¹Mann–Whitney U test, ²values given in parentheses are the normal reference ranges. Hb: Hemoglobin, Hct: hematocrit, MCV: mean corpuscular volume, CRP: C reactive protein, F: female, M: male.

partially explain the disease's pathogenesis (15). In the literature, there are data indicating that prohepcidin levels are not correlated with urinary and serum hepcidin levels (16). In a study conducted by Tarkun et al., the serum hepcidin levels of the MPN group including PV and ET patients and the HC group were similar (17). Similarly in the present study, hepcidin levels of the MPN group and HC group were similar. Although in PV patients serum hepcidin levels were lower when compared to the HC group, they were not significantly different. In light of this information, it can be suggested that there is a defense mechanism in order to decrease the iron supply and suppress hematopoiesis in MPN patients.

GDF15 production is associated with cellular stress, damage, and apoptosis. In diseases where ineffective erythropoiesis is observed, GDF15 is released from

mature erythroblasts in high amounts. High GDF15 levels suppress the hepcidin production. This suppression requires high amounts of GDF15 levels (7,18–20). In the literature, there are several studies indicating the potential role of GDF15 in hepcidin release regulation. Tanno et al. detected that in patients with β -thalassemia where ineffective erythropoiesis was observed, serum GDF15 levels were high and hepcidin levels were low. Again in this study, in human hepatocyte cultures, serum GDF15 and recombinant GDF15 in patients with thalassemia were shown to suppress hepcidin production. High levels (>5000 pg/mL) of GDF15 were required for this suppression (18). In another study that Mast et al. conducted, in high-intensity blood donors where effective hematopoiesis is expected, hepcidin levels were low; however, there was no significant increase in GDF15 levels (21). In the study by

Table 3. Serum hepcidin, GDF15, and mitoferrin-1 levels of patients and HCs.

	Hepcidin (ng/mL) Mean ± SD	GDF15 (pg/mL) Mean ± SD	Mitoferrin-1 (pg/mL) Mean ± SD
MPNs (n = 27 patients)	22.50 ± 17.38	280.44 ± 138.44	37.00 ± 10.19
HCs (n = 27 patients)	23.51 ± 12.14	178.20 ± 77.03	42.73 ± 9.63
P-value	0.806	0.002*	0.039*
PV (n = 10 patients)	15.31 ± 10.91	301.64 ± 159.84	40.35 ± 12.94
HC (n = 27 patients)	23.51 ± 12.14	178.20 ± 77.03	42.73 ± 9.63
P-value	0.291	0.013*	0.791
ET (n = 17 patients)	26.74 ± 19.30	267.97 ± 127.78	35.03 ± 7.95
HC (n = 27 patients)	23.51 ± 12.14	178.20 ± 77.03	42.73 ± 9.63
P-value	0.756	0.034*	0.038*
PV (n = 10 patients)	15.31 ± 10.91	301.64 ± 159.84	40.35 ± 12.94
ET (n = 17 patients)	26.74 ± 19.30	267.97 ± 127.78	35.03 ± 7.95
P-value	0.131	0.734	0.371

*P < 0.05, statistically significant.

Table 4. Relationships between hepcidin, GDF15, and mitoferrin-1 levels and presence of *JAK2* mutation and splenomegaly.

	Hepcidin (ng/mL) Mean ± SD	GDF15 (pg/mL) Mean ± SD	Mitoferrin-1 (pg/mL) Mean ± SD
<i>JAK2</i> mutation (+) (n = 24)	19.44 ± 13.78	281.42 ± 143.80	36.51 ± 10.63
<i>JAK2</i> mutation (-) (n = 3)	47.06 ± 26.95	272.55 ± 106.13	40.96 ± 4.68
P-value	0.007*	0.919	0.487
Splenomegaly (+) (n = 11)	18.92 ± 20.12	291.30 ± 168.16	39.48 ± 11.67
Splenomegaly (-) (n = 16)	24.97 ± 15.42	272.92 ± 119.27	35.29 ± 9.02
P-value	0.385	0.742	0.303

*P < 0.05, statistically significant.

Tarkun et al., in the MPN group, serum GDF15 levels were detected to be significantly high; however, no difference was detected in serum hepcidin levels of MPN patients when compared to the control group. In addition, no correlation was observed between GDF15 and hepcidin levels. These results were linked to the fact that GDF15 was not high enough to suppress hepcidin. Moreover, despite the increase in erythropoietic activity and GDF15 levels, it was hypothesized that the normal levels of hepcidin measured in MPN patients could be a defense mechanism to reduce iron utilization in order to prevent erythropoietic activity from increasing (17). In our study, there was no correlation between serum GDF15 and hepcidin levels.

In our study, in the MPN patients, serum mitoferrin-1 levels were found to be significantly lower when compared to the HC group. Similar to the MPN group, in ET patients, significantly low serum mitoferrin-1 levels were obtained.

In the PV patients, although the serum mitoferrin-1 levels were low, there was no statistical difference. In the comparison of PV and ET patients, no statistical difference was observed in terms of mitoferrin-1 values. A few theories can be speculated for low serum mitoferrin-1 levels in MPN patients. First of all, the serum levels may be low because mitoferrin-1 helps iron molecules enter mitochondria and it is used in order to provide iron entry into the mitochondria. In this case, the decrease in serum mitoferrin-1 level can be an indirect indicator of the increase of hematopoiesis in MPN cases. Another reason for low mitoferrin-1 in MPN patients could be a defense mechanism just as there is no decrease in serum hepcidin levels. We believe that low levels of serum mitoferrin-1 may reduce mitochondrial iron uptake and control the myeloproliferation. For this to be conclusive, there is a need for studies investigating the mitoferrin-1 gene expression

and synthesis at the cellular level. This cellular defense mechanism can occur through additional undetected factors that inhibit mitoferrin-1 expression. A third mechanism can occur through Abcb 10 forming a complex with mitoferrin-1, ferrochelatase proteins, or a member of this yet-to-be-discovered complex inhibiting mitoferrin directly. All these theories suggest a defense mechanism in order to prevent thromboembolic events, which are a part of the clinical picture and occur as a result of increased hyperviscosity in MPN cases. In PV patients where erythropoiesis is dominant, low serum mitoferrin-1 values were not obtained and this is linked to the insufficient number of patients. In the literature, there is no study about serum mitoferrin-1 measurements in MPN patients. As our study is the first study showing serum mitoferrin-1 levels in these patients, a data comparison could not be carried out.

Hepcidin binds and inhibits ferroportin and stops iron transport from macrophages and iron absorption from intestines (22). When hepcidin is bound to ferroportin, ferroportin phosphorylation occurs (23). Domenico et al. showed that JAK2 is necessary for this phosphorylation; and JAK2's binding to ferroportin is hepcidin-dependent (24). In the following years, Ross et al. showed that for hepcidin-mediated ferroportin internalization, JAK2 is not necessary and the JAK-STAT pathway is not included in ferroportin phosphorylation (25). In the study conducted by Tarkun et al., hepcidin levels of MPN patients were similar in patients with or without *JAK2* mutation. In the same study, GDF15 levels were significantly higher in patients with *JAK2* V617F. Despite these findings, no

negative correlation could be shown between hepcidin and GDF15. There is a need for further studies (17). Due to these contradictory literature data, the direct role of *JAK2* mutation on iron metabolism in MPN patients is inconclusive. In our study, within the MPN group, serum GDF15 and mitoferrin-1 values between the patients with and without *JAK2* mutation were similar. We found that in patients with *JAK2* mutation, hepcidin levels were significantly lower, similar to patients with iron deficiency. However, there were only three patients in the *JAK2* negative group, so we think that it is not appropriate to make a conclusion.

There are several shortcomings of the present study. First of all, the number of patients was low. In addition, we studied the levels of proteins in serum with ELISA and we could not demonstrate the levels of expression of the molecules with genetic methods due to technical insufficiency.

In conclusion, in this study GDF15, hepcidin, and mitoferrin-1 levels of MPN patients were evaluated and this is the only study demonstrating the mitoferrin-1 levels in MPN patients. GDF15 levels in the MPN group were higher than in the HC group, which can be attributed to increased erythropoiesis in these patients. This study revealed that hepcidin was not suppressed despite the increased GDF15 levels and erythropoiesis in these patients. Moreover, mitoferrin-1 levels were lower in MPN patients when compared to HC. It can be hypothesized that similar hepcidin levels in patients and controls and low mitoferrin-1 levels in patients may be a defense mechanism against erythroid activity and thromboembolic complications.

References

- Hoffman R, Baker KR, Prchal JT. The polycythemias. In: Hoffman R, Benz EJ, Shattil SJ, editors. Hematology: Basic Principles and Practice. 6th ed. Philadelphia, PA, USA: Saunders, Elsevier; 2005. pp. 1209-1291.
- Tefferi A, Vainchenker W. Myeloproliferative neoplasms: molecular pathology, essential clinical understanding, and treatment strategies. J Clin Oncol 2011; 29: 573-582.
- Tefferi A. JAK and MPL mutations in myeloid malignancies. Leukemia Lymphoma 2008; 49: 388-397.
- Tefferi A. JAK2 mutations and clinical practice in myeloproliferative neoplasms. Cancer J 2007; 13: 366-371.
- Josef T, Prchal FJ. Polycythemia vera. In: Lichtman M, Kaushansky K, Beutler E, Seligsohn U, Kipps T, Prchal FJ, editors. Williams Hematology. 8th ed. New York, NY, USA: McGraw-Hill; 2010. pp. 1919-1943.
- Nemeth E, Ganz T. The role of hepcidin in iron metabolism. Acta Haematol 2009; 122: 78-86.
- Tanno T, Noel P, Miller JL. Growth differentiation factor 15 in erythroid health and disease. Curr Opin Hematol 2010; 17: 184-90.
- Nancy CA. ABCs of erythroid mitochondrial iron uptake. P Natl Acad Sci USA 2009; 106: 16012-16013.
- Shaw GC, Cope JJ, Li L, Corson K, Hersey C, Ackermann GE, Gwynn B, Lambert AJ, Wingert RA, Traver D et al. Mitoferrin is essential for erythroid iron assimilation. Nature 2006; 440: 96-100.
- Paradkar PN, Zumbrennen KB, Paw BH, Ward DM, Kaplan J. Regulation of mitochondrial iron import through differential turnover of mitoferrin 1 and mitoferrin 2. Mol Cell Biol 2009; 29: 1007-1016.
- Chen W, Paradkar PN, Li L, Pierce EL, Langer NB, Takahashi-Makise N, Hyde BB, Shirihai OS, Ward DM, Kaplan J et al. Abcb10 physically interacts with mitoferrin-1 (Slc25a37) to enhance its stability and function in the erythroid mitochondria. P Natl Acad Sci USA 2009; 106: 16263-16268.

12. Chen W, Dailey HA, Paw BH. Ferro chelatase forms an oligomeric complex with mitoferrin-1 and Abcb10 for erythroid heme biosynthesis. *Blood* 2010; 116: 628-630.
13. Park CH, Valore EV, Waring AJ, Ganz T. Hepcidin, a urinary antimicrobial peptide synthesized in the liver. *J Biol Chem* 2001; 276: 7806-7810.
14. Winder A, Lefkowitz R, Ghoti H, Leiba M, Ganz T, Nemeth E, Rachmilewitz EA. Urinary hepcidin excretion in patients with myelodysplastic syndrome and myelofibrosis. *Br J Haematol* 2008; 142: 668-680.
15. Kwapisz J, Zekanowska E, Jasiniewska J. Decreased serum prohepcidin concentration in patients with polycythemia vera. *J Zhejiang Univ Sci B* 2009; 10: 791-795.
16. Frazer DM, Anderson GJ. Heparin compared with prohepcidin: an absorbing story. *Am J Clin Nutr* 2009; 89: 475-476.
17. Tarkun P, Mehtap Ö, Ateşoğlu EB, Gedük A, Musul MM, Hacıhanefioğlu A. Serum hepcidin and growth differentiation factor-15 (GDF-15) levels in polycythemia vera and essential thrombocythemia. *Eur J Haematol* 2013; 91: 228-235.
18. Tanno T, Bhanu NV, Oneal PA, Goh SH, Staker P, Lee YT, Moroney JW, Reed CH, Luban NL, Wang RH et al. High levels of GDF15 in thalassemia suppress expression of the iron regulatory protein hepcidin. *Nat Med* 2007; 13: 1096-1101.
19. Tamary H, Shalev H, Perez-Avraham G, Zoldan M, Levi I, Swinkels DW, Tanno T, Miller JL. Elevated growth differentiation factor 15 expression in patients with congenital dyserythropoietic anemia type I. *Blood* 2008; 112: 5241-5244.
20. Musallam KM, Taher AT, Duca L, Cesaretti C, Halawi R, Cappellini MD. Levels of growth differentiation factor-15 are high and correlate with clinical severity in transfusion independent patients with β thalassemia intermedia. *Blood Cells Mol Dis* 2011; 47: 232-234.
21. Mast AE, Foster TM, Pinder HL, Beczkiewicz CA, Bellissimo DB, Murphy AT, Kovacevic S, Wroblewski VJ, Witcher DR. Behavioral, biochemical, and genetic analysis of iron metabolism in high-intensity blood donors. *Transfusion* 2008; 48: 2197-2204.
22. Nemeth E. Iron regulation and erythropoiesis. *Curr Opin Hematol* 2008; 15: 169-175.
23. Withuhn BA, Quelle FW, Silvennoinen O, Yi T, Tang B, Miura O, Ihle JN. JAK2 associates with the erythropoietin receptor and is tyrosinephosphorylated and activated following stimulation with erythropoietin. *Cell* 1993; 74: 227-236.
24. Domenico ID, Lo E, Ward DM, Kaplan J. Heparin-induced internalization of ferroportin requires binding and cooperative interactions with Jak2. *P Natl Acad Sci USA* 2009; 106: 3800-3805.
25. Ross SL, Tran L, Winters A, Lee KJ, Plewa C, Foltz I, King C, Miranda LP, Allen J, Beckman H et al. Molecular mechanism of hepcidin-mediated ferroportin internalization requires ferroportin lysines, not tyrosines or JAK-STAT. *Cell Metab* 2012; 15: 905-917.