

1-1-2002

## Radiation Transfer for Exponential $c(x)$

FATMA ERDOĞAN

Follow this and additional works at: <https://journals.tubitak.gov.tr/physics>



Part of the [Physics Commons](#)

---

### Recommended Citation

ERDOĞAN, FATMA (2002) "Radiation Transfer for Exponential  $c(x)$ ," *Turkish Journal of Physics*: Vol. 26: No. 3, Article 3. Available at: <https://journals.tubitak.gov.tr/physics/vol26/iss3/3>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Physics by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact [academic.publications@tubitak.gov.tr](mailto:academic.publications@tubitak.gov.tr).

# Radiation Transfer for Exponential $c(x)$

Fatma ERDOĞAN

Ankara University, Department of Physics Engineering,  
06100 Tandoğan, Ankara-TURKEY  
e-mail: ferdogan@science.ankara.edu.tr

Received 16.02.2001

## Abstract

The emergent flux for radiation transfer in inhomogeneous half-space for exponential  $c(x) = \frac{1+ke^{-mx}}{1+be^{-mx}}$  is obtained by using Modified-Eddington method, where  $m$ ,  $b$ ,  $k$  are constants.

As a result of this work, angular flux is obtained in terms of Jacobi polynomials.

**Key Words:** The emergent flux, Radiation transfer, Modified-Eddington

## 1. Introduction

The radiation problems for the homogeneous medium have been solved using various methods [1, 2, 3]. These kind of problems are then extended to problems of the form

$$\mu\partial_x\varphi(x, \mu) + \varphi(x, \mu) = \frac{c(x)}{2} \int_{-1}^1 \varphi(x, \mu') d\mu'; 0 \leq x \leq \infty \quad (1)$$

$$\varphi(0, \mu) = g(\mu), 0 \leq \mu \leq 1 \quad (2)$$

$$\lim_{x \rightarrow \infty} \varphi(x, \mu) < \infty, \quad (3)$$

where  $c(x)$  is a continuous function of position [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17]. In most of these works  $c(x)$  is considered to have the forms,

$$c(x) = c_0 e^{-x/s} \quad (4)$$

or

$$c(x) = \int_0^\infty \alpha(t) e^{-x/t} dt, \quad (5)$$

where

$$\int_0^{\infty} |t(\alpha(t))|^2 dt < \infty, \quad (6)$$

For these choices of  $c(x)$ , the singular eigenfunction method of solution is used [4, 5].

Pomraining and Larsen [6] obtained the solutions of the transport equation using Case's method. G.C. Pomraining has calculated [11] the radiation diffuse problem and hence the emergent flux for the homogeneous medium. Wakil, Saad and Hendi [16] used the Modified-Eddington approximation for the exponential medium.

In our work we use the Modified-Eddington method to determine

$$c(x) = \frac{1 + ke^{-mx}}{1 + be^{-mx}} \quad (7)$$

with which the emergent flux is calculated.

## 2. The Modified-Eddington Method

The angular flux in the Modified-Eddington method is defined as

$$\varphi(x, \mu) = M(x)E(x, \mu) + H(x)O(x, \mu), \quad (8)$$

where the zeroth and first moments of the flux are given by

$$M(x) = \frac{1}{2} \int_{-1}^1 \varphi(x, \mu) d\mu \quad (9)$$

and

$$H(x) = \frac{1}{2} \int_{-1}^1 \varphi(x, \mu) \mu d\mu \quad (10)$$

$E(x, \mu)$  and  $O(x, \mu)$  in the definition of the angular flux are even and odd functions of  $\mu$ , respectively, and normalized as

$$\frac{1}{2} \int_{-1}^1 E(x, \mu) d\mu = 1; \quad (11)$$

$$\frac{1}{2} \int_{-1}^1 O(x, \mu) d\mu = 0. \quad (12)$$

Multiplying Eq.(1) by  $\mu^n$ ,  $n = 0, 1$ , respectively and integrating over  $\mu$  in the range  $(-1, 1)$  we get, respectively for each  $n$ ,

$$\frac{d}{dx}H(x) + [1 - c(x)]M(x) = 0 \quad (13)$$

$$\frac{d}{dx}[D(x)M(x)] + H(x) = 0, \quad (14)$$

where

$$D(x) = \frac{1}{2} \int_{-1}^1 E(x, \mu) \mu^2 d\mu. \quad (15)$$

In order to eliminate spatial derivatives Eqs. (1, 8, 14, 15) are used and

$$E(x, \mu) = \frac{c(x)}{1 - \nu^2 \mu^2} \quad (16)$$

$$O(x, \mu) = \mu \frac{E(x, \mu)}{D(x)} \quad (17)$$

are obtained, where

$$\nu^2(x) = \frac{1 - c(x)}{D(x)} \quad (18)$$

and

$$2\nu(x)c(x) = \ln [(1 + \nu(x)) / (1 - \nu(x))]. \quad (19)$$

Equations (13, 14) give

$$D \frac{d^2}{dx^2} [D(x)M(x)] - (1 - c(x))M(x) = 0. \quad (20)$$

Form  $\ll 1$ ,  $D(x)$  can be assumed as a constant.

### 3. The Calculations

We can solve Eq.(20) for  $c(x)$  given in Eq.(17) using the Modified-Eddington method:

$$D \frac{d^2}{dx^2} M(x) - \left( 1 - \frac{1 + ke^{-mx}}{1 + be^{-mx}} \right) M(x) = 0. \quad (21)$$

When the variable  $x$  is changed to  $s$  as

$$s = -e^{-mx}, \quad (22a)$$

the following equation is found:

$$\frac{d^2 M}{ds^2} + \frac{1}{s} \frac{dM(s)}{ds} + \frac{1}{s^2(1-bs)} \left\{ \frac{(b-k)s}{Dm^2} \right\} M(s) = 0. \quad (22b)$$

Lets choose

$$M(s) = (1-bs)y(s). \quad (23)$$

Placing it into Eq.(22b), we get:

$$\frac{d^2 y}{ds^2} + \frac{(1-3bs)}{s(1-bs)} \frac{dy}{ds} + \frac{1}{s(1-bs)} \left[ -b + \frac{b-k}{Dm^2} \right] y(s) = 0 \quad (24)$$

Using  $t=bs$  ;

$$\frac{d^2 y}{dt^2} + \frac{(1-3t)}{t(1-t)} \frac{dy}{dt} + \frac{1}{t(1-t)} \left[ -1 + \frac{b-k}{Dm^2 b} \right] y = 0 \quad (25)$$

and with  $t = \frac{1}{2}(1-z)$  we obtain

$$(1-z^2) \frac{d^2 y}{dz^2} + (1-3z) \frac{dy}{dz} + \left[ -1 + \frac{b-k}{Dm^2 b} \right] y = 0. \quad (26)$$

This differential equation is in the form similar to the Jacobi differential equation which is given as

$$(1-z^2) \frac{d^2 y}{dz^2} + \{ \alpha - \beta + (\alpha + \beta - 2)z \} \frac{dy}{dz} + (n+1)(n+\alpha+\beta)y = 0, \quad (27)$$

where  $\alpha, \beta$  are constants [18, 19].

$(1-z)^\alpha (1+z)^\beta P_n^{(\alpha, \beta)}(z)$  satisfies this differential equation. Comparing the two equations (26, 27) we have

$$\alpha = 0 \quad (28a)$$

$$\beta = -1 \quad (28b)$$

$$n = \mp \sqrt{\frac{b-k}{Dm^2 b}}. \quad (28c)$$

The solution to Eq.(21) can now be written in the form

$$M(x) = P_{\mp \sqrt{\frac{b-k}{Dm^2 b}}}^{(0, -1)} (1 + 2be^{-mx}). \quad (29)$$

From Eq.(14), we get

$$H(x) = -D \frac{dM(x)}{dx} = 2bm DP_{\pm \sqrt{\frac{b-k}{Dm^2 b}}}^{(0, -1)'} (1 + 2be^{-mx}). \quad (30)$$

Using Eq.(8) and the expressions in Eq.(16, 17, 29, 30) the angular flux can be obtained as

$$\begin{aligned} \varphi(x, \mu) = & \left( \frac{1 + ke^{-mx}}{1 + be^{-mx}} \right) \left( \frac{1}{1 - \nu^2 \mu^2} \right) P_{\mp \sqrt{\frac{b-k}{Dm^2b}}}^{(0,-1)} (1 + 2be^{-mx}) \\ & + \left( \frac{1 + ke^{-mx}}{1 + be^{-mx}} \right) \left( \frac{2bm\mu}{1 - \nu^2 \mu^2} \right) P_{\mp \sqrt{\frac{b-k}{Dm^2b}}}^{(0,-1)'} (1 + 2be^{-mx}), \end{aligned} \quad (31)$$

where b, D, m, k are constants.

Using the solution given in Eq.(31), one can calculate the constants C and D

$$C = 1 - 2 \int_0^1 \varphi(0, -\mu) \mu d\mu, \quad (32)$$

$$D = \frac{1}{3} + 2 \int_0^1 \varphi(0, -\mu) \mu^2 d\mu. \quad (33)$$

as

$$C = 1 + \frac{1}{\nu^2} \left( \frac{1+k}{1+b} \right) \left\{ \ln(1 - \nu^2) P_{\mp \sqrt{\frac{b-k}{Dm^2b}}}^{(0,-1)} (1 + 2b) + \frac{4bm}{\nu} (-\nu + \arctan h\nu) P_{\mp \sqrt{\frac{b-k}{Dm^2b}}}^{(0,-1)'} (1 + 2b) \right\} \quad (34)$$

and

$$D = \frac{1}{3} + \frac{2}{\nu^3} \left( \frac{1+k}{1+b} \right) \left\{ (-\nu + \arctan h\nu) P_{\mp \sqrt{\frac{b-k}{Dm^2b}}}^{(0,-1)} (1 + 2b) + \frac{\nu^2 + \ln(1 - \nu^2)}{\nu} P_{\mp \sqrt{\frac{b-k}{Dm^2b}}}^{(0,-1)'} (1 + 2b) \right\}. \quad (35)$$

## 4. Conclusion

The emergent flux for radiation transfer in inhomogeneous half-space is calculated using  $c(x) = (1 + ke^{-mx})/(1 + be^{-mx})$ . This calculation has been previously handled by Pomraning using the linear transport equation. In our work the Modified. Eddington method is used and analytical expressions in terms of Jacobi polynomials are obtained.

## References

- [1] S. Chandrasekhar, Radiative Transfer (Dover Publ.,New York, 1960).
- [2] K. M. Case and P. F. Zweifel, Linear Transport Theory (Addition-Wesley, 1967).
- [3] V. V. Sobolev, Light Scattering in Planetary Atmospheres (Pergamon Press, London, 1975).
- [4] B. J. Martin, *J. Appl. Math.*, **20** (1971) 703.
- [5] T. W. Mullikin and C. E. Siewert, *Ann. Nucl. Energy.*, **7** (1980) 205.
- [6] G. C. Pomraning and E. W. Larsen, *J. Math. Phys.*, **21** (1980) 1603.
- [7] E. W. Larsen, G. C. Pomraning and V. C. Badham, *J. Math. Phys.*, **21** (1980) 2448.
- [8] T. W. Mullikin, Sandia National Laboratories Report Sand., 80-1065 (1980)

- [9] E. W. Larsen and T. W. Mullikin, *J. Math. Phys.*, **22** (1981) 856.
- [10] D. M. Garcia and C. E. Siewert, *J. Quant. Spectrosc. Radiat. Transfer.*, **25** (1981) 277.
- [11] G. C. Pomraning, *J. Math. Phys.*, **22** (1981) 1623.
- [12] E. W. Larsen, *J. Math. Phys.*, **22** (1981) 1463.
- [13] E. W. Larsen, *Prog. Nucl. Energy*, **8** (1981) 203.
- [14] D. M. Garcia and C. E. Siewert, *J. Quant Spectrosc. Radiat. Transfer.*, **27** (1982) 141.
- [15] G. C. Pomraning, E. W. Larsen and V. C. Badnam, *J. Math. Phys.*, **23** (1982) 815.
- [16] E. Wakil, E. A. Saad and A. A. Hendi, *Astrophysics and Space Science.*, **100** (1984) 381.
- [17] F. Erdoğan and C. Tezcan, *J. Quant. Spectrosc. Radiat. Transfer.*, (in the press)
- [18] W. Bell, *Special Functions* (D.Van Nostrand Company Ltd., London, 1968).
- [19] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover Publications, Inc., New York, 1970).