

1-1-2017

The Hahn-Banach theorem for $\$A\$$ -linear operators

BAHRİ TURAN

FATMA BİLİCİ

Follow this and additional works at: <https://dctubitak.researchcommons.org/math>



Part of the [Mathematics Commons](#)

Recommended Citation

TURAN, BAHRİ and BİLİCİ, FATMA (2017) "The Hahn-Banach theorem for $\$A\$$ -linear operators," *Turkish Journal of Mathematics*: Vol. 41: No. 5, Article 25. <https://doi.org/10.3906/mat-1610-24>
Available at: <https://dctubitak.researchcommons.org/math/vol41/iss5/25>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Mathematics by an authorized editor of TÜBİTAK Academic Journals.

The Hahn–Banach theorem for A -linear operators

Bahri TURAN*, Fatma BİLİCİ

Department of Mathematics, Faculty of Science, Gazi University, Teknikokullar, Ankara, Turkey

Received: 07.10.2016

Accepted/Published Online: 02.01.2017

Final Version: 28.09.2017

Abstract: In this short paper we present a generalization of the Hahn–Banach extension theorem for A -linear operators. Some theoretical applications and results are given.

Key words: Riesz space, positive operator, A -linear operator, Hahn–Banach extension theorem

1. Introduction

We assume the reader to be familiar with the elementary theory of Riesz spaces and order bounded operators. In this regard, we use [1, 3, 7] as sources of unexplained terminology and notation. Moreover, all Riesz spaces under consideration are assumed to be Archimedean.

We denote by $L_b(E, F)$ the class of all order bounded operators from the Riesz space E into F and by $L_b(E)$ the order bounded operators from E into itself. Recall that $\pi \in L_b(E)$ is called an orthomorphism of E if $x \perp y$ in E implies that $\pi(x) \perp y$. Orthomorphisms of E will be denoted by $Orth(E)$. The principal order ideal generated by the identity operator I in $Orth(E)$ is called the ideal center of E and is denoted by $Z(E)$. Let A be a Riesz algebra (lattice ordered algebra), i.e. A is a Riesz space that is simultaneously an associative algebra with the additional property that $a, b \in A_+$ implies that $a \cdot b \in A_+$. An f -algebra A is a Riesz algebra that satisfies the extra requirement that $a \wedge b = 0$ implies $a \cdot c \wedge b = c \cdot a \wedge b = 0$ for all $c \in A_+$. If A is an Archimedean f -algebra, then A is necessarily associative and commutative. The collections $Orth(E)$ are, with respect to composition as multiplication, Archimedean f -algebras with the identity mapping I as a unit element. Another well-known example of f -algebras is $C(K)$ of all real continuous functions on a topological space K .

Let A be an f -algebra and E be a Riesz space. E is said to be a left o -module (order module) over A if there exists a map $A \times E \rightarrow E : (a, x) \rightarrow ax$ satisfying the following:

1. E is a left module over A ,
2. for each $a \in A_+$ and $x \in E_+$ we have $ax \in E_+$.

An o -module over A is called f -module if it has the following property:

3. if $x \perp y$ in E , then for each $a \in A$ we have $ax \perp y$.

If A has a unit element e ,

4. $ex = x$ for each $x \in E$,

*Correspondence: bturan@gazi.edu.tr

2010 AMS Mathematics Subject Classification: 47B60, 47B65, 46A40

and then E is said to be a unitary f -module. A right f -module over A is defined similarly. We shall only consider left f -modules from now on and these will simply be referred to as f -modules. If E is an f -module over A , then there exists a map $p : A \rightarrow \text{Orth}(E)$ defined by $p(a) = \pi_a$ where $\pi_a(x) = ax$ for each $x \in E$. p is a unital algebra and Riesz homomorphisms. Conversely, if there exists such a map $p : A \rightarrow \text{Orth}(E)$ then the map $(a, x) \rightarrow ax = p(a)(x)$ of $A \times E \rightarrow E$ defines an f -module structure on E over A .

Example 1 (a) A Riesz space E is an f -module over $A = \mathbb{R}$, the f -algebra of the real numbers. The embedding $p : A \rightarrow \text{Orth}(E)$ is the identification of \mathbb{R} with the subalgebra of $\text{Orth}(E)$ consisting of the multiples of the identity.

(b) A unital f -algebra A is an f -module over itself in a natural way. The mapping $p : A \rightarrow \text{Orth}(A)$ is the well-known identification (surjective algebra and Riesz isomorphism) of $a \in A$ with the orthomorphism $p(a) = \pi_a$ where $\pi_a(b) = a \cdot b$, for each $b \in A$ [1, 7].

(c) Let E be an f -module over A . The mapping $p : A \rightarrow \text{Orth}(E)$ is a positive algebra homomorphism and hence p is a Riesz homomorphism [4].

Definition 2 Let E and F be two modules over A . A mapping $\Gamma : E \rightarrow F$ will be called A -linear if $\Gamma(x + y) = \Gamma(x) + \Gamma(y)$ and $\Gamma(ax) = a\Gamma(x)$ for each $a \in A$ and $x, y \in E$. A mapping $q : E \rightarrow F$ will be called A -sublinear if $q(x + y) \leq q(x) + q(y)$ and $q(ax) = aq(x)$ for each $a \in A_+$ and $x, y \in E$.

The set of order bounded and A -linear maps will be denoted by $L_b(E, F; A)$. Order properties of the space of A -linear operators were given by Turan in [5].

Example 3 (a) Let E be a Riesz space. The bilinear map $Z(E) \times E \rightarrow E : (\pi, x) \rightarrow \pi x = \pi(x)$ shows that E is an f -module over its center. As E is assumed to be Archimedean, $\text{Orth}(E)$ is also Archimedean and therefore commutative. Thus, we have:

$$\text{Orth}(E) \subseteq L_b(E, E; Z(E)) \subseteq L_b(E).$$

(b) Let E be a Riesz space of all continuous piecewise linear functions on $[0, 1]$. Then $Z(E) = \{\lambda I : \lambda \in \mathbb{R}\}$ and so $L_b(E, E; Z(E)) = L_b(E)$.

(c) Let E be a Riesz space. If E has the principal projection property, then $L_b(E, E; Z(E)) = \text{Orth}(E)$ [1, Theorem 8.3].

(d) Let $A = l_\infty$, $E = l_p$; $1 \leq p < \infty$; then $L_b(E, E; A) = \text{Orth}(E)$.

Many versions of the Hahn–Banach theorem have been given. In this work, we shall gather an extension theorem for A -linear operators and we shall give some applications of the extension theorem.

2. Generalization of the Hahn–Banach theorem

The following theorem is a general version of what is known as the Hahn–Banach extension theorem.

Theorem 4 Let E be a module over unital Dedekind complete f -algebra A , F an Archimedean Dedekind complete o -module over A , and $q : E \rightarrow F$ an A -sublinear map. If N is a submodule of E and $\Phi : N \rightarrow F$

is an A -linear map satisfying $\Phi(x) \leq q(x)$ for all $x \in N$, then there exists an A -linear map Γ from E to F such that $\Gamma(x) = \Phi(x)$ for all $x \in N$ and $\Gamma(x) \leq q(x)$ for all $x \in E$.

Proof The proof of the one-step extension is different from the other proof of Hahn–Banach theorems. If this is done, then an application of Zorn’s lemma guarantees an extension of Φ to all of E with the desired properties. Let N be a submodule of E . If $N = E$ holds, then the proof is trivial. Assume that $N \neq E$ and take $z \in E \setminus N$. Let N_1 be the submodule of E generated by N and z , i.e. $N_1 = \{x + az : x \in N, a \in A\}$. This representation is not unique, since it is possible that $az \in N$ for some nonzero a in A . This leads to difficulties in showing the “one-step extension”. It is clear that for each $y, u \in N$ we have

$$\Phi(y) - q(y - z) \leq q(u + z) - \Phi(u).$$

This inequality coupled with the Dedekind completeness of F guarantees that both

$$k = \sup \{ \Phi(y) - q(y - z) : y \in N \}$$

and

$$t = \inf \{ q(u + z) - \Phi(u) : u \in N \}$$

exist in F , and they satisfy $k \leq t$. Now, for any $w \in F$ satisfying $k \leq w \leq t$ (for instance, $w = k$), let us define

$$\Gamma_1 : N_1 \rightarrow F; (x + az) \rightarrow \Gamma_1(x + az) = \Phi(x) + aw.$$

We shall now show that Γ_1 is well defined. As A is Dedekind complete, A is e -uniform complete [3, Theorem 42.6]. If $0 \leq a$ in A then $(a + \frac{1}{n}e)^{-1}$ is in A for all n ($n = 1, 2, 3, \dots$) by [2, Theorem 11.1]. For all n , we have

$$w \leq t \leq q \left[\left(a + \frac{1}{n}e \right)^{-1} x + z \right] - \Phi \left[\left(a + \frac{1}{n}e \right)^{-1} x \right]$$

and it follows that

$$\Phi(x) + \left(a + \frac{1}{n}e \right) w \leq q \left[x + \left(a + \frac{1}{n}e \right) z \right] \leq q(x + az) + \frac{1}{n}eq(z).$$

As F is Archimedean, we have

$$\Phi(x) + aw \leq q(x + az).$$

For general $a \in A$, write $a = a^+ - a^-$. Let $u : A \rightarrow B_{a^+}$ be the order projection, where B_{a^+} is the band generated by a^+ . We have $u \in Orth(A)$, $u^2 = u$, $u(a) = a^+$. Since A is Riesz and algebra isomorphic to $Orth(A)$ [7, Theorem 141.1], there exists $0 \leq b \in A$ such that $b^2 = b$, $ba = a^+$, $ba^+ = a^+$, $ba^- = 0$, and $(b - e)a = a^-$. From the first case we have

$$\Phi(x) + a^+w \leq q(x + a^+z)$$

and hence

$$\begin{aligned} b(\Phi(x) + a^+w) &= b\Phi(x) + b(a^+w) = b\Phi(x) + (ba^+)w = b\Phi(x) + a^+w \\ &\leq bq(x + a^+z) = q(bx + b(a^+z)) = q(bx + b^2(a^+z)) = bq(x + az). \end{aligned}$$

Similarly, since $k \leq w$ we get

$$(e - b)(\Phi(x) - a^-w) \leq (e - b)q(x + az).$$

Adding these inequalities, we obtain

$$\Phi(x) + aw \leq q(x + az) \tag{1}$$

for all $a \in A$. We suppose $x, y \in N$ and $a, c \in A$ are such that $x + az = y + cz$. Then, from (1), we have

$$\Phi(x - y) + (a - c)w \leq q((x - y) + (a - c)z) = q(0) = 0$$

and

$$\Phi(y - x) + (c - a)w \leq q((y - x) + (c - a)z) = q(0) = 0,$$

and therefore $\Gamma_1(x + az) = \Gamma_1(y + cz)$. Evidently, Γ_1 is dominated by q and is an A -linear map. An application of Zorn's lemma guarantees an extension of Φ to all of E with the desired properties. \square

Let K be an extremally disconnected compact Hausdorff space (sometimes called a Stonian space). Then $C(K)$ is a Dedekind complete f -algebra. Taking $A = C(K)$, we can get the following result that was proved by Vincent-Smith as Theorem 1 in [6].

Corollary 5 *Let E be a module over Dedekind complete $A = C(K)$ and $q : E \rightarrow A$ be an A -sublinear map. If N is a submodule of E and $\Phi : N \rightarrow A$ is an A -linear map satisfying $\Phi(x) \leq q(x)$ for all $x \in N$, then there exists some A -linear map Γ on E that extends Φ and is such that $\Gamma(x) \leq q(x)$ for all $x \in E$.*

If we take $A = \mathbb{R}$, then we obtain the following corollary that was given as Theorem 83.13 [7] or Theorem 2.1 [1].

Corollary 6 *Let E be a (real) vector space, F a Dedekind complete Riesz space, and $q : E \rightarrow F$ a sublinear map. If N is a vector subspace of E and $\Phi : N \rightarrow F$ is an operator satisfying $\Phi(x) \leq q(x)$ for all $x \in N$, then there exists some linear map Γ on E that extends Φ and is such that $\Gamma(x) \leq q(x)$ for all $x \in E$.*

It is known that there are many applications of the Hahn–Banach extension theorem. Some of these applications were given for linear positive operators in [1]. Similar applications of Theorem 1 can be given for A -linear positive operators. Two of them will be given below. Let E be a unitary module over A . Then every submodule of E is a vector subspace of E . A submodule N of a unitary module E is called a Riesz submodule whenever N is closed under the lattice operations of E (i.e. whenever N is a Riesz subspace of E).

Corollary 7 *Let E be a unitary f -module over unital Dedekind complete f -algebra A , F an Archimedean Dedekind complete o -module over A , and $\Gamma : E \rightarrow F$ a positive A -linear operator. If N is a Riesz submodule of E and $\Phi : N \rightarrow F$ is a positive A -linear map satisfying $\Phi(x) \leq \Gamma(x)$ for all $x \in N_+$, then there exists some positive A -linear map H on E that extends Φ and is such that $H(x) \leq \Gamma(x)$ for all $x \in E_+$.*

Proof It is similar to the proof of Theorem 2.2, which is given in [1]. \square

Corollary 8 *Let E be a unitary f -module over unital Dedekind complete f -algebra A and F an Archimedean Dedekind complete o -module over A . If N is a Riesz submodule of E and $\Phi : N \rightarrow F$ is a positive A -linear map, then the following statements are equivalent:*

1. Φ extends to a positive A -linear map.
2. Φ extends to an order bounded A -linear map.
3. There exists a monotone A -sublinear mapping $q : E \rightarrow F$ satisfying $\Phi(x) \leq q(x)$ for all $x \in N$.

Proof The proof is the same as the proof of Theorem 2.3 [1] with a slight difference. □

References

- [1] Aliprantis CD, Burkinshaw O. Positive Operators. London, UK: Academic Press, 1985.
- [2] De Pagter B. f -Algebras and orthomorphism. PhD, Leiden University, Leiden, the Netherlands, 1981.
- [3] Luxemburg WĀJ, Zaanen AC. Riesz Space I. Amsterdam: North Holland, 1971.
- [4] Turan B. On f -linearity and f -orthomorphisms. *Positivity* 2000; 4: 293-301.
- [5] Turan B. Order properties of the space of A -linear operators. *Demonstratio Mathematica* 2006; 2: 429-438.
- [6] Vincent-Smith G. The Hahn-Banach theorem for modules. *P Lond Math Soc* 1967; s3-17: 72-90.
- [7] Zaanen AC. Riesz Spaces II. Amsterdam, the Netherlands: North Holland, 1983.