

1-1-2009

C-Banded Karyotype and Nucleolar Organizer Regions (NORs) of Wild Boar, *Sus scrofa* (Artiodactyla: Suidae) from Anatolia

ATILLA ARSLAN

İRFAN ALBAYRAK

Follow this and additional works at: <https://journals.tubitak.gov.tr/biology>



Part of the [Biology Commons](#)

Recommended Citation

ARSLAN, ATILLA and ALBAYRAK, İRFAN (2009) "C-Banded Karyotype and Nucleolar Organizer Regions (NORs) of Wild Boar, *Sus scrofa* (Artiodactyla: Suidae) from Anatolia," *Turkish Journal of Biology*. Vol. 33: No. 1, Article 5. <https://doi.org/10.3906/biy-0805-35>
Available at: <https://journals.tubitak.gov.tr/biology/vol33/iss1/5>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Biology by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

C-Banded Karyotype and Nucleolar Organizer Regions (NORs) of Wild Boar, *Sus scrofa* (Artiodactyla: Suidae) from Anatolia

Atila ARSLAN¹, İrfan ALBAYRAK²

¹Department of Biology, Faculty of Science and Arts, Selçuk University, 42031 Konya - TURKEY

²Department of Biology, Faculty of Science and Arts, Kırıkkale University, 71450 Kırıkkale - TURKEY

Received: 28.05.2008

Abstract: The present study reports the karyotype, C-banding, and nucleolar organizer regions (NORs) of 6 *Sus scrofa* (Linnaeus, 1758) males from Anatolia. The karyotype of *S. scrofa* comprised (2n) 38 chromosomes, the number of chromosomal arms (FN) was 64, and the number of autosomal arms (FNa) was 60. C-positive regions appeared to be restricted to the centromeric regions of autosomes 1, and 13-18, and the entire long arm of the Y chromosome. Some autosomes had very slight C-bands. The X chromosome appeared to be entirely euchromatic. NORs were identified by the silver-staining technique and were observed on secondary constriction sites of 2 metacentric chromosomes pairs.

Key Words: *Sus scrofa*, wild boar, cytogenetic, Anatolia

Anadolu'daki Yaban Domuzu, *Sus scrofa* (Artiodactyla: Suidae)'nin C-bantlı Karyotipi ve Nükleolar Organizatör Bölgeleri (NORs)

Özet: Bu çalışmada Anadolu'daki altı erkek *Sus scrofa* (Linnaeus, 1758) örneğinin karyotip, C-bant ve nükleolar organizatör bölgelerinin (NORs) özellikleri araştırıldı. *S. scrofa*'nın karyotipinin (2n) 38 kromozomdan oluştuğu, temel kromozom kol sayısı (FN) 64 ve otozomal kromozom kol sayısı (FNa) 60 olduğu gözlemlendi. C-pozitif bölgeler, 1, 13-18 nolu kromozomların sentromerik bölgelerinde ve Y kromozomun uzun kolunun tamamında görüldü. Bazı otozomlar çok hafif C-bantlara sahiptir. X kromozomun tamamının ökromatik tipte olduğu belirlendi. Gümüş boyama tekniği ile tanımlanan nükleolar organizatör bölgeleri (NORs), iki çift metasentrik kromozomdaki ikincil boğum yerlerinde bulundu.

Anahtar Sözcükler: *Sus scrofa*, yaban domuzu, sitogenetik, Anadolu

Introduction

Turan (1) stated that *Sus scrofa* Linnaeus, 1758 is found throughout Turkey. Mursaloğlu (2) reported that the wild boar in Anatolia is represented by *Sus scrofa libycus* Gray, 1868. Karyological studies around the world show that the diploid chromosome number of the domestic pig (*Sus scrofa domestica*) and the wild boar (*Sus scrofa scrofa*) ranges from 36 to 38 (3-12). The karyology of *S. scrofa* from Turkey was reported by Albayrak and İnci (13), but they did not report any

chromosomal polymorphisms in *S. scrofa*. Nonetheless, chromosome number polymorphisms have been frequently recorded in wild boar populations as a result of centric fusion (3,8,12). The distribution of constitutive heterochromatin (C-band) in the chromosomes of domestic pigs and wild boar has been studied by Sysa (14) and Hansen (15). Nucleolar organizer regions (NORs) of this species have also been determined by some authors using fluorescence *in situ* hybridization (FISH) and silver staining techniques (16-21).

Herein, we present the C-banding and Ag-NORs data for *S. scrofa* from Turkey, and a comparison of these results with those from other parts of the world.

Materials and Methods

The study animals (6 males) were collected from Kırkkale Province in 2007. Karyotype preparations were obtained from the bone marrow of the colchicined animals (22), followed by conventional Giemsa staining. Constitutive heterochromatin and NORs were detected to identify pairs of each autosomal and sexual chromosome via C-banding (23) and Ag-NOR staining (24), respectively. Chromosome morphology was established by calculating centromeric indexes. In total, 10-20 slides were prepared from each animal for analysis, and a minimum of 20 well-spread metaphase plates were also analyzed from each animal. Some standard voucher specimens (skins and skulls) were deposited in the Department of Biology, Faculty of Science and Arts, Selçuk University, Konya, Turkey.

Results and Discussion

The karyotype of Turkish *S. scrofa* comprises 38 chromosomes. The number of autosomal arms (FNa) is 60 and the number of fundamental arms (FN) is 64. Twelve autosomal pairs are bi-armed (nos. 1-12) and 6 pairs are acrocentric (nos. 13-18). The X chromosome is a medium-sized submetacentric and the Y chromosome is a small metacentric (Figure 1). We know that for populations studied elsewhere, diploid chromosome numbers of 36 and 37 were recorded from the USA, Italy, the Netherlands, and Poland (7,11,12,25). According to previous reports (3,8,12), the causes of this chromosomal polymorphism could perhaps be attributed to a Robertsonian translocation (i.e. centric fusion) occurring between 1 or 2 pairs of acrocentric chromosomes. All previous reports indicated (2n = 36, 37, and 38) the same number (60) of autosomal arms (FNa). Centric fusion was not observed neither in the present study nor in the study performed by Albayrak and İnci (13) on Turkish wild boar.




Figure 1. Metaphase spread and karyotype of Turkish *Sus scrofa*. Arrow indicates the secondary constrictions (NOR).

C-bands are located at the centromeric regions of submetacentric and acrocentric chromosomes nos. 1 and 13-18, as well as the entire long arm of the Y chromosome. Some autosomes have only very slight centromeric C-bands. The X chromosome appears to be entirely euchromatic. The C-bands of acrocentric chromosomes were larger than those on chromosome no. 1 (Figure 2). In the domestic pig the centromeric regions of all chromosomes have been reported to be C-positive (14,15). Furthermore, interstitial C-bands were observed in a chromosome pair by Hansen (15). Telomeric C-bands have also been identified by Adegá et al. (26) in the domestic pig; however, neither interstitial nor telomeric C-bands were determined in our specimens. These results show that (with the exclusion of chromosome no. 1) the centromere regions of bi-armed chromosomes have been completed by a reduction in heterochromatin blocks. According to Adegá et al. (26), the classic C-banding technique used in cytogenetic studies detects the major constitutive heterochromatin blocks in pig chromosomes; however, the classical method has also been used in studies by Sysa (22) and Hansen (23).

Using silver-nitrate staining we determined that the NORs are localized at the secondary constriction sites of 2 metacentric chromosome pairs (nos. 4 and 8). Active NOR signals were observed to be different in each chromosome, in both the conventional karyotype and Ag-stained metaphase (Figures 1 and 3). In the present study metacentric and submetacentric chromosomes were not karyotyped separately; therefore, chromosomes were numbered as mixed so that in our results chromosomes 4 and 8 corresponded to chromosomes 8 and 10, respectively, in other reports. Most researchers have reported a preferential staining of the NORs on chromosome 10. Mellink et al. (27), however, observed a high incidence of Ag-stained NORs on chromosome no. 8 in wild *Sus* species (*S. scrofa scrofa*, *S. scrofa vittatus*, *S. verrucosus*, *S. celebensis*, and *S. salvanius*). It has been revealed by the FISH technique that domestic pig rDNA is located either in the secondary constrictions of both chromosome pairs 10 and 8 (27) or only in the secondary constrictions of chromosome 10 (28). Mellink et al. (21) determined that the signal on chromosome 10 can be stronger than that in chromosome 8, as in the present study.




Figure 2. Metaphase spread and C-banded karyotype of Turkish *Sus scrofa*.




Figure 3. Silver-stained metaphases of Turkish *Sus scrofa*. Arrow indicates the Ag-NOR.

Consequently, the localization of NORs obtained from Turkish *Sus scrofa* is similar to nominative and non-nominative species. Although Turkish *Sus scrofa libycus* is known to be an ancestor to all domestic pigs, it is clearly different than others, because centromeric C-bands were not seen in most bi-armed chromosomes of the Turkish specimens that have been studied to date.

Acknowledgments

We would like to thank Dr. A. C. Kitchener for linguistic revision of the manuscript.

Corresponding Author:

Atilla ARSLAN

Department of Biology,

Faculty of Science and Arts,

University of Selçuk,

Konya - TURKEY

E-mail: aarslan@selcuk.edu.tr

References

1. Turan N. Türkiye'nin av ve yaban hayvanları (Memeliler). Ogun Matbaacılık Sanayi, Ankara, 1984.
2. Mursaloğlu B. Türkiye'nin azalan memelileri hakkında. Turk Biol Derg 14: 65-70, 1964.
3. Mcfee AF, Banner MW, Rary JM. Variation in chromosome number among European wild pigs. Cytogenetics 5: 75-81, 1966.
4. Zivkovic S, Jovanovic V, Isakovic I et al. Chromosome complement of the European wild pig (*Sus scrofa* L.). Experientia 27: 224-226, 1971.
5. Gustavsson I, Hageltorn M, Zech L et al. Identification of the chromosomes in a centric fusion/fission polymorphic system of the pig (*Sus scrofa* L.). Hereditas 75: 153-155, 1973.
6. Tikhonov VN, Troshina AI. Identification of chromosomes and their aberrations in karyotypes of subspecies of *Sus scrofa* L. by differential staining. Doklady Akademii Nauk SSR 214: 932-935, 1974.
7. Bosma AA. Chromosomal polymorphism and G banding patterns in the wild boar (*Sus scrofa* L.) from the Netherlands. Genetica 46: 391-399, 1976.
8. Hansen-Melander E, Melander Y. The karyotype of the pig. Hereditas 77: 149-158, 1974.
9. Gustavsson I. Standard karyotype of the domestic pig. Hereditas 109: 151-157, 1988.
10. Bosma AA, De Haan NA, Macdonald AA. The current status of cytogenetics of the Suidae: a review. Bongo, Berlin 18: 258-272, 1991.
11. Macchi E, Tarantola M, Perrone A et al. Cytogenetic variability in the wild boar (*Sus scrofa scrofa*) in Piedmont (Italy): Preliminary Data. Ibx J Mountain Ecol 3: 17-18, 1995.
12. Rejduch B, Slota E, Rozycki M et al. Chromosome of polymorphism in a litter of European wild boar (*Sus scrofa scrofa* L.). Anim Sci 21: 57-62, 2003.
13. Albayrak İ, İnci S. The karyotype of the wild boar *Sus scrofa* Linnaeus, 1758 in Turkey (Mammalia: Artiodactyla). Turk J Zool 31: 65-68, 2007.
14. Sysa PS. Polymorphism of metaphase chromosomes in swine (*Sus scrofa* L.). Genetica 52-53: 321-315, 1980.
15. Hansen KM. Sequential Q- and C-band staining of pig chromosomes, and some comments on C-band polymorphism and C-band technique. Hereditas 96: 183-189, 1982.
16. Czaker R, Mayr B. Comparative studies on the polymorphism of nucleolus organizer regions (NORs) in four breeds of domestic pigs (*Sus scrofa domestica* L.) with special emphasis on the development of breeds. J Zool Syst Evol Res 20: 233-241, 1982.
17. Mayr B, Schweizer D, Geber G. NOR activity, heterochromatin differentiation, and the Robertsonian polymorphism in *Sus scrofa* L.. Hereditas 75: 79-80, 1984.
18. Bosma AA, De Haan NA, Mellink CHM. Ribosomal RNA genes located on chromosome 16 of the domestic pig. Cytogenet. Cell Genet 58: 21-24, 1992.

19. Popescu CP, Boscher J, Malynicz GL. Chromosome R-banding patterns and NOR homologies in the European wild pig and four breeds of domestic pig. *Annales de Genetiq* 32: 136-140, 1989.
20. Mellink CHM, Bosma AA, De Haan NA. et al. Numerical variation of nucleolar organizer regions after silver staining in domestic and wild Suidae (Mammalia). *Anim Genet* 23: 231-239, 1992.
21. Mellink CHM, Bosma AA, De Haan NA. Variation in size of Ag-NORs and fluorescent rDNA in situ hybridization signals in six breeds of domestic pig. *Hereditas* 120: 141-149, 1994.
22. Ford CE, Hamerton JL. Colchicine-hypotonic citrate squash sequence for mammalian chromosomes. *Stain Tech* 31: 247-251 1956.
23. Sumner AT. A simple technique for demonstrating centromeric heterochromatin. *Exp. Cell Res* 75: 304-306, 1972.
24. Howell WM, Black DA. Controlled silver staining of nucleolar organizer regions with a protective colloidal developer: a 1-step method. *Experientia* 36: 1014-1015, 1980.
25. Groves C. Ancestors for the pigs: taxonomy and phylogeny of the genus *Sus*. Technical Bulletin No 3 Department of Prehistory, Research School of Pacific Studies, Australian National Univ., 1981.
26. Adegá F, Chaves R, Guedes-Pinto H. Chromosome restriction enzyme digestion in domestic pig (*Sus scrofa*) constitutive heterochromatin arrangement. *Gene Genet Syst* 80: 49-56, 2005.
27. Mellink CHM, Bosma AA, De Haan NA et al. Distribution of rRNA genes in breeds of domestic pig studied by non-radioactive in situ hybridization and selective silver staining. *Genet Sel Evol* 23, suppl. 1: 168-172, 1991.
28. Miyake ML, O'Brien SJ, Kaneda Y. Regional localization of rDNA genes on pig chromosome 10 by in situ hybridization. *Japan J Vet Sci* 50: 341-345, 1988.